



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

cGMP and cAMP pathways rearrange ARHGAP17 and ARHGEF6 protein complexes to control Rac1 in platelets

Citation for published version:

Nagy, Z, Wynne, K, Von Kriegsheim, A, Gambaryan, S & Smolenski, A 2015, 'cGMP and cAMP pathways rearrange ARHGAP17 and ARHGEF6 protein complexes to control Rac1 in platelets' *BMC pharmacology & toxicology*, vol. 16, no. Suppl 1, pp. A16. DOI: 10.1186/2050-6511-16-S1-A16

Digital Object Identifier (DOI):

[10.1186/2050-6511-16-S1-A16](https://doi.org/10.1186/2050-6511-16-S1-A16)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

BMC pharmacology & toxicology

Publisher Rights Statement:

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



MEETING ABSTRACT

Open Access

cGMP and cAMP pathways rearrange ARHGAP17 and ARHGEF6 protein complexes to control Rac1 in platelets

Zoltan Nagy^{1,2}, Kieran Wynne², Alexander von Kriegsheim³, Stepan Gambaryan⁴, Albert Smolenski^{1,2*}

From 7th International Conference on cGMP Generators, Effectors and Therapeutic Implications Trier, Germany. 19-21 June 2015

Endothelial cells inhibit blood platelets by releasing nitric oxide (NO) and prostacyclin (PGI₂). NO and PGI₂ activate cGMP and cAMP synthesis, leading to the activation of cGMP- and cAMP-dependent protein kinases (PKG, PKA), phosphorylation of substrate proteins, and inhibition and reversal of platelet activation and aggregation. A known target of PKA and PKG is the small G-protein Rac1. The active form of Rac1, Rac1-GTP, contributes to cytoskeletal reorganization, platelet spreading, adhesion, granule release, and aggregation. PKG and PKA activation have been shown to reduce Rac1-GTP levels. Rac1-GTP is controlled by GTPase-activating proteins (GAPs - reduce Rac1-GTP) and Guanine-nucleotide exchange factors (GEFs - elevate Rac1-GTP). We hypothesized that specific GAPs and/or GEFs might mediate the inhibitory effects of PKG and PKA on Rac1. Screening for potential substrates among the GAPs and GEFs expressed in platelets resulted in the identification of the GAP ARHGAP17 (also known as Rich1 or Nadrin) as well as the GEF ARHGEF6 (also known as alpha-PIX or Cool2) as new substrates of PKG and PKA in human platelets. We mapped the phosphorylation sites on ARHGAP17 and ARHGEF6, and we investigated the functional consequences of phosphorylation. Phosphorylation of ARHGAP17 by PKA resulted in a dissociation of ARHGAP17 and its interaction partner CIP4. Phosphorylation of ARHGEF6 by both kinases resulted in the association of the ARHGEF6/GIT1 complex with the phospho-serine/threonine binding protein 14-3-3.

Our data suggest that PKG- and PKA-mediated phosphorylations of ARHGAP17 and ARHGEF6 lead to a reorganization of protein complexes involved in the control of

Rac1. These events are likely to contribute to Rac1 inhibition by cyclic nucleotides. Similar findings regarding the regulation of a critical small G-protein by GAP and GEF phosphorylation have been described for Rap1 in platelets before. We conclude that phosphorylation of GAPs and GEFs of small G-proteins might represent a general principle of platelet regulation by cGMP and cAMP.

Authors' details

¹School of Medicine, University College Dublin, Dublin, Ireland. ²Conway Institute, University College Dublin, Dublin, Ireland. ³Systems Biology Ireland, University College Dublin, Dublin, Ireland. ⁴Sechenov Institute of Evolutionary Physiology and Biochemistry, Russian Academy of Sciences, St. Petersburg, 194223, Russia.

Published: 2 September 2015

doi:10.1186/2050-6511-16-S1-A16

Cite this article as: Nagy et al.: cGMP and cAMP pathways rearrange ARHGAP17 and ARHGEF6 protein complexes to control Rac1 in platelets. *BMC Pharmacology and Toxicology* 2015 **16**(Suppl 1):A16.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



* Correspondence: albert.smolenski@ucd.ie

¹School of Medicine, University College Dublin, Dublin, Ireland

Full list of author information is available at the end of the article