



# THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### **Isokinetic shoulder extension strength of trained female uni-lateral arm amputee swimmers**

**Citation for published version:**

Payton, C, Naemi, R, Machtsiras, G & Sanders, R 2008, 'Isokinetic shoulder extension strength of trained female uni-lateral arm amputee swimmers' 16th FINA World Sports Medicine Congress, Manchester, United Kingdom, 7/04/08 - 8/04/08, .

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Early version, also known as pre-print

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.





Payton, C. J.<sup>1</sup>, Naemi, R.<sup>2</sup>, Machtsiras, G.<sup>2</sup> and Sanders, R.<sup>2</sup>

<sup>1</sup>Department of Exercise & Sport Science, Manchester Metropolitan University, Cheshire, UK

<sup>2</sup>Centre for Aquatics Research and Education, Edinburgh University, Edinburgh, UK

## INTRODUCTION

Land-based strength and conditioning work is an important component of a swimmer's training, with development of the shoulder complex usually being an important consideration. Swimmers with an amputation at elbow level, who perform strength training exercises, do so with both their affected and unaffected arms. What has not been well established is the extent of the difference in strength between their two shoulders, and how this varies between arm amputee swimmers.

**The purpose of this study was to establish the within and between-swimmer differences in shoulder extension strength of unilateral forearm amputee swimmers.**

## METHODS

- Participants were nine well-trained female swimmers ( $16.1 \pm 3.2$  years;  $1.64 \pm 0.04$  m;  $57.7 \pm 6.5$  kg). All had a single congenital arm amputation at the level of the elbow.
- Best times for the 50 m front crawl ranged between 30.4-39.6 s.
- Shoulder extension strength was measured on a BIODEX System 3 Pro isokinetic dynamometer.
- Five maximal effort shoulder flexion-extension repetitions were performed with the affected and then unaffected arms at a slow speed ( $60^\circ \cdot s^{-1}$ ) and a fast speed ( $180^\circ \cdot s^{-1}$ ).
- Mean extension torque generated over the mid-range of the movement ( $80-100^\circ$  extension) was found for each repetition, in each condition.
- Shoulder strength (N·m) - the highest value recorded for this mean extension torque.
- Two-way repeated-measures ANOVA was used to evaluate differences between the conditions.



## RESULTS






Figure 1. Shoulder strength (mean  $\pm$  SD) at velocity of a)  $60^\circ \cdot s^{-1}$  and b)  $180^\circ \cdot s^{-1}$ .

Figure 2. Typical extension torque curve.

- The shoulder of the unaffected side was significantly stronger ( $p < 0.05$ ) than the shoulder on the affected side, at both the slow and fast test speeds, although the degree of strength asymmetry varied considerably between the swimmers.
- The swimmers exhibited significantly greater ( $p < 0.05$ ) shoulder strength at the slow test speed, compared to the fast test speed, on both arms. This finding can be explained by the muscle force-velocity relationship.
- Swimmers who ranked highly when strength was expressed in absolute terms were not necessarily as highly ranked when their strength scores were normalised for body mass (Jaric, 2002).
- No significant relationship was found between the shoulder strength (relative or absolute) or strength asymmetry of the swimmers, and their performance level.

## CONCLUSION

Unilateral forearm amputee swimmers generally have greater shoulder extension strength on their unaffected side, compared to their affected side, although there is considerable variation in strength asymmetry between swimmers.

## REFERENCE

Jaric, S. (2002). Muscle strength testing: use of normalisation for body size. *Sports Medicine*, 32 (10), 615-631.

## ACKNOWLEDGEMENTS

The authors would like to acknowledge British Swimming for funding this project and the swimmers for their participation.