

Cognitive, Affective, & Behavioral Neuroscience

Neural responses to others' pain vary with psychopathic traits in healthy adult males

Journal:	<i>Cognitive, Affective, and Behavioral Neuroscience</i>
Manuscript ID:	CABN-RA-14-209.R2
Manuscript Type:	Research Article
Date Submitted by the Author:	24-Feb-2015
Complete List of Authors:	Seara-Cardoso, Ana; University College London, Division of Psychology and Language Sciences Viding, Essi; UCL, Lickley, Rachael; Royal Holloway, University of London, Department of Psychology Sebastian, Catherine; Royal Holloway, University of London, Department of Psychology
Keywords:	empathy, psychopathic personality, fMRI, anterior insula, midcingulate gyrus, inferior frontal gyrus

SCHOLARONE™
Manuscripts

Only

1
2
3 **Neural responses to others' pain vary with psychopathic traits in healthy adult males**
4

5 **Running title:** Psychopathy and response to others' pain
6
7
8
9

10 Ana Seara-Cardoso^{a, b}, Essi Viding^{a, c}, Rachael A. Lickley^a, Catherine L. Sebastian^d
11
12
13

14
15 ^a Division of Psychology and Language Sciences, University College London, United Kingdom
16

17 ^b CIPsi, School of Psychology, University of Minho, Portugal
18

19 ^c Institute of Cognitive Neuroscience, University College London, United Kingdom
20

21 ^d Department of Psychology, Royal Holloway, University of London, United Kingdom
22
23
24
25
26

27 * Corresponding author:
28

29 Ana Seara-Cardoso
30

31 Division of Psychology and Language Sciences, University College London
32

33 26 Bedford Way, London WC1H 0AP, UK
34
35

36 +44(0)2076791051; ana.cardoso.09@ucl.ac.uk
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Disrupted empathic processing is a core feature of psychopathy. Neuroimaging data suggest that individuals with high levels of psychopathic traits show atypical responses to others' pain in a network of brain regions typically recruited during empathic processing (anterior insula, inferior frontal gyrus, and mid- and anterior cingulate cortex). Here, we investigated whether neural responses to others' pain vary with psychopathic traits within the general population in a similar manner to that found in individuals at the extreme end of the continuum. As predicted, variation in psychopathic traits was associated with variation in neural responses to others' pain in the network of brain regions typically engaged during empathic processing. Consistent with previous research, our findings indicate the presence of suppressor effects in the association between levels of affective-interpersonal and lifestyle-antisocial dimensions of psychopathy, and neural responses to others' pain. That is, after controlling for the influence of the other dimension, higher affective-interpersonal psychopathic traits were associated with reduced neural responses to other's pain; whilst higher lifestyle-antisocial psychopathic traits were associated with increased neural responses to others' pain. Our findings provide further evidence that atypical function in this network might represent neural markers of disrupted emotional and empathic processing; that the two dimensions of psychopathy might tap into distinct underlying vulnerabilities; and, most importantly, that the relationships observed at the extreme end of the psychopathy spectrum apply to the non-clinical distribution of these traits, providing further evidence for continuities in the mechanisms underlying psychopathic traits across the general population.

Keywords: empathy; psychopathic personality; fMRI; anterior insula; midcingulate gyrus; inferior frontal gyrus

Introduction

Empathy is a multidimensional phenomenon that involves the capacity to resonate with and understand affective states of others (e.g. Singer & Lamm, 2009). It likely comprises both cognitive and affective components. One affective component, termed ‘affective resonance’, involves experiencing an affective state elicited by the observation or imagination of another person’s affective state. This experience, particularly in response to others’ distress, is thought to play a crucial role in appropriate social interaction. For example, it is thought that experiencing an affective response to others’ distress can elicit prosocial behavior (Nichols, 2001), whilst the absence of such a response can lead to a failure to inhibit aggression towards others (Blair, 2013; Blair, Mitchell, & Blair, 2005). Ultimately, a blunted empathic response system may lead to the development of inappropriate moral behavior (Blair et al., 2005).

Neuroimaging studies have utilized a wide range of different experimental tasks and stimuli (e.g. watching another person in painful situations, seeing a loved one about to receive an electric shock, or viewing another person expressing disgust) to probe the neural bases of empathy (see Fan et al., 2011 for a comprehensive review). Recent meta-analyses of these studies (Fan, Duncan, de Greck, & Northoff, 2011; Lamm, Decety, & Singer, 2011) indicate that the observation of others’ experiences of distress, and more specifically of others’ experiences of pain, consistently elicits robust activation in anterior insula (AI), inferior frontal gyrus (IFG), and a region spanning the border between mid-cingulate cortex (midCC) and anterior cingulate cortex (ACC).

1
2
3 Callous and un-empathic behavior is the hallmark of psychopathy, a personality disorder
4
5 characterized by a constellation of traits including affective-interpersonal traits, such as lack of
6
7 consideration for others' feelings and a tendency to manipulate others; and lifestyle-antisocial
8
9 behavior characteristics, such as impulsiveness and persistent antisocial behavior (Hare, 1993;
10
11 Hare & Neumann, 2008). It has been proposed that the absence of a robust spontaneous empathic
12
13 response to others' distress explains why individuals with psychopathy find it easier to commit
14
15 acts of antisocial behavior towards others (Blair, 2013; Blair et al., 2005). Indeed, both
16
17 behavioral and neuroimaging data are consistent with the notion that these individuals do not
18
19 find other people's distress as salient as their peers do (see Blair, 2013, for a recent review). For
20
21 example, individuals with extreme levels of psychopathic traits present a profile of blunted
22
23 emotional reactivity to aversive stimuli including pictures of mutilated bodies and physical
24
25 assault (Levenston, Patrick, Bradley, & Lang, 2000; Patrick, Bradley, & Lang, 1993), impaired
26
27 recognition of distress cues in others (Blair, Colledge, Murray, & Mitchell, 2001; Blair et al.,
28
29 2004; Blair et al., 2002) and atypical neural responses to stimuli depicting others experiencing
30
31 pain in the network of brain regions typically recruited during empathic processing (i.e. anterior
32
33 insula, IFG and midCC/dACC; Decety, Chen, Harenski, & Kiehl, 2013; Decety, Skelly, & Kiehl,
34
35 2013; Lockwood et al., 2013; Marsh et al., 2013; Meffert, Gazzola, Den Boer, Bartels, &
36
37 Keysers, 2013).

38
39 The study of psychopathy in the general population has been the subject of considerable attention
40
41 recently. There seems to be an increasing interest in the subject, be it on the influence of these
42
43 traits in the workplace, or on the prevalence of high levels of these traits in people who hold key
44
45 positions in society, such as in politics or banking. Research has now shown that the structure of
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 psychopathic personality is dimensional rather than categorical; that is, psychopathic traits are
4 normally distributed in the general population, and individuals with a diagnosis of psychopathy
5 represent an extreme end of that distribution (see Hare and Neumann, 2008, for a review).
6
7 Findings from studies inspecting the behavioral and neurophysiological correlates of
8 psychopathic traits in the general population seem to mirror those observed in clinical/forensic
9 samples and suggest that there may exist continuities in the mechanisms underlying psychopathy
10 (see Koenigs, Baskin-Sommers, Zeier, & Newman, 2011; Seara-Cardoso & Viding, 2014, for
11 recent reviews).
12
13
14
15
16
17
18
19
20
21
22
23

24 With regards to empathic processing, evidence suggests that high levels of psychopathic traits in
25 the general population are associated with reduced emotional reactivity to aversive stimuli (e.g.
26 Benning, Patrick, & Iacono, 2005; Justus & Finn, 2007), as well as weaker self-reported
27 affective responses to others' emotional faces (Ali, Amorim, & Chamorro-Premuzic, 2009;
28 Seara-Cardoso, Dolberg, Neumann, Roiser, & Viding, 2013; Seara-Cardoso, Neumann, Roiser,
29 McCrory, & Viding, 2012). At the neural level, evidence suggests that in the general population
30 psychopathic traits are associated with atypical responses in brain regions including inferior
31 frontal gyrus, ventromedial prefrontal cortex and amygdala when processing emotional facial
32 expressions (Carré, Hyde, Neumann, Viding, & Hariri, 2013; Gordon, Baird, & End, 2004;
33 Hyde, Byrd, Votruba-Drzal, Hariri, & Manuck, 2014), when punishing others with electric
34 shocks (Molenberghs et al., 2014), as well as when rating one's own affective response to others'
35 emotional faces (Seara-Cardoso, Sebastian, Viding, & Roiser, under review). These findings
36 suggest that links between psychopathy and poor empathic responding extend throughout the
37 continuum of psychopathic traits both at the behavioral and neural level.
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

There is also clear evidence that youth and adults with extreme levels of psychopathic traits show atypical neural responses to others' pain when compared with healthy controls (Decety, Chen, et al., 2013; Decety, Skelly, et al., 2013; Lockwood et al., 2013; Marsh et al., 2013; Meffert et al., 2013). However, there is little evidence to suggest whether neural responses to others experiencing pain similarly vary continuously with psychopathic traits in the general adult population. Here, we employed the imaging paradigm and analysis strategy previously described in Lockwood et al. (2013), to study whether individual variability in neural responses to others' pain is associated with psychopathic traits in the general population. Lockwood et al. (2013) measured fMRI responses to pictures of others' hands and feet either in pain or in no pain (control condition) in a large sample of children with conduct problems and typically developing controls. As predicted, the children with conduct problems exhibited significantly reduced neural responses in regions previously associated with empathic processing, namely AI, IFG and ACC, in comparison to the typically developing control group. However, considerable heterogeneity of neural response was seen within the conduct problems group. When callous traits (similar to adult affective-interpersonal psychopathic traits) and conduct disorder symptoms (similar to adult lifestyle-antisocial behavior characteristics) were analyzed together as continuous independent variables in regression analyses, neural responses to others' pain were negatively associated with callous traits (in AI and ACC), but positively associated with conduct disorder symptoms (in ACC). These relationships only became apparent when the unique contribution of each of these variables was inspected, controlling for the other.

This pattern of opposing relationships between the two dimensions of psychopathic traits and measures of affective processing, with relationships only emerging after shared variance is

1
2
3 controlled for, is consistent with research to date suggesting that these two dimensions exert
4
5 suppressor effects on each other (e.g. Hicks & Patrick, 2006; Blonigen et al. 2010; Vanman et
6
7 al., 2003). Suppression, in this case cooperative suppression, occurs when two correlated
8
9 variables (as is the case for the two dimensions of psychopathic traits) present opposing
10
11 relationships with a given criterion variable, such that the inclusion of both concurrently in a
12
13 regression model increases the association of each with the criterion variable (Watson et. al,
14
15 2013). In other words, the association of each dimension of psychopathy is greater when the
16
17 variance shared with the other dimension is accounted for, because variance shared with the
18
19 other dimension does not present the same relationship with the criterion variable and therefore
20
21 suppresses the association (Blonigen et al., 2010). In psychopathy research, these suppressor
22
23 effects seem to indicate that, although affective-interpersonal and lifestyle-antisocial features
24
25 often co-occur and present shared components, there are also unique aspects of each dimension
26
27 (i.e. those not shared with the other dimension) that are related to distinct types of atypical
28
29 emotional and cognitive processing.
30
31
32
33
34
35
36
37
38

39 With respect to emotional processing, behavioral studies in both general and psychopathic
40
41 samples have shown that, when holding the other dimension constant, the affective-interpersonal
42
43 dimension (characterized by blunt affect and shallowness) is indeed associated with reduced
44
45 reactivity to emotional stimuli, whilst the lifestyle-antisocial dimension (characterized by
46
47 impulsivity and irresponsibility) is associated with increased reactivity to emotional stimuli (e.g.
48
49 Hicks & Patrick, 2006; Seara-Cardoso et al., 2012; Uzieblo, Verschuere, van den Bussche, &
50
51 Crombez, 2010; Blonigen et al., 2010; Vanman et al., 2003). This pattern of divergent
52
53 associations between the two dimensions of psychopathy has also been found at the neural level
54
55
56
57
58
59
60

1
2
3 in the amygdala (a region implicated in affective processing) in response to non-pain related
4 emotional stimuli in children with conduct problems (Lozier, Cardinale, VanMeter, & Marsh,
5 2014; Sebastian et al., 2012) and typical adults (Carré et al., 2013; Hyde et al., 2014), as well as
6
7
8 in AI and ACC during empathy processing in children with conduct problems, as discussed
9
10
11 above (Lockwood et al., 2013).
12
13
14

15
16
17 In sum, extant evidence indicates that individuals with extreme levels of psychopathy present a
18 pattern of reduced behavioral and neural response to others' suffering which may, in part,
19 explain some of their characteristic inappropriate social interactions. However, we do not yet
20 know if neural processing of others' pain relates to variability in psychopathic traits in those
21 individuals who function in the community. We used the methodology described in Lockwood et
22 al. (2013), to study whether neural responses to others' pain vary with psychopathic traits within
23 the general population in a similar manner to individuals with extreme levels of these traits. If
24 the neurobiological correlates of psychopathy vary along a continuum in the general population,
25 we would expect to find a pattern of neural responses in brain regions typically recruited during
26 empathic processing (i.e. anterior insula, IFG and mid/dACC) consistent with previous research
27 with individuals with extreme levels of psychopathic traits (Decety, Chen, et al., 2013; Decety,
28 Skelly, et al., 2013; Lockwood et al., 2013; Marsh et al., 2013; Meffert et al., 2013). More
29 specifically, consistent with the literature showing that affective-interpersonal and lifestyle-
30 antisocial dimensions of psychopathy may reflect distinct underlying vulnerabilities, we
31 predicted that these two dimensions of psychopathy would exert suppressor effects on each other
32 in relation to activity in these regions while viewing others' pain.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Methods

Participants

Fifty-three right-handed male participants from the community with no reported history of psychiatric illness were recruited for this study. Of these, six were excluded before pre-processing due to: failure to complete the task (2 participants); excessive response times (2 participants); one incidental finding (1 participant); and corrupted fMRI data due to excessive movement (1 participant). Analyses of the residuals from the multiple regression models inspecting the relationships between neural responses and psychopathic traits revealed one extreme outlier. This participant was excluded, leaving 46 participants in the analyses [mean age: 27.93, range: 19-40]. According to GPower software (Faul et al., 2007), a sample size between 38 (for one-tailed analyses) and 49 (for two-tailed analyses) is appropriate to detect an effect size of $R^2 = 0.17$, similar to the average effect size reported in Lockwood et al. (2013), at an alpha significance level of 0.05 with 80% power. Thus, an appropriate sample size was recruited. All participants provided written informed consent according to the guidelines approved by UCL Division of Psychology and Language Sciences Ethics Committee who provided ethical approval for this study.

Experimental Task

Stimuli were 192 digital photographs showing another person's hand or foot in painful or non-painful situations (taken from Gu et al., 2010). "Pain" and "No Pain" stimuli (96 pictures per condition) were matched on physical properties and were validated as eliciting empathy-related activations in a previous study (Gu et al., 2010). Stimuli were presented in 24 pain and no-pain

1
2
3 blocks each lasting 20 s and consisting of eight images, each displayed for 2,000 ms with a 500
4
5 ms interstimulus interval. Blocks were pseudorandomized, with the same block type never
6
7 presented more than twice in a row. A fixation cross was presented for 15 s every six blocks. To
8
9 ensure attention, participants performed a hand/foot key press judgment on every trial.
10
11 Participants practiced outside the scanner with painful and non-painful images not seen in the
12
13 main experiment, until $\geq 80\%$ accuracy was reached.
14
15
16
17

18 19 20 *Psychometric Measures*

21
22 Participants completed the *Self-Report Psychopathy Scale Short Form* (SRP-SF; Paulhus,
23
24 Neumann, & Hare, in press), a 29-item scale designed to measure psychopathic attributes in non-
25
26 institutionalized samples. The SRP-SF assesses psychopathic traits, organized in four facets –
27
28 interpersonal, affective, lifestyle and antisocial – consistent with recent research on the
29
30 Psychopathy Checklist – Revised (PCL-R; Hare, 2003). Higher scores on the SRP questionnaire
31
32 reflect higher levels of psychopathic traits. Like the PCL-R, the four facets can be modelled in
33
34 terms of the traditional two-factor dimensions: affective-interpersonal and lifestyle-antisocial.
35
36 The SRP has been shown to have a clear latent structure and good construct validity (Mahmut,
37
38 Menictas, Stevenson, & Homewood, 2011; Neumann & Pardini, 2012; Neumann, Schmitt,
39
40 Carter, Embley, & Hare, 2012), and is strongly correlated with the PCL-R (Lilienfeld & Fowler,
41
42 2006; Paulhus et al., in press). In the present sample, Cronbach's alpha for the total SRP scale
43
44 was .91, for the affective-interpersonal scale was .88 and for the lifestyle-antisocial scale was .84.
45
46 Affective-interpersonal scores varied between 15 and 61 ($M=29.85$; $SD=9.11$), lifestyle-
47
48 antisocial scores varied between 15 and 47 ($M=29.15$; $SD=8.89$), and the two scales presented a
49
50 correlation coefficient of $r = .66$ ($P < .001$), thus presenting a similar distribution to previously
51
52
53
54
55
56
57
58
59
60

1
2
3 reported distributions from larger samples of adults from the general population (Seara-Cardoso
4 et al., 2012; Foulkes et al., 2013; Foulkes et al., 2014; Paulhus et al., in press). Participants also
5 completed the State-Trait Anxiety Inventory (STAI; Spielberger et al., 1984), which comprises
6 two subscales measuring trait and state anxiety. The Matrix Reasoning subscale of the Wechsler
7 Abbreviated Scale of Intelligence (Wechsler, 1999) was administered to estimate general
8 intellectual ability.
9
10
11
12
13
14
15
16
17
18
19

20 *Magnetic resonance imaging acquisition*

21
22 Images were acquired using a Siemens Avanto 1.5 T MRI scanner at the Birkbeck-UCL Centre
23 for Neuroimaging with a 32-channel head coil. One-hundred-eighty-nine multislice T2*
24 weighted echo planar images (EPIs) with blood oxygenation-level-dependent (BOLD) contrast
25 were acquired in a single run of 9 min. The T2* EPI sequence was based on Weiskopf, Hutton,
26 Josephs, & Deichmann (2006) and used the following acquisition parameters: 35 2 mm slices
27 acquired in a descending trajectory with a 1 mm gap; echo time = 50 ms; repetition time = 2.975
28 s; slice tilt = -30°; flip angle = 90°; field of view = 192 mm; matrix size = 64 x 64. A 5.5 min T1-
29 weighted MPRAGE scan was acquired for coregistration, normalization, and overlay.
30
31
32
33
34
35
36
37
38
39
40
41
42
43

44 *Image processing and analyses*

45
46 EPI data were analysed using SPM8 (www.fil.ion.ucl.ac.uk/spm) in Matlab. The first five and
47 last two volumes were discarded. Data were realigned to the sixth volume, normalized to the
48 Montreal Neurological Institute template resampling to a voxel size of 2x2x2 mm, and smoothed
49 with an 8 mm full width at half-maximum Gaussian filter. Data were high-pass filtered at 128 s
50
51
52
53
54
55
56
57
58
59
60

1
2
3 to remove low-frequency drifts, and the statistical model included an AR(1) autoregressive
4
5 function to account for autocorrelations.
6
7

8
9
10 After preprocessing, a block analysis compared neural activity associated with pain and no-pain
11
12 conditions. Regressors included Pain and No-Pain (blocks of 20 s duration) and fixation (15 s),
13
14 modelled as boxcar functions convolved with a canonical hemodynamic response function. The
15
16 six realignment parameters were also modelled as effects of no interest. At the first level, Pain >
17
18 No-Pain and No-Pain > Pain contrasts were created. Contrast images were entered into second-
19
20 level analyses, where both SRP dimensions were either entered separately or simultaneously as
21
22 covariates in multiple regression models. Relationships between total SRP scores and BOLD
23
24 response were also examined.
25
26
27
28
29
30

31
32 Whole-brain analyses for the Pain > No Pain contrast are reported using a cluster forming
33
34 threshold of $P < .001$ (uncorrected, cluster size >10), with cluster-level family-wise error (FWE)
35
36 correction. Region-of-interest (ROI) analyses were conducted in four a priori regions of interest
37
38 (bilateral anterior insula, IFG, ACC and midCC). The first three were taken from Lockwood et
39
40 al. (2013) and the midCC was added because it regularly features in meta-analyses of empathy
41
42 for pain, with clusters bordering midCC and ACC (Fan et al., 2011; Lamm et al., 2011). ROI
43
44 analyses were conducted as described in Lockwood et al. (2013). ROIs were anatomically
45
46 defined using masks from the automated anatomical labelling (AAL) atlas (Maldjian, Laurienti,
47
48 Kraft, & Burdette, 2003), and the MarsBaR toolbox (<http://marsbar.sourceforge.net>) was used to
49
50 calculate average contrast estimates across bilateral ROIs and to conduct t-tests at a standard
51
52 statistical threshold of $p < .05$ (Eisenberger et al., 2010; Masten et al., 2011). Contrast estimates
53
54
55
56
57
58
59
60

1
2
3 extracted with Marsbar were also used in IBM SPSS and MS Excel to conduct regression
4
5 analyses and to generate the illustrative partial regression plots presented in Figure 1.
6
7
8
9

10 **Results**

11 *Behavioral Data*

12
13
14
15
16
17 Mean reaction times (RTs) and percentage error rates were calculated. Consistent with previous
18
19 studies (Gu et al., 2010; Lockwood et al., 2013), RTs during Pain were significantly slower than
20
21 No-Pain ($t(45)=5.76$, $p<.001$; Pain: $M=767.11$, $SD=106.62$; No-Pain: $M=738.10$, $SD=104.52$).
22
23

24
25 There were no significant differences in percentage error rates between conditions (Pain:
26
27 $M=2.97$; $SD=2.50$; No-Pain: $M=2.72$; $SD=2.36$).
28
29
30
31

32
33 On the basis of previous studies showing that unique variance associated with affective-
34
35 interpersonal and lifestyle-antisocial traits show opposing associations with emotional reactivity
36
37 (e.g. Carré et al., 2013; Hicks & Patrick, 2006; Lockwood et al., 2013; Lozier et al., 2014; Seara-
38
39 Cardoso et al., 2012), we conducted two regression analyses where both dimensions of
40
41 psychopathy were entered as predictors of the difference in mean RTs for Pain > No-Pain and
42
43 difference in percentage error rates for Pain > No-Pain, respectively. After controlling for levels
44
45 of the other dimension, lifestyle-antisocial traits presented a significant positive association with
46
47 difference in percentage error rates ($t=2.253$; $P=.03$, whilst affective-interpersonal traits
48
49 presented an at-trend negative association ($t=-1.72$; $P=.09$). That is, when holding the other
50
51 dimension constant, higher levels of affective-interpersonal traits were associated (at trend level)
52
53 with fewer errors during the Pain relative to No-Pain condition, whilst higher levels of lifestyle-
54
55
56
57
58
59
60

1
2
3 antisocial traits were associated with increased error rates in the Pain relative to No-Pain
4
5 condition. There were no significant associations with difference in RTs in Pain relative to No-
6
7 Pain condition. There were no significant bivariate associations between total SRP score or
8
9 either dimension of psychopathic traits and mean RT/percentage error rate differences between
10
11 conditions (all $P > .30$).
12
13
14
15
16
17

18 *Imaging results*

19
20 Results from the whole-brain analyses for the main effect of Pain > No-Pain are displayed in
21
22 Table 1 (see supplementary table S1 for No-Pain>Pain). Main effects were found in regions
23
24 previously associated with empathy for pain, and largely replicated previous studies using the
25
26 same stimuli (Gu et al., 2010; Lockwood et al., 2013), including bilateral IFG extending to
27
28 anterior insula ($P < .001$, FWE corrected at cluster level). ROI analyses for the main effect of Pain
29
30 > No Pain also revealed the predicted pattern of significant BOLD response in the bilateral AI
31
32 ($t[45] = 1.68$, $P = .05$) and IFG ($t[45] = 3.61$, $P < .001$), but not in midCC ($t[45] = .70$, $P = .24$)
33
34 and ACC ($t[45] = -.10$, $P = .34$). Additionally, entering difference (Pain > No-Pain) in error rate
35
36 and difference in RT between conditions as predictors of BOLD response in two SPM models
37
38 also showed that BOLD response in all ROIs presented significant positive relationships with
39
40 difference in error rate [AI ($t[44] = 3.08$, $P < 0.01$), IFG ($t[44] = 2.08$, $P = .02$), midCC ($t[44] =$
41
42 2.69 , $P < 0.01$) and ACC ($t[44] = 2.94$, $P < 0.01$)] and difference in RT between pain vs no-pain
43
44 conditions [(AI ($t[44] = 2.98$, $P < 0.01$), midCC ($t[44] = 1.69$, $P = 0.05$ and ACC ($t[44] = 1.97$, P
45
46 $= 0.03$); with the exception of the IFG which association was at-trend ($t[44] = 1.05$, $P = .15$)].
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1. Whole brain analyses showing main-effects for Pain > No-Pain BOLD response

Brain regions	Peak						Cluster	
	L/R	X	y	z	T	Z	Extent (k)	<i>P</i> (FWE)
Middle temporal gyrus	L	-44	-68	-2	11.75	>8	3540	<.001
Occipital gyrus	R	32	-84	1	10.83	7.56	3367	<.001
Supramarginal gyrus	L	-56	-30	32	10.10	7.26	1326	<.001
Supramarginal gyrus	R	66	-24	38	8.45	6.50	854	<.001
Precentral gyrus	L	-50	4	30	6.97	5.71	818	<.001
Cerebellum	R	16	-76	-50	6.21	5.25	191	.03
Inferior frontal gyrus	R	56	38	6	5.51	4.79	369	<.001
Insula	L	-38	-4	-10	4.79	4.48	81	.33
Precentral gyrus	R	50	6	30	4.48	4.27	388	<.001
Inferior frontal gyrus	L	-52	38	6	4.26	3.88	51	.60
Inferior frontal gyrus	L	-40	28	0	4.25	3.88	206	.03
Ext. Insula	L	-32	28	4	4.17	3.82		
Postcentral gyrus	L	32	-34	42	4.25	3.88	29	.85
Amygdala	R	22	-4	-14	4.21	3.84	42	.70
Cerebellum	R	16	-74	-50	3.76	3.49	50	.61

Notes: Whole-brain analysis reported at a threshold level of $P < .001$ (uncorrected, cluster size > 10

voxels). Spatial coordinates (x, y, z) are in Montreal Neurological Institute space. R = Right; L = Left.

On the basis of previous studies showing that unique variance associated with affective-interpersonal and lifestyle-antisocial traits show opposing associations with emotional reactivity (e.g. Carré et al., 2013; Hicks & Patrick, 2006; Lockwood et al., 2013; Lozier et al., 2014; Seara-Cardoso et al., 2012), we entered both dimensions of psychopathy as predictors in a single

1
2
3 multiple regression model at the second level in SPM, and tested whether neural response in our
4 ROIs was associated with each dimension individually after controlling for the other (see Figure
5
6
7
8 1). As predicted, ROI analyses for Pain > No-Pain revealed that, after controlling for lifestyle-
9
10 antisocial traits, unique variance associated with affective-interpersonal traits was negatively
11
12 related to BOLD response in AI ($t[43] = 1.87, P = 0.03$), IFG ($t[43] = 2.68, P < .01$), and midCC
13
14 ($t[43] = 2.38, P = 0.01$), and at-trend in ACC ($t[43] = 1.24, P = 0.11$). That is, when holding
15
16 levels of lifestyle-antisocial behavior constant, increased levels of affective-interpersonal traits
17
18 were associated with a decrease in neural responses to others' pain in these regions. After
19
20 controlling for affective-interpersonal traits, unique variance associated with lifestyle-antisocial
21
22 traits was positively related to differential BOLD response in AI ($t[43] = 2.51, P < 0.01$), IFG
23
24 ($t[43] = 3.16, P < 0.01$), midCC ($t[43] = 2.64, P < 0.01$) and ACC ($t[43] = 1.92, P = 0.03$). That
25
26 is, when holding levels of affective-interpersonal traits constant, increased levels of lifestyle-
27
28 antisocial behavior traits were associated with an increase in neural responses to others' pain in
29
30 these regions.
31
32
33
34
35
36
37
38

39 To exclude potential confounds of trait anxiety and cognitive ability, we included trait anxiety
40
41 and estimated IQ as additional covariates in follow-up regression models. Including these
42
43 variables did not change the pattern of results (all significant results remained at $P < .05$, with the
44
45 exception of the association of lifestyle-antisocial traits and BOLD response in ACC ($P = .12$).
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1. Partial regression plots showing opposing relationships between response to Pain > No-Pain in bilateral AI, IFG and midCC, and unique variance associated with affective-interpersonal and lifestyle-antisocial psychopathic traits after controlling for each other (a similar pattern was also seen in the ACC, adjacent to midCC). Left: negative relationships between BOLD response to Pain > No-Pain and affective-interpersonal traits after controlling for the effect of lifestyle-antisocial traits. Right: positive relationships between BOLD response to Pain > No-Pain and lifestyle-antisocial traits after controlling for the effect of affective-interpersonal traits. R^2 reflects partial correlation coefficient of determination. Insets show horizontal and mid-sagittal sections of bilateral AI ($z=0$), IFG ($z=15$) and midCC ($x=0$) ROIs overlaid on an average T1 structural image from all participants.

1
2
3 To test whether these opposing results were genuine suppressor effects, we inspected the
4
5 bivariate associations between psychopathic dimensions and total score and differential BOLD
6
7 response in three separate regression models. These analyses revealed weaker and largely non-
8
9 significant bivariate associations between neural responses in our ROIs and affective-
10
11 interpersonal traits [AI ($t[44] = .27, P = .39$); IFG ($t[44] = .72, P = .24$); midCC ($t[44] = .80, P =$
12
13 $.21$); ACC ($t[44] = .04, P = .97$)], lifestyle-antisocial traits [AI ($t[44] = 1.66, P = .05$); IFG ($t[44]$
14
15 $= 1.74, P = .44$); midCC ($t[44] = 1.36, P = .09$); ACC ($t[44] = 1.46, P = .15$)] and total
16
17 psychopathy score [AI ($t[44] = .72, P = .24$); IFG ($t[44] = .60, P = .55$); midCC ($t[44] = .27, P =$
18
19 $.40$); ACC ($t[44] = .44, P = .22$)].
20
21
22
23
24
25
26

27 For completeness, due to previous research linking amygdala dysfunction to psychopathic traits,
28
29 we inspected whether neural responses in amygdala varied as a function of psychopathic traits.
30
31 No significant associations were found.
32
33
34
35

36 Discussion

37
38
39
40
41 Neuroimaging studies have shown that the observation of other people experiencing distress, in
42
43 particular pain, elicits robust activation in AI, IFG and midCC/ACC (Fan et al., 2011; Lamm et
44
45 al., 2011). Consistent with the idea that individuals with extreme levels of psychopathy do not
46
47 find other people's distress as salient as their peers do (Blair, 2005), it has been reported that
48
49 these individuals show atypical neural activity in these regions in response to others' pain, when
50
51 compared with healthy controls (Decety, Chen, et al., 2013; Decety, Skelly, et al., 2013;
52
53 Lockwood et al., 2013; Marsh et al., 2013; Meffert et al., 2013). However, although affective
54
55
56
57
58
59
60

1
2
3 dysfunction is considered to be a critical, defining feature of psychopathy (Blair et al., 2005)
4
5 there is little evidence as to whether empathic neural responses to others' pain vary continuously
6
7 with psychopathic traits in typical adults. That is, as to whether the pattern of **relationships**
8
9 **between psychopathic personality traits and** neural response to others' pain observed at the
10
11 extreme end of the psychopathy distribution is also observed in a non-clinical distribution of
12
13 these traits in functioning members of the general population.
14
15
16

17
18
19
20 In line with predictions, we found that psychopathic traits were significantly associated with
21
22 neural responses to stimuli depicting others experiencing pain in AI, IFG, and midCC/ACC.
23
24 More specifically, we found suppressor effects between the two dimensions of psychopathy in
25
26 terms of their relationships with neural responses to others' pain in these regions. Unique
27
28 variance associated with affective-interpersonal traits was negatively associated with neural
29
30 responses to others' pain in these regions, whilst at the same time unique variance in lifestyle-
31
32 antisocial traits was positively associated with neural responses.
33
34
35
36
37
38

39 It has been proposed that the AI, IFG and midCC/ACC play separate but complementary roles in
40
41 empathic processing. The AI is proposed to be critical for sensory integration (Critchley, Wiens,
42
43 Rotshtein, Öhman, & Dolan, 2004), for the representation and integration of feeling states
44
45 (Craig, 2009) and for effectively discriminating between emotionally salient and non-salient
46
47 information (Gu et al., 2012). The midCC/ACC, with extensive connections from the
48
49 somatosensory cortices, and to and from the insula, amygdala, ventral striatum and periaqueductal
50
51 gray, is thought to be a hub region in affective, cognitive and motor control and, ultimately, to
52
53 influence motor centers responsible for expressing affect and executing goal-directed behavior
54
55
56
57
58
59
60

1
2
3 (Bernhardt & Singer, 2012; Shackman et al., 2011). Whereas the AI is thought to serve as an
4 input region of the system, translating sensations into subjective feelings and awareness, the
5
6 cingulate may function as an output region, exerting volitional control (Gu et al., 2012). The
7
8 IFG, on the other hand, is thought to play a role in emotional contagion and emotional
9
10 recognition (Shamay-Tsoory, 2011), as well as in emotion regulation and pain suppression
11
12 (Ochsner & Gross, 2005; Salomons, Johnstone, Backonja, Shackman, & Davidson, 2007; Wager,
13
14 Davidson, Hughes, Lindquist, & Ochsner, 2008).
15
16
17
18
19
20
21

22 Psychopathic traits are characterized by lack of empathy, disregard for other people's feelings,
23
24 impulsiveness and antisocial behavior. It would therefore be unsurprising that individuals with
25
26 extreme levels of these traits presented atypical engagement of the regions outlined above when
27
28 they are faced with others' distress. We did not find significant bivariate associations between
29
30 psychopathic traits and neural responses in these regions, **which could be due to sample size**
31
32 **and lack of statistical power or to limited range of scores in the extreme end of**
33
34 **psychopathic traits in our sample, and** associations only became apparent once the two
35
36 dimensions of psychopathic traits were inspected and their shared variance was controlled for.
37
38 **However, and in line with previous research with clinical/forensic samples (e.g. Hicks &**
39
40 **Patricks, 2006; Vanman et al., 2003; Lockwood et al., 2012),** we found a cooperative
41
42 suppression effect between the affective-interpersonal and lifestyle-antisocial dimensions of
43
44 psychopathy and neural responses to others' pain. This cooperative suppression effects occurs
45
46 because the two dimensions of psychopathic traits are correlated with each other but present
47
48 opposing relationships with neural responses to pain-related stimuli in these regions. The
49
50 association of each dimension of psychopathy with neural responses becomes apparent when the
51
52
53
54
55
56
57
58
59
60

1
2
3 shared variance is accounted for, that is, when the other dimension is held constant. The variance
4
5 shared with the other dimension does not present the same relationship with the criterion variable
6
7 and therefore suppresses the association (Blonigen et al., 2010).
8
9

10
11
12 We also observed a pattern of cooperative suppression between the two dimensions of
13
14 psychopathic traits and the difference in error rate between pain and no-pain conditions. More
15
16 specifically, we found that when holding the other dimension constant, lifestyle-antisocial traits
17
18 presented a positive association with difference in error rate between conditions whilst affective-
19
20 interpersonal traits present a negative association with difference in error rate. That is, when
21
22 holding affective-interpersonal traits constant, higher lifestyle-antisocial traits corresponded to a
23
24 higher rate of errors made in the pain vs no-pain condition; whilst when holding lifestyle-
25
26 antisocial traits constant higher affective-interpersonal traits corresponded to a reduced rate of
27
28 errors made in the pain vs no-pain condition (at trend levels). An increased error rate in the Pain
29
30 condition is thought to result from increased reactivity to the emotional content of the stimuli
31
32 (i.e. depicting others' pain in comparison to no-pain) and consequent interference in task
33
34 performance (hand/feet judgment) (Gu et al., 2013). These results are in line with the notion that
35
36 higher levels of affective-interpersonal traits are accompanied by less reactivity to stimuli
37
38 depicting others' pain (reflected by less 'interference' by others' pain and lower error rate) and
39
40 higher levels of lifestyle-antisocial are accompanied by higher reactivity to the same stimuli
41
42 (reflected by higher interference and higher error rate). Furthermore, we found significant
43
44 positive associations between difference in error rate and neural responses in all above
45
46 mentioned regions, that is, those participants who presented higher difference in error rates in
47
48 pain vs no-pain also presented higher neural responses in these regions. This corroborates the
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 notion that individual differences in reactivity to others' pain influences neural responses in the
4
5
6 neural circuitry thought to be involved in empathy for pain.
7
8
9

10 We found that unique variance associated with the two dimensions of psychopathic traits,
11
12 affective-interpersonal and lifestyle-antisocial, presented opposing associations with neural
13
14 response to pain (relative to no pain) in anterior insula, IFG, midCC and ACC. After shared
15
16 variance with lifestyle-antisocial traits was removed, affective-interpersonal traits (characterized
17
18 by lack of consideration for others' well-being) presented negative associations with neural
19
20 response in AI, IFG, and midCC, which is consistent with the characteristic lack of arousal to
21
22 others' distress and general blunted affect associated with this set of traits. In contrast, after
23
24 removing variance associated with affective-interpersonal traits, lifestyle-antisocial
25
26 characteristics (marked by poor inhibitory control), were positively associated with response in
27
28 these regions, consistent with evidence showing that these traits are associated with hyper-
29
30 reactivity to emotional stimuli and poor emotional and behavioral regulation in both extreme and
31
32 typical samples (Carré et al., 2013; Hyde et al., 2014; Patrick, Hicks, Krueger, & Lang, 2005;
33
34 Seara-Cardoso et al., 2012). Our results are in line with and provide further evidence for the
35
36 notion that, although the two dimensions of psychopathy co-occur, they may tap into two
37
38 distinctive underlying constructs. These constructs share components but also present unique
39
40 aspects (i.e. those not shared with the other) that are related to distinct types of atypical
41
42 emotional processing (Patrick et al., 2007). For example, variance in lifestyle-antisocial
43
44 behaviors in the general population may stem from multiple different sources. Individuals who
45
46 present with these behaviors may do so because they lack empathy and concern for others (low
47
48 emotional reactivity) or they may show reactive aggression to threat (increased emotional
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 reactivity). Once variance shared with affective-interpersonal traits is controlled for, what is left
4
5 is variance in lifestyle-antisocial behavior that is driven by factors other than those which are in
6
7 common with affective-interpersonal traits. **Likewise, individuals with high levels of affective-**
8
9 **interpersonal traits may differ on their levels of antisocial behavior. Individuals with high**
10
11 **levels of affective-interpersonal traits but low levels of lifestyle-antisocial behavior seem to**
12
13 **present significantly higher education and intelligence than those with both high levels of**
14
15 **affective-interpersonal traits and antisocial behavior (Hervé, 2007; Mokros et al., 2015).**
16
17 **These two groups have been referred to as ‘manipulative’ and ‘aggressive psychopaths’,**
18
19 **respectively, illustrating their distinct behavioral profile (Hervé, 2007). The neurocognitive**
20
21 **profiles of these two groups have not been explored and it would be of interest to assess**
22
23 **whether their distinct patterns of behavior rest upon distinct patterns of emotional**
24
25 **reactivity.**
26
27
28
29
30
31
32
33

34 It should be noted, however, that although the use of partial correlations is a powerful and
35
36 informative technique to identify associations between different variables, it also poses some
37
38 difficulties in the interpretation of results (Lynam, Hoyle, & Newman, 2006). The most
39
40 important one is the difficulty in knowing exactly what construct is left once the variance of
41
42 another correlated construct is removed (Lynam et al., 2006). The replication of the present
43
44 findings using larger sample with a group comparison approach, with groups defined by high and
45
46 low levels on the two dimensions, would provide important further validation of these results.
47
48 However, it is worth noting that this approach has its own limitations, for example owing to the
49
50 moderate positive correlation between the two dimensions, individuals high on one dimension
51
52
53
54
55
56
57
58
59
60

1
2
3 but not the other may be difficult to recruit and somewhat unrepresentative of how these traits
4
5 are distributed.
6
7

8
9
10 Finally, it is worth noting the pattern of associations found in midCC/ACC. We found
11
12 significant associations between midCC response and both dimensions of psychopathic traits;
13
14 whilst the association between ACC response was significant with lifestyle-antisocial traits, and
15
16 at-trend with affective-interpersonal traits. According to the meta-analyses conducted by Fan et
17
18 al. (2011) and Lamm et al. (2011), the region implicated in empathy for pain spans the border
19
20 between these two regions. However, in spite of these associations with individual differences,
21
22 we did not detect a main effect of Pain > No-Pain in these regions. Correlations and main effects
23
24 are statistically distinct and, for any given region and any given process, each can be observed in
25
26 isolation or both can occur (Calder, Ewbank, & Passamonti, 2011). When a robust correlation
27
28 with a personality trait is found in the absence of a group main effect, it is likely because lower
29
30 and higher scores on the personality dimension are associated with relative reductions and
31
32 increases in the neural response. This produces an overall effect that does not significantly differ
33
34 from zero, thus rendering the main-effect of task manipulation in that region non-significant.
35
36
37
38
39
40
41
42

43 **Conclusions**

44
45
46
47
48 In summary, we demonstrate that neural responses to others' pain in AI, IFG and midCC, regions
49
50 typically associated with empathic processing, are associated with variation in psychopathic
51
52 traits in the general population. Strikingly, the two dimensions of psychopathy presented
53
54 opposite associations with neural response in these regions. These results provide further
55
56
57
58
59
60

1
2
3 evidence for the notions that atypical function in these regions might represent neural markers of
4
5 disrupted emotional and empathic processing for individuals with high levels of psychopathic
6
7 traits; that the two dimensions of psychopathy tap into two separable constructs, with distinct
8
9 underlying vulnerabilities; and, finally, that the relationships observed at the extreme end of the
10
11 psychopathy distribution apply to non-clinical distribution of these traits in the general
12
13 population, i.e. that there are continuities in the mechanisms underlying psychopathy.
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Acknowledgements

This work was supported by a Portuguese Foundation for Science and Technology Doctoral Grant (Fundação para a Ciência e Tecnologia; SFRH/BD/60279/2009) awarded to A.S.C.; a British Academy Small Research Grant (SG101362) awarded to C.L.S., an Economic and Social Research Council (RES-062-23-2202) award to E.V., and a Wellcome Trust Biomedical Vacation Scholarship awarded to R.A.L. E.V. is a Royal Society Wolfson Research Merit Award holder. C.L.S. was partially supported during the writing of this paper by an Economic and Social Research Council award (ES/K008951/1). We would like to thank Prof. Craig Neumann for his help and advice with regard to the Self-Report Psychopathy Scale.

Conflict of interest

None.

Ethical Standards

The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008.

References

- 1
2
3
4
5
6 Ali, F., Amorim, I. S., & Chamorro-Premuzic, T. (2009). Empathy deficits and trait emotional
7
8 intelligence in psychopathy and Machiavellianism. *Personality and Individual Differences*, 47(7),
9
10 758-762.
11
- 12 Benning, S. D., Patrick, C. J., & Iacono, W. G. (2005). Psychopathy, startle blink modulation, and
13
14 electrodermal reactivity in twin men. *Psychophysiology*, 42(6), 753-762. doi: 10.1111/j.1469-
15
16 8986.2005.00353.x
17
- 18 Bernhardt, B. C., & Singer, T. (2012) The neural basis of empathy. *Vol. 35* (pp. 1-23).
19
- 20 Blair, R. J. R. (2013). The neurobiology of psychopathic traits in youths. *Nature Reviews Neuroscience*,
21
22 14(11), 786-799. doi: 10.1038/nrn3577
23
- 24 Blair, R. J. R., Colledge, E., Murray, L., & Mitchell, D. G. V. (2001). A selective impairment in the
25
26 processing of sad and fearful expressions in children with psychopathic tendencies. *Journal of*
27
28 *Abnormal Child Psychology*, 29(6), 491-498.
29
- 30 Blair, R. J. R., Mitchell, D., & Blair, K. (2005). *The psychopath: Emotion and the brain*. Malden, MA:
31
32 Blackwell Publishing.
33
- 34 Blair, R. J. R., Mitchell, D. G. V., Peschardt, K. S., Colledge, E., Leonard, R. A., Shine, J. H., . . . Perrett,
35
36 D. I. (2004). Reduced sensitivity to others' fearful expressions in psychopathic individuals.
37
38 *Personality and Individual Differences*, 37(6), 1111-1122. doi: 10.1016/j.paid.2003.10.008
39
40
- 41 Blair, R. J. R., Mitchell, D. G. V., Richell, R. A., Kelly, S., Leonard, A., Newman, C., & Scott, S. K.
42
43 (2002). Turning a deaf ear to fear: Impaired recognition of vocal affect in psychopathic
44
45 individuals. *Journal of Abnormal Psychology*, 111(4), 682-686. doi: 10.1037//0021-
46
47 843x.111.4.682
48
49
- 50 Blonigen, D. M., Patrick, C. J., Douglas, K. S., Poythress, N. G., Skeem, J. L., Lilienfeld, S. O., Edens, J.
51
52 F., & Krueger, R. F. (2010). Multi-method assessment of psychopathy in relation to factors of
53
54
55
56
57
58
59
60

- 1
2
3 internalizing and externalizing from the personality assessment inventory: The impact of method
4 variance and suppressor effects. *Psychological Assessment*, 22(1), 96-107.
5
6
7
8 Calder, A. J., Ewbank, M., & Passamonti, L. (2011). Personality influences the neural responses to
9 viewing facial expressions of emotion. *Philosophical Transactions of the Royal Society B:
10 Biological Sciences*, 366(1571), 1684-1701. doi: 10.1098/rstb.2010.0362
11
12
13 Carré, J. M., Hyde, L. W., Neumann, C. S., Viding, E., & Hariri, A. R. (2013). The neural signatures of
14 distinct psychopathic traits. *Social Neuroscience*, 8(2), 122-135.
15
16
17
18 Craig, A. D. (2009). How do you feel - now? The anterior insula and human awareness. *Nature Reviews
19 Neuroscience*, 10(1), 59-70.
20
21
22
23 Critchley, H. D., Wiens, S., Rotshtein, P., Öhman, A., & Dolan, R. J. (2004). Neural systems supporting
24 interoceptive awareness. *Nature Neuroscience*, 7(2), 189-195.
25
26
27 Decety, J., Chen, C., Harenski, C., & Kiehl, K. A. (2013). An fMRI study of affective perspective taking
28 in individuals with psychopathy: Imagining another in pain does not evoke empathy. *Frontiers in
29 Human Neuroscience* (SEP).
30
31
32
33 Decety, J., Skelly, L. R., & Kiehl, K. A. (2013). Brain response to empathy-eliciting scenarios involving
34 pain in incarcerated individuals with psychopathy. *JAMA Psychiatry*, 70(6), 638-645.
35
36
37
38 Eisenberger, N. I., Berkman, E. T., Inagaki, T. K., Rameson, L. T., Mashal, N. M., & Irwin, M. R. (2010).
39 Inflammation-induced anhedonia: Endotoxin reduces ventral striatum responses to reward.
40
41
42
43 *Biological Psychiatry*, 68(8), 748-754.
44
45
46
47 Fan, Y., Duncan, N. W., de Greck, M., & Northoff, G. (2011). Is there a core neural network in empathy?
48 An fMRI based quantitative meta-analysis. *Neuroscience and Biobehavioral Reviews*, 35(3), 903-
49 911.
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160

1
2
3
4
5 Foulkes, L., Seara-Cardoso, A., Neumann, C.S., Rogers, J.S.C., & Viding, E. (2014). Looking after
6 number one: Associations between psychopathic traits and measures of social motivation and
7 functioning in a community sample of males. *Journal of Psychopathology and Behavioural*
8 *Assessment*, 36(1), 22-29.

14 Foulkes, L., McCrory, E., Neumann, C.S., & Viding, E. (2014). Inverted social reward: associations
15 between psychopathic traits and self-report and experimental measures of social reward. *PLoS*
16 *One*, 9(8), e106000.

20 Gordon, H. L., Baird, A. A., & End, A. (2004). Functional differences among those high and low on a
21 trait measure of psychopathy. *Biological Psychiatry*, 56(7), 516-521.

24 Gu, X., Gao, Z., Wang, X., Liu, X., Knight, R. T., Hof, P. R., & Fan, J. (2012). Anterior insular cortex is
25 necessary for empathetic pain perception. *Brain*, 135(9), 2726-2735.

28 Gu, X., Liu, X., Guise, K. G., Naidich, T. P., Hof, P. R., & Fan, J. (2010). Functional dissociation of the
29 frontoinsular and anterior cingulate cortices in empathy for pain. *Journal of Neuroscience*,
30 30(10), 3739-3744.

33 Gu, X., Liu, X., Van Dam, N.T., Hof, P.R., & Fan, J. (2013). Cognition-emotion integration in the
34 anterior insula cortex. *Cerebral Cortex*, 23(1), 20-27.

37 Hare, R. D. (1993). *Without conscience: The disturbing world of the psychopaths among us*. New York,
38 NY: Simon & Schuster.

41 Hare, R. D. (2003). *The Hare Psychopathy Checklist - Revised* (2nd edition ed.). Toronto, ON: Multi-
42 Health Systems.

45 Hare, R. D., & Neumann, C. S. (2008). Psychopathy as a clinical and empirical construct. *Annual Review*
46 *of Clinical Psychology*, 4, 217-246.

49 Hervé, H. (2007). Psychopathic subtypes: Historical and contemporary perspectives. In J. C. Yuille & H.
50 Hervé (Eds.). *The psychopath: Theory, research and practice*, 431-460. Mahwah, NJ: Erlbaum.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- Hicks, B. M., & Patrick, C. J. (2006). Psychopathy and negative emotionality: Analyses of suppressor effects reveal distinct relations with emotional distress, fearfulness, and anger-hostility. *Journal of Abnormal Psychology, 115*(2), 276-287. doi: 10.1037/0021-843x.115.2.276
- Hyde, L. W., Byrd, A. L., Votruba-Drzal, E., Hariri, A. R., & Manuck, S. B. (2014). Amygdala reactivity and negative emotionality: Divergent correlates of antisocial personality and psychopathy traits in a community sample. *Journal of Abnormal Psychology, 123*(1), 214-224. doi: 10.1037/a0035467
- Justus, A. N., & Finn, P. R. (2007). Startle modulation in non-incarcerated men and women with psychopathic traits. *Personality and Individual Differences, 43*(8), 2057-2071.
- Koenigs, M., Baskin-Sommers, A., Zeier, J., & Newman, J. P. (2011). Investigating the neural correlates of psychopathy: A critical review. *Molecular Psychiatry, 16*(8), 792-799. doi: 10.1038/mp.2010.124
- Lamm, C., Decety, J., & Singer, T. (2011). Meta-analytic evidence for common and distinct neural networks associated with directly experienced pain and empathy for pain. *NeuroImage, 54*(3), 2492-2502.
- Levenston, G. K., Patrick, C. J., Bradley, M. M., & Lang, P. J. (2000). The psychopath as observer: Emotion and attention in picture processing. *Journal of Abnormal Psychology, 109*(3), 373-385. doi: 10.1037//0021-843x.109.3.373
- Lilienfeld, S. O., & Fowler, K. A. (2006). The self-report assessment of psychopathy: Problems, pitfalls, and promises. In C. J. Patrick (Ed.), *Handbook of Psychopathy* (pp. 107-132). New York: Guilford Press.
- Lockwood, P. L., Sebastian, C. L., McCrory, E. J., Hyde, Z. H., Gu, X., De Brito, S. A., & Viding, E. (2013). Association of callous traits with reduced neural response to others' pain in children with conduct problems. *Current Biology, 23*(10), 901-905.
- Lozier, L. M., Cardinale, E. M., VanMeter, J. W., & Marsh, A. A. (2014). MEdition of the relationship between callous-unemotional traits and proactive aggression by amygdala response to fear among children with conduct problems. *JAMA Psychiatry*. doi: 10.1001/jamapsychiatry.2013.4540

- 1
2
3 Lynam, D. R., Hoyle, R. H., & Newman, J. P. (2006). The perils of partialling: Cautionary tales from
4 aggression and psychopathy. *Assessment, 13*(3), 328-341.
5
6
7 Mahmut, M. K., Menictas, C., Stevenson, R. J., & Homewood, J. (2011). Validating the Factor Structure
8 of the Self-Report Psychopathy Scale in a Community Sample. *Psychological Assessment, 23*(3),
9 670-678.
10
11
12
13
14 Maldjian, J. A., Laurienti, P. J., Kraft, R. A., & Burdette, J. H. (2003). An automated method for
15 neuroanatomic and cytoarchitectonic atlas-based interrogation of fMRI data sets. *NeuroImage,*
16 *19*(3), 1233-1239. doi: 10.1016/s1053-8119(03)00169-1
17
18
19
20
21 Marsh, A. A., Finger, E. C., Fowler, K. A., Adalio, C. J., Jurkowitz, I. T. N., Schechter, J. C., . . . Blair, R.
22 J. R. (2013). Empathic responsiveness in amygdala and anterior cingulate cortex in youths with
23 psychopathic traits. *Journal of Child Psychology and Psychiatry and Allied Disciplines.*
24
25
26
27 Masten, C. L., Colich, N. L., Rudie, J. D., Bookheimer, S. Y., Eisenberger, N. I., & Dapretto, M. (2011).
28 An fMRI investigation of responses to peer rejection in adolescents with autism spectrum
29 disorders. *Developmental Cognitive Neuroscience, 1*(3), 260-270.
30
31
32
33
34 Meffert, H., Gazzola, V., Den Boer, J. A., Bartels, A. A. J., & Keysers, C. (2013). Reduced spontaneous
35 but relatively normal deliberate vicarious representations in psychopathy. *Brain, 136*(8), 2550-
36 2562. doi: 10.1093/brain/awt190.
37
38
39
40
41 Mokros, A., Hare, R.D., Neumann, C.S., Santtila, p., Habermeyer, E., & Nistchke, J. (2015). Variants of
42 psychopathy in adult male offenders: A latent profile analysis. *Journal of Abnormal Psychology.*
43
44
45 Molenberghs, P., Bosworth, R., Nott, Z., Louis, W. R., Smith, J. R., Amiot, C. E., . . . Decety, J. (2014).
46 The influence of group membership and individual differences in psychopathy and perspective
47 taking on neural responses when punishing and rewarding others. *Human Brain Mapping, n/a-*
48 *n/a.* doi: 10.1002/hbm.22527
49
50
51
52
53 Neumann, C. S., & Pardini, D. A. (2012). Factor Structure and Construct Validity of the Self-Report
54 Psychopathy (SRP) Scale and the Youth Psychopathic Traits Inventory (YPI) in Young Men.
55 *Journal of Personality Disorders, 1-15.* doi: 10.1521/pedi_2012_26_063
56
57
58
59
60

- 1
2
3 Neumann, C. S., Schmitt, D. S., Carter, R., Embley, I., & Hare, R. D. (2012). Psychopathic Traits in
4 Females and Males across the Globe. *Behavioral Sciences and the Law*, 30(5), 557-574. doi:
5 10.1002/bsl.2038
6
7
8
9
10 Nichols, S. (2001). Mindreading and the cognitive architecture underlying altruistic motivation. *Mind and*
11 *Language*, 16, 425-455.
12
13
14 Ochsner, K. N., & Gross, J. J. (2005). The cognitive control of emotion. *Trends in Cognitive Sciences*,
15 9(5), 242-249.
16
17
18 Patrick, C. J., Bradley, M. M., & Lang, P. J. (1993). Emotion in the criminal psychopath: Startle reflex
19 modulation. *Journal of Abnormal Psychology*, 102(1), 82-92. doi: 10.1037//0021-843x.102.1.82
20
21
22 Patrick, C. J., Hicks, B. M., Krueger, R. F., & Lang, A. R. (2005). Relations between psychopathy facets
23 and externalizing in a criminal offender sample. *Journal of Personality Disorders*, 19(4), 339-
24 356.
25
26
27
28 Patrick, C. J., Hicks, B. M., Nichol, P. E., & Krueger, O. F. (2007). A bifactor approach to modeling the
29 structure of the psychopathy checklist-revised. *Journal of Personality Disorders*, 21(2), 118-141.
30
31
32 Paulhus, D. L., Neumann, C. S., & Hare, R. D. (in press). *Manual for the Self-Report Psychopathy Scale*
33 (4th ed.). Toronto: Multi-Health Systems.
34
35
36
37 Salomons, T. V., Johnstone, T., Backonja, M. M., Shackman, A. J., & Davidson, R. J. (2007). Individual
38 differences in the effects of perceived controllability on pain perception: Critical role of the
39 prefrontal cortex. *Journal of Cognitive Neuroscience*, 19(6), 993-1003.
40
41
42
43 Seara-Cardoso, A., Dolberg, H., Neumann, C., Roiser, J. P., & Viding, E. (2013). Empathy, morality and
44 psychopathic traits in women. *Personality and Individual Differences*, 55(3), 328-333.
45
46
47
48 Seara-Cardoso, A., Neumann, C., Roiser, J., McCrory, E., & Viding, E. (2012). Investigating associations
49 between empathy, morality and psychopathic personality traits in the general population.
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 Seara-Cardoso, A., Sebastian, C. L., Viding, E., & Roiser, J. (under review). Affective resonance in
4
5 response to others' emotional faces varies with affective ratings and psychopathic traits in
6
7 amygdala and anterior insula.
8
9
10 Seara-Cardoso, A., & Viding, E. (2014). Functional Neuroscience of Psychopathic Personality in Adults.
11
12 *Journal of Personality*, n/a-n/a. doi: 10.1111/jopy.12113
13
14 Sebastian, C. L., McCrory, E. J. P., Cecil, C. A. M., Lockwood, P. L., De Brito, S. A., Fontaine, N. M. G.,
15
16 & Viding, E. (2012). Neural responses to affective and cognitive theory of mind in children with
17
18 conduct problems and varying levels of callous-unemotional traits. *Archives of General*
19
20 *Psychiatry*, 69(8), 814-822. doi: 10.1001/archgenpsychiatry.2011.2070
21
22
23 Shackman, A. J., Salomons, T. V., Slagter, H. A., Fox, A. S., Winter, J. J., & Davidson, R. J. (2011). The
24
25 integration of negative affect, pain and cognitive control in the cingulate cortex. *Nature Reviews*
26
27 *Neuroscience*, 12(3), 154-167. doi: 10.1038/nrn2994
28
29
30 Shamay-Tsoory, S. G. (2011). The neural bases for empathy. *Neuroscientist*, 17(1), 18-24.
31
32 Singer, T., & Lamm, C. (2009). The social neuroscience of empathy. *Annals of the New York Academy of*
33
34 *Sciences*, 1156, 81-96.
35
36
37 Spielberg, C. D. (1984). *State-Trait Anxiety Inventory*. Consulting Psychologists Press, Inc.
38
39 Uzieblo, K., Verschuere, B., van den Bussche, E., & Crombez, G. (2010). The validity of the
40
41 psychopathic personality inventory-revised in a community sample. *Assessment*, 17(3), 334-346.
42
43
44 Vanman, E.J., Mejia, V.Y., Dawson, M.E., Schell, A.M., & Raine, A. (2003). Modification of the startle
45
46 reflex in a community sample: do one or two dimensions of psychopathy underlie emotional
47
48 processing? *Personality and Individual Differences*, 35, 2007-2021.
49
50 Wager, T. D., Davidson, M. L., Hughes, B. L., Lindquist, M. A., & Ochsner, K. N. (2008). Prefrontal-
51
52 Subcortical Pathways Mediating Successful Emotion Regulation. *Neuron*, 59(6), 1037-1050.
53
54 Weiskopf, N., Hutton, C., Josephs, O., & Deichmann, R. (2006). Optimal EPI parameters for reduction of
55
56 susceptibility-induced BOLD sensitivity losses: A whole-brain analysis at 3 T and 1.5 T.
57
58 *NeuroImage*, 33(2), 493-504.
59
60

1
2
3 Wexchler, D. (1999). *Wechsler abbreviated scale of intelligence*. San Antonio, TX: Psychological
4
5 Corporation.
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Review Only

Table S1. Whole brain analyses showing main-effects across all participants for No-Pain> Pain**BOLD response**

Brain regions	Peak						Cluster	
	L/R	x	y	z	t	Z	Extent (k)	P (FWE)
Calcarine sulcus	L	-8	-88	12	5.42	4.73	1416	<.001
White matter	R	28	-48	14	5.20	4.58	73	.39
Superior frontal gyrus	R	18	58	2	4.57	4.12	363	<.001
White matter	L	-30	-44	0	4.42	4.00	21	.92
Superior frontal gyrus	L	-14	50	-10	3.91	3.61	36	.77
Caudate nucleus	L	-18	24	4	3.96	3.65	12	.98
Superior frontal gyrus	R	24	-44	-34	3.96	3.65	15	.96
Supramarginal gyrus	R	44	-52	34	3.84	3.55	24	.90
White matter	L	-24	38	10	3.59	3.35	16	.96

Notes: Whole-brain analyses reported at a threshold level of $P < .001$ (uncorrected, cluster size > 10 voxels). Spatial coordinates (x, y, z) are in Montreal Neurological Institute space. R = Right; L = Left.