



ELSEVIER

Available online at www.sciencedirect.com

ScienceDirect

Current Opinion in
Microbiology

cAMP signalling and its role in host cell invasion by malaria parasites

Abigail J Perrin¹, Avnish Patel², Christian Flueck²,
Michael J Blackman^{1,2} and David A Baker²

Cyclic adenosine monophosphate (cAMP) is an important signalling molecule across evolution, but until recently there was little information on its role in malaria parasites. Advances in gene editing – in particular conditional genetic approaches and mass spectrometry have paved the way for characterisation of the key components of the cAMP signalling pathway in malaria parasites. This has revealed that cAMP signalling plays a critical role in invasion of host red blood cells by *Plasmodium falciparum* merozoites through regulating the phosphorylation of key parasite proteins by the cAMP-dependent protein kinase (PKA). These insights will help us to investigate parasite cAMP signalling as a target for novel antimalarial drugs.

Addresses

¹ Malaria Biochemistry Laboratory, The Francis Crick Institute, London, United Kingdom

² Faculty of Infectious and Tropical Diseases, London School of Hygiene & Tropical Medicine, London, United Kingdom

Corresponding author: Baker, David A (David.Baker@lshtm.ac.uk)

Current Opinion in Microbiology 2020, 58:69–74

This review comes from a themed issue on **Host–microbe interactions: parasites**

Edited by **Honorine Ward** and **Kami Kim**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 5th October 2020

<https://doi.org/10.1016/j.mib.2020.09.003>

1369-5274/Crown Copyright © 2020 Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Protozoan parasites of the genus *Plasmodium* are responsible for causing malaria, a disease with a huge global impact. The malaria parasite invades several cell types during the course of its complex life cycle in the vertebrate host and mosquito vector. Following the bite of an infected mosquito, injected sporozoites rapidly migrate to the liver, traversing several host cells before settling inside a hepatocyte. Here, each parasite divides and differentiates to produce thousands of merozoites which are released into the bloodstream [1]. All the clinical symptoms of malaria are caused by repeated cycles of red blood cell (RBC) invasion, modification and merozoite release (egress) with associated destruction of the host

cell [2]. Some blood-stage parasites differentiate into sexual forms called gametocytes which mature and undergo fertilization in the female mosquito midgut following ingestion of an infected blood meal [3]. Motile ookinete forms then develop, which invade the epithelial layer, crossing the insect midgut [4] before developing into oocysts and eventually releasing thousands of new sporozoites. These migrate to and invade the mosquito salivary glands within which they become infectious before reintroduction into the vertebrate host [5]. Each of the cellular invasion events during the lifecycle are essential for the replication and transmission of malaria parasites, and as such are of great interest as targets of anti-malarial therapeutic agents [6].

Molecular characterisation of host cell invasion by *Plasmodium* has traditionally been very challenging; invasion occurs rapidly and invasive zoites are technically difficult to isolate and maintain for *in vitro* studies. However, increasingly sophisticated genetic and biochemical tools as well as recent breakthroughs in live cell imaging have allowed detailed dissection of the roles played by key proteins and signalling molecules, particularly in the invasion of human RBCs by merozoites of *P. falciparum*, the *Plasmodium* species that causes most malaria-associated fatalities. Very recent studies have highlighted a pivotal role for the second messenger cAMP in this process [7^{**},8^{**},9^{*}]. Whilst cAMP is fundamental to a huge range of signal transduction processes, from human metabolism [10] to the behaviour of social amoeba [11], its role in malaria parasites was previously unclear. In this review we describe and summarise the newly elucidated crucial function of cAMP-dependent signalling in RBC invasion [7^{**},8^{**},9^{*}].

Key components of *Plasmodium* cAMP signalling

Cellular cAMP concentrations are controlled by the interplay of two classes of enzymes, adenylyl cyclases (ACs) and phosphodiesterases (PDEs), that respectively synthesise and degrade cAMP. The *Plasmodium* genome encodes two ACs (AC α and AC β) and four PDEs (PDE α , PDE β , PDE γ and PDE δ). Transcription of *AC α* occurs only in the sexual and pre-erythrocytic life cycle stages [12,13] whereas *AC β* expression is restricted primarily to mature intra-erythrocytic parasites (schizonts) as they approach egress, implicating AC β as the essential source of cAMP production in blood stage parasites [13–16].


PDE γ and *PDE δ* are expressed in mature gametocyte and mosquito stages where the enzymes are thought to play roles that depend on their capacity to hydrolyse cGMP [17–21]. Both *PDE α* and *PDE β* are transcribed maximally in mature blood stage schizonts. In an important distinction between these enzyme isoforms, *PDE α* hydrolyses cGMP only [22,23] whereas *PDE β* is a dual-specific PDE able to hydrolyse both cGMP and cAMP [8**]. *PDE β* is therefore the sole PDE that regulates cAMP levels in asexual blood stage parasites and is also the only essential PDE during this clinically important stage in the *P. falciparum* life cycle [8**,22].

Most of the critical functions of cAMP in eukaryotes are mediated via the activity of the cAMP-dependent protein kinase (PKA). This enzyme exists as a protein heterodimer comprising a regulatory subunit (PKAr) and a catalytic subunit (PKAc). Binding of PKAr to PKAc inhibits kinase activity, but upon binding of cAMP, PKAr dissociates from PKAc, relieving the inhibition and allowing PKAc to phosphorylate protein substrates [24,25]. Whilst PKA is thought to be the main effector of cAMP-dependent signalling in *P. falciparum*, the parasite genome also encodes a protein previously designated as Epac (exchange protein activated by cAMP), with predicted cyclic nucleotide-binding domains [26]. The biochemistry, localisation and function of the *Plasmodium* proteins involved in cyclic nucleotide signalling are described in detail in a recent review [27] and summarised in Figures 1 and 2 (Table 1).

cAMP is critical for merozoite invasion of RBCs

Invasion of RBCs by *P. falciparum* merozoites is a multi-step process which occurs following the generation and maturation of a new generation of daughter merozoites within an infected RBC. In preparation for invasion these intracellular merozoites must exit from their host RBC in a cGMP-dependent process known as egress [29,30], releasing proteins from apical secretory organelles called exonemes, micronemes and rhoptries that are critical for egress and invasion. Free merozoites associate tightly with a target RBC within seconds, pulling host RBC membrane around themselves as they drive into the cell powered by an actinomyosin motor [31]. A role for cAMP signalling in invasion was initially suggested by the ability of pharmacological inhibitors of mammalian cAMP regulatory and responsive proteins to interfere with invasion [14,26,32] as well as phosphoproteome data suggesting that PKA is highly active in mature schizonts [33,34]. A more detailed analysis of cAMP signalling was recently made possible by novel genetic techniques [35–37] that allowed robust, inducible disruption of *AC β* [7**], *PKAc* [7**,9*] and *PDE β* [8**] in *P. falciparum*. These studies showed unambiguously that in the absence of either cAMP synthesis or PKA activity, merozoites form normally but are completely unable to invade RBCs [7**,9*].

Figure 1




Key components of cAMP signalling in the *Plasmodium* merozoite. *AC β* is thought to associate with the cytosolic surface of the rhoptry membrane. Here it produces cAMP that regulates the activity of PKA, which has a diffuse localisation within the parasite cytoplasm. *PDE β* is trafficked via the ER to an apical location, from which it appears to re-localise to the parasite plasma membrane in mature schizonts. *AMA1* is stored in micronemes before it is discharged onto and across the merozoite surface just before egress. Most of this *AMA1* is then proteolytically removed except for a small fraction at the apical end that binds to RON proteins on the RBC surface during invasion. These processes are regulated in a poorly understood manner by PKA-dependent phosphorylation of the cytoplasmic tail of *AMA1*.

Artificially increasing cAMP levels by disruption of *PDE β* also significantly impaired invasion and moreover prevented further development of those parasites that did go on to invade [8**]. Taken together, these findings demonstrate that finely tuned regulation of intracellular cAMP levels is crucial to control the invasion process and subsequent parasite development.

Key events before invasion do not depend on cAMP

It has been suggested that before RBC invasion the production of cAMP causes an Epac-dependent rise in merozoite cytosolic Ca^{2+} and that this in turn leads to the secretion of critical proteins from micronemes [26]. However, recent conditional genetic knockout studies unequivocally show that Ca^{2+} signalling, microneme secretion and egress do not require cAMP [7**], PKA [7**,9*] or Epac [7**,15,28]. Instead, chemical genetic approaches suggest that these processes critically depend on cGMP production and the activity of the parasite cGMP-dependent protein kinase (PKG) [29,30,38]. Epac does not play an essential role at any point in merozoite invasion *in vitro* [7**]; indeed, it appears to be completely dispensable in asexual blood stages and is readily lost or mutated in laboratory isolates [7**,15,28].

Figure 2



cAMP signalling and its interactions with cGMP signalling in mature schizonts.

AC β catalyses the conversion of ATP to cAMP, which binds to the PKA regulatory subunit, PKAr, releasing inhibition of PKAc kinase activity. PKAc then phosphorylates a broad range of substrates, including the cytoplasmic tail of AMA1. This induces a conformational change in AMA1 that is important for merozoite invasion. cGMP, thought to be produced by GC α , activates PKG. This leads to the intracellular release of Ca²⁺, the activation of calcium-dependent protein kinases (CDPKs), secretion of exoneme and microneme proteins (including AMA1), and ultimately egress. cAMP and cGMP are both hydrolysed by PDE β . CDPK1 is phosphorylated following PKG activation and is thought to enhance the activity of PKAc, which may in turn negatively regulate the function of CDPK1. Dotted lines indicate phosphorylation events.

Table 1

Key proteins involved in cAMP signalling in asexual blood stages of *Plasmodium*

Protein	Name	Identifier (PlasmoDB)	Null phenotype
AC β	Adenylyl cyclase β	PF3D7_0802600	Essential for merozoite invasion [7**]
Epac	Exchange protein activated by cAMP	PF3D7_1417400	Non-essential in asexual blood stage parasites [7**,15,28]
PDE β	Phosphodiesterase β	PF3D7_1321500	Essential for efficient merozoite invasion and subsequent ring-stage development [8**]
PKAc	cAMP-dependent protein kinase catalytic subunit	PF3D7_0934800	Essential for merozoite invasion [7**,9*]
PKAr	cAMP-dependent protein kinase regulatory subunit	PF3D7_1223100	No data available

Observation of cAMP signalling-defective merozoites by time-lapse and electron microscopy immediately following egress has shown that, despite an inability to invade, the parasites are still able to interact with RBCs [7**,9*]. AC β -null and PKA-null merozoites induce deformations

of the RBC surface [7**,9*], indicating that the parasite actinomyosin motor is active [31,39] and can drag membrane around the merozoite to some degree as it attempts to initiate invasion. The mutant merozoites can also induce echinocytosis [7**,9*], a transient, striking change in RBC architecture that is indicative of successful discharge of parasite rhoptries [40]. Collectively, these observations suggest that cAMP-dependent signalling is not essential for early stages in the invasion pathway but is required for a relatively late step in the process.

PKA mediates a large number of phosphorylation events, many of which may be critical for invasion

One plausible model that might explain why the cAMP dependency of merozoite invasion is restricted to a late stage in the process came from the identification of the PKA-dependent phosphorylation of a merozoite protein called apical membrane antigen 1 (AMA1) [7**,9*,41], a type I integral membrane protein that is secreted from micronemes onto the merozoite surface just before egress. Much of this AMA1 is rapidly shed from the merozoite surface upon egress, but some is retained (Figure 1). Upon interaction with a target RBC, the AMA1 ectodomain binds to parasite rhoptry neck (RON) proteins that are also discharged into the RBC membrane, forming a tight interaction between the two cells before the merozoite enters. Phosphorylation of the short cytoplasmic tail of AMA1 at Ser₆₁₀, mediated directly by PKA [7**,9*,41], causes a significant conformational change in the tail structure [7**] and is critical for invasion [41]. Interactions between PKAc-null or AC β -null merozoites and RBCs show similarities to those observed when AMA1-RON interactions are blocked [7**,9*,40–44]. Furthermore, parasites deficient in cAMP signalling do not shed AMA1 efficiently from the merozoite surface [7**] and fewer merozoites attach to RBCs in the absence of PKAc [9*]. By contrast, PDE β -null parasites in which cAMP levels are elevated show premature shedding of AMA1 [8**]. Together these observations present the possibility that the invasion defect observed in the absence of cAMP signalling could be largely or wholly explained by a lack of PKA-dependent regulation of AMA1 function.

Comparative phosphoproteomic studies have shown that, in addition to AMA1, a large number of proteins are phosphorylated in mature schizonts in response to cAMP-dependent signalling [7**,8**,9*]. At least 60 phosphorylation sites within ~40 schizont proteins can be confidently identified as cAMP-dependent, based on their significantly decreased phosphorylation in PKAc-knockout and AC β -knockout parasites and significantly enhanced phosphorylation in PDE β knockouts [7**,8**]. At least 20 of these proteins are thought to be essential for blood stage parasite development [15], but further investigation of the effect of PKA-dependent phosphorylation

of these sites will be required to determine what contribution they make to merozoite invasion.

Unanswered questions and avenues for further study

What are the roles of cAMP signalling in RBC invasion and subsequent parasite development?

Loss of cAMP signalling has no observable impact on merozoite maturation or function until after egress and interaction of merozoites with a new host RBC [7^{**},9^{*}]. Around this stage of the invasion pathway we know that PKA-mediated phosphorylation of AMA1 modulates its proteolytic shedding from the merozoite surface. However, the mechanistic link between these two processes, and the function of AMA1 shedding remain unclear. It is plausible that a phosphorylation-dependent conformational change in the cytosolic domain of AMA1 marks the protein for efficient proteolysis which may in turn be required for correct tight junction formation and entry of the parasite into the RBC. However, the critical role of cAMP at and beyond this point in the erythrocytic life cycle is challenging to dissect with current experimental tools. The large number of cAMP-dependent phosphorylation events in merozoites [7^{**},8^{**}], coupled with the observation that PDE β -deficient merozoites that enter RBCs cannot develop further [8^{**}], make it tempting to speculate that PKA plays an important role in regulating the function not just of AMA1 but also of other key components of the invasion machinery as well as proteins involved in early intracellular development.

How do cGMP and cAMP signalling interact?

The precise temporal regulation of cyclic nucleotide-dependent signalling events in mature schizonts and merozoites appears critical for successful egress and invasion; blocking these signalling events prevents invasion and/or egress [7^{**},9^{*},29,30] whilst artificially elevating cyclic nucleotide levels leads to premature egress and severe impairments to invasion [8^{**},29]. Current knowledge suggests that cGMP-dependent processes are important for controlling key events just before egress and that cAMP then becomes important just before invasion. However, it is still unclear precisely how the transition from cGMP-dependent to cAMP-dependent control is timed and co-ordinated. Tight control of the expression of the cyclases, PDEs and kinases likely plays an important role but it is possible that post-translational modifications, such as the phosphorylation of PKA [45], PDE β , AC β and GC α [46] or the sensing of small molecules also contribute to the timely regulation of their activities. The homology between AC β and the bicarbonate-sensitive mammalian soluble adenylyl cyclase has led to the suggestion that bicarbonate ions may activate cAMP production in asexual blood stage parasites, but this has yet to be conclusively demonstrated [26].

Is cAMP signalling important for invasion events in mosquito and pre-erythrocytic life cycle stages?

Almost all of our insights into the role of cAMP in invasion apply to asexual blood stage malaria parasites, for which genetic tools and culture systems are most advanced. The role of cAMP may be quite different during sporozoite invasion of hepatocytes; early work suggested that AC α -mediated cAMP production is required for this process [13] but more recent findings indicate that PKAc is non-essential at this life cycle stage [47^{*}]. Whilst little is known about the role of cAMP signalling in mosquito stage invasion events, we are hopeful that this situation will soon change since it has recently become possible to apply similar conditional genetic approaches that have transformed our understanding of merozoite invasion to all stages of the *Plasmodium* life cycle [48^{*}].

Could components of cAMP signalling make good drug targets?

The essentiality of cAMP signalling for malaria parasite development in the human hosts makes this pathway an attractive target for therapeutics. Alongside its role in invasion, regulation of cAMP signalling is also critical for gametocyte survival [49], raising the attractive prospect that pharmacological targeting of cAMP signalling could impact both blood stage replication and transmission to the mosquito vector. The cyclases, PDEs and PKA are all potential target enzymes that are being explored, although designing drugs with sufficient selectivity over human isoforms will be an important challenge to overcome during development.

Conflicts of interest statement

Nothing declared.

Acknowledgements

This work was supported by Wellcome Trust grant 106239/Z/14/A (AJP and MJB), Wellcome Trust grant 106240/Z/14/Z (AP, CF and DAB) and Wellcome ISSF2 funding to the London School of Hygiene & Tropical Medicine. Funding to AJP and MJB also came from the Francis Crick Institute (<https://www.crick.ac.uk/>) which receives its core funding from Cancer Research UK (FC001043; <https://www.cancerresearchuk.org>), the UK Medical Research Council (FC001043; <https://www.mrc.ac.uk/>), and the Wellcome Trust (FC001043; <https://wellcome.ac.uk/>). Figures Created with BioRender.com.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
 - of outstanding interest
1. Vaughan AM, Kappe SHI: **Malaria parasite liver infection and exoerythrocytic biology.** *Cold Spring Harb Perspect Med* 2017, **7**.
 2. Cowman AF, Tonkin CJ, Tham W-H, Duraisingh MT: **The molecular basis of erythrocyte invasion by malaria parasites.** *Cell Host Microbe* 2017, **22**:232-245.
 3. Bennink S, Kiesow MJ, Pradel G: **The development of malaria parasites in the mosquito midgut.** *Cell Microbiol* 2016, **18**:905.

4. Baton L, Ranford Cartwright L: **How do malaria ookinetes cross the mosquito midgut wall?** *Trends Parasitol* 2005, **21**:22-28.
 5. Kojin BB, Adelman ZN: **The Sporozoite's journey through the mosquito: a critical examination of host and parasite factors required for salivary gland invasion.** *Front Ecol Evol* 2019, **7**:284.
 6. Burns AL, Dans MG, Balbin JM, de Koning-Ward TF, Gilson PR, Beeson JG, Boyle MJ, Wilson DW: **Targeting malaria parasite invasion of red blood cells as an antimalarial strategy.** *FEMS Microbiol Rev* 2019, **43**:223-238.
 7. Patel A, Perrin AJ, Flynn HR, Bisson C, Withers-Martinez C, Treeck M, Flueck C, Nicastro G, Martin SR, Ramos A et al.: **Cyclic AMP signalling controls key components of malaria parasite host cell invasion machinery.** *PLoS Biol* 2019, **17**:e3000264.
- Uses conditional genetics to pinpoint the roles of AC β and PKA in host RBC invasion.
8. Flueck C, Drought LG, Jones A, Patel A, Perrin AJ, Walker EM, Nofal SD, Sniijders AP, Blackman MJ, Baker DA: **Phosphodiesterase beta is the master regulator of cAMP signalling during malaria parasite invasion.** *PLoS Biol* 2019, **17**:e3000154.
- Describes the functional characterisation of PDE β in asexual blood stages of *P. falciparum*.
9. Wilde M-L, Triglia T, Marapana D, Thompson JK, Kouzmitchev AA, Bullen HE, Gilson PR, Cowman AF, Tonkin CJ: **Protein kinase A is essential for invasion of Plasmodium falciparum into human erythrocytes.** *mBio* 2019, **10**.
- Describes the generation and phenotype of parasites deficient in PKA, confirming the conclusions of [7••].
10. Ravnskaer K, Madiraju A, Montminy M: **Role of the cAMP pathway in glucose and lipid metabolism.** *Handb Exp Pharmacol* 2015:29-49.
 11. Saran S, Meima ME, Alvarez-Curto E, Weening KE, Rozen DE, Schaap P: **cAMP signaling in Dictyostelium. Complexity of cAMP synthesis, degradation and detection.** *J Muscle Res Cell Motil* 2002, **23**:793-802.
 12. Muhia DK, Swales CA, Eckstein-Ludwig U, Saran S, Polley SD, Kelly JM, Schaap P, Krishna S, Baker DA: **Multiple splice variants encode a novel adenyl cyclase of possible plastid origin expressed in the sexual stage of the malaria parasite Plasmodium falciparum.** *J Biol Chem* 2003, **278**:22014-22022.
 13. Ono T, Cabrita-Santos L, Leitao R, Bettiol E, Purcell LA, Diaz-Pulido O, Andrews LB, Tadakuma T, Bhanot P, Mota MM et al.: **Adenylyl cyclase alpha and cAMP signaling mediate Plasmodium sporozoite apical regulated exocytosis and hepatocyte infection.** *PLoS Pathog* 2008, **4**:e1000008.
 14. Salazar E, Bank EM, Ramsey N, Hess KC, Deitsch KW, Levin LR, Buck J: **Characterization of Plasmodium falciparum adenylyl cyclase- β and its role in erythrocytic stage parasites.** *PLoS One* 2012, **7**:e39769.
 15. Zhang M, Wang C, Otto TD, Oberstaller J, Liao X, Adapa SR, Udenze K, Bronner IF, Casandra D, Mayho M et al.: **Uncovering the essential genes of the human malaria parasite Plasmodium falciparum by saturation mutagenesis.** *Science* 2018, **360**:eaap7847.
 16. Bushell E, Gomes AR, Sanderson T, Anar B, Girling G, Herd C, Metcalf T, Modrzynska K, Schwach F, Martin RE et al.: **Functional profiling of a plasmodium genome reveals an abundance of essential genes.** *Cell* 2017, **170**:260.
 17. Taylor CJ, McRobert L, Baker DA: **Disruption of a Plasmodium falciparum cyclic nucleotide phosphodiesterase gene causes aberrant gametogenesis.** *Mol Microbiol* 2008, **69**:110-118.
 18. McRobert L, Taylor CJ, Deng W, Fivelman QL, Cummings RM, Polley SD, Billker O, Baker DA: **Gametogenesis in malaria parasites is mediated by the cGMP-dependent protein kinase.** *PLoS Biol* 2008, **6**:e139.
 19. Muhia DK, Swales CA, Deng W, Kelly JM, Baker DA: **The gametocyte-activating factor xanthurenic acid stimulates an increase in membrane-associated guanylyl cyclase activity in the human malaria parasite Plasmodium falciparum.** *Mol Microbiol* 2001, **42**:553-560.
 20. Hirai M, Arai M, Kawai S, Matsuoka H: **PbGC β Is essential for plasmodium ookinete motility to invade midgut cell and for successful completion of parasite life cycle in mosquitoes.** *J Biochem* 2006, **140**:747-757.
 21. Lakshmanan V, Fishbaugher ME, Morrison B, Baldwin M, Macarulay M, Vaughan AM, Mikolajczak SA, Kappe SHI: **Cyclic GMP balance is critical for malaria parasite transmission from the mosquito to the mammalian host.** *mBio* 2015, **6**:e02330.
 22. Wentzinger L, Bopp S, Tenor H, Klar J, Brun R, Beck HP, Seebeck T: **Cyclic nucleotide-specific phosphodiesterases of Plasmodium falciparum: PfPDE α , a non-essential cGMP-specific PDE that is an integral membrane protein.** *Int J Parasitol* 2008, **38**:1625-1637.
 23. Howard BL, Harvey KL, Stewart RJ, Azevedo MF, Crabb BS, Jennings IG, Sanders PR, Manallack DT, Thompson PE, Tonkin CJ et al.: **Identification of potent phosphodiesterase inhibitors that demonstrate cyclic nucleotide-dependent functions in apicomplexan parasites.** *ACS Chem Biol* 2015, **10**:1145-1154.
 24. Kemp BE, Bylund DB, Huang TS, Krebs EG: **Substrate specificity of the cyclic AMP-dependent protein kinase.** *Proc Natl Acad Sci U S A* 1975, **72**:3448.
 25. Taylor SS, Ilouz R, Zhang P, Kornev AP: **Assembly of allosteric macromolecular switches: lessons from PKA.** *Nat Rev Mol Cell Biol* 2012, **13**:646-658.
 26. Dawn A, Singh S, More KR, Siddiqui FA, Pachikara N, Ramdani G, Langsley G, Chitnis CE: **The central role of cAMP in regulating plasmodium falciparum merozoite invasion of human erythrocytes.** *PLoS Pathog* 2014, **10**:e1004520.
 27. Baker DA, Drought LG, Flueck C, Nofal SD, Patel A, Penzo M, Walker EM: **Cyclic nucleotide signalling in malaria parasites.** *Open Biol* 2017, **7**.
 28. Claessens A, Affara M, Assefa SA, Kwiatkowski DP, Conway DJ: **Culture adaptation of malaria parasites selects for convergent loss-of-function mutants.** *Sci Rep* 2017, **7**:41303.
 29. Collins CR, Hackett F, Strath M, Penzo M, Withers-Martinez C, Baker DA, Blackman MJ: **Malaria parasite cGMP-dependent protein kinase regulates blood stage merozoite secretory organelle discharge and egress.** *PLoS Pathog* 2013, **9**:e1003344.
 30. Koussis K, Withers-Martinez C, Baker DA, Blackman MJ: **Simultaneous multiple allelic replacement in the malaria parasite enables dissection of PKG function.** *Life Sci Alliance* 2020, **3**.
 31. Perrin AJ, Collins CR, Russell MRG, Collinson LM, Baker DA, Blackman MJ: **The actinomyosin motor drives malaria parasite red blood cell invasion but not egress.** *mBio* 2018, **9**:e00905-18.
 32. Syn C, Parzy D, Traincard F, Boccaccio I, Joshi MB, Lin DT, Yang X-M, Assemet K, Doerig C, Langsley G: **The H89 cAMP-dependent protein kinase inhibitor blocks Plasmodium falciparum development in infected erythrocytes.** *Eur J Biochem* 2001, **268**:4842-4849.
 33. Lasonder E, Green JL, Grainger M, Langsley G, Holder AA: **Extensive differential protein phosphorylation as intraerythrocytic Plasmodium falciparum schizonts develop into extracellular invasive merozoites.** *Proteomics* 2015, **15**:2716-2729.
 34. Lasonder E, Green JL, Camarda G, Talabani H, Holder AA, Langsley G, Alano P: **The Plasmodium falciparum schizont phosphoproteome reveals extensive phosphatidylinositol and cAMP-protein kinase A signaling.** *J Proteome Res* 2012, **11**:5323-5337.
 35. Collins CR, Das S, Wong EH, Andenmatten N, Stallmach R, Hackett F, Herman J-P, Müller S, Meissner M, Blackman MJ: **Robust inducible Cre recombinase activity in the human malaria parasite Plasmodium falciparum enables efficient gene deletion within a single asexual erythrocytic growth cycle.** *Mol Microbiol* 2013, **88**:687-701.
 36. Jones ML, Das S, Belda H, Collins CR, Blackman MJ, Treeck M: **A versatile strategy for rapid conditional genome engineering**

- using loxP sites in a small synthetic intron in *Plasmodium falciparum*. *Sci Rep* 2016, **6**:21800.
37. Knuepfer E, Napiorkowska M, van Ooij C, Holder AA: **Generating conditional gene knockouts in Plasmodium - a toolkit to produce stable DiCre recombinase-expressing parasite lines using CRISPR/Cas9**. *Sci Rep* 2017, **7**:3881.
 38. Brochet M, Collins MO, Smith TK, Thompson E, Sebastian S, Volkmann K, Schwach F, Chappell L, Gomes AR, Berriman M *et al.*: **Phosphoinositide metabolism links cGMP-dependent protein kinase G to essential Ca²⁺ signals at key decision points in the life cycle of malaria parasites**. *PLoS Biol* 2014, **12**: e1001806.
 39. Das S, Lemgruber L, Tay CL, Baum J, Meissner M: **Multiple essential functions of Plasmodium falciparum actin-1 during malaria blood-stage development**. *BMC Biol* 2017, **15**:70.
 40. Weiss GE, Gilson PR, Taechalertpaisarn T, Tham W-H, de Jong NWM, Harvey KL, Fowkes FJL, Barlow PN, Rayner JC, Wright GJ *et al.*: **Revealing the sequence and resulting cellular morphology of receptor-ligand interactions during Plasmodium falciparum invasion of erythrocytes**. *PLoS Pathog* 2015, **11** e1004670.
 41. Leykauf K, Treeck M, Gilson PR, Nebl T, Braulke T, Cowman AF, Gilberger TW, Crabb BS: **Protein kinase A dependent phosphorylation of apical membrane Antigen 1 plays an important role in erythrocyte invasion by the malaria parasite**. *PLoS Pathog* 2010, **6**:e1000941.
 42. Treeck M, Zacherl S, Herrmann S, Cabrera A, Kono M, Struck NS, Engelberg K, Haase S, Frischknecht F, Miura K *et al.*: **Functional analysis of the leading malaria vaccine candidate AMA-1 reveals an essential role for the cytoplasmic domain in the invasion process**. *PLoS Pathog* 2009, **5** e1000322.
 43. Richard D, MacRaild CA, Riglar DT, Chan J-A, Foley M, Baum J, Ralph SA, Norton RS, Cowman AF: **Interaction between Plasmodium falciparum apical membrane antigen 1 and the rhoptry neck protein complex defines a key step in the erythrocyte invasion process of malaria parasites**. *J Biol Chem* 2010, **285**:14815-14822.
 44. Yap A, Azevedo MF, Gilson PR, Weiss GE, O'Neill MT, Wilson DW, Crabb BS, Cowman AF: **Conditional expression of apical membrane antigen 1 in Plasmodium falciparum shows it is required for erythrocyte invasion by merozoites**. *Cell Microbiol* 2014, **16**:642-656.
 45. Kumar S, Kumar M, Ekka R, Dvorin JD, Paul AS, Madugundu AK, Gilberger T, Gowda H, Duraisingh MT, Keshava Prasad TS *et al.*: **PfCDPK1 mediated signaling in erythrocytic stages of Plasmodium falciparum**. *Nat Commun* 2017, **8**:63.
 46. Alam MM, Solyakov L, Bottrill AR, Flueck C, Siddiqui FA, Singh S, Mistry S, Viskaduraki M, Lee K, Hopp CS *et al.*: **Phosphoproteomics reveals malaria parasite Protein Kinase G as a signalling hub regulating egress and invasion**. *Nat Commun* 2015, **6**:7285.
 47. Choudhary HH, Gupta R, Mishra S: **PKAc is not required for the preerythrocytic stages of Plasmodium berghei**. *Life Sci Alliance* 2019, **2**.
Shows that sporozoite invasion of hepatocytes is independent of cAMP signalling via PKA.
 48. Tibúrcio M, Yang ASP, Yahata K, Suárez-Cortés P, Belda H, Baumgarten S, van de Vegte-Bolmer M, van Gemert G-J, van Waardenburg Y, Levashina EA *et al.*: **A novel tool for the generation of conditional knockouts to study gene function across the plasmodium falciparum life cycle**. *mBio* 2019, **10**.
Describes a novel parasite line that will enable the study of cAMP signalling across diverse life cycle stages.
 49. Ramdani G, Naissant B, Thompson E, Breil F, Lorthiois A, Dupuy F, Cummings R, Duffier Y, Corbett Y, Mercereau-Puijalon O *et al.*: **cAMP-signalling regulates gametocyte-infected erythrocyte deformability required for malaria parasite transmission**. *PLoS Pathog* 2015, **11** e1004815.