

MIT Joint Program on the Science and Policy of Global Change



Effects of Air Pollution Control on Climate

Ronald G. Prinn, John Reilly, Marcus Sarofim, Chien Wang and Benjamin Felzer

Report No. 118

January 2005

The MIT Joint Program on the Science and Policy of Global Change is an organization for research, independent policy analysis, and public education in global environmental change. It seeks to provide leadership in understanding scientific, economic, and ecological aspects of this difficult issue, and combining them into policy assessments that serve the needs of ongoing national and international discussions. To this end, the Program brings together an interdisciplinary group from two established research centers at MIT: the Center for Global Change Science (CGCS) and the Center for Energy and Environmental Policy Research (CEEPR). These two centers bridge many key areas of the needed intellectual work, and additional essential areas are covered by other MIT departments, by collaboration with the Ecosystems Center of the Marine Biology Laboratory (MBL) at Woods Hole, and by short- and long-term visitors to the Program. The Program involves sponsorship and active participation by industry, government, and non-profit organizations.

To inform processes of policy development and implementation, climate change research needs to focus on improving the prediction of those variables that are most relevant to economic, social, and environmental effects. In turn, the greenhouse gas and atmospheric aerosol assumptions underlying climate analysis need to be related to the economic, technological, and political forces that drive emissions, and to the results of international agreements and mitigation. Further, assessments of possible societal and ecosystem impacts, and analysis of mitigation strategies, need to be based on realistic evaluation of the uncertainties of climate science.

This report is one of a series intended to communicate research results and improve public understanding of climate issues, thereby contributing to informed debate about the climate issue, the uncertainties, and the economic and social implications of policy alternatives. Titles in the Report Series to date are listed on the inside back cover.

Henry D. Jacoby and Ronald G. Prinn,
Program Co-Directors

For more information, please contact the Joint Program Office

Postal Address: Joint Program on the Science and Policy of Global Change
77 Massachusetts Avenue
MIT E40-428
Cambridge MA 02139-4307 (USA)

Location: One Amherst Street, Cambridge
Building E40, Room 428
Massachusetts Institute of Technology

Access: Phone: (617) 253-7492
Fax: (617) 253-9845
E-mail: globalchange@mit.edu
Web site: <http://MIT.EDU/globalchange/>

Effects of Air Pollution Control on Climate

Ronald Prinn*, John Reilly*, Marcus Sarofim*, Chien Wang* and Benjamin Felzer†

Abstract

Urban air pollution and climate are closely connected due to shared generating processes (e.g., combustion) for emissions of the driving gases and aerosols. They are also connected because the atmospheric lifecycles of common air pollutants such as CO, NO_x and VOCs, and of the climatically important methane gas (CH₄) and sulfate aerosols, both involve the fast photochemistry of the hydroxyl free radical (OH). Thus policies designed to address air pollution may impact climate and vice versa. We present calculations using a model coupling economics, atmospheric chemistry, climate and ecosystems to illustrate some effects of air pollution policy alone on global warming. We consider caps on emissions of NO_x, CO, volatile organic carbon, and SO_x both individually and combined in two ways. These caps can lower ozone causing less warming, lower sulfate aerosols yielding more warming, lower OH and thus increase CH₄ giving more warming, and finally, allow more carbon uptake by ecosystems leading to less warming. Overall, these effects significantly offset each other suggesting that air pollution policy has a relatively small net effect on the global mean surface temperature and sea level rise. However, our study does not account for the effects of air pollution policies on overall demand for fossil fuels and on the choice of fuels (coal, oil, gas), nor have we considered the effects of caps on black carbon or organic carbon aerosols on climate. These effects, if included, could lead to more substantial impacts of capping pollutant emissions on global temperature and sea level than concluded here. Caps on aerosols in general could also yield impacts on other important aspects of climate beyond those addressed here, such as the regional patterns of cloudiness and precipitation.

Contents

1. Introduction	1
2. A chemistry primer	2
3. Integrated Global System Model.....	4
4. Numerical experiments	6
4.1 Effects on concentrations.....	8
4.2 Effects on ecosystems.....	9
4.3 Economic effects.....	10
4.4 Effects on temperature and sea level.....	11
5. Summary and Conclusions	12
6. References	14

1. INTRODUCTION

Urban air pollution has a significant impact on the chemistry of the atmosphere and thus potentially on regional and global climate. Already, air pollution is a major issue in an increasing number of megacities around the world, and new policies to address urban air pollution are likely to be enacted in many developing countries irrespective of the participation of these countries in any explicit future climate policies. The emissions of gases and microscopic particles (aerosols) that are important in air pollution and climate are often highly correlated due to shared generating processes. Most important among these processes is combustion of fossil fuels and biomass which

* Joint Program on the Science and Policy of Global Change, MIT, Cambridge MA 02139, USA.

† Ecosystems Center, Marine Biological Laboratory, Woods Hole, MA 02543, USA.

Article in review for inclusion in: *Human-Induced Climate Change: An Interdisciplinary Assessment*, Snowmass Workshop 10th Anniversary Volume, M. Schlesinger (editor), Cambridge University Press.

produces carbon dioxide (CO₂), carbon monoxide (CO), nitrogen oxides (NO_x), volatile organic compounds (VOCs), black carbon (BC) aerosols, and sulfur oxides (SO_x, comprised of some sulfate aerosols, but mostly SO₂ gas which subsequently forms white sulfate aerosols). In addition, the atmospheric lifecycles of common air pollutants such as CO, NO_x and VOCs, and of the climatically important methane (CH₄) and sulfate aerosols, both involve the fast photochemistry of the hydroxyl free radical (OH). Hydroxyl radicals are the dominant “cleansing” chemical in the atmosphere, annually removing about 3.7 gigatons (1 gigaton = 10¹⁵ gm) of reactive trace gases from the atmosphere; this amount is similar to the total mass of carbon removed annually from the atmosphere by the land and ocean combined (Ehhalt, 1999; Prinn, 2003).

In this paper we report exploratory calculations designed to show some of the major effects of specific global air pollutant emission caps on climate. In other words, could future air pollution policies help to mitigate future climate change or exacerbate it? For this purpose, we will need to consider carefully the connections between the chemistry of the atmosphere and climate. These connections are complex and their nonlinearity is exemplified by the fact that concentrations of ozone in urban areas for a given level of VOC emissions tend to increase with increasing NO_x emissions until a critical CO-dependent or VOC-dependent NO_x emission level is reached. Above that critical level, ozone concentrations actually decrease with increasing NO_x emissions emphasizing the need for policies to consider CO, VOC and NO_x emission reductions jointly rather than independently.

In order to interpret the results of our calculations presented later, it is necessary to understand some of the reasons for the above complexity and nonlinearity in air chemistry. Hence, the next section provides a review of the key issues, aimed especially at the non-expert. In two sections following that, we introduce the global model that we use for our calculations and present and interpret the results. We end with a summary and concluding remarks.

2. A CHEMISTRY PRIMER

The ability of the lower atmosphere (troposphere) to remove most air pollutants depends on complex chemistry driven by the relatively small amount of the sun’s ultraviolet light that penetrates through the upper atmospheric (stratospheric) ozone layer (see: Ehhalt, 1999; Prinn, 2003). This chemistry is also driven by emissions of NO_x, CO, CH₄ and VOCs and leads to the production of O₃ and OH. **Figure 1** reviews, with much simplification, the chemical reactions involved (Prinn, 1994). The importance of this chemistry to climate change occurs because it involves both climate-forcing greenhouse gases (H₂O, CH₄, O₃) and air pollutants (CO, NO, NO₂). It also involves aerosols (H₂SO₄, HNO₃, BC) that influence climate (through reflecting or absorbing sunlight), productivity of ecosystems (through their exposure to O₃, and to H₂SO₄ and HNO₃ in acid rain), and human health (through inhalation). Also important are free radicals and atoms in two forms: very reactive species like O(¹D) and OH, and less reactive ones like HO₂, O(³P), NO and NO₂.




Figure 1. Summary of the chemistry in the troposphere important in the linkage between urban air pollution and climate (after Prinn, 1994, 2003). VOCs (not shown) are similar to CH₄ in their reactions with OH, but they form acids, aldehydes and ketones in addition to CO.

Referring to Figure 1, when OH reacts with CH₄ the CH₄ is converted mostly to CO in steps that consume OH and also produce HO₂. The OH in turn converts CO to CO₂, NO₂ to HNO₃, and SO₂ to H₂SO₄. The primary OH production pathway occurs when H₂O reacts with the O(¹D) atoms that come from dissociation of O₃ by ultraviolet (UV) light. Within about a second of its formation, on average, OH reacts with other gases, either by donating its O atom (*e.g.*, to CO to form CO₂ and H) or by removing H (*e.g.*, from CH₄ to form CH₃ and H₂O). The H and CH₃ formed in these ways attach rapidly to O₂ to form hydroperoxy (HO₂) or methylperoxy (CH₃O₂) free radicals which are relatively unreactive. If there is no way to rapidly recycle HO₂ back to OH, then levels of OH are kept relatively low. The addition of NO_x emissions into the mix significantly changes the chemistry. Specifically, a second pathway is created in which NO reacts with HO₂ to form NO₂ and to reform OH. Ultraviolet light then decomposes NO₂ to produce O atoms (which attach to O₂ to form O₃) and reform NO. Hence NO_x (the sum of NO and NO₂) is a catalyst which is not consumed in these reactions. The production rate of OH by this secondary path in polluted air is about five times faster than the above primary pathway involving O(¹D) and H₂O (Ehhalt, 1999). The reaction of NO with HO₂ does not act as a sink for HO_x (the sum of OH and HO₂) but instead determines the ratio of OH to HO₂. Calculations for

polluted air suggest that HO₂ concentrations are about 40 times greater than OH (Ehhalt, 1999). This is due mainly to the much greater reactivity of OH compared to HO₂.

If emissions of air pollutants that react with OH, such as CO, VOCs, CH₄, and SO₂, are increasing, then keeping all else constant, OH levels should decrease. This would increase the lifetime and hence concentrations of CH₄. However, increasing NO_x emissions should increase tropospheric O₃ (and hence the primary source of OH), as well as increase the recycling rate of HO₂ to OH (the second source of OH). This OH increase should lower CH₄ concentrations. Thus changing the level of OH causes greenhouse gas, and thus climate, changes. Climate change will also influence OH. Higher ocean temperatures should increase H₂O in the lower troposphere and thus increase OH production through its primary pathway. Higher atmospheric temperatures also increase the rate of reaction of OH with CH₄, decreasing the concentrations of both. Greater cloud cover will reflect more solar ultraviolet light, thus decreasing OH, and vice versa.

Added to these interactions involving gases, are those involving aerosols. For example, increasing SO₂ emissions and/or OH concentrations should lead to greater concentrations of sulfate aerosols which are a cooling influence. Accounting for all of these interactions, and other related ones (see *e.g.*, Prinn, 2003), requires that a detailed interactive atmospheric chemistry and climate model be used to assess the effects of air pollution reductions on climate.

3. INTEGRATED GLOBAL SYSTEM MODEL

For our calculations, we utilize the MIT Integrated Global System Model (IGSM). The IGSM consists of a set of coupled submodels of economic development and its associated emissions, natural biogeochemical cycles, climate, air pollution, and natural ecosystems (Prinn *et al.*, 1999; Reilly *et al.*, 1999; Webster *et al.*, 2002, 2003). It is specifically designed to address key questions in the natural and social sciences that are amenable to quantitative analysis and are relevant to environmental policy. The current structure of the IGSM is shown in **Figure 2**.

Chemically and radiatively important trace gases and aerosols are emitted as a result of human activity. The Emissions Prediction and Policy Analysis (EPPA) submodel incorporates the major relevant demographic, economic, trade, and technical issues involved in these emissions at the national and global levels. Natural emissions of these gases are also important and are computed in the Natural Emissions Model (NEM) which is driven by IGSM predictions of climate and ecosystem states around the world.

The coupled atmospheric chemistry and climate submodel is in turn driven by the combination of these anthropogenic and natural emissions. This submodel includes atmospheric and oceanic chemistry and circulation, and land hydrological processes. The atmospheric chemistry component has sufficient detail to include its sensitivity to climate and different mixes of emissions, and to address the effects on climate of policies proposed for control of air pollution and vice versa (Wang *et al.*, 1998; Mayer *et al.*, 2000). Of particular importance to the calculations presented here, the urban air pollution (UAP) submodel is based upon, and designed




Figure 2. Schematic illustrating the framework, submodels, and processes in the MIT Integrated Global System Model (IGSM). Feedbacks between the component models that are currently included, or proposed for inclusion in later versions, are shown as solid or dashed lines respectively (adapted from Prinn *et al.*, 1999).

to simulate, the detailed chemical and dynamical processes in current 3D urban air chemistry models (Mayer *et al.*, 2000). For this purpose, the emissions calculated in the EPPA submodel are divided into two parts: urban emissions which are processed by the UAP submodel before entering the global chemistry/climate submodel, and non-urban emissions which are input directly into the large-scale model. The UAP enables simultaneous consideration of control policies applied to local air pollution and global climate. It also provides the capability to assess the effects of air pollution on ecosystems, and to predict levels of irritants important to human health in the growing number of megacities around the world. The atmospheric and oceanic circulation components in the IGSM are simplified compared to the most complex models available, but they capture the major processes and, with appropriate parameter choices, can mimic quite well the zonal-average behavior of the complex models (Sokolov and Stone, 1998; Sokolov *et al.*, 2003). We use the version of the IGSM with 2D atmospheric and 2D oceanic

submodels here, although the latest version has a 3D ocean to capture better the deep ocean circulations that serve as heat and CO₂ sinks (Kamenkovich *et al.*, 2002, 2003). The 2D/2D version we use here resolves separately the land and ocean (LO) processes at each latitude and so is referred to as the 2D-LO-2D version.

The outputs from the coupled atmospheric chemistry and climate model then drive a Terrestrial Ecosystems Model (TEM; Xiao *et al.*, 1998) which calculates key vegetation properties including production of vegetation mass, land-atmosphere CO₂ exchanges, and soil nutrient contents in 18 globally distributed ecosystems. TEM then feeds back its computed CO₂ fluxes to the climate/atmospheric chemistry submodel, and its soil nutrient contents to NEM, to complete the IGSM interactions. The current IGSM does not include treatment of black carbon (BC) aerosols (see Figure 1). Detailed studies with a global 3D chemistry and climate model indicate multiple, regionally variable and partially-offsetting, effects of BC on absorption and reflection of sunlight, reflectivity of clouds, and the strength of lower tropospheric convection (Wang, 2004). These detailed studies also suggest important BC-induced changes in the geographic pattern of precipitation, not surprisingly since aerosols have important and complex effects on cloud formation, and on whether clouds will even produce precipitation. Methods to capture these effects in the IGSM are currently being explored. In light of the difficulty in simulating these and other regional effects, the numerical results presented here are limited to temperature and sea level effects, primarily at the global and hemispheric level.

4. NUMERICAL EXPERIMENTS

To investigate, at least qualitatively, some of the important potential impacts of controls of air pollutants on temperature, we have carried out runs of the IGSM in which individual pollutant emissions, or combinations of these emissions, are held constant from 2005 to 2100. These are compared to a reference run (denoted “ref”) in which there is no explicit policy to reduce greenhouse gas emissions (see Reilly *et al.*, 1999; Webster *et al.*, 2002).

Specifically, in five runs of the IGSM, we consider caps at 2005 levels of emissions of the following air pollutants:

- (1) NO_x only (denoted “NO_x cap”),
- (2) CO plus VOCs only (denoted “CO/VOC cap”),
- (3) SO_x only (denoted “SO_x cap”),
- (4) Cases (1) and (2) combined (denoted “3 cap”),
- (5) Cases (1), (2) and (3) combined (denoted “all cap”).

Cases (1) and (2) are designed to show the individual effects of controls on NO_x and reactive carbon gases (CO, VOC), although such individual actions are very unlikely. Case (3) addresses further controls on emissions of sulfur oxides from combustion of fossil fuels and biomass, and from industrial processes. Cases (4) and (5) address combinations more likely to be representative of a real comprehensive air pollution control approach.

One important caveat in interpreting our results is that we are neglecting the effects of air pollutant controls on: (a) the overall demand for fossil fuels (*e.g.*, leading to greater efficiencies in energy usage and/or greater demand for non-fossil energy sources), and (b), the relative mix of fossil fuels used in the energy sector (*i.e.* coal versus oil versus gas). Consideration of these effects, which may be very important, will require calculation in the EPPA model of the impacts of NO_x, CO, VOC and SO_x emission reductions on the cost of using coal, oil, and gas. Such calculations have not yet been included in the current global economic models (including EPPA) used to address the climate issue. Such inclusion requires relating results from existing very detailed studies of costs of meeting near-term air pollution control to the more aggregated structure, and longer time horizon, of models used to examine climate policy.

In **Figure 3** we show the ratios of the emissions of NO_x, CO/VOC, and SO_x in the year 2100 to the reference case in 2100 when their emissions are capped at 2005 levels. Because these chemicals are short-lived (hours to several days for NO_x, VOCs, and SO_x, few months for CO), the effects of their emissions are largely restricted to the hemispheres in which they are emitted (and for the shortest-lived pollutants restricted to their source regions). Figure 3 therefore shows hemispheric as well as global emission ratios. For calibration, the reference global emissions of NO_x, CO/VOC, and SO_x in 2100 are about 5, 2.5, and 1.5 times their 2000 levels.




Figure 3. Global, northern hemispheric (NH) and southern hemispheric (SH) emissions in the year 2100 of CO/VOC, NO_x and SO_x, when they are capped at 2005 levels (CAP), are shown as ratios to emissions in the reference (REF) case (no caps).

4.1 Effects on concentrations

In **Figure 4**, the global and hemispheric average lower tropospheric concentrations of CH₄, O₃, sulfate aerosols, and OH in each of the above five capping cases are shown as percentage changes from the relevant global or hemispheric reference. From Figure 4a, the major global effects of capping SO_x are to decrease sulfate aerosols and slightly increase OH (due to lower SO₂ which is an OH sink). Capping of NO_x leads to decreases in O₃ and OH and an increase in CH₄ (caused by the lower OH which is a CH₄ sink). The CO and VOC cap increases OH and thus increases sulfate (formed by OH and SO₂) and decreases CH₄. Note that CO and VOC changes have opposing effects on O₃ so the net changes when they are capped together are small. Combining NO_x, CO and VOC caps leads to an O₃ decrease (driven largely by the NO_x decrease) and a slight increase in CH₄ (the enhancement due to the NO_x caps being partially offset by the opposing CO/VOC caps). Finally, capping all emissions causes substantial lowering of sulfate aerosols and O₃ and a small increase in CH₄.

The two hemispheres generally respond somewhat differently to these caps due to the short air pollutant lifetimes and dominance of northern over southern hemispheric emissions (Figs. 4b and 4c). The northern hemisphere contributes the most to the global averages and therefore responds similarly (compare Figs. 4a and 4c). The southern hemisphere shows very similar decreases in sulfate aerosol from its reference when compared to the northern hemisphere when either SO_x or all emissions are capped (compare Figs. 4b and 4c).

When compared to the southern hemisphere, the northern hemispheric ozone levels decrease by much larger percentages below their northern hemisphere reference when either NO_x, NO_x/CO/VOC, or all emissions are capped. Capping NO_x emissions leads to significant decreases in OH and thus increases in methane in both hemispheres (Figs. 4b and 4c). Because methane has a long lifetime (about 9 years, Prinn *et al.*, 2001) relative to the interhemispheric




Figure 4. Concentrations of climatically and chemically important species (CH₄, O₃, aerosols, OH) in the five cases with capped emissions are shown as percent changes from their relevant global or hemispheric average values in the reference case for the year 2100: (a) global-average; (b) southern hemispheric; and (c) northern hemispheric concentrations.

mixing time (about 1 to 2 years), its global concentrations are influenced by OH changes in either hemisphere alone, or in both. Hence CH₄ also increases in both hemispheres when NO_x/CO/VOC or all emissions are capped even though the OH decreases only occur in the northern hemisphere in these two cases (see Figs. 4b and 4c).

4.2 Effects on ecosystems

Effects of air pollution on the land ecosystem sink for carbon can be significant due to reductions in ozone-induced plant damage (**Figure 5**, see also Felzer *et al.*, 2004). Net primary production (NPP, the difference between plant photosynthesis and plant respiration), as well as net ecosystem production (NEP, which is the difference between NPP and soil respiration plus decay, and represents the net land sink), both increase when ozone decreases. This is evident in the case illustrated in Figure 5 where all pollutants are capped and ozone decreased by about 13% globally (Figure 4a). The effect is even greater when we assume that cropland and managed forests receive optimal levels of nitrogen fertilizer (“with Fertilizer” case; Felzer *et al.*, 2004a,b). The land sink (NEP) is increased by 30 to 49% or 0.6 to 0.9 gigatons of carbon (in CO₂) in 2100 through the illustrated pollution caps (Figure 5, 1 gigaton=10¹⁵ gm).

These ecosystem calculations do not include the additional positive effects on NPP and NEP of decreased acid deposition and decreased exposure to SO₂ and NO₂ gas, that would result from the pollution caps considered. They also do not include the negative effects on NPP and NEP of decreasing nutrient nitrate and possibly sulfate deposition that also arise from these caps.




Figure 5. Net annual uptake of carbon by vegetation alone (net primary production) and vegetation plus soils (net ecosystem production, the land carbon sink) for the NO_x/SO_x/CO plus VOC capped (allcap) case is shown for the year 2100 as a percentage change from the reference case. The results show the effects with optimal nitrogen use through fertilization on cropland (with Fertilizer) or with levels of nitrogen in croplands assumed to be the same as those in equivalent natural ecosystems (without Fertilizer).

4.3 Economic effects

If we could confidently value damages associated with climate change, we could estimate the avoided damages in dollar terms resulting from reductions in temperature due to the lowered level of atmospheric CO₂ caused by the above increases in the land carbon sink achieved with the ozone caps. We could similarly value the temperature changes due to caps in other pollutants besides ozone. However monetary damage estimates suffer from numerous shortcomings (*e.g.*, Jacoby, 2004). Felzer *et al.* (2004a,b) valued increases in carbon storage in ecosystems due to decreased ozone exposure in terms of the avoided costs of fossil fuel CO₂ reductions needed to achieve an atmospheric stabilization target. The particular target they examined was 550 ppm CO₂. The above extra annual carbon uptake (due to avoided ozone damage) of 0.6 to 0.9 gigatons of carbon is only 2 to 4% of year 2100 reference projections of anthropogenic fossil CO₂ emissions (which reach nearly 25 gigatonsC/year in 2100 according to Felzer *et al.* (2004b)). However, as these authors point out, this small level of additional uptake can have a surprisingly large effect on the cost of achieving a climate policy goal. Here we conduct a similar analysis using a 5% discount rate, and adopting the policy costs associated with 550 ppm CO₂ stabilization, to estimate the policy cost savings that would result from the increased carbon uptake through 2100 in the “allcap” compared to the “ref” scenarios shown in Figure 5. The savings are \$2.5 (“without Fertilizer”) to \$4.7 (“with Fertilizer”) trillion (1997 dollars). These implied savings are 12 to 22% of the total cost of a 550ppm stabilization policy.

The disproportionately large economic value of the additional carbon uptake has two reasons. One reason is that the fossil carbon reduction savings are cumulative; the total reference 2000-2100 carbon uptake is 36 (without Fertilizer) and 75 (with Fertilizer) gigatons, or about 6 to 13 years of fossil carbon emissions at current annual rates. A second reason is that the additional uptake avoids the highest marginal cost options. This assumes that the implemented policies would be cost effective in the sense that the least costly carbon reduction options would be used first, and more costly options would only be used later if needed. An important caveat here is that, as shown in Felzer *et al.* (2004a,b), a carbon emissions reduction policy also reduces ozone precursors so that an additional cap on these precursors associated with air pollution policy results in a smaller additional reduction, and less avoided ecosystem damage. A pollution cap as examined here, assuming there was also a 550ppm carbon policy in place, leads to only a 0.1 to 0.8 gigaton increase in the land sink in 2100 (compare 0.6 to 0.9 gigatons in Figure 5) and a cumulative 2000-2100 increase of carbon uptake of 13 to 40 gigatons of carbon, which is about one-half of the above increased cumulative uptake when the pollution cap occurs assuming there is no climate policy.

4.4 Effects on temperature and sea level

The impact of these various pollutant caps on global and hemispheric mean surface temperature and sea level changes from 2000 to 2100 are shown in **Figure 6** as percentages relative to the global-average reference case changes of 2.7°C and 0.4 meters respectively.

The largest increases in temperature and sea level occur when SO_x alone is capped due to the removal of reflecting (cooling) sulfate aerosols. Because most SO_x emissions are in the northern hemisphere, the temperature increases are greatest there. For the NO_x caps, temperature increases in the southern hemisphere (driven by the CH₄ increases), but decreases in the northern hemisphere (due to the cooling effects of the O₃ decreases exceeding the warming driven by the CH₄ increases). For CO and VOC reductions, there are small decreases in temperature driven by the accompanying aerosol increases and CH₄ reductions, with the greatest effects being in the northern hemisphere where most of the CO and VOC emissions (and aerosol production) occur.

When NO_x, CO, and VOCs are all capped, the nonlinearity in the system is evidenced by the fact that the combined effects are not simple sums of the effects from the individual caps. Ozone




Figure 6. Effects of air pollution caps in the five capping cases on the global, northern hemispheric and southern hemispheric average temperature increases, and the global sea level rise, between 2000 and 2100 are shown as percent changes from their average values (global or hemispheric) in the reference case. Also shown are the equivalent results for the case where the enhanced sink due to the ozone cap is included along with the caps on all pollutants. For this case, we assume the average of the fertilized and non-fertilized sink enhancements from Figure 5.

decreases and aerosol increases (offset only slightly by CH₄ increases) lead to even less warming and sea level rise than obtained by adding the CO/VOC and NO_x capping cases. Finally the capping of all emissions yields temperature and sea level rises that are smaller but qualitatively similar to the case where only SO_x is capped, but the rises are greater than expected from simple addition of the SO_x-capped and CO/VOC/NO_x-capped cases. Nevertheless, the capping of CO, VOC and NO_x serves to reduce the warming induced by the capping of SO_x.

Note that these climate calculations in Figure 6 omit the cooling effects of the CO₂ reductions caused by the lessening of the inhibition of the land sink by ozone (Figure 5). This omission is valid if we presume that anthropogenic CO₂ emissions, otherwise restricted by a climate policy, are allowed to increase to compensate for these reductions. This was the basis for our economic analysis in the previous section. To illustrate the lowering of climate impacts if we allowed the sink-related CO₂ reductions to occur, we show a sixth case in Figure 6 (“allcap+sink”) which combines the capping of all air pollutant emissions with the enhanced carbon sink from Figure 5. Now we see that the sign of the warming and sea level rise seen in the “allcap” case is reversed in the “allcap+sink” case. If we could value this lowering of climate impacts, it would provide an alternative to the economic analysis in section 4.3.

5. SUMMARY AND CONCLUSIONS

To illustrate some of the impacts of air pollution policy on climate change, we examined five highly idealized but informative scenarios for placing caps on emissions of SO_x, NO_x, CO plus VOCs, NO_x plus CO plus VOCs, and all of these pollutants combined. These caps kept global emissions at 2005 levels through 2100 and their effects on climate were compared to a reference run with no caps applied. Our purpose was not to claim that these scenarios are in any way realistic or likely, but rather that they served to illustrate quite well the complex interactions between air pollutant emissions and changes in temperature and sea level.

In general, placing caps on NO_x alone, or NO_x, CO and VOCs together, leads to lower ozone levels, and thus less radiative forcing of climate change by this gas, and to less inhibition by ozone of carbon uptake by ecosystems which also leads to less radiative forcing (this time by CO₂). Less radiative forcing by these combined effects means less warming and less sea level rise.

Placing caps on NO_x alone also leads to decreases in OH and thus increases in CH₄. These OH decreases and CH₄ increases are lessened (but not reversed) when there are simultaneous NO_x, CO and VOC caps. Increases in CH₄ lead to greater radiative forcing. Placing caps on SO_x leads to lower sulfate aerosols, less reflection of sunlight back to space by these aerosols (direct effect) and by clouds seeded with these aerosols (indirect effect), and thus to greater radiative forcing of climate change due to solar radiation. Enhanced radiative forcing by these aerosol and CH₄ changes combined leads to more warming and sea level rise. Hence these impacts on climate of the pollutant caps partially cancel each other. Specifically, depending on the capping case, the 2000-2100 reference global average climate changes are altered only by +4.8 to -2.6%

(temperature) and +2.2 to -2.2 % (sea level). Except for the NO_x alone case, the alterations of temperature are of the same sign but significantly greater in the northern hemisphere (where most of the emissions and emission reductions occur) than in the southern hemisphere. Note that for the NO_x alone caps, the temperature decrease caused by ozone reductions is greater than the temperature increase driven by methane increases in the northern hemisphere while the opposite is true in the southern hemisphere (Figure 6).

It is well established that urban air pollution control policies are beneficial for human health and downwind ecosystems. As far as ancillary benefits are concerned, our calculations suggest that air pollution policies may have only a small influence, either positive or negative, on mitigation of global-scale climate change. However, even small contributions to climate change mitigation can be disproportionately important in economic terms. This occurs because, as we show in the case of increased carbon uptake, these effects mean that the highest cost climate change mitigation measures, those occurring at the margin, can be avoided. To further check on the validity of our conclusions, future work should include:

- (1) the effects of air pollution policy on overall demand for fossil fuels and individual demands for coal, oil and gas;
- (2) the effects of caps on black carbon (as a regulated air pollutant) on climate;
- (3) the effects on ecosystems of changes in deposition rates of acids, nitrates, and sulfates and levels of exposure to SO₂ and NO₂ resulting from air pollution reductions.

Acknowledgments

This research was supported by the U.S Department of Energy, U.S. National Science Foundation, and the Industry Sponsors of the MIT Joint Program on the Science and Policy of Global Change: Alstom Power (France), American Electric Power (USA), BP p.l.c. (UK/USA), ChevronTexaco Corporation (USA), DaimlerChrysler AG (Germany), Duke Energy (USA), J-Power (Electric Power Development Co., Ltd.) (Japan), Electric Power Research Institute (USA), Electricité de France, ExxonMobil Corporation (USA), Ford Motor Company (USA), General Motors (USA), Mirant (USA), Murphy Oil Corporation (USA), Oglethorpe Power Corporation (USA), RWE/Rheinbraun (Germany), Shell International Petroleum (Netherlands/UK), Statoil (Norway), Tennessee Valley Authority (USA), Tokyo Electric Power Company (Japan), TotalFinaElf (France), Vetlesen Foundation (USA).

6. REFERENCES

- Ehhalt, D.H., 1999: Gas phase chemistry of the troposphere. *Topics in Physical Chemistry*, **6**: 21-109.
- Felzer, B., Kicklighter, D., Melillo, J., Wang, C., Zhuang, Q., and Prinn, R., 2004a: Effects of ozone on net primary production and carbon sequestration in the conterminous United States using a biogeochemistry model. *Tellus B*, **56**: 230-248.
- Felzer, B., Reilly, J., Melillo, J., Kicklighter, D., Wang, C., Prinn, R., Sarofim, M., and Zhuang, Q.: 2004b: *Past and future effects of ozone on net primary production and carbon sequestration using a global biogeochemical model*. MIT JPSPGC Report 103 (http://mit.edu/globalchange/www/MITJPSPGC_Rpt103.pdf); *Climatic Change*, in press
- Jacoby, H.D., 2004: Informing climate policy given incommensurable benefits estimates. *Global Environmental Change Part A*, **14**(3): 287-279; MIT JPSPGC Reprint 2004-7.
- Kamenkovich, I.V., Sokolov, A.P. and Stone, P., 2002: An efficient climate model with a 3D ocean and statistical-dynamical atmosphere. *Climate Dynamics*, **1**: 585-598.
- Kamenkovich, I.V., Sokolov, A.P. and Stone, P., 2003: Feedbacks affecting the response of the thermohaline circulation to increasing CO₂: A study with a model of intermediate complexity. *Climate Dynamics*, **21**: 119-130.
- Mayer, M., Wang, C. Webster, M., and Prinn, R.G.: 2000. Linking local air pollution to global chemistry and climate. *J. Geophysical Research*, **105**: 20,869-20,896.
- Prinn, R.G., 1994, The interactive atmosphere: Global atmospheric-biospheric chemistry. *Ambio*, **23**: 50-61.
- Prinn, R.G., 2003: The cleansing capacity of the atmosphere. *Annual Reviews Environment and Resources*, **28**: 29-57.
- Prinn, R.G., Huang, J., Weiss, R., Cunnold, D., Fraser, P., Simmonds, P., McCulloch, A., Harth, C., Salameh, P., O'Doherty, S., Wang, R., Porter, L., and Miller, B., 2001: Evidence for substantial variations of atmospheric hydroxyl radicals in the past two decades. *Science*, **292**:1882-1888.
- Prinn, R.G., Jacoby, H., Sokolov, A., Wang, C., Xiao, X., Yang, Z., Eckaus, R., Stone, P., Ellerman, A.D., Melillo, J., Fitzmaurice, J., Kicklighter, D., Holian, G. and Liu, Y., 1999: Integrated Global System Model for climate policy assessment: feedbacks and sensitivity studies. *Climatic Change*, **41**: 469-546.
- Reilly, J., Prinn, R., Harnisch, J., Fitzmaurice, J., Jacoby, H., Kicklighter, D., Melillo, J., Stone, P., Sokolov, A. and Wang, C., 1999: Multi-gas assessment of the Kyoto Protocol. *Nature*, **401**: 549-555.
- Sokolov, A., and Stone, P., 1998: A flexible climate model for use in integrated assessments. *Climate Dynamics*, **14**: 291-303.
- Sokolov, A., Forest, C.E. and Stone, P., 2003: Comparing oceanic heat uptake in AOGCM transient climate change experiments. *J. Climate*, **16**: 1573-1582.
- Wang, C., Prinn, R. and Sokolov, A., 1998: A global interactive chemistry and climate model: Formulation and testing. *J. Geophysical Research*, **103**: 3399-3417.
- Wang, C., 2004: A modeling study on the climate impacts of black carbon aerosols. *J. Geophysical Research*, **109**: D03106, doi: 10.1029/2003JD004084.
- Webster, M.D., Babiker, M., Mayer, M., Reilly, J.M., Harnisch, J., Sarofim, M.C., and Wang, C., 2002: Uncertainty in emissions projections for climate models. *Atmospheric Environment*, **36**: 3659-3670.
- Webster, M.D., Forest, C.E., Reilly, J.M., Babiker, M., Kicklighter, D., Mayer, M., Prinn, R.G., Sarofim, M., Sokolov, A., Stone, P.H., and Wang, C., 2003: Uncertainty analysis of climate change and policy response. *Climatic Change*, **61**: 295-320.
- Xiao, X., Melillo, J., Kicklighter, D., McGuire, A., Prinn, R., Wang, C., Stone, P. and Sokolov, A., 1998: Transient climate change and net ecosystem production of the terrestrial biosphere. *Global Biogeochemical Cycles*, **12**: 345-360.

REPORT SERIES of the MIT Joint Program on the Science and Policy of Global Change

1. **Uncertainty in Climate Change Policy Analysis** *Jacoby & Prinn* December 1994
2. **Description and Validation of the MIT Version of the GISS 2D Model** *Sokolov & Stone* June 1995
3. **Responses of Primary Production and Carbon Storage to Changes in Climate and Atmospheric CO₂ Concentration** *Xiao et al.* Oct 1995
4. **Application of the Probabilistic Collocation Method for an Uncertainty Analysis** *Webster et al.* Jan. 1996
5. **World Energy Consumption and CO₂ Emissions: 1950-2050** *Schmalensee et al.* April 1996
6. **The MIT Emission Prediction and Policy Analysis (EPPA) Model** *Yang et al.* May 1996
7. **Integrated Global System Model for Climate Policy Analysis** *Prinn et al.* June 1996 (*superseded* by No. 36)
8. **Relative Roles of Changes in CO₂ and Climate to Equilibrium Responses of Net Primary Production and Carbon Storage** *Xiao et al.* June 1996
9. **CO₂ Emissions Limits: Economic Adjustments and the Distribution of Burdens** *Jacoby et al.* July 1997
10. **Modeling the Emissions of N₂O & CH₄ from the Terrestrial Biosphere to the Atmosphere** *Liu* August 1996
11. **Global Warming Projections: Sensitivity to Deep Ocean Mixing** *Sokolov & Stone* September 1996
12. **Net Primary Production of Ecosystems in China and its Equilibrium Responses to Climate Changes** *Xiao et al.* November 1996
13. **Greenhouse Policy Architectures and Institutions** *Schmalensee* November 1996
14. **What Does Stabilizing Greenhouse Gas Concentrations Mean?** *Jacoby et al.* November 1996
15. **Economic Assessment of CO₂ Capture and Disposal** *Eckaus et al.* December 1996
16. **What Drives Deforestation in the Brazilian Amazon?** *Pfaff* December 1996
17. **A Flexible Climate Model For Use In Integrated Assessments** *Sokolov & Stone* March 1997
18. **Transient Climate Change and Potential Croplands of the World in the 21st Century** *Xiao et al.* May 1997
19. **Joint Implementation: Lessons from Title IV's Voluntary Compliance Programs** *Atkeson* June 1997
20. **Parameterization of Urban Sub-grid Scale Processes in Global Atmospheric Chemistry Models** *Calbo et al.* July 1997
21. **Needed: A Realistic Strategy for Global Warming** *Jacoby, Prinn & Schmalensee* August 1997
22. **Same Science, Differing Policies; The Saga of Global Climate Change** *Skolnikoff* August 1997
23. **Uncertainty in the Oceanic Heat and Carbon Uptake & their Impact on Climate Projections** *Sokolov et al.* September 1997
24. **A Global Interactive Chemistry and Climate Model** *Wang, Prinn & Sokolov* September 1997
25. **Interactions Among Emissions, Atmospheric Chemistry and Climate Change** *Wang & Prinn* Sept. 1997
26. **Necessary Conditions for Stabilization Agreements** *Yang & Jacoby* October 1997
27. **Annex I Differentiation Proposals: Implications for Welfare, Equity and Policy** *Reiner & Jacoby* Oct. 1997
28. **Transient Climate Change and Net Ecosystem Production of the Terrestrial Biosphere** *Xiao et al.* November 1997
29. **Analysis of CO₂ Emissions from Fossil Fuel in Korea: 1961–1994** *Choi* November 1997
30. **Uncertainty in Future Carbon Emissions: A Preliminary Exploration** *Webster* November 1997
31. **Beyond Emissions Paths: Rethinking the Climate Impacts of Emissions Protocols** *Webster & Reiner* November 1997
32. **Kyoto's Unfinished Business** *Jacoby, Prinn & Schmalensee* June 1998
33. **Economic Development and the Structure of the Demand for Commercial Energy** *Judson et al.* April 1998
34. **Combined Effects of Anthropogenic Emissions & Resultant Climatic Changes on Atmospheric OH** *Wang & Prinn* April 1998
35. **Impact of Emissions, Chemistry, and Climate on Atmospheric Carbon Monoxide** *Wang & Prinn* April 1998
36. **Integrated Global System Model for Climate Policy Assessment: Feedbacks and Sensitivity Studies** *Prinn et al.* June 1998
37. **Quantifying the Uncertainty in Climate Predictions** *Webster & Sokolov* July 1998
38. **Sequential Climate Decisions Under Uncertainty: An Integrated Framework** *Valverde et al.* Sept. 1998
39. **Uncertainty in Atmospheric CO₂ (Ocean Carbon Cycle Model Analysis)** *Holian* Oct. 1998 (*superseded* by No. 80)
40. **Analysis of Post-Kyoto CO₂ Emissions Trading Using Marginal Abatement Curves** *Ellerman & Decaux* October 1998

Contact the Joint Program Office to request a copy. The Report Series is distributed at no charge.

REPORT SERIES of the MIT Joint Program on the Science and Policy of Global Change

41. **The Effects on Developing Countries of the Kyoto Protocol and CO₂ Emissions Trading** *Ellerman et al.* November 1998
42. **Obstacles to Global CO₂ Trading: A Familiar Problem** *Ellerman* November 1998
43. **The Uses and Misuses of Technology Development as a Component of Climate Policy** *Jacoby* Nov. 1998
44. **Primary Aluminum Production: Climate Policy, Emissions and Costs** *Harnisch et al.* December 1998
45. **Multi-Gas Assessment of the Kyoto Protocol** *Reilly et al.* January 1999
46. **From Science to Policy: The Science-Related Politics of Climate Change Policy in the U.S.** *Skolnikoff* January 1999
47. **Constraining Uncertainties in Climate Models Using Climate Change Detection Techniques** *Forest et al.* April 1999
48. **Adjusting to Policy Expectations in Climate Change Modeling** *Shackley et al.* May 1999
49. **Toward a Useful Architecture for Climate Change Negotiations** *Jacoby et al.* May 1999
50. **A Study of the Effects of Natural Fertility, Weather and Productive Inputs in Chinese Agriculture** *Eckaus & Tso* July 1999
51. **Japanese Nuclear Power and the Kyoto Agreement** *Babiker, Reilly & Ellerman* August 1999
52. **Interactive Chemistry and Climate Models in Global Change Studies** *Wang & Prinn* September 1999
53. **Developing Country Effects of Kyoto-Type Emissions Restrictions** *Babiker & Jacoby* October 1999
54. **Model Estimates of the Mass Balance of the Greenland and Antarctic Ice Sheets** *Bugnion* October 1999
55. **Changes in Sea-Level Associated with Modifications of Ice Sheets over 21st Century** *Bugnion* Oct. 1999
56. **The Kyoto Protocol and Developing Countries** *Babiker, Reilly & Jacoby* October 1999
57. **Can EPA Regulate Greenhouse Gases Before the Senate Ratifies the Kyoto Protocol?** *Bugnion & Reiner* November 1999
58. **Multiple Gas Control Under the Kyoto Agreement** *Reilly, Mayer & Harnisch* March 2000
59. **Supplementarity: An Invitation for Monopsony?** *Ellerman & Sue Wing* April 2000
60. **A Coupled Atmosphere-Ocean Model of Intermediate Complexity** *Kamenkovich et al.* May 2000
61. **Effects of Differentiating Climate Policy by Sector: A U.S. Example** *Babiker et al.* May 2000
62. **Constraining Climate Model Properties Using Optimal Fingerprint Detection Methods** *Forest et al.* May 2000
63. **Linking Local Air Pollution to Global Chemistry and Climate** *Mayer et al.* June 2000
64. **The Effects of Changing Consumption Patterns on the Costs of Emission Restrictions** *Lahiri et al.* Aug. 2000
65. **Rethinking the Kyoto Emissions Targets** *Babiker & Eckaus* August 2000
66. **Fair Trade and Harmonization of Climate Change Policies in Europe** *Viguier* September 2000
67. **The Curious Role of "Learning" in Climate Policy: Should We Wait for More Data?** *Webster* October 2000
68. **How to Think About Human Influence on Climate** *Forest, Stone & Jacoby* October 2000
69. **Tradable Permits for Greenhouse Gas Emissions: A primer with reference to Europe** *Ellerman* Nov. 2000
70. **Carbon Emissions and The Kyoto Commitment in the European Union** *Viguier et al.* February 2001
71. **The MIT Emissions Prediction and Policy Analysis Model: Revisions, Sensitivities and Results** *Babiker et al.* February 2001
72. **Cap and Trade Policies in the Presence of Monopoly and Distortionary Taxation** *Fullerton & Metcalf* March 2001
73. **Uncertainty Analysis of Global Climate Change Projections** *Webster et al.* March 2001 (*superseded by No. 95*)
74. **The Welfare Costs of Hybrid Carbon Policies in the European Union** *Babiker et al.* June 2001
75. **Feedbacks Affecting the Response of the Thermohaline Circulation to Increasing CO₂** *Kamenkovich et al.* July 2001
76. **CO₂ Abatement by Multi-fueled Electric Utilities: An Analysis Based on Japanese Data** *Ellerman & Tsukada* July 2001
77. **Comparing Greenhouse Gases** *Reilly, Babiker & Mayer* July 2001
78. **Quantifying Uncertainties in Climate System Properties using Recent Climate Observations** *Forest et al.* July 2001
79. **Uncertainty in Emissions Projections for Climate Models** *Webster et al.* August 2001

Contact the Joint Program Office to request a copy. The Report Series is distributed at no charge.

REPORT SERIES of the MIT Joint Program on the Science and Policy of Global Change

80. **Uncertainty in Atmospheric CO₂ Predictions from a Global Ocean Carbon Cycle Model** *Holian et al.*
September 2001
81. **A Comparison of the Behavior of AO GCMs in Transient Climate Change Experiments** *Sokolov et al.*
December 2001
82. **The Evolution of a Climate Regime: Kyoto to Marrakech** *Babiker, Jacoby & Reiner* February 2002
83. **The "Safety Valve" and Climate Policy** *Jacoby & Ellerman* February 2002
84. **A Modeling Study on the Climate Impacts of Black Carbon Aerosols** *Wang* March 2002
85. **Tax Distortions and Global Climate Policy** *Babiker, Metcalf & Reilly* May 2002
86. **Incentive-based Approaches for Mitigating GHG Emissions: Issues and Prospects for India** *Gupta* June 2002
87. **Deep-Ocean Heat Uptake in an Ocean GCM with Idealized Geometry** *Huang, Stone & Hill* September 2002
88. **The Deep-Ocean Heat Uptake in Transient Climate Change** *Huang et al.* September 2002
89. **Representing Energy Technologies in Top-down Economic Models using Bottom-up Info** *McFarland et al.* October 2002
90. **Ozone Effects on Net Primary Production and C Sequestration in the U.S. Using a Biogeochemistry Model** *Felzer et al.* November 2002
91. **Exclusionary Manipulation of Carbon Permit Markets: A Laboratory Test** *Carlén* November 2002
92. **An Issue of Permanence: Assessing the Effectiveness of Temporary Carbon Storage** *Herzog et al.* Dec. 2002
93. **Is International Emissions Trading Always Beneficial?** *Babiker et al.* December 2002
94. **Modeling Non-CO₂ Greenhouse Gas Abatement** *Hyman et al.* December 2002
95. **Uncertainty Analysis of Climate Change and Policy Response** *Webster et al.* December 2002
96. **Market Power in International Carbon Emissions Trading: A Laboratory Test** *Carlén* January 2003
97. **Emissions Trading to Reduce Greenhouse Gas Emissions in the U.S.: The McCain-Lieberman Proposal**
Paltsev et al. June 2003
98. **Russia's Role in the Kyoto Protocol** *Bernard et al.* June 2003
99. **Thermohaline Circulation Stability: A Box Model Study** *Lucarini & Stone* June 2003
100. **Absolute vs. Intensity-Based Emissions Caps** *Ellerman & Sue Wing* July 2003
101. **Technology Detail in a Multi-Sector CGE Model: Transport Under Climate Policy** *Schafer & Jacoby* July 2003
102. **Induced Technical Change and the Cost of Climate Policy** *Sue Wing* September 2003
103. **Effects of Ozone on NPP and Carbon Sequestration Using a Global Biogeochemical Model** *Felzer et al.*
January 2004
104. **A Modeling Analysis of Methane Exchanges Between Alaskan Ecosystems and the Atmosphere** *Zhuang et al.* November 2003
105. **Analysis of Strategies of Companies under Carbon Constraint** *Hashimoto* January 2004
106. **Climate Prediction: The Limits of Ocean Models** *Stone* February 2004
107. **Informing Climate Policy Given Incommensurable Benefits Estimates** *Jacoby* February 2004
108. **Methane Fluxes Between Ecosystems & Atmosphere at High Latitudes During the Past Century** *Zhuang et al.* March 2004
109. **Sensitivity of Climate to Diapycnal Diffusivity in the Ocean** *Dalan et al.* May 2004
110. **Stabilization and Global Climate Policy** *Sarofim et al.* July 2004
111. **Technology and Technical Change in the MIT EPPA Model** *Jacoby et al.* July 2004
112. **The Cost of Kyoto Protocol Targets: The Case of Japan** *Paltsev et al.* July 2004
113. **Air Pollution Health Effects: Toward an Integrated Assessment** *Yang et al.* July 2004
114. **The Role of Non-CO₂ Greenhouse Gases in Climate Policy: Analysis Using the MIT IGSM** *Reilly et al.* Aug 2004
115. **Future United States Energy Security Concerns** *Deutch* September 2004
116. **Explaining Long-Run Changes in the Energy Intensity of the U.S. Economy** *Sue Wing* September 2004
117. **Modeling the Transport Sector: The Role of Existing Fuel Taxes in Climate Policy** *Paltsev et al.* Nov. 2004
118. **Effects of Air Pollution Control on Climate** *Prinn et al.* January 2005

Contact the Joint Program Office to request a copy. The Report Series is distributed at no charge.