

## Original Research

# Post-Trial Feedback Alters Landing Performance in Adolescent Female Athletes Using a Portable Feedback System

Thomas W Kernozek, PhD, FACSM<sup>1 a</sup>, Drew Rutherford, MS<sup>1</sup>, Becky Heinert, PT, MSPT, SCS<sup>2</sup>, Jessica Onsager, PT, DPT<sup>1</sup>, Maria Lee, PT, DPT<sup>1</sup>, Jeremie Schiedermayer, PT, DPT<sup>1</sup>, Stephanie Dietrich, SPT<sup>1</sup>, Renee Dade, SPT<sup>1</sup>, Thomas Gus Almonroeder, PT, DPT, PhD<sup>1</sup>

<sup>1</sup> Department of Health Professions, Physical Therapy Program, University of Wisconsin – La Crosse, 1300 Badger Street, La Crosse, WI, USA; La Crosse Institute for Movement Science (LIMS), University of Wisconsin – La Crosse, 1300 Badger Street, La Crosse, WI, USA, <sup>2</sup> La Crosse Institute for Movement Science (LIMS), University of Wisconsin – La Crosse, 1300 Badger Street, La Crosse, WI, USA; Gundersen Health System, Sports Medicine, La Crosse, WI, USA

Keywords: training, movement system, motor learning, kinematics, kinetics, anterior cruciate ligament

<https://doi.org/10.26603/001c.18808>

---

## International Journal of Sports Physical Therapy

Vol. 16, Issue 1, 2021

---

### Background

Post-performance verbal and visual feedback based on data collected via lab-based instruments have been shown to improve landing patterns related to non-contact ACL injury. Biomechanical methods are often complex, difficult to transport and utilize in field settings, and costly, which limits their use for injury prevention. Developing systems that can readily provide feedback outside of the lab setting may support large scale use of feedback training for ACL injury prevention.

### Purpose/Hypothesis

The purpose of this study was to investigate the effectiveness of a single training session using a custom portable feedback training system that provides performance cues to promote changes in impact kinetics and lower extremity position during landing in female athletes.

### Study Design

Repeated measures

### Methods

One hundred fifty female athletes (ages 13-18 years old) landed from a 50 cm platform with and without feedback related to vertical ground reaction force (vGRF), vGRF symmetry and lower extremity position. Feedback was provided via a portable, low-cost system that included two custom-built force plates interfaced with a digital camera. Each athlete performed six pre-test trials followed by two blocks of six trials where they received visual feedback from the training system and individualized verbal cues from an investigator. Following training blocks, athletes completed six post-test trials without feedback and then six dual-task trials where a ball was randomly thrown to the performer during the landing (transfer task). vGRF and knee to ankle (K:A) separation ratio were measured and the average responses were reported for each trial block.

### Results

Differences in vGRF between baseline, post-test and transfer task trial blocks were observed ( $F(2,298)=181.68$ ,  $p < .0001$ ). Mean (SD) peak vGRF (body weight) were 4.43 (0.90), 3.28 (0.61), and 3.80 (0.92), respectively. Differences in K:A ratio between baseline, post-test and transfer task trial blocks were shown ( $F(2,298)=68.47$ ,  $p < .0001$ ). Mean (SD) K:A ratio were 0.87 (0.21), 0.98 (0.19), and 0.92 (0.19), respectively.

---

<sup>a</sup> **Corresponding author:** Thomas W. Kernozek, PhD, FACSM Department of Health Professions Physical Therapy Program University of Wisconsin-La Crosse La Crosse, WI 54601 e-mail: kernozek.thom@uwlax.edu

## Conclusion

A portable feedback system may be effective in reducing peak vGRFs and promoting a more desirable K:A ratio during landing and transfer task landing in adolescent female athletes.

## Level of Evidence

3b

## INTRODUCTION

Non-contact injury to the anterior cruciate ligament (ACL) in female athletes has been investigated quite extensively. Despite the plethora of investigations, females are two to eight times more likely to suffer this traumatic injury as compared to males.<sup>1</sup> Although the proposed etiology of non-contact ACL ruptures appears multifactorial, improper landing mechanics in female athletes has been cited as associated with increased injury risk.<sup>2-4</sup> Furthermore, a second ACL rupture is more likely in athletes within two years after ligament reconstruction following return to sport that involve jumping and landing activities where aberrant landing patterns were shown.<sup>5-7</sup>

The best strategy for identifying these aberrant landing patterns and the most effective intervention for their remediation is debatable. Augmented feedback using external sources (ie. use of video display and verbal cueing and instruction based on expertise from an instructor) is most often employed in the clinical setting due to their relative ease of implementation. Traditionally, clinicians attempt to modify athletes' landing mechanics by providing instructions to promote more optimal body postures and joint alignment; however, these types of instructions often only produce transient changes in landing mechanics.<sup>8</sup> As a result, there appears to be a need for more advanced methods of athlete feedback. Post-performance video feedback has been shown to promote changes in landing patterns that may reduce ACL injury in female athletes (e.g. lower vertical ground reaction forces, less knee abduction).<sup>3,4,9-16</sup> This is encouraging, as video can be recorded in clinical or field settings using widely available, low-cost equipment such as smartphones, tablets, etc. However, the information obtained from such videos is subjective and does not provide insight into the forces experienced during landing. In addition, most standard video cameras sample data at rates that are too low for dynamic movements such as drop landings.

Assessments that incorporate three-dimensional (3-D) analysis of drop landings have also been used to identify kinematic and kinetic measures that may be related to a higher risk of non-contact ACL injuries in female athletes.<sup>3,8,15,17,18</sup> Post-trial feedback has been shown to be effective in modifying neuromuscular risk factors in drop landing and can provide more accurate and objective kinematic and kinetic information which may serve to better identify risk factors for injury or re-injury.<sup>15</sup> However, many of these investigations have used expensive and elaborate laboratory-based methods to provide such feedback<sup>4,19-21</sup> that would be impractical to implement large scale especially in field settings. Therefore, this type of performance-based feedback may not have the ability to reach the target population (ie. adolescent female high school athletes).

There appears to be a need to develop landing assessment and training tools that can provide immediate post-trial feedback relevant to an athlete's performance that is both objective and qualitative. The purpose of this study was to investigate the effectiveness of a single training session using a custom portable feedback training system that provides performance cues to promote changes in impact kinetics and lower extremity position during landing in female athletes. Feedback provided during training was based on kinetic/kinematic data recorded via the system and individualized cues from an investigator.

## METHODS

### SUBJECTS

This study was approved by the University of Wisconsin La Crosse's Institutional Review Board. Prior to participation in this study, all subjects or parent provided their written informed consent in accordance with University guidelines.

One hundred fifty females at regional high schools between the ages of 13-18 years old participated in this single session study. Age, height, body mass index, and Tegner level<sup>22</sup> was recorded. All were actively participating female athletes. Exclusion criteria for the study included (1) any current lower extremity injury, (2) knee pain at rest or during running or jumping, (3) pregnancy, (4) any cardiovascular abnormalities or medical condition that limited training as indicated by a physical activity readiness questionnaire (PAR-Q).<sup>23</sup> All participants utilized their own athletic footwear and wore comfortable athletic clothing during the study.

### PROCEDURES

Participants were asked to complete a drop landing from a 50 centimeter (cm) platform. Prior to testing, athletes were given verbal instructions for the landing task including: 1) jump forward with both feet off the raised platform and land with both feet while having one foot near the center of each force platform, (2) jump bilaterally rather than stepping off the box, (3) land as to not fall forward off of the force platforms, (4) return to standing from landing position and maintain that position for two seconds. During data collection, trials were discounted if those requirements were not met. Participants were allowed up to three practice trials to familiarize themselves with the task. Data were collected generally in a gymnasium or common space with a ceiling height that could accommodate the experimental setup.

Kinetic data were obtained from two custom, high impact force plates designed for these in situ data collections. Each force platform was positioned adjacent to one another

and 25 cm in front of the 50 cm platform. Each force platform was custom made with 4 calibrated load cells (ntep-1klb shear beam load cell) capable of measuring vertical force. Bilateral vertical ground reaction force (vGRF) data were sampled at 2000 Hz and normalized to each participant's body weight. Validation of the system was performed in a pilot investigation of 20 participants where the custom force platforms were placed directly on top of two commercially available force platforms (Model 4080, Bertec Corporation, Columbus, OH) and sampled at 2000 Hz on both systems. The peak vGRF was within 5% of that obtained with the commercially available force platforms. This accuracy criteria are likely considered reasonable regarding force plate accuracy for a portable system during a dynamic impact situation such as landing.<sup>24</sup> Frontal plane video of each participant focused on the lower extremities were recorded from a high-speed camera at 100 Hz (DFK 23UV024, The Imaging Source, LLC, Charlotte, NC, USA) during each performance trial. The camera was positioned on a tripod at a height of 65 cm and at a distance of 130 cm from the force plates. Custom scripts were implemented within commercial software (Innovative Sports Training, Inc., Chicago, IL, USA) and used on a digital monitor in front of the performer to provide immediate post-trial feedback. (Figure 1)

Athletes completed 30 landing trials divided into five blocks of six trials with one minute of rest provided between each trial block. During the initial block (Pre-test), participants were blinded to any form of feedback to determine their baseline landing performance. Prior to beginning the training blocks, the athletes were provided with a brief overview of the information they would receive on the visual feedback display post each trial. This information included peak vGRF displayed in body weight, symmetry of lower extremity vGRF demonstrated through a seesaw/teeter totter display, and frontal plane video that was replayed by the investigator to depict a qualitative impression of the performer's lower extremity alignment and overall body position during landing (Figure 2)

The baseline test was followed by the first training block (Training 1) where participants received post-trial feedback based on these data coupled with cues for improving landing performance from an investigator. Athletes were provided with both externally focused feedback first (e.g. "try to reduce the vGRF value", "try to land quieter or more softly") and then internally focused feedback (e.g. "try to keep your knees over your toes," "try to land with your knees out"). The feedback provided was individualized and dependent upon what was observed based on the peak vGRF, vGRF symmetry and general impression from the video of the athletes' frontal plane landing kinematics. The general order of the feedback was based on the order of processing of these data on the display (peak vGRF, vGRF symmetry and then video of performance). These data were available within 20 seconds after each landing. The video was available in second window initially behind the force display and was available last due to computer processing requirements associated with recording these images at high speed. Athletes were given time to review the data and then relevant cues were provided by the investigator. The next trial was then immediately performed such that the entire testing



Figure 1: Typical experimental setup taken to each school. Photo depicts the drop platform, two portable force platforms, high speed video camera focused on the frontal plane, and display showing the vertical ground reaction force (vGRF), vGRF symmetry and when cued the ability to review high speed frontal view of the performer landing. After each landing, the researcher and participant discussed the feedback and internal and external cues were provided prior to the next performance trial during the training blocks. During the pre-test, post-test and transfer task the display was turned off.




Figure 2: Feedback display showing peak vertical ground reaction force (vGRF), vGRF symmetry and in another window the ability to replay and pause high speed frontal plane video.

and training session took approximately 20 minutes. The general training and feedback used was systematic where once that variable such as vGRF showed substantial improvement, the next variable was selected for feedback. During the second training set (Training 2), the feedback from the investigator was gradually withdrawn and the athletes were asked to self-evaluate their peak vGRF, loading symmetry and body alignment and position projected to the display. Participants were encouraged to incorporate strategies that they felt were most helpful for performance im-

**Table 1: Means and standard deviations of demographic data for 150 female athletes tested**

Age (years)	Height (m)	Weight (kg)	Body Mass Index
14.94 (1.61)	1.7 (0.08)	60.95 (11.09)	20.86 (3.58)

Standard deviations are provided in within brackets.

provement based on their first training block. Following completion of Training 1 and Training 2, a post-test (six trial block) was completed without any verbal or visual feedback. Finally, a transfer task was examined where a dual task landing was performed without feedback. During the dual transfer task, participants were required to attend to catching a ball while landing. Six trials were performed with a tester either throwing the ball or faking a throw. The order of the testing condition (throw or fake throw) was randomized for each participant. Outcome data from the transfer condition performance trials were pooled.

**DATA PROCESSING**


Kinetic data were exported and processed in Excel where peak vGRF in body weight for each trial were determined. Scaled video data were analyzed within Kinovea (<https://www.kinovea.org>) to determine knee to ankle separation ratio (K:A ratio).<sup>25</sup> Kinovea is a software program that allows various video formats to be opened, scaled and points where points on the image can be used to calculate various kinematic measurements. This approach to measuring knee abduction has shown high intra and interrater reliability (0.97 and 0.92 respectively).<sup>25</sup> From a force threshold of 10 N, a single video frame was selected 100 ms<sup>26</sup> after impact for analysis of frontal plane knee motion. K:A ratio was the distance from the estimated knee joint center to ankle joint center.<sup>25,27,28</sup> K:A ratios of less than 1.0 are indicative of more knee valgus positioning during landing.<sup>25,27,28</sup>

**STATISTICAL METHODS**


The dependent variables of interest were peak vGRF and K:A ratio. Means for peak total vGRF in multiples of body weight and K:A ratio were calculated for the three blocks of six trials (baseline, post-test, and transfer task). A repeated measures analysis of variance (ANOVA) (alpha set to 0.05) was then performed to examine differences in trial blocks (baseline, post-test, and transfer task) in vGRF and then on K:A ratio. Post hoc comparisons were performed using the Bonferroni approach. Effect sizes (Cohen’s *d*) were calculated between the same trial blocks. Statistical analysis was completed utilizing SPSS, version 25 (IBM Corporation, Armonk, NY, USA).

**RESULTS**

Demographic data on participants are provided in [Table 1](#). Means and standard deviations for these participants were: age of 14.94 ± 1.61 years, height of 1.67 ± 0.08 m, weight of 60.95 ± 11.09 kg, and body mass index of 20.86 ± 3.58. Ath-



**Figure 3: Depiction of the means and standard deviations for vertical ground reaction force (vGRF) in body weight for the pre-test, post-test and transfer task.**



**Figure 4: Depiction of the means and standard deviation for knee to ankle (K:A) ratio for the pre-test, post-test and transfer task.**

letes reported a Tegner level<sup>22</sup> of at least 5/10 and were currently competing in competitive volleyball and/or basketball programs at the high school or club level since the goal was to test actively participating female athletes.

Mean vGRF were different from baseline, post-test and transfer task trial blocks ( $F(2,298)=181.68, p < .0001$ ). The mean (standard deviation) peak vGRF in multiples of body weight for the pre-test, post-test and transfer task respectively were 4.43 (0.90), 3.28 (0.61), and 3.80 (0.92). Post hoc

comparisons showed that baseline vGRF was 29.96% higher than post-test and 14.22% higher during the transfer task but the transfer task was 15.85% greater than the post-test (all  $p < .0001$ ). The effect size for vGRF from baseline to post-test was 1.52, baseline to transfer was .69 and post-test to transfer was .68.

Mean K:A ratio were different from baseline, post-test and transfer task trial blocks ( $F(2,298)=63.47$ ,  $p < .0001$ ). The mean (standard deviation) K:A ratio for the baseline, post-test and transfer task respectively were 0.87 (0.21), 0.98 (0.19), and 0.92 (0.19). Post hoc comparisons showed K:A ratio during the baseline was 12.64% greater than post-test and 6.12% greater during the transfer task but the transfer task was 5.75% reduced compared to the post-test (all  $p < .0001$ ). The effect size for K:A ratio from baseline to post-test was 0.55, baseline to transfer was .25 and post-test to transfer was .32.

## DISCUSSION

Findings of the current study indicate that using a portable force plate system interfaced with a digital camera is an effective tool for providing feedback to promote immediate positive changes in vGRF and K:A ratio of adolescent female athletes. The participants in the study demonstrated a reduction in peak vGRF and higher K:A ratio (indicative of less knee abduction) following a training session which included visual feedback related to vGRF, vGRF symmetry and video of frontal plane landing kinematics, as well as cueing for an investigator. Effect sizes were large and moderate for baseline to post-test for vGRF and K:A ratio. Despite these variables regressing toward baseline values, these improvements persisted during the transfer task however both of their effect sizes were between small to medium. Positive transfer of an improved movement pattern to an untrained task, in this case, the dual-task where a ball was randomly passed to the performer is considered an indicator of motor learning.<sup>29</sup> Despite previous investigations suggesting that augmented feedback is effective in eliciting improvements in landing mechanics, and this is the first time a portable system has been used that provides salient data that participants appear to respond to that can be used to alter landing mechanics for neuromuscular training on-site with athletic teams.<sup>4,10,13-15,28,30-33</sup>

The present findings differ in part from Munro and Herrington<sup>14</sup> who also evaluated changes in vGRF and knee abduction angle using augmented feedback. In their study, the authors incorporated a combination of strategies including video demonstration of correct landing form, individualized post-trial expert video assessment and a checklist for self-analysis of the performance. Subjects were rated on kinematic variables such as trunk lean, knee valgus and knee flexion during landing. Using video analysis to determine the knee frontal plane projection angle (FPPA), the authors found a 23.9° reduction in knee abduction angle post feedback but no change in vGRF.<sup>14</sup> In the current study, a decrease in K:A ratio was also found immediately after training, but these findings differed from Munro and Herrington<sup>14</sup> in that the subjects in the present study displayed a nearly 30% decrease in peak vGRF after a single session. However, the present study's findings do become more tem-

pered as baseline vGRF were only 14.22% higher compared to the transfer task. One reason for these differences may be related to the type of feedback that was utilized for the training session. Both studies used feedback strategies that emphasized key kinematic risk factors with expert feedback for performance improvements; both also utilized self-directed assessments of landing mechanics. However, in the current study, implicit feedback that focused on the performance outcome also incorporated using vGRF and vGRF symmetry. As the participants demonstrated an understanding and mastery of targeted biomechanical variables, verbal cuing by the expert evaluator was withdrawn and the reliance on self-correction using external feedback increased. Studies have suggested that using external cues for skill acquisition may accelerate the learning process by facilitating movement automaticity and enhance the production of effective and efficient movement patterns.<sup>33-36</sup> It has also been postulated that a new skill acquired using an external focus of attention is more resilient under psychological and physiological fatigue.<sup>33,37-41</sup> Biomechanically, greater knee flexion angles and lower peak ground reaction forces have been reported when landing instructions invoke an external focus of attention.<sup>31,42,43</sup> Therefore, in the present study, the utilization of the externally focused performance variables projected on the screen during the landing training may have resulted in reduced vGRFs during landing with feedback withdrawal and during the transfer task.

Previous research has reported that changes in landing mechanics achieved through augmented feedback may be transferable to a new task. Stroube et al.<sup>44</sup> reported that high risk landing mechanics that were identified during a tuck jump with subsequent feedback and training provided over 8 weeks resulted in 37-40% reductions in knee valgus angles during a drop landing. They did not measure ground reaction force nor force symmetry directly but did report changes to "excessive landing noise" and "foot contact timing not equal". Etnoyer et al.<sup>10</sup> showed that participants that received augmented feedback during a box drop-jump task were able to maintain greater peak knee flexion angles during a running stop-jump task compared to controls during a single training session; however, no kinematic differences were reported during a sidestep cutting maneuver. In the current study, reductions in both K:A ratio and vGRF values were found during performance of a transfer task of catching a ball during landing compared to baseline. Compared to baseline, the athletes in the present study demonstrated a 5.75% decrease in K:A ratio during the transfer task. Similarly, the vGRF changes were maintained throughout the transfer task with a 14.22% decrease in force attenuation after training compared to baseline. These findings are important as a transfer task may be more representative of an athletic activity in which the focus of attention is on multiple environmental cues often occurring simultaneously. Instructional sessions that rely solely on internally focused feedback may interfere with an athlete's ability to apply the newly acquired movement strategies to a less controlled or unstructured task. It has been suggested that incorporating unknown patterns of movement to a training program may enhance the athlete's motivation during training and stimulate the premotor cortex to find more optimal solutions to unanticipated events during per-

formance.<sup>51</sup> Although the results of the transfer task activity show promise, it is speculative to apply these findings to actual athletic competition. Because only a single session of training was provided, the maintenance of any change in performance or how multiple sessions of training can further influence skill learning are currently unknown. Future studies should focus on longer term retention of the biomechanical changes and how landing performance changes during more complex movement patterns that more closely mimics athletic activities.

The feedback training system utilized in this study overcomes many of the barriers associated with conventional laboratory-based equipment, as it is relatively easy to use, inexpensive to develop, and portable. This contributed to the ability to assess/train a large number of athletes in the field. There have been preliminary attempts to explore the potential utility of lower cost gaming systems (e.g. Wii Balance Board or Microsoft Kinect) for movement assessment and feedback training.<sup>45,46</sup> These systems overcome the barriers associated with laboratory-based equipment; however, they have technical limitations (e.g. low sampling rates, limited sensor range) that restricts their accuracy for dynamic tasks such as landing. The researchers believe that there is a need to continue to develop/utilize systems such as this which are conducive to testing/training in the field, but also overcome the technical limitations of other systems.

Findings from this investigation appear to indicate that a portable force plate interfaced with a digital camera provide promise to promoting biomechanical changes that may reduce an athlete's relative risk for ACL injury. Several researchers have reported that greater vertical ground reaction forces at impact and increased knee valgus predispose female athletes to non-contact ACL injuries.<sup>3,47-51</sup> Greater impact forces have been associated with anterior tibial accelerations.<sup>32</sup> In vitro and modeling studies have reported greater ACL strain and tibial shear with landing patterns that lack adequate knee flexion to absorb impact forces.<sup>26,52-55</sup> Increased knee valgus angle during landing has been associated with increased risk for ACL injury in females.<sup>3</sup> These kinetic and kinematic findings have led to the

development of neuromuscular programs aimed at mitigating these risk factors.<sup>4,13,15,32,56</sup> However, most preventative programs do not utilize visual feedback training. The effect of combining of this type of feedback system into a more traditional neuromuscular training program offers promise and appears to warrant further investigation. It is plausible that the use of an augmented feedback system may provide a more effective avenue for neuromuscular training that incorporates the principles of motor learning in skill development.

The present study has limitations. Although the use of a drop landing is widely used in research studies to evaluate kinematics and kinetics that may be related to knee injury, the task may lack external validity. Similarly, the transfer task utilized in this study of a random "catch" or "no catch" may not provide a close enough parallel to the multiple simultaneous environmental cues and motor planning required for sports. Lastly, this study was conducted over a single testing session therefore the long-term retention effects of these performance changes is unknown.

## CONCLUSIONS

The results of this study indicate that using a portable clinical feedback system may be an effective tool in reducing peak vGRFs and knee abduction angles during a drop landing and transfer task in adolescent females. Future studies should focus on the retention effects of using augmented feedback systems on performance modifications.

---

## CONFLICTS OF INTEREST

Authors present no conflict of interest associated with this work.

Submitted: January 20, 2020 CST, Accepted: July 24, 2020 CST



## REFERENCES

1. Arendt EA, Agel J, Dick R. Anterior cruciate ligament injury patterns among collegiate men and women. *J Athl Train*. 1999;34(2):86-92.
2. Myer GD, Ford KR, Khoury J, Succop P, Hewett TE. Clinical correlates to laboratory measures for use in non-contact anterior cruciate ligament injury risk prediction algorithm. *Clin Biomech (Bristol, Avon)*. 2010;25(7):693-699. doi:10.1016/j.clinbiomech.2010.04.016
3. Hewett TE, Myer GD, Ford KR, et al. Biomechanical measures of neuromuscular control and valgus loading of the knee predict anterior cruciate ligament injury risk in female athletes: a prospective study. *Am J Sports Med*. 2005;33(4):492-501. doi:10.1177/0363546504269591
4. Onate JA, Guskiewicz KM, Sullivan RJ. Augmented feedback reduces jump landing forces. *J Orthop Sports Phys Ther*. 2001;31(9):511-517. doi:10.2519/jospt.2001.31.9.511
5. Paterno MV, Schmitt LC, Ford KR, et al. Biomechanical measures during landing and postural stability predict second anterior cruciate ligament injury after anterior cruciate ligament reconstruction and return to sport. *Am J Sports Med*. 2010;38(10):1968-1978. doi:10.1177/0363546510376053
6. Wiggins AJ, Grandhi RK, Schneider DK, Stanfield D, Webster KE, Myer GD. Risk of secondary injury in younger athletes after anterior cruciate ligament reconstruction: A systematic review and meta-analysis. *Am J Sports Med*. 2016;44(7):1861-1876. doi:10.1177/0363546515621554
7. Ardern CL. Anterior cruciate ligament reconstruction-Not exactly a one-way ticket back to the preinjury level: A review of contextual factors affecting return to sport after surgery. *Sports Health*. 2015;7(3):224-230. doi:10.1177/1941738115578131
8. Sugimoto D, Myer GD, Bush HM, Klugman MF, Medina McKeon JM, Hewett TE. Compliance with neuromuscular training and anterior cruciate ligament injury risk reduction in female athletes: a meta-analysis. *J Athl Train*. 2012;47(6):714-723. doi:10.4085/1062-6050-47.6.10
9. Benjaminse A, Lemmink KAPM, Diercks RL, Otten B. An investigation of motor learning during side-step cutting: design of a randomised controlled trial. *BMC Musculoskelet Disord*. 2010;11:235. doi:10.1186/1471-2474-11-235
10. Etnoyer J, Cortes N, Ringleb SI, Van Lunen BL, Onate JA. Instruction and jump-landing kinematics in college-aged female athletes over time. *J Athl Train*. 2013;48(2):161-171. doi:10.4085/1062-6050-48.2.09
11. Parsons JL, Alexander MJL. Modifying spike jump landing biomechanics in female adolescent volleyball athletes using video and verbal feedback. *J Strength Cond Res*. 2012;26(4):1076-1084. doi:10.1519/JSC.0b013e31822e5876
12. Welling W, Benjaminse A, Gokeler A, Otten B. Enhanced retention of drop vertical jump landing technique: A randomized controlled trial. *Hum Mov Sci*. 2016;45:84-95. doi:10.1016/j.humov.2015.11.008
13. Prapavessis H, McNair PJ. Effects of instruction in jumping technique and experience jumping on ground reaction forces. *J Orthop Sports Phys Ther*. 1999;29(6):352-356. doi:10.2519/jospt.1999.29.6.352
14. Munro A, Herrington L. The effect of videotape augmented feedback on drop jump landing strategy: Implications for anterior cruciate ligament and patellofemoral joint injury prevention. *Knee*. 2014;21(5):891-895. doi:10.1016/j.knee.2014.05.011
15. Myer GD, Stroube BW, DiCesare CA, et al. Augmented feedback supports skill transfer and reduces high-risk injury landing mechanics: a double-blind, randomized controlled laboratory study. *Am J Sports Med*. 2013;41(3):669-677. doi:10.1177/0363546512472977
16. Oñate JA, Guskiewicz KM, Marshall SW, Giuliani C, Yu B, Garrett WE. Instruction of jump-landing technique using videotape feedback: altering lower extremity motion patterns. *Am J Sports Med*. 2005;33(6):831-842. doi:10.1177/0363546504271499
17. Myer GD, Faigenbaum AD, Chu DA, et al. Integrative training for children and adolescents: techniques and practices for reducing sports-related injuries and enhancing athletic performance. *Phys Sportsmed*. 2011;39(1):74-84. doi:10.3810/psm.2011.02.1854
18. Myers CA, Torry MR, Peterson DS, et al. Measurements of tibiofemoral kinematics during soft and stiff drop landings using biplane fluoroscopy. *Am J Sports Med*. 2011;39(8):1714-1722. doi:10.1177/0363546511404922

19. Barrios JA, Crossley KM, Davis IS. Gait retraining to reduce the knee adduction moment through real-time visual feedback of dynamic knee alignment. *J Biomech.* 2010;43(11):2208-2213. doi:10.1016/j.jbiomech.2010.03.040
20. Herman DC, Oñate JA, Weinhold PS, et al. The effects of feedback with and without strength training on lower extremity biomechanics. *Am J Sports Med.* 2009;37(7):1301-1308. doi:10.1177/0363546509332253
21. Olbrantz C, Bergelin J, Asmus J, Kernozek T, Rutherford D, Gheidi N. Effect of posttrial visual feedback and fatigue during drop landings on patellofemoral joint stress in healthy female adults. *J Appl Biomech.* 2018;34(1):82-87. doi:10.1123/jab.2017-0074
22. Tegner Y, Lysholm J. Rating systems in the evaluation of knee ligament injuries. *Clin Orthop Relat Res.* 1985;(198):43-49.
23. Adams R. Revised physical activity readiness questionnaire. *Can Fam Physician.* 1999;45:992, 995, 1004-1005.
24. Silva MG, Moreira PVS, Rocha HM, Silva MG, Moreira PVS, Rocha HM. Development of a low cost force platform for biomechanical parameters analysis. *Res Biomed Engineer.* 2017;33(3):259-268. doi:10.1590/2446-4740.01217
25. Mizner RL, Chmielewski TL, Toepke JJ, Tofte KB. Comparison of 2-dimensional measurement techniques for predicting knee angle and moment during a drop vertical jump. *Clin J Sport Med.* 2012;22(3):221-227. doi:10.1097/JSM.0b013e31823a46ce
26. Kernozek TW, Ragan RJ. Estimation of anterior cruciate ligament tension from inverse dynamics data and electromyography in females during drop landing. *Clin Biomech (Bristol, Avon).* 2008;23(10):1279-1286. doi:10.1016/j.clinbiomech.2008.08.001
27. Noyes FR, Barber-Westin SD, Fleckenstein C, Walsh C, West J. The drop-jump screening test: difference in lower limb control by gender and effect of neuromuscular training in female athletes. *Am J Sports Med.* 2005;33(2):197-207. doi:10.1177/0363546504266484
28. Ericksen HM, Lefevre C, Luc-Harkey BA, Thomas AC, Gribble PA, Pietrosimone B. Females decrease vertical ground reaction forces following 4-week jump-landing feedback intervention without negative affect on vertical jump performance. *J Sport Rehabil.* February 2019:1-5. doi:10.1123/jsr.2018-0140
29. Schmitt R, Lee T, Winstein CJ, Wulf G, Zelazak H. *Motor Control and Learning, 6E.* Human Kinetics; 2018.
30. Popovic T, Caswell SV, Benjaminse A, Siragy T, Ambegaonkar J, Cortes N. Implicit video feedback produces positive changes in landing mechanics. *J Exp Orthop.* 2018;5(1):12. doi:10.1186/s40634-018-0129-5
31. Gokeler A, Benjaminse A, Welling W, Alferink M, Eppinga P, Otten B. The effects of attentional focus on jump performance and knee joint kinematics in patients after ACL reconstruction. *Phys Ther Sport.* 2015;16(2):114-120. doi:10.1016/j.ptsp.2014.06.002
32. McNair PJ, Prapavessis H, Callender K. Decreasing landing forces: effect of instruction. *Br J Sports Med.* 2000;34(4):293-296. doi:10.1136/bjism.34.4.293
33. Benjaminse A, Gokeler A, Dowling AV, et al. Optimization of the anterior cruciate ligament injury prevention paradigm: novel feedback techniques to enhance motor learning and reduce injury risk. *J Orthop Sports Phys Ther.* 2015;45(3):170-182. doi:10.2519/jospt.2015.4986
34. Wulf G, Dufek JS. Increased jump height with an external focus due to enhanced lower extremity joint kinetics. *J Mot Behav.* 2009;41(5):401-409. doi:10.1080/00222890903228421
35. Wulf G, Lewthwaite R. Conceptions of ability affect motor learning. *J Mot Behav.* 2009;41(5):461-467. doi:10.3200/35-08-083
36. Zachry T, Wulf G, Mercer J, Bezodis N. Increased movement accuracy and reduced EMG activity as the result of adopting an external focus of attention. *Brain Res Bull.* 2005;67(4):304-309. doi:10.1016/j.brainresbull.2005.06.035
37. Benjaminse A, Otten E. ACL injury prevention, more effective with a different way of motor learning? *Knee Surg Sports Traumatol Arthrosc.* 2011;19(4):622-627. doi:10.1007/s00167-010-1313-z
38. Masters RSW, Poolton JM, Maxwell JP. Stable implicit motor processes despite aerobic locomotor fatigue. *Conscious Cogn.* 2008;17(1):335-338. doi:10.1016/j.concog.2007.03.009
39. Poolton JM, Maxwell JP, Masters RSW, Raab M. Benefits of an external focus of attention: common coding or conscious processing? *J Sports Sci.* 2006;24(1):89-99. doi:10.1080/02640410500130854
40. Hardy L, Mullen R, Jones G. Knowledge and conscious control of motor actions under stress. *Br J Psychol.* 1996;87 ( Pt 4):621-636. doi:10.1111/j.2044-8295.1996.tb02612.x



41. Ong NT, Bowcock A, Hodges NJ. Manipulations to the timing and type of instructions to examine motor skill performance under pressure. *Front Psychol*. 2010;1:196. doi:10.3389/fpsyg.2010.00196
42. Myer GD, Paterno MV, Ford KR, Hewett TE. Neuromuscular training techniques to target deficits before return to sport after anterior cruciate ligament reconstruction. *J Strength Cond Res*. 2008;22(3):987-1014. doi:10.1519/JSC.0b013e31816a86cd
43. Myer GD, Ford KR, Brent JL, Hewett TE. The effects of plyometric vs. dynamic stabilization and balance training on power, balance, and landing force in female athletes. *J Strength Cond Res*. 2006;20(2):345-353. doi:10.1519/R-17955.1
44. Stroube BW, Myer GD, Brent JL, Ford KR, Heidt RS, Hewett TE. Effects of task-specific augmented feedback on deficit modification during performance of the tuck-jump exercise. *J Sport Rehabil*. 2013;22(1):7-18. doi:10.1123/jsr.22.1.7
45. Clark RA, Howells B, Feller J, Whitehead T, Webster KE. Clinic-based assessment of weight-bearing asymmetry during squatting in people with anterior cruciate ligament reconstruction using Nintendo Wii Balance Boards. *Arch Phys Med Rehabil*. 2014;95(6):1156-1161. doi:10.1016/j.apmr.2014.02.024
46. Mentiplay BF, Hasanki K, Perraton LG, Pua Y-H, Charlton PC, Clark RA. Three-dimensional assessment of squats and drop jumps using the Microsoft Xbox One Kinect: Reliability and validity. *J Sports Sci*. 2018;36(19):2202-2209. doi:10.1080/02640414.2018.1445439
47. Padua DA, DiStefano LJ, Beutler AI, de la Motte SJ, DiStefano MJ, Marshall SW. The landing error scoring system as a screening tool for an anterior cruciate ligament injury-prevention program in elite-youth soccer athletes. *J Athl Train*. 2015;50(6):589-595. doi:10.4085/1062-6050-50.1.10
48. Chaudhari AM, Andriacchi TP. The mechanical consequences of dynamic frontal plane limb alignment for non-contact ACL injury. *J Biomech*. 2006;39(2):330-338. doi:10.1016/j.jbiomech.2004.11.013
49. Leppänen M, Pasanen K, Kujala UM, et al. Stiff landings are associated with increased ACL injury risk in young female basketball and floorball players. *Am J Sports Med*. 2017;45(2):386-393. doi:10.1177/0363546516665810
50. Dufek JS, Bates BT. Biomechanical factors associated with injury during landing in jump sports. *Sports Med*. 1991;12(5):326-337. doi:10.2165/00007256-199112050-00005
51. Norcross MF, Blackburn JT, Goerger BM, Padua DA. The association between lower extremity energy absorption and biomechanical factors related to anterior cruciate ligament injury. *Clin Biomech (Bristol, Avon)*. 2010;25(10):1031-1036. doi:10.1016/j.clinbiomech.2010.07.013
52. Laughlin WA, Weinhandl JT, Kernozek TW, Cobb SC, Keenan KG, O'Connor KM. The effects of single-leg landing technique on ACL loading. *J Biomech*. 2011;44(10):1845-1851. doi:10.1016/j.jbiomech.2011.04.010
53. Levine JW, Kiapour AM, Quatman CE, et al. Clinically relevant injury patterns after an anterior cruciate ligament injury provide insight into injury mechanisms. *Am J Sports Med*. 2013;41(2):385-395. doi:10.1177/0363546512465167
54. Shin CS, Chaudhari AM, Andriacchi TP. The effect of isolated valgus moments on ACL strain during single-leg landing: a simulation study. *J Biomech*. 2009;42(3):280-285. doi:10.1016/j.jbiomech.2008.10.031
55. Withrow TJ, Huston LJ, Wojtys EM, Ashton-Miller JA. The relationship between quadriceps muscle force, knee flexion, and anterior cruciate ligament strain in an in vitro simulated jump landing. *Am J Sports Med*. 2006;34(2):269-274. doi:10.1177/0363546505280906
56. Milner CE, Fairbrother JT, Srivatsan A, Zhang S. Simple verbal instruction improves knee biomechanics during landing in female athletes. *Knee*. 2012;19(4):399-403. doi:10.1016/j.knee.2011.05.005