

## Effect of an indigenous AM and PGPR combination on chilli growth and productivity in lateritic soil

Debashis Kuila<sup>1,2</sup> <https://orcid.org/0000-0001-9353-1289>

Bijoy Mal<sup>2</sup> <https://orcid.org/0000-0001-6282-1996>

Sudip Mondal<sup>2</sup> <https://orcid.org/0000-0003-1717-6355>

Somdatta Ghosh<sup>3\*</sup> <https://orcid.org/0000-0002-2558-8520>

Gunjan Biswas<sup>2</sup> <https://orcid.org/0000-0001-8776-7762>

<sup>1</sup>Mycorrhiza & Microbiology Research Section, UG & PG Department of Botany, Midnapore College (Autonomous). WB. India.

<sup>2</sup>Department of Botany and Forestry, Vidyasagar University. Midnapore 721102. WB. India.

<sup>3</sup>PG & UG Department of Botany, Midnapore College (Autonomous). Midnapore. WB. India.

\*Corresponding Author e-mail: [somdattaghosh@yahoo.co.in](mailto:somdattaghosh@yahoo.co.in)

### ABSTRACT

Infertile lateritic soil is particularly deficient in phosphorus (P) and Nitrogen (N). Arbuscular Mycorrhiza (AM) has a key role to uptake bound P from the soil and provide to the plants growing under P-poor conditions and improve water and nutrient uptake. *Azotobacter* fixes free nitrogen and phosphate solubilizing bacteria (PSB) release bound phosphate, are the important groups of plant growth-promoting rhizobacteria (PGPR), sometimes they may act as mycorrhiza helper and applied with AM as biofertilizer. This pot experiment was conducted to determine the primary impact of singly and combined application of native bio-inoculants, the AM, *Acaulospora*, and the PGPR, *Azotobacter* and *Pseudomonas* sp. (PSB) on growth and yield of chilli (*Capsicum frutescens* L.), growing in acid lateritic soil. Inoculated treatments were compared for growth and productivity of chilli in terms of height, leaf number, leaf area, root collar diameter, number of flowers and number of fruits, final fresh and dry yield. The productivity of chilli showed a maximum in combined treatment of *Acaulospora*, *Azotobacter*, and PSB. Also the AM spore count and root colonization found maximum in that treatment. Hence the application of indigenous AM inoculation along with native PGPR, *Azotobacter* and PSB may present better productivity in low fertile lateritic soil.

Keywords: *Acaulospora*, *Azotobacter*, infertile soil, mycorrhiza, PSB

## Efecto de una combinación indígena de AM y PGPR sobre el crecimiento y la productividad de los chiles en suelos lateríticos

### RESUMEN

El suelo laterítico infértil es particularmente deficiente en fósforo (P) y nitrógeno (N). Las micorrizas arbusculares (AM) tiene un papel clave para absorber el P unido del suelo y proporcionar a las plantas que crecen en condiciones de P pobre y mejorar la absorción de agua y nutrientes. *Azotobacter* que fija el nitrógeno libre y las bacterias solubilizadoras de fosfato (PSB), son grupos importantes de rizobacterias que promueven el crecimiento de las plantas (PGPR). A veces pueden actuar de conjunto con micorrizas y aplicarse con AM como biofertilizante. Este experimento en maceta se realizó para determinar el impacto primario de la aplicación individual y combinada de bio-inoculantes nativos, AM *Acaulospora* y PGPR *Azotobacter* y *Pseudomonas* sp. (PSB) sobre el crecimiento y el rendimiento del chile (*Capsicum frutescens* L.), que crece en suelo ácido laterítico. Los tratamientos inoculados se compararon para el crecimiento y la productividad del chile en términos de altura, número de hojas, área foliar, diámetro de raíz, número de flores, número de frutos, rendimiento final fresco y seco. La productividad de los

chiles mostró un máximo en el tratamiento combinado de *Acaulospora*, *Azotobacter* y PSB. También el recuento de esporas de AM y la colonización de raíces encontraron el máximo en ese tratamiento. De ahí la aplicación de la inoculación de AM indígena junto con PGPR nativo, *Azotobacter* y PSB pueden presentar una mejor productividad en suelos lateríticos de baja fertilidad.

Palabras clave: *Acaulospora*, *Azotobacter*, suelo infértil, micorriza, PSB

## INTRODUCTION

The fertility level of dry acid lateritic soil is very low. The soil is naturally deficient in phosphorus, nitrogen, and other essential nutrients like calcium, magnesium, etc. In this type of soil, nutrients remain unavailable to the plant, resulting in unfavourable conditions for plant growth (Koley, 2000). In conventional agricultural practices, uses of overdose agrochemicals day by day increasing of low fertility and toxicity in the soil in agricultural fields and pesticide-resistant pests throughout India. As a result of these real problems, adverse effects on health are being reflected (Mittal *et al.*, 2013; Rahman and Debnath *et al.*, 2015). Therefore, in modern agriculture strategy, the huge demand for producing organically toxic-free vegetables is increasing, which also cost-effective and environment-friendly (Madhusudhan, 2016; Pandey and Singh, 2012).

Mycorrhiza is the mutualistic root–fungus relationship between non-pathogenic soil fungi and the plant roots (Frank, 1885; van der Heijden *et al.*, 2015). Arbuscular mycorrhiza (AM), belonging to the phylum Glomeromycota (Redecker *et al.*, 2000; Kapoor *et al.*, 2008), are the most common symbiotic association of mycorrhiza found in 90% of land plants (Schüßler *et al.*, 2001; Brundrett, 2002; Smith and Read, 2008). AM colonizes in the cortical tissue of roots during active plant growth (Kirk *et al.*, 2001) and procures nutrient form beyond the nutrients depletion zone of roots (Li *et al.*, 1991). Organic acids and phosphatases produced by AM help to release of available phosphorous from unavailable complexes (Bagyaraj, 1984; Entry *et al.*, 2002; Fomina *et al.*, 2005). They also increase nitrogen and carbon content in soil (Almas *et al.*, 2004). The fine extra-radical hyphae of arbuscular mycorrhizal fungi (AMF) involve in the nutrient process and uptake phosphorous (Bowen, 1973; Bucher, 2007), provide nitrogen (Madder *et al.*, 2000; Chalot *et al.*, 2006; Nuccio *et al.*, 2013) by absorbing

ammonium, nitrate and amino acids (Hodge *et al.*, 2001), and deliver other plant micronutrients like copper, zinc, etc. (Smith and Read, 1997), and in exchange, withdraw organic sugar from plant roots (Kottke, 2002).

AM hyphae can absorb water from lower water potential than the plant roots (Bethlenfalvay *et al.*, 1988; Püschel *et al.*, 2020) and form a mycelial mat which promotes retaining soil moisture. AM symbiosis is effective to overcome low pH in acidic soil which restrict plant growth (Clark, 1997; Muthukumar *et al.*, 2014). The increasing water and nutrient uptake ability of AMF in low fertile and dry condition (Smith and Read, 1997; Aúge, 2001) provide improved plant growth in less-nutrient soil, such as acid lateritic soil (Weber *et al.*, 1992; Brundrett, 2009). They also protect plants from pathogenic fungi and nematode disease (Bagyaraj, 1984). Nowadays, arbuscular mycorrhizal inoculation to the crops has become an appreciable alternative that increases the growth and yield (Mishra and Verma, 1982; Johansen *et al.*, 1993; Ghosh and Verma, 2006; Samanta and Verma, 2006; Sengupta *et al.*, 2006; Robinson *et al.*, 2016; Chukwuka *et al.*, 2017).

*Acaulospora* is the AM fungal genus belonging to the family Acaulosporaceae, has been distributed in more than 30 species (Simanungkalit, 2006). *Acaulospora* sp. are a common dominant group of arbuscular mycorrhiza at the lateritic ecosystem (Ghosh and Verma, 2011). A special feature, having the sporiferous saccule with the saccule subtending hypha present in *Acaulospora* (INVAM, 2018).

The plant-growth-promoting rhizobacteria (PGPR), present in plant mycorrhizosphere, promote the growth of the plant (Andrade, 2004; Giri *et al.*, 2005). *Azotobacter* and phosphate solubilizing bacteria are the important plant growth promoters (PGP) and act as mycorrhiza helper, having an important role in the growth of hyphae from germinating

AM spores, colonization of plant roots by AM fungi and growth of external AM hyphae and dehydrogenase activity of the AM fungus (Burla *et al.*, 1996). *Azotobacter* is free-living soil bacteria, fix free nitrogen directly from the environment. Inoculation of *Azotobacter* sp. to the plant has showed improved beneficial effects in seed germination, shoot, root biomass, and yield (Kumar *et al.*, 2000; Kumar *et al.*, 2009; Malik *et al.*, 2009; Reddy *et al.*, 2003). Phosphate solubilizing bacteria (PSB) can release bound phosphate to the available P form (Bhattacharyya and Jain, 2000; Ponmurugan and Gopi, 2006). Many research works related to plant growth promotion by the application of phosphate solubilizers have been reported earlier (Khiari and Parent, 2002; Saxena and Sharma, 2003; Yazdani *et al.*, 2009).

Chilli (*Capsicum frutescens* L.) is the part of the Solanaceae family, is a very essential ingredient in our daily vegetables. It is rich in volatile oils, fatty oils, capsaicinoids, carotenoids, vitamins, proteins, fibers, and minerals and acts as a natural bactericidal, having medicinal uses to treat muscle pain, cough, asthma, sore throat, etc. (Bosland and Votava, 2000). As it is an economically important crop, the average yield of chilli in India is lower as compared to other developed countries (APEDA, 2019; FAOSTAT, 2017). Cropping in unused barren land containing infertile soil such as lateritic soil may become a suitable solution.

The present research was conducted to determine the primary impact of singly and combined application of native bio-inoculants, the AM, *Acaulospora*, and the PGPR, *Azotobacter* and *Pseudomonas* sp. (PSB) on growth and yield of chilli (*Capsicum frutescens* L.), growing in acid lateritic soil.

## MATERIALS AND METHODS

### *Plant material*

Certified seed of chilli (*Capsicum frutescens* L.) were collected from District Agricultural Head office, Paschim Medinipur, West Bengal, India. Healthy seeds were selected manually and surface disinfected by immersion in an aqueous solution of 0.1% (w/v) mercuric chloride (HgCl<sub>2</sub>) for 3-4 min followed by washing three times with autoclaved deionized

water under aseptic conditions in a laminar flow chamber.

### *Experimental conditions*

The experiment was conducted with chilli (*C. frutescens*) planted in sterilized lateritic soil (Sylvia, 1994). It was done in 22.30° N Latitude and 87.20° E Longitude in Midnapore subdivision of West Bengal in the pre-winter season. The primary soil characteristics were tested according to Jackson (1973), having pH 5.61, electrical conductivity (EC) of 0.18 m mohs/cm<sup>2</sup>, moisture content of 2.7%, organic carbon (OC) of 0.63 g kg<sup>-1</sup>, total nitrogen (N) of 0.04% and phosphate (P) of 0.03%.

### *Isolation of the AM propagules*

Separation and isolation of indigenous AM propagules from the native ecosystem was done by following the wet sieving and decanting technique (Gerdermann and Nicolson, 1963). Spores of single species were separated morphologically using a LABOMED CSM2 microscope, and surface disinfected by 200 µg ml<sup>-1</sup> streptomycin and 2% Chloramine-T solution (w/v) (Mosse, 1973).

The pure culture of each inoculum was cultured in sterilized sand and soil mixture. The soil: sand (1: 1, v/v) mixture was sterilized by oven drying at 85 °C for 8 h with a gap of 48 h and again dried for 8 h (Sylvia, 1994). Pure cultures were done in funnels and mass cultures in surface-disinfected earthen pots, grown in an automated growth chamber with sorghum (*Sorghum vulgare* L.) plants. The mass cultures were maintained up to 90 days in the growth chamber at 26±2 °C under a photoperiod of 16 h light and 8 h dark, an irradiance provided by 110 W fluorescent lamps (Philips, India).

The successful mass culture of the indigenous AM inocula was identified morphologically by following Schenck and Perez (1990) and the INVAM web photo guide (INVAM, 2018) and confirmed as *Acaulospora* sp.

### *Bacterial cultures*

*Azotobacter* was isolated from lateritic rhizospheric soil in Ashbey mannitol agar media for free-living nitrogen-fixing bacteria (Pelczar *et al.*, 1957; Subba, 1977). Pikovskaya media for native phosphate

solubilizing bacteria (Pikovskaya, 1948; Sundara and Sinha, 1963; Subba, 1977) were used. *Azotobacter* sp. and *Pseudomonas* sp. (PSB) were confirmed according to Bergey manual (Holt and Krieg, 1984; Krieg *et al.*, 1994).

#### Seedling inoculation

Surface disinfected seeds of chilli were germinated in aseptic conditions and seeded with a 10 cm gap from each other in a seedling tray within a growing chamber under the environment at temperature 25-30 °C, humidity 65-70% and 15-16 h of lighting. Sterilized soil and sand mixture in 2:1 (v/v) ratio was used as growing substrate.

After germination, the healthy chilli seedlings were transferred to polythene bags (18 X 24 cm) filled with 3 kg sterilized soil (one seedling per pot) to grow for 90 days in the greenhouse. AM inoculation was done by soil-based inoculum obtained from mass culture (20 spores /g, 25 g /pot), applied under 4 cm from the soil surface. 10 ml of the bacterial cultures contained about 10<sup>8</sup> CFU ml<sup>-1</sup> liquid medium were applied per pot.

The experiment was designed in a randomized block with eight treatments and three

replicates. The treatments of this experiment consisted with T<sub>0</sub> = No inoculation (Control), T<sub>1</sub> = *Azotobacter* (AZO), T<sub>2</sub> = Phosphate solubilizing bacteria (PSB), T<sub>3</sub> = *Acaulospora* sp. (Ac), T<sub>4</sub> = *Azotobacter* + PSB (AZO+PSB), T<sub>5</sub> = *Acaulospora* sp. + *Azotobacter* (Ac+AZO), T<sub>6</sub> = *Acaulospora* sp. + PSB (Ac+PSB), T<sub>7</sub> = *Acaulospora* sp. + *Azotobacter* + PSB (Ac+AZO+PSB).

Plant parameters were measured at 30, 60 and 90 days after plantation (dap) in three replicates per treatment in terms of shoot height (cm), leaf number per plant, leaf area (cm<sup>2</sup>), the total number of flower and fruit appeared per plant, and total fresh weight (g) and dry mass (g) of fruit yield per plant. The fresh yield was calculated by summing the total fresh weight of fruits obtained. The dry mass was measured by drying the fruits in a hot air oven at 65 °C for 48 hours until a constant weight was obtained.

Rhizospheric soil samples were tested for the AM spore population by wet sieving and decanting technique (Gerdermann and Nicolson, 1963). Plant root samples were treated with 10% KOH and stained with trypan blue (Phillips and Hayman, 1970) to study the colonization percentage by the equation 1:

$$\text{Root colonization \%} = \left( \frac{\text{Number of root pieces colonized}}{\text{Total number of root pieces observed}} \right) \times 100$$

#### Statistical analysis

Statistical analysis of data was done by comparison of means using the least significant difference (LSD) at p < 0.05 probability, after the performing analysis of variance (ANOVA) using IBM SPSS 20 and MS Excel 2013. Pearson correlation coefficient between variables was analyzed at p < 0.05 and p < 0.01 level of significance.

#### RESULTS AND DISCUSSION

The inoculation of chilli seedlings with an indigenous AM and PGPR combination increased the growth since the early stage, with prominent effect of AM.

At 30 dap, the treatment T<sub>7</sub> having *Acaulospora* sp. + *Azotobacter* + PSB (Ac+AZO+PSB) showed height, leaf number,

leaf area and root collar diameter slightly superior. Nevertheless, after 60 days of culture, the plants in treatment Ac+AZO+PSB (T7) were 23.66% higher than the control, followed by Ac+AZO (21.3%), Ac+PSB (15.5%) and Ac (10.6%). The same pattern was observed in the total leaf number and root collar diameter. Leaf area was in Ac+AZO+PSB 14.3%, in Ac+AZO 11.1% and in Ac 10.0% higher than the control.

The first appearance of flowering have few days of difference among the treatments. It was observed on 26<sup>th</sup> day after plantation in AZO+PSB and Ac+AZO+PSB, at 28<sup>th</sup> day in Ac+AZO and at 29<sup>th</sup> day in AZO, Ac, and Ac+PSB. On the other hands, first fruiting was observed at 33<sup>rd</sup> dap in AZO+PSB followed by Ac+AZO+PSB and Ac+AZO at the 36<sup>th</sup> day; and Ac and Ac+PSB at 41<sup>st</sup> day.

After 90 dap, in treatment with the combination of AM and PGPR (Ac+AZO+PSB, T7) all the parameters reached the maximum, without significantly differences with Ac+AZO (T5) in plant height, leaf number and root collar diameter. As well as T7 treatment was similar to T6 (Ac+PBS) in root collar diameter (Table 1). The shoot height

in Ac+AZO+PSB was 31.64% superior to control and the leaf number 37.8%. The total fruit number was found maximum in Ac+AZO+PSB, 127.5% better than control and significantly higher ( $p < 0.05$ ) all the rest. A similar result was observed in the total fresh and dry mass of fruits yield per plant (Figure 1).

Table 1. Effect AM and PGPR on chilli (*Capsicum frutescens* L.) plants parameters after 90 days after plantation.

| Treatments                  | Height (cm) | Leaf Number | Leaf Area (cm <sup>2</sup> ) | Root Collar Diameter (cm) | Flower Number | Fruit Number |
|-----------------------------|-------------|-------------|------------------------------|---------------------------|---------------|--------------|
| T <sub>0</sub> Control      | 17.38 c     | 28.3 b      | 2.58 b                       | 0.98 c                    | 5 b           | 3.66 d       |
| T <sub>1</sub> (AZO)        | 19.5 b      | 30.6 b      | 2.68 b                       | 1.13 b                    | 7 b           | 5 c          |
| T <sub>2</sub> (PSB)        | 18.74 c     | 29 b        | 2.63 b                       | 1.07 c                    | 6 b           | 5 c          |
| T <sub>3</sub> (Ac)         | 20.52 b     | 31.3 b      | 2.79 b                       | 1.29 a                    | 7 b           | 6.33 b       |
| T <sub>4</sub> (AZO+PSB)    | 20.12 b     | 31 b        | 2.77 b                       | 1.2 b                     | 6.6 b         | 5 c          |
| T <sub>5</sub> (Ac+AZO)     | 21.18 a     | 36.6 a      | 2.95 b                       | 1.35 a                    | 8 a           | 7 b          |
| T <sub>6</sub> (Ac+PSB)     | 20.76 b     | 34.6 b      | 2.82 b                       | 1.3 a                     | 7.6 a         | 6.33 b       |
| T <sub>7</sub> (Ac+AZO+PSB) | 22.88 a     | 39 a        | 3.05 a                       | 1.38 a                    | 8.6 a         | 8.33 a       |

Data with different letters in the same column indicate significant differences at  $p < 0.05$  according to LSD test. AZO-Azotobacter sp., Ac- Acaulospora sp., PBS-Pseudomonas sp.




Figure 1. Fresh yield and dry mass of *Capsicum frutescens* L. plants inoculated with AM and PGPR. Treatments T<sub>1</sub> (AZO), T<sub>2</sub> (PSB), T<sub>3</sub> (Ac), T<sub>4</sub> (AZO+PSB), T<sub>5</sub> (Ac+AZO), T<sub>6</sub> (Ac+PSB), T<sub>7</sub> (Ac+AZO+PSB). AZO-Azotobacter sp., Ac- Acaulospora sp., PBS-Pseudomonas sp.

In single inoculation, the effect of AM was superior than *Azotobacter* and PSB in root collar diameter and fruit number. In dual inoculation, the combination of AM and *Azotobacter* was better than AM and PSB in height and leaf number. Plants inoculated with *Azotobacter* sp. have shown an increase in shoot and yield effects. It have been tested on different crops (Reddy *et al.*, 2003; Singh and Rana, 2005; Kumar *et al.*, 2009; Malik *et al.*, 2009). The dual effect of inoculation of AMF and *Azotobacter* was found encouraging in earlier research work also (Paul *et al.*, 2011; Dal *et al.*, 2018; Shaimaa and Massoud, 2017). *Azotobacter* adds only nitrogen in soil and PSB release available phosphate, AM mainly translocate nutrients including micronutrients through the extensive mycelial network (Kayama and Yamanaka, 2014; Rouphael *et al.*, 2015) and also release phosphates by phosphatase (Zhang *et al.*, 2014).

Hence, AM alone is found almost sufficient for nutrition rather than *Azotobacter* or PSB in single inoculation. In dual treatments, also uptake ability of AM was in advance than the only supply of nutrients, and *Azotobacter* as a source of N and AM as P substitute worked best. Interaction between the AM, *Acaulospora* with *Azotobacter* and PSB, in this

triple application may provide the development of plant growth and yield (Nadeem *et al.*, 2014; Hashem *et al.*, 2016). Enhanced supply of N and P in the rhizosphere by PGPR boosts up AM uptake in the triple inoculation resulting establishment of better mycorrhizal activity (Selvakumar *et al.*, 2012; Vafadar *et al.*, 2014; Raklami *et al.*, 2019).

The mycorrhizal infection and spore population were found maximum in Ac+AZO+PSB (T7) treatment followed by MA combined with PGPR (Ac+AZO, Ac+PSB) (Figure 2). Mycorrhizal colonization was mostly mycelial and arbuscular.

Mycorrhizal colonization with class – III intensity produced in Ac+AZO+PSB and Ac+AZO. In these treatments, colonization percentage were observed higher than the inoculation formed in Ac alone. A strong significant positive correlation ( $r=0.997$ ,  $p<0.05$  and  $p<0.01$ ) was observed between AM root colonization and spore density. They were also showed a significant positive correlation ( $p<0.05$  and  $p<0.01$ ) with all the measured plant parameters, including fresh yield and dry yield. The last one were also positively and significantly correlated ( $p<0.05$  and  $p<0.01$ ) with plant height.




Figure 2. Mycorrhizal colonization and spore population in root and rhizosphere of chilli plant inoculated with AZO-*Azotobacter* sp., Ac- *Acaulospora* sp., PSB-*Pseudomonas* sp.

In all the AM related treatments, arbuscular colonization and spore formation occurred. Formation of arbuscles and high-intensity colonization in the plant roots indicates active symbiosis between *Acaulospora* and *C. frutescens* in those treatments. As the natural propagules in lateritic soil are low, the mycorrhizal application to the plant increased the growth and yield (Ghosh and Verma, 2006; Hempel *et al.*, 2013). AM are more active in nutrient-poor dry soils, plants depended more on mycorrhizae for nutrients and moisture in this condition (Brundrett, 2009; Begum *et al.*, 2019). The mycorrhization facilitates an effective hyphal network that spreads throughout the rhizosphere and improves the nutrient absorbing ability of the host plant (Avio *et al.*, 2006; Püschel *et al.*, 2020). Active mycorrhizal colonization in the plants alters some physiological processes and increases water and nutrient uptake which enhances crop growth and yield (Oyetunji *et al.*, 2003; Mardukhi *et al.*, 2011; Douds *et al.*, 2005), which also influenced the chilli plant (Gaur *et al.*, 1998; Vyas and Vyas, 2014).

The test plant, *C. frutescens* establish showed AM dependency on the dominant indigenous AM genus, *Acaulospora*, in this lateritic zone (Sieverding, 1991; Ghosh *et al.*, 2008), which shows active adaptation and efficacy in native soil (Oliveira *et al.*, 2005; Samanta and Verma, 2006; Akib *et al.*, 2018). The triple application of AM, *Azotobacter*, and PSB were found to influence the growth of the crops too (Shwetha and Lakshman, 2013; Lallawmkima *et al.*, 2018), though with other AM species, not indigenous or in lateritic soil.

Enhancement in productivity of chilli through the mycorrhization established in earlier works (Selvakumar and Thamizhiniyan, 2011; Bhuvaneshwari *et al.*, 2014). The mycorrhizal inoculation can enhance the yield of chillies significantly and reduced the application of chemical fertilizers by up to 50 percent (Bagyaraj and Sreeramulu, 1982). Application of AM with such mycorrhiza helping rhizobacteria develops a synergistic microbial interaction (Artursson *et al.*, 2006) and maybe a substitute for chemical fertilizer particularly in this dry acidic soil.

## CONCLUSIONS

Application of indigenous AM, *Acaulospora* sp. along with beneficial and native plant growth-

promoting rhizobacteria, *Azotobacter* and PSB facilitate *C. frutescens* L. better growth and enhanced the yield grown in infertile dry and acid lateritic soil. Such AM and microbial consortium could be benefit other crops also in this soil type and organic agriculture and horticultural practice particularly in other poor nutrient soil and in surely reduced cost in comparison to chemical fertilizers.

## ACKNOWLEDGMENT

The first author is indebted to the University Grant Commission (UGC), Government of India for the financial assistance to conduct the research work. The authors are also thankful to Prof. N. K. Verma for his comments and advice on the study.

## Conflict of interest

This work was done all together and no potential conflict of interest present among the authors.

## Author contributions

Conceptualization SG, Data curation DK and BM, Formal analysis DK, Funding acquisition SG, Investigation SG, Methodology DK, BM and SM, Project administration SG, Resources SG and GB, Supervision SG, Software DK and SM, Validation SG, Visualization GB, Writing—original draft DK, Writing—review & editing SG.

## REFERENCES

- Akib MA, Mustari K, Kuswinanti T, Syaiful SA, (2018) Identification and abundance of indigenous endomycorrhiza isolated from nickel post-mining plantation in Sorowako. *Int J Curr Res Biosci Plant Biol* 5(4): 8-16; doi: 10.20546/ijcrbp.2018.504.002
- Almas AR, Bakken LR, Mulder J (2004) Changes in tolerance of soil microbial communities in Zn and Cd contaminated soils. *Soil Biol Biochem* 36(5): 805-813
- Andrade G (2004) Role of functional groups of microorganisms on the rhizosphere microcosm dynamics. In: Varma A, Abbott L, Werne D, Hampp R (eds). *Plant Surface Microbiology*, pp. 51-68. NY Springer, New York
- APEDA (2019) Agri Exchange, Agricultural & Processed Food Products Export Development

- Authority, Ministry of Commerce and industry, government of India. Available in: <http://apeda.gov.in/> Accessed 11/05/2019
- Artursson V, Finlay RD, Jansson JK (2006) Interactions between arbuscular mycorrhizal fungi and bacteria and their potential for stimulating plant growth. *Environmental Microbiology* 8: 1-10
- Aúge RM (2001) Water relations, drought and vesicular – arbuscular mycorrhizal symbiosis. *Mycorrhiza* 11: 3-42
- Avio L, Pellegrino E, Bonari E, Giovannetti M (2006) Functional diversity of arbuscular mycorrhizal fungal isolates in relation to extraradical mycelial networks. *New Phytologist* 172: 347-357; doi: 10.1111/j.1469-8137.2006.01839.x
- Bagyaraj DJ, Sreeramulu KR (1982) Preinoculation with VA mycorrhiza improves growth and yield of chilli transplanted in the field and saves phosphatic fertilizer. *Plant Soil* 69: 375-381
- Bagyaraj DJ (1984) Biological interactions with VA mycorrhizal fungi. In: Powell CL, Bagyaraj DJ (eds). *VA Mycorrhiza*, pp. 131-154. CRC Press, Florida
- Begum N, Qin C, Ahanger MA, Raza S, Khan MI, Ashraf M, Ahmed N, Zhang L (2019) Role of Arbuscular Mycorrhizal Fungi in Plant Growth Regulation: Implications in Abiotic Stress Tolerance. *Front Plant Sci* 10: 1068; doi: 10.3389/fpls.2019.01068
- Bethlenfalvay GJ, Brown MS, Ames RN, Thomas RS (1988) Effects of drought on host and endophyte development in mycorrhizal Soyabeans in relation to water use and phosphate uptake. *Plant Physiol* 72: 565-571
- Bhattacharyya P, Jain RK (2000) Phosphorus solubilizing biofertilizers in the whirlpool of rock phosphate-challenges and opportunities. *Fertilizer News* 45: 45-52
- Bhuvaneshwari G, Reetha S, Sivaranjani R, Ramakrishnan K (2014) Effect of AM fungi and *Trichoderma* species as stimulations of growth and morphological character of chilli (*Capsicum annum* .L). *International Journal of Current Microbiology and Applied Science* 3(3): 447-455
- Bosland PW, Votava EJ (2000) Peppers: Vegetable and spice capsicums. CABI Publishing, Oxon
- Bowen GD (1973) Mineral nutrition in mycorrhizas. In: Marks GC, Kozlowski TT (eds). *Ectomycorrhizae*, pp. 151-201. Academic Press, New York
- Brundrett MC (2009) Mycorrhizal associations and other means of nutrition of vascular plants: understanding the global diversity of host plants by resolving conflicting information and developing reliable means of diagnosis. *Plant Soil* 320: 1-41
- Brundrett MC (2002) Coevolution of roots and mycorrhizas of land plants. *New Phytol* 154: 275-304
- Bucher M (2007) Functional biology of plant phosphate uptake at root and mycorrhiza interfaces. *New Phytol* 173: 11-26
- Burla M, Goverde M, Schwinn FJ, Wiemken A (1996) Influence of biocontrol organisms on root pathogenic fungi and on the plant symbiotic microorganisms *Rhizobium phaseoli* and *Glomus mosseae*. *J Plant Dis Prot* 103: 156-163
- Chalot M, Blaudez D, Brun A (2006) Ammonia: a candidate for nitrogen transfer at the mycorrhizal interface. *Tren Plant Sci* 11: 263-266
- Chukwuka KS, Okechukwu RU, Umukoro BO (2017) Arbuscular mycorrhiza fungi, NPK (15-15-15) and cow dung interaction in sustainable cassava production and food security. *Adv Plants Agric Res* 7(4): 328-335; doi: 10.15406/apar.2017.07.00262
- Clark RB (1997) Arbuscular mycorrhizal adaptation, spore germination root colonization and lost. *Plant Soil* 192(1): 15-22
- Dal C, Barion G, Ferrari M, Visioli G, Dramis L, Panozzo A, Vamerali T (2018) Effects of field inoculation with VAM and bacteria consortia on root growth and nutrients uptake in common wheat. *Sustainability* 10: 3286; doi: 10.3390/su10093286
- Douds DD, Nagahashi G, Pfeffer PE, Kayser WM, Reider C (2005) On-farm production and



- utilization of arbuscular mycorrhizal fungus inoculum. *Can J Plant Sci* 85: 15-21
- Entry JA, Rygiewiez PT, Watrud LS, Donnelly PK (2002) Influence of adverse soil condition on the formation and functioning of arbuscular mycorrhizas. *Adv Environ Res* 7: 123-138
- FAOSTAT (2017) Chili production in 2016, Crops-World Regions-Production Quantity-Green Chillies and Peppers from pick lists, UN Food and Agriculture Organization, Statistics Division. Available in: <http://faostat.fao.org/> Accessed 25/04/2018
- Fomina MA, Alexander IJ, Colpaert JV, Gadd GM (2005) Solubilization of toxic metal minerals and metal tolerance of mycorrhizal fungi. *Soil Biology and Biochemistry* 37: 297-299
- Frank AB (1885) Ueber die aug warzel symbiose beruhende, Ernährung gewisser Baume durch Unterirdische Plize Ber Dtsch. *Bot Ges* 3: 128-145
- Gaur A, Adholeya A, Mukerji KG (1998) A comparison of AM fungi inoculants using *Capsicum* and *Polianthes* in marginal soil amended with organic matter. *Mycorrhiza* 7(6): 307-312
- Gerdermann JW, Nicolson TH (1963) Spores of mycorrhizal endogone species extracted from soil by wet sieving and decanting. *Transactions of the British Mycological Society* 46: 235-244
- Ghosh S, Verma NK (2006) Growth and mycorrhizal dependency of *Acacia mangium* Willd. inoculated with three vesicular arbuscular mycorrhizal fungi in lateritic soil. *New Forests* 31: 75-81
- Ghosh S, Verma NK (2011) Impact of rhizospheric conditions on AM diversity, succession, and colonization in two plantations of *Acacia auriculiformis* and *Eucalyptus tereticornis*. *Mycorrhiza News* 22(4): 5-7
- Ghosh S, Knap UK, Verma NK (2008) Effect of four arbuscular mycorrhizae on *Acacia mangium* Willd. Seedlings in lateritic soil. *Indian J Plant Physiol* 13(4): 375-380
- Giri B, Giang PH, Kumari R, Prasad R, Varma A (2005) Microorganisms in soils: roles in genesis and functions. In: Buscotand F, Varma A (eds). *Soil Biology*, pp. 19-43. NY Springer-Verlag, New York
- Hashem A, Abd Allah EF, Alqarawi AA, Al-Huqail AA, Wirth S, Egamberdieva D (2016) The interaction between arbuscular mycorrhizal fungi and endophytic bacteria enhances plant growth of *Acacia gerrardi* under salt stress. *Front Microbiol* 7: 1089; doi: 10.3389/fmicb.2016.01089
- Hempel S, Hötzenberger L, Kühn I, Michalski SG, Rillig MC, Zobel M, Moora M (2013) Mycorrhizas in the Central European Flora: relationships with plant life history traits and ecology. *Ecology* 94: 1389-1399
- Hodge A, Campbell CD, Fitter AH (2001) An arbuscular mycorrhizal fungus accelerates decomposition and acquires nitrogen directly from organic material. *Nature* 413: 297-299
- Holt JG, Krieg NR (1984) *Bergey's Manual of Systematic Bacteriology*. Williams and Wilkins, Baltimore
- INVAM (2018) Classification, International Culture Collection of Vesicular Arbuscular Mycorrhizal Fungi. Available in: <https://invam.wvu.edu> Accessed 23/05/2019
- Jackson ML (1973) *Soil chemical analyses*, Prentice Hall, New Delhi
- Johansen A, Jakobsen I, Jensen ES (1993) Hyphal N transport by a vesicular-arbuscular mycorrhizal fungus associated with cucumber grown at three nitrogen levels. *Plant and Soil* 160: 1-9; doi: 10.1007/BF00150340
- Kapoor R, Sharma D, Bhatnagar AK (2008) Arbuscular mycorrhizae in micropropagation systems and their potential applications. *Sci Hortic* 116: 227-239
- Kayama M, Yamanaka T (2014) Growth characteristics of ectomycorrhizal seedlings of *Quercus glauca*, *Quercus salicina* and *Castanopsis cuspidata* planted on acidic soil. *Trees* 28: 569-583; doi: 10.1007/s00468-013-0973-y
- Khiari L, Parent V (2002) Phosphorus transformations in acid light-textured soils

- treated with dry swine manure. *Canad J Microbiol* 85: 75-87
- Kirk PM, Cannon PF, David JC, Stalfers JA (2001) *Ainswrth and Bisby's Dictionary of the fungi*, (9th ed). CAB International, Wallingford, UK
- Koley AK (2000) *Basic concepts of soil science*. New age international publishers, India
- Kottke I (2002) Mycorrhizae-rhizosphere determinants of plant communities. In: Waisel Y, Eshel A, Kafkafi U (eds). *Plant Roots, the hidden half* (3rd ed), pp. 919-932. Marcel Dekker, New York
- Krieg NR, Holt JG, Sneath PHA, Staley JT, Williams ST (1994) *Bergey's Manual of Determinative Bacteriology*, (9th ed). Williams & Wilkins, Baltimore
- Kumar V, Aggarwal NK, Singh BP (2000) Influence of P - solubilizing analogue resistant mutants of *Azotobacter chroococcum* on yield and quality parameters of *Helianthus annuus*. *Folia Microbiologica* 45: 347-352
- Kumar V, Solanki AS, Sharma S (2009) Yield and economics of *Withania somnifera* influenced by dual inoculation of *Azotobacter chroococcum* and *Pseudomonas putida*. *Turkish Journal of Biology* 33: 219-223
- Lallawmkima I, Singh SK, Sharma M (2018) Application of *Azotobacter*, Vesicular Arbuscular Mycorrhiza and Phosphate Solubilizing Bacteria for potato cultivation in Central Plain Zone (Pb-3) of Punjab. *Journal of Environmental Biology* 39: 985-989; doi: 10.22438/jeb/39/6/MRN-463
- Li Xia - Lin, George E, Marschiner H (1991) Extension of the phosphorus depletion zone in VA – mycorrhizal white clover in calcareous soil. *Plant soil* 136(1): 41-48
- Madder P, Vierheilig H, Boller T, Streitwalf B, Engle Freg P, Chritie, Wietnken A (2000) Transport of 15N from a soil compartment separated by a polytetrafluoroethylene membrane to plant roots via the hyphae of arbuscular mycorrhizal fungi. *New Phytologist* 146: 155-161
- Madhusudhan L (2016) Organic Farming-Ecofriendly Agriculture. *J Ecosys Ecograph* 6: 209; doi: 10.4172/2157-7625.1000209
- Malik BS, Paul S, Ahlawat AK, Singh AM, Shivay YS (2009) Productivity and quality of wheat spp. grown with different fertilization condition. *Indian Journal of Agricultural Sciences* 79: 636-40
- Mardukhi B, Rejali F, Daei G, Ardakani Md R, Malakouti Md J, Miransari Md (2011) Arbuscular mycorrhizas enhance nutrient uptake in different wheat genotypes at high salinity levels under field and greenhouse conditions. *Comptes Rendus Biologies* 334(7): 564-571; doi: 10.1016/j.crv.2011.05.001
- Mishra RR, Verma NK (1982) Effect of different mycorrhizal treatments on the growth of onion. *Acta Botanica India* 11: 49-52
- Mittal S, Kaur G, Vishwakarma G (2013) Effects of Environmental Pesticides on the Health of Rural Communities in the Malwa Region of Punjab, India: A Review. *Human and Ecological Risk Assessment An International Journal* 20: 366-387; doi: 10.1080/10807039.2013.788972
- Mosse B (1973) Advances in the study of vesicular arbuscular mycorrhiza. *Ann Rev Phytopath* 11: 171-196
- Muthukumar T, Priyadharsini P, Uma E, Jaison S, Pandey RR (2014) Role of arbuscular mycorrhizal fungi in alleviation of acidity stress on plant growth. In: Miransari Md (ed). *Use of Microbes for the Alleviation of Soil Stresses*, pp. 43-72. Springer, New York
- Nadeem SM, Ahmad M, Zahir ZA, Javaid A, Ashraf M (2014) The role of mycorrhizae and plant growth promoting rhizobacteria (PGPR) in improving crop productivity under stressful environments. *Biotechnol Adv* 32: 429-448; doi: 10.1016/j.biotechadv.2013.12.005
- Nuccio EE, Hodge A, Pett-Ridge J, Herman DJ, Weber PK, Firestone MK (2013) An arbuscular mycorrhizal fungus significantly modifies the soil bacterial community and nitrogen cycling during litter decomposition. *Environmental Microbiology* 15: 1870-1881; doi: 10.1111/1462-2920.12081

- Oliveira RS, Vosatka M, Dodd JC, Castro PML (2005) Studies on the diversity of arbuscular mycorrhizal fungi and the efficacy of two native isolates in highly alkaline anthropogenic sediment. *Mycorrhiza* 16: 23-31
- Oyetunji OJ, Ekanayake IJ, Osonubi O (2003) The influence of arbuscular mycorrhizae fungus, mulch and fertilizer application on the yield of yams in an agroforestry system in south western Nigeria. *Maurik Bull* 6: 75-82
- Pandey J, Singh A (2012) Opportunities and constraints in organic farming: An Indian perspective. *Journal of Scientific Research Banaras Hindu University* 56: 47-72
- Paul S, Singh R, Tyagi M (2011) Interactive effect with AM fungi and *Azotobacter* inoculated seed on germination, plant growth and yield in cotton (*Gossypium hirsutum*). *Indian Journal of Agricultural Sciences* 81(11): 1041-55
- Pelczar MJ, Bard RC, Burnett GW, Conn HJ, Demoss RD, Euans EE, Weiss FA, Jennison MW, Meckee AP, Riker AJ, Warren J, Weeks OB (1957) *Manual of microbiological methods*. Society of American Bacteriology McGraw Hill Book Company Inc, New York
- Phillips JM, Hayman DS (1970) Improved procedure for clearing roots and staining parasitic and vesicular arbuscular mycorrhizal fungi for rapid assessment of infection. *Trans Br Mycol Soc* 55: 158-161
- Pikovskaya RI (1948) Mobilization in phosphate in soil in concentration with vital activities of some microbial species. *Microbiologia* 17: 362-370
- Ponmurugan P, Gopi C (2006) *In vitro* production of growth regulators and phosphate activity by phosphate solubilizing bacteria. *African J Biotechnol* 5: 348-350
- Püschel D, Bitterlich M, Rydlová J (2020) Facilitation of plant water uptake by an arbuscular mycorrhizal fungus: a Gordian knot of roots and hyphae. *Mycorrhiza* 30: 299-313; doi: 10.1007/s00572-020-00949-9
- Raklami A, Bechtaoui N, Tahiri AI, Anli M, Meddich A, Oufdou K, (2019) Use of Rhizobacteria and Mycorrhizae Consortium in the Open Field as a Strategy for Improving Crop Nutrition, Productivity and Soil Fertility. *Front Microbiol* 10: 1106; doi: 10.3389/fmicb.2019.01106
- Rahman KM, Debnath SC (2015) Agrochemical use, environmental and health hazards in Bangladesh. *International Research Journal of Interdisciplinary & Multidisciplinary Studies* 1: 75-79
- Reddy PS, Rao TVSS, Venkatramana P, Suryanarayaana N (2003) Response of mulberry varieties to VAM and *Azotobacter* biofertilizers inoculation. *Indian J Plant Physiol* 8(2): 171-174
- Redecker D, Morton JB, Bruns TD (2000) Ancestral lineages of arbuscular mycorrhizal fungi (Glomales). *Mol Phylogen Evol* 14: 276-284
- Robinson L, Feng W, Gulbis N, Hajdu K, Harrison RJ, Jeffries P, Xu X (2016) The Use of Arbuscular Mycorrhizal Fungi to Improve Strawberry Production in Coir Substrate. *Front Plant Sci* 7: 1237; doi: 10.3389/fpls.2016.01237
- Rouphael Y, Franken P, Schneider C, Schwarz D, Giovannetti M, Agnolucci M (2015) Arbuscular mycorrhizal fungi act as bio-stimulants in horticultural crops. *Sci Hort* 196: 91-108; doi: 10.1016/j.scienta.2015.09.002
- Samanta S, Verma NK (2006) Effect of VA mycorrhiza on the growth and protein content in fruits of *Capsicum annuum* grown in acid lateritic soil. *J Mycopathol Res* 44(2): 197-200
- Saxena J, Sharma V (2003) Phosphate solubilizing activity of microbes and their role as biofertilizer. In: Trivedi PC (ed). *Advances in Microbiology*, pp. 59-73. Scientific Publ, Jodhpur
- Schenck NC, Perez Y (1990) *Manual for the identification of VA mycorrhizal fungi (INVAM)* (3rd ed). University of Florida, Gainesville
- Schüßler A, Schwarziff D, Walker C (2001) A new fungal phylum, the Glomeromycota: Phylogeny and evolution. *Mycological Research* 105: 1413-1421
- Selvakumar G, Chandrasekaran M, Charlotte S, Kiyoon K, Tongmin S (2012) Spore associated bacteria (SAB) of arbuscular mycorrhizal fungi (amf) and plant growth promoting rhizobacteria (PGPR) increase nutrient uptake and plant growth under stress conditions. *Korean Journal of Soil Science and Fertilizer* 45(4): 582-592; doi: 10.7745/KJSSF.2012.45.4.582

- Selvakumar G, Thamizhiniyan P (2011) The effect of the arbuscular mycorrhizal (am) fungus *Glomus intraradices* on the growth and yield of chilli (*Capsicum annuum* L.) under salinity stress. *World Applied Sciences Journal* 14(8): 1209-1214
- Sengupta D, Verma NK, Ghosh BC (2006) Effect of vesicular arbuscular mycorrhiza and *Rhizobium* on field grown ground nut in acid lateritic soil. In: Prakash A, Mehrotra VS (Eds). *Mycorrhiza*, pp. 215-218. Scientific Publishers (India), Jodhpur
- Shaimaa AMd, Massoud ON (2017) Impact of Inoculation with Mycorrhiza and *Azotobacter* under Different N and P Rates on Growth, Nutrient status, Yield and Some Soil Characteristics of Washington Navel Orange Trees. *Middle East J Agric Res* 6(3): 617-638
- Shwetha C, Lakshman HC (2013) Effect of AM fungi, *Azotobacter* and phosphate solubilizing bacteria in improvement of *Amaranthus paniculatus* L. - A leafy vegetable. *Research Journal of Biotechnology* 8: 36-39
- Sieverding E (1991) Vesicular-arbuscular mycorrhizal management in tropical agro system. German Technical Co-operation (GZT), Eschborn
- Simanungkalit RDM (2006) Arbuscular Mycorrhiza Fungi. Available in: <http://www.balittanah.litbang.pertanian.go.id> Accessed 14/06/2019
- Singh CS, Rana JPS (2005) Arbuscular mycorrhizal fungi. In: Kaushik BD (ed). *Advances in microbiology at IARI 1961-2004*, pp. 123-34. Mouno publishing house, New Delhi
- Smith SE, Read DJ (1997) *Mycorrhizal Symbiosis*. Academic Press, London
- Smith SE, Read DJ (2008) *Mycorrhizal Symbiosis*, (3rd Edition). Academic Press, London
- Subba NS (1977) *Soil Microorganisms and Plant Growth*. Oxford and IBH Publishing Co, New Delhi
- Sundara WVB, Sinha MK (1963) Phosphate dissolving micro-organisms in the soil and rhizosphere. *Indian J Agric Sci* 33: 272-278
- Sylvia DM (1994) Vesicular-arbuscular mycorrhizal fungi. In: Weaver RW, Angle S, Bottomley P, Bezdicek D, Smith S, Tabatabai A, Wollum A (Eds). *Methods of soil Analysis, (Part 2) Microbiological and Biochemical properties*, pp. 351-378. Soil Science Society of America, Madison; ISBN: 9780891188650
- Vafadar F, Amooaghaie R, Otroshy Md (2014) Effects of plant-growth-promoting rhizobacteria and arbuscular mycorrhizal fungus on plant growth, stevioside, NPK, and chlorophyll content of *Stevia rebaudiana*. *Journal of Plant Interactions* 9(1): 128-136; doi: 10.1080/17429145.2013.779035
- van der Heijden MGA, Martin FM, Selosse MA, Sanders IR (2015) Mycorrhizal ecology and evolution: the past, the present, and the future. *New Phytologist* 205: 1406-1423
- Vyas M, Vyas A (2014) Field Response of *Capsicum annuum* Dually Inoculated with AM Fungi and PGPR in Western Rajasthan. *International Journal of Research Studies in Biosciences* 2(3): 21-26
- Weber E, George E, Beck DP, Saxena MC, Marschner H (1992) Vesicular arbuscular mycorrhizal and phosphorus uptake in chickpea grown in Northern Syria. *Expl Agric* 28: 433-442
- Yazdani M, Bahmanyar MA, Pirdashti H, Esmaili MA (2009) Effect of Phosphate solubilization microorganisms (PSM) and plant growth promoting rhizobacteria (PGPR) on yield and yield components of Corn (*Zea mays* L.). *Proc World Acad Science Eng Technol* 37: 90-92
- Zhang L, Fan JQ, Ding XD, He XH, Zhang FS, Feng G (2014) Hyphosphere interactions between an arbuscular mycorrhizal fungus and a phosphate solubilizing bacterium promote phytate mineralization in soil. *Soil Biol Biochem* 74: 177-183; doi: 10.1016/j.soilbio.2014.03.004

Recibido: 16-04-2020

Aceptado: 22-05-2020

This is an open access article distributed under a Creative Commons Attribution-NonCommercial 4.0 International (CC BY-NC 4.0) <https://creativecommons.org/licenses/by-nc/4.0/> The use, distribution or reproduction is permitted citing the original source and authors.