

Biogeosciences, 18, 3689–3700, 2021
<https://doi.org/10.5194/bg-18-3689-2021>

© Author(s) 2021. This work is distributed under the Creative Commons Attribution 4.0 License.



Reviews and syntheses: Heterotrophic fixation of inorganic carbon – significant but invisible flux in environmental carbon cycling

Alexander Braun¹, Marina Spona-Friedl¹, Maria Avramov¹, Martin Elsner^{1,2}, Federico Baltar³, Thomas Reinthaler³, Gerhard J. Herndl^{3,4}, and Christian Griebler^{1,3}

¹Institute of Groundwater Ecology, Helmholtz Zentrum München, Ingolstaedter Landstrasse 1, 85764 Neuherberg, Germany

²Department of Analytical Chemistry and Water Chemistry, Technical University of Munich, Munich, Germany

³Department of Functional and Evolutionary Ecology, University of Vienna, Althanstrasse 14, 1090 Vienna, Austria

⁴Department of Marine Microbiology and Biogeochemistry, Royal Netherlands Institute for Sea Research, Utrecht University, P.O. Box 59, 1790 AB Den Burg, the Netherlands

Correspondence: Christian Griebler (christian.griebler@univie.ac.at)

Received: 10 December 2020 – Discussion started: 17 December 2020

Revised: 3 May 2021 – Accepted: 17 May 2021 – Published: 21 June 2021

Abstract. Heterotrophic CO₂ fixation is a significant yet underappreciated CO₂ flux in environmental carbon cycling. In contrast to photosynthesis and chemolithoautotrophy – the main recognized autotrophic CO₂ fixation pathways – the importance of heterotrophic CO₂ fixation remains enigmatic. All heterotrophs – from microorganisms to humans – take up CO₂ and incorporate it into their biomass. Depending on the availability and quality of growth substrates, and drivers such as the CO₂ partial pressure, heterotrophic CO₂ fixation contributes at least 1 %–5 % and in the case of methanotrophs up to 50 % of the carbon biomass. Assuming a standing stock of global heterotrophic biomass of 47–85 Pg C, we roughly estimate that up to 5 Pg C might be derived from heterotrophic CO₂ fixation, and up to 12 Pg C yr^{−1} originating from heterotrophic CO₂ fixation is funneled into the global annual heterotrophic production of 34–245 Pg C yr^{−1}. These first estimates on the importance of heterotrophic fixation of inorganic carbon indicate that this pathway should be incorporated in present and future carbon cycling budgets.

ble for more than 60 % of the “enhanced greenhouse effect” resulting in global warming (Beer et al., 2010; Berg, 2011; Houghton, 2007; Le Quéré et al., 2016).

While photosynthesis and chemosynthesis are the most important processes of carbon fixation, non-autotrophic carbon fixation, i.e., the carbon fixation mediated by heterotrophic organisms, might also be relevant, albeit uncommonly quantified. While heterotrophs are, per definition, organisms that respire organic compounds to gain energy and build up biomass, CO₂ fixation plays also an essential role in heterotrophic carbon metabolism. The diversity of carboxylating enzymes in nature reaches far beyond autotrophy and virtually all heterotrophs harbor numerous enzymes fixing dissolved inorganic carbon. Even though the first carboxylase in heterotrophs was discovered more than 80 years ago (Wood and Werkman, 1936), the role of heterotrophs in carbon cycling has so far largely focused on the oxidation of organic substrates using oxygen or alternative electron acceptors (e.g. nitrate, ferric iron, sulfate) and the production of CO₂. Similar to the CO₂ fixation by autotrophs, “heterotrophic CO₂ fixation” might, however, constitute a significant carbon flux in specific habitats. The relevance of this process has hardly been quantified due to the lack of reliable estimates of heterotrophic CO₂ fixation for most organisms and habitats, and the presumption that CO₂ fixation in natural environments is restricted to autotrophic organisms.

To fill this gap, we review the current knowledge on (i) the role of heterotrophic CO₂ fixation for cellular

1 Introduction

Fixation of CO₂ is a fundamental biosynthetic process in nature (Beer et al., 2010; Berg et al., 2007), providing the main source of metabolic energy on Earth (Giovannoni and Stingl, 2005). At the same time, it acts as a sink for atmospheric CO₂, the most important greenhouse gas, which is responsi-

metabolism, (ii) respiration and non-autotrophic CO₂ fixation, and (iii) CO₂ fixation in habitats dominated by heterotrophs, and we provide (iv) quantitative estimates of heterotrophic CO₂ fixation in different environments.

2 Role of heterotrophic CO₂ fixation in cellular metabolism

The non-autotrophic uptake of inorganic carbon has been reported for a wide range of organisms from prokaryotes and fungi to vertebrates (Wood and Werkman, 1938; Kleiber et al., 1952; Cochrane, 1958; Hartman et al., 1972; Perez and Matin, 1982; Schinner et al., 1982; Parkinson et al., 1990; Roslev et al., 2004; Hesselsoe et al., 2005; Feisthauer et al., 2008; Spona-Friedl et al., 2020) and plants (Melzer and O’Leary, 1987). Currently, more than 20 carboxylases are known, forming an integral part of the central and peripheral metabolic pathways of heterotrophic metabolism (Fig. 1), e.g., in gluconeogenesis, the synthesis of fatty acids, amino acids, vitamins and nucleotides, the assimilation of leucine, and anaplerosis (Evans and Slotin, 1940; Krebs, 1941; Wood and Werkman, 1941; Werkman and Wood, 1942; Kornberg and Krebs, 1957; Wood and Stjernholm, 1962; Kornberg, 1965; Scrutton, 1971; Hartman and Keen, 1973; Dijkhuizen and Harder, 1984; Parkinson et al., 1991; Attwood, 1995; Han et al., 2000; Sauer and Eikmanns, 2005; Erb et al., 2009; Schink, 2009; Erb, 2011; Bar-Even et al., 2012). Carboxylation in heterotrophs not just compensates for the dependence on organic matter, but rather CO₂ fulfills the role of a “co-substrate”, providing an effective and simple way to extend an existing organic carbon substrate by a single C1 unit as part of the secondary production (Erb, 2011).

The most important CO₂ fixation pathway in all organisms is anaplerosis. Anaplerosis replenishes intermediates in the tricarboxylic acid (TCA) cycle, which have been released for biosynthesis. TCA metabolites are used as building blocks for macromolecular compounds, e.g. almost half of all amino acids in prokaryotes are directly synthesized from oxaloacetate and α -ketoglutarate (Fuchs, 1999). For this purpose, heterotrophs use the enzymes pyruvate carboxylase present in a large variety of organisms, including prokaryotes, archaea, yeasts, fungi, and higher organisms (e.g. mammals), and phosphoenol pyruvate (PEP) carboxylase, widely distributed in bacteria (Attwood, 1995; Jitrapakdee and Wallace, 1999; Sauer and Eikmanns, 2005; Jitrapakdee et al., 2008) (Fig. 1). The replenishment of metabolites continuously withdrawn from the TCA cycle via the anaplerotic reaction of PEP carboxylase entails an assimilation of CO₂ corresponding to 25 % of the initial substrate’s carbon content. In a systematic stable isotope labeling experiment with *Bacillus subtilis*, a gram-positive heterotrophic bacterium widespread in the environment, the interdependency of pathways and rates of CO₂ fixation on the concurrent utilization of organic substrate(s) was explored (Spona-Friedl et

al., 2020). Over the course of the experiments *B. subtilis* assimilated 6 % and 5 % of carbon biomass from the external H¹³CO₃ pool when growing on glucose and lactate, respectively (Spona-Friedl et al., 2020). Growth on malate, an intermediate of the TCA cycle, expected to serve directly to refill the oxaloacetate pool of the TCA cycle, still revealed a contribution to biomass production from inorganic carbon of 3 % (Spona-Friedl et al., 2020). In fact, PEP carboxylase was still actively transforming pyruvate to oxaloacetate. Heterotrophic CO₂ fixation continued to a lower extent even in the absence of cell growth during the stationary phase (Spona-Friedl et al., 2020), indicating that anaplerotic reactions are important in low-productivity habitats (see below).

Overall, heterotrophic CO₂ fixation via anaplerosis in microorganisms contributes around 1 % to 8 % to the carbon biomass (Romanenko, 1964; Perez and Matin, 1982; Doronina and Trotsenko, 1984; Miltner et al., 2004; Roslev et al., 2004; Hesselsoe et al., 2005; Šantrůčková et al., 2005; Feisthauer et al., 2008; Akinyede et al., 2020; Spona-Friedl et al., 2020). Under particular environmental conditions even higher contributions were reported (Perez and Matin, 1982). The advantage that CO₂ is readily available to the cell either as atmospheric gas or, more commonly, in its hydrated form HCO₃⁻ obviously outcompetes the disadvantage that carboxylation is generally an endergonic reaction (Faber et al., 2015). This thermodynamic obstacle may be less important when carboxylation supports the assimilation of organic substrates more reduced than the organism’s biomass, resulting in carbon-limited but excess-energy conditions (Heijnen and Roels, 1981; Ensign et al., 1998; von Stockar et al., 2006; Battley, 2013). In this case, in addition to anaplerosis further carboxylation reactions are induced (Fig. 1) to add oxidized C (from CO₂) to the reduced organic substrate for adjusting the degree of reduction to that of the biomass (Fig. 2). For example, the assimilation of leucine and propionate into biomass entails carboxylation of the initial C-6 and C-3 carbon bodies, respectively, and thus triggers an assimilation of dissolved inorganic carbon (DIC) that corresponds to 17 % and 33 % of the initial substrate’s carbon content, respectively (Erb, 2011). In aerobic methane oxidation, the full oxidation potential of one molecule of CO₂ is needed to adjust the high degree of reduction of methane to that of biomass during its assimilation. Consequently, methanotrophs derive up to 50 % of their carbon biomass from CO₂ (Strong et al., 2015; Battley, 2013).

Besides the degree of reduction of organic carbon sources, the partial pressure of CO₂ plays a role. Carboxylases may catalyze carboxylation as well as decarboxylation of organic compounds, and the equilibrium of the reaction depends on the concentrations of all compounds involved. An increase in the CO₂ concentration may move the equilibrium of the reaction toward the product of the carboxylation, and thus lead to an increase in CO₂ fixation.

In a physiological context, the amount of inorganic carbon fixed by heterotrophs, from either an endogenous or ex-




Figure 1. Selected heterotrophic CO_2 fixation reactions and pathways. PEP: phosphoenolpyruvate; DAPA: 7,8-diaminononanoate; AIR: 1-(5'-phosphoribosyl)-5-aminoimidazole; CAIR: 1-(5-phospho-D-ribose)-5-amino-4-imidazolecarboxylate; CoA: Coenzyme-A.

ogenous source, may be dependent on the metabolic state of the organisms and the specific environmental conditions. In their early work, Romanenko et al. (1972) suggested that the rate of heterotrophic anaplerotic fixation of DIC is strictly proportional to the heterotrophic bacterial carbon production. Since then, a number of factors have been identified potentially influencing the relative contribution of anaplerotic and other non-autotrophic CO_2 fixation reactions on biomass production. In laboratory experiments with the bacterial strain *Thiobacillus novellus*, for example, a higher amount of CO_2 was fixed under nutrient-limited conditions (Perez and Matin, 1982). Moreover, mixotrophic bacterial strains fixed more DIC compared to those grown autotrophically (Perez and Matin, 1982). Fungi fixed relatively more CO_2 at lower organic carbon (glucose and maltose) concentrations (Schinner et al., 1982). In conclusion, the degree of heterotrophic CO_2 fixation highly depended on the availability of easily degradable organic carbon sources (Schinner et al., 1982).

Studies on the possible relationship between heterotrophic DIC fixation and the activity of prokaryotic cells revealed

contradicting results. While Roslev et al. (2004) mentioned actively growing cells fix more DIC than resting cells, Merlin et al. (2003) report enhanced uptake of DIC by heterotrophic bacteria during slow growth and starvation. A relationship between DIC and heterotrophic bacterial production has been reported frequently as exemplified below.

3 Respiration and non-autotrophic CO_2 fixation

The production of CO_2 via respiration and the parallel fixation of CO_2 in heterotrophs take place simultaneously. The heterotrophic fixation of CO_2 is thus generally considered a back-reaction; i.e., part of the originally produced CO_2 from respiration is re-assimilated. Following this line of arguments, the more reduced an organic substrate is the less CO_2 is released (Fig. 2). The heterotrophic fixation of DIC does not necessarily lead to a net carbon biomass production; however, if microbes oxidize geogenic methane, this would result in a net carbon biomass production. Experimentally it is difficult to differentiate respiratory CO_2 flux from concurrent anaplerotic CO_2 fixation. As a consequence, there




Figure 2. Anaplerotic CO_2 fixation contributes 1%–8% of carbon biomass (indicated by the grey band) in heterotrophic cells. Dependent to the organism and in relation to the uptake of the individual organic compounds and their entry into the TCA cycle and central metabolic pathways the relative amount of inorganic carbon assimilated varies, as highlighted by the red arrows. See examples for malic and lactic acid. With organic carbon sources more reduced than the organism's biomass (dashed line) further carboxylation reactions are induced (indicated by black line), increasing the overall carbon contribution from CO_2 beyond anaplerosis (grey band). In methanotrophs, 50% of the cell's carbon may originate from CO_2 fixation. For further explanations, see text.

are numerous experiments and field studies determining dark CO_2 fixation, but only a few studies quantified the assimilation of DIC by non-autotrophs.

Respiration in aquatic systems is frequently determined via the consumption of dissolved oxygen (Robinson and Williams, 2005), potentially underestimating the carbon use efficiency of heterotrophs. Depending on the substrate, the respiration quotient ($\Delta\text{CO}_2 / -\Delta\text{O}_2$) varies between 0.7–1.3 (Robinson, 2019), leading to an error between 20% and 40% with regard to CO_2 production from respiration. Moreover, the respiration quotient also varies because other oxygen-consuming processes are potentially taking place simultaneously (e.g. nitrification) (Robinson, 2019). For instance, it is 138 O_2 for 106 CO_2 for ideal Redfield-type organic matter and 150 O_2 for 106 CO_2 for more realistic marine organic matter (Fraga et al., 1998; Paulmier et al., 2009). Calculations based on a study on temperate forest soils revealed a reduction of overall CO_2 emissions due to dark CO_2 fixation by mainly heterotrophic microbes (Akinyede et al., 2020). Collectively, with respect to C cycling, heterotrophic CO_2 fixation and the carbon flux from the inorganic pool into heterotrophic biomass can be regarded as a process more important than hitherto assumed.

4 CO_2 fixation in habitats dominated by heterotrophs

In contrast to sunlit habitats, where photoautotrophs make up a significant portion of the total biomass and photosynthesis is of major importance in carbon cycling, heterotrophs and chemolithoautotrophs represent the prevailing biota in the “dark habitats”, i.e., soils, subsurface environments, and the deep sea. These dark environments, although characterized by disproportionately lower biological activity, exceed their photic counterparts in both volume and biomass. In the oceans, the deep sea (below 200 m) exceeds the sunlit surface layer by a factor of 18 in volume and, remarkably, by a factor of 2 in biomass (Aristegui et al., 2009). Therefore, the so-called “dark CO_2 fixation” not only occurs in specific “hot spots” on the seafloor (hydrothermal vents, cold seeps, and mud volcanoes), or in anoxic waters, but also throughout the entire oxygenated “dark” water column (Reinthal et al., 2010; Yakimov et al., 2014). In limnic environments, the dark groundwater ecosystems outnumber surface waters 100-fold in terms of water volume (Danielopol et al., 2003), and similarly, soils are also, with the exception of their surface, exclusively dark habitats.

Yet, heterotrophic CO_2 fixation does not occur only in the dark environments since heterotrophs are also found in the photic zone. This is particularly relevant in the ocean because the photic zone is where the highest biomass concentrations are found. Recently, it has been estimated that the inclusion of dark CO_2 fixation (integrated over the euphotic layer, 0–150 m depth) would increase oceanic primary production estimates by 2.5%–22% (Baltar et al., 2019). A similar situation might be assumed for surface inland waters; however, global estimations are missing so far.

Dark DIC fixation has been reported for all types of ecosystems, including marine habitats (Wuchter et al., 2003; Middelburg, 2011; DeLorenzo et al., 2012; Molari et al., 2013; Baltar and Herndl, 2019; Lengger et al., 2019; Smith et al., 2019; Vasquez-Cardenas et al., 2020), brackish and freshwater systems (Bräuer et al., 2013; Santoro et al., 2013; Noguerola et al., 2015; Signori et al., 2017; Vick-Majors and Priscu, 2019; Zhao et al., 2020), cave waters and groundwater ecosystems (Pedersen and Ekendahl, 1992a, b; Kotelnikova and Pedersen, 1998; Kellermann et al., 2012; Lazar et al., 2017), and soil habitats (Ehleringer et al., 2000; Miltner et al., 2004, 2005; Šantrůčková et al., 2005, 2018; Akinyede et al., 2020, and references therein). In the absence of solar radiation, particularly in the dark ocean, CO_2 fixation rates of up to $\sim 125 \text{ mg C m}^{-3} \text{ d}^{-1}$ have been measured, amounting to 30% (on a per volume basis) of the phototrophic CO_2 fixation in ocean surface waters (Zopfi et al., 2001; Detmer et al., 1993; Casamayor et al., 2001; Baltar et al., 2010). In a eutrophic lagoon, dark DIC fixation accounted for 31% of total DIC fixation in the water column (Lliros et al., 2011). Recently it was shown that the ratio between dark / light CO_2 fixation in oceanic surface waters which is usually around 0.1 increases with depth, reaching a ratio of

1 at 120–160 m depth (Baltar et al., 2019). In the past, however, dark DIC fixation has frequently been attributed to the activity of chemoautotrophs only. Only a few studies so far have provided strong quantitative evidence for heterotrophic CO₂ fixation in aquatic and terrestrial ecosystems (Table 1).

As indicated, part of the dark CO₂ fixation in oceans has been attributed to chemolithoautotrophic archaea (Wuchter et al., 2003; Ingalls et al., 2006), obtaining the energy required for the endergonic carboxylation through the oxidation of reduced inorganic compounds, such as ammonia or hydrogen sulfide (Swan et al., 2011; Zhang et al., 2020). A total annual chemolithoautotrophic CO₂ fixation rate of 0.77 Pg C was calculated for the oceans (Middelburg, 2011). The observed fluxes of the reduced inorganic compounds available as energy sources, however, seem largely insufficient to explain the relatively high dark CO₂ fixation rates (Overbeck, 1979; Tuttle and Jannasch, 1979; Baltar et al., 2010; Reinthaler et al., 2010; Herndl and Reinthaler, 2013). In some cases, the supply rates of the reduced inorganic compounds used as an energy source explain less than 40 % of the observed dark CO₂ fixation rates (Zopfi et al., 2001). Recently, chemoautotrophic nitrification was estimated to explain < 13 % of the dark CO₂ fixation (integrated over the euphotic zone), with the rest coming from either heterotrophic DIC fixation or other chemoautotrophic processes (Baltar and Herndl, 2019).

The potential energy sources for the unexplained proportion of the dark CO₂ fixation remain enigmatic. Possible explanations could be either an underestimation of the supply rates of reduced inorganic compounds or the uptake of CO₂ by heterotrophic organisms (Zopfi et al., 2001; Baltar et al., 2019). In the surface ocean in particular, DIC incorporation via anaplerotic reactions might play an important role in compensating for metabolic imbalances in marine bacteria under oligotrophic conditions, contributing > 30 % of the carbon incorporated into biomass (González et al., 2008; Palovaara et al., 2014). Evidence for the latter comes from experiments with Arctic seawater, which exhibited high DIC fixation rates (0.5–2.5 µg C L⁻¹ d⁻¹) correlating with heterotrophic bacterial production (Alonso-Sáez et al., 2010). Using different molecular tools, DIC uptake was attributed mainly to heterotrophic *Gamma*- and *Betaproteobacteria* rather than to typical chemoautotrophs, thus showing that chemolithoautotrophs were not the main drivers of CO₂ fixation in this habitat (Alonso-Sáez et al., 2010). Further evidence comes from the genome of *Polaribacter* sp. MED152, a representative of Bacteroidetes, which typically comprise about 10 %–20 % of the prokaryotic abundance in seawater (González et al., 2008). A unique combination of membrane transporters and carboxylases in these organisms indicates the importance of anaplerosis in addition to other DIC fixation pathways (González et al., 2008). If the heterotrophic metabolism of bacteria is suddenly intensified (e.g., after an input of organic matter), dark DIC fixation rates and the expression of transcripts associated with key anaplerotic enzymes increase proportionally (Baltar et al., 2016). As men-

tioned above, contradicting results were obtained on the relationship between heterotrophic CO₂ fixation and the availability of organic matter. A few studies suggest a relative increase in dark DIC fixation in oligotrophic habitats harboring slow-growing or starving bacterial populations (Perez and Matin, 1982; Schinner et al., 1982; Merlin et al., 2003; Alonso-Sáez et al., 2010; Santoro et al., 2013). Considering the slow community-wide specific growth rates of heterotrophic bacteria in oligotrophic and/or cold waters, such as the marine aphotic zone, the Arctic Ocean, deep sea sediments, groundwater systems and the terrestrial subsurface, alpine limnic systems, and deep-lake sediments, enhanced anaplerotic DIC uptake can be expected. However, there is also evidence for the stimulation of dark DIC fixation in response to organic matter enrichment in different types of soils (Miltner et al., 2005; Šantrůčková et al., 2018). Hence, these contradictory findings require further, more systematic research.

Other environmental factors that may influence dark DIC fixation include the concentrations of CO₂ and bicarbonate as inorganic carbon sources. An increase in the CO₂ concentration may shift the equilibrium of the carboxylation–decarboxylation reactions increasing CO₂ fixation. Elevated partial pressure of CO₂ might stimulate dark DIC fixation. In temperate forest soils, rates of dark microbial CO₂ fixation were positive correlated with the CO₂ concentration (Spohn et al., 2019). Similarly, with increasing CO₂ concentrations, higher dark DIC fixation was observed in wetland soils affected by subcrustal CO₂ degassing (Beulig et al., 2015). Here, besides known chemoautotrophs, CO₂ fixation via anaplerotic reactions was shown for putatively heterotrophs, i.e., subdivision 1 Acidobacteriaceae, lacking enzymatic pathways for autotrophic CO₂ fixation (Beulig et al., 2015). In experiments with two marine heterotrophic bacterial isolates, elevation of CO₂ concentration provoked an increase in CO₂ fixation along with a decrease in respiration (Teiro et al., 2012). Thus, we may assume that a rise in CO₂ concentrations and CO₂-induced geochemical changes will alter carbon turnover in affected ecosystems, with dark DIC fixation and anaplerotic reactions becoming more important.

5 Quantitative estimates of heterotrophic CO₂ fixation in different environments

5.1 Quantification of heterotrophic DIC fixation

It is difficult to properly quantify heterotrophic fixation of inorganic carbon in the environment. Not surprisingly, quantitative data almost exclusively originate from laboratory experiments using cultures and tissues in combination with carbon isotopic labeling (e.g. Spona-Friedl et al., 2020). Field studies generally report on dark carbon fixation, including the activity of chemoautotrophs and heterotrophs. So far, evidence for the significant contribution of heterotrophic CO₂

Table 1. Dissolved inorganic carbon (DIC) assimilation rates from a range of aquatic (marine and limnic) and soil environments. Dark carbon fixation (DCF) is shown as fraction of either bacterial heterotrophic production (BP) or respiration. Original data were converted to similar units whenever possible to allow comparison.

Aquatic ecosystems	Depth [m]	DIC fixation [$\mu\text{g C L}^{-1} \text{d}^{-1}$]	BP [$\mu\text{g C}^{-1} \text{d}^{-1}$]	DCF/BP [%]	Source	Remarks
Arctic	Seawater cultures	0.5–2.3	0.4–2.5	100 %	Alonso-Saez et al. (2010)	Only potential for DCF
Mediterranean Sea	4900	0.096 ± 0.02	0.048	200 %	Yakimov et al. (2014)	Only anaerobic
Tropical South China Sea	200–1500	0.72–1.68	0.48–4.8	40 %–105 %	Zhou et al. (2017)	Probably a large fraction anaerobic
Tropical estuary	1–18	4.8–14.4	55.2–1142	1.3 %–9 %	Signori et al. (2017)	Probably mostly anaerobic
Eutrophic lagoon	1–5	206			Llirios et al. (2011)	Probably mostly anaerobic
Boreal lake sediments	1–3	13.2–48 $\text{mg C m}^{-2} \text{d}^{-1}$	BP 96–216 $\text{mg C m}^{-2} \text{d}^{-1}$	8.4 %–37.4 %	Santoro et al. (2013)	Probably a large fraction anaerobic
Tropical lake sediments	1–3	0.12–20.4 $\text{mg C m}^{-2} \text{d}^{-1}$	BP 14.4–583 $\text{mg C m}^{-2} \text{d}^{-1}$	0.4 %–80.4 %	Santoro et al. (2013)	Probably a large fraction anaerobic
Deep granitic ground-water biofilms	812–1240	0.2–2 $\mu\text{g C m}^{-2} \text{d}^{-1}$	n.d.	n.d.	Ekendahl and Pedersen (1994)	Probably a large fraction anaerobic
Terrestrial ecosystems		DIC fixation [$\mu\text{g C g}^{-1} \text{d}^{-1}$]	R [$\mu\text{g CO}_2\text{-C g}^{-1} \text{d}^{-1}$]	DCF/R [%]		
Temperate forest soil	0–0.7	0.036–0.32	0.95–19.1	1.2 %–3.9 %	Spohn et al. (2019)	^{13}C label mainly in AA, indicating anaerobiosis
Temperate agricultural soil	0–1	0.06–0.86	n.d.	n.d.	Akinyede et al. (2020)	Dominance of heterotrophs
Temperate agricultural soil	0–0.3	0.26	0.63	2.7 %	Milner et al. (2004)	Probably a large fraction anaerobic
Temperate agricultural soil	0–0.3	0.19	9.82	1 %–5 %	Milner et al. (2005)	DCF mainly driven by aerobic heterotrophs
Range of temperate forest & field soils	0.05–0.15	1.82–23.6*	0.65–9.16	3 %–39 %	Šantrůčková et al. (2005)	Probably a large fraction anaerobic
Arctic tundra soils	0–0.15	0.035–0.4	n.d.	n.d.	Nel and Cramer (2019)	Probably mostly anaerobic
Arctic tundra soils		0.04–0.08	0.79–10.7	0.04 %–16 %	Šantrůčková et al. (2018)	Anaerobic enzymes comprised the majority of carboxylase genes

* Values taken from Table 2 in Akinyede et al. (2020), n.d. – not determined.

fixation, as highlighted for selected studies in Table 1, is based on additional measures complementing the quantification of dark carbon fixation, i.e. molecular microbial community analysis occasionally including the quantitative assessment of functional genes involved in carbon fixation and characterization of environmental conditions. In fact, significant dark carbon fixation (i) in the obvious absence of chemoautotrophs and related genes or (ii) in well-oxygenated environments that lack potential electron donors (e.g. H₂, H₂S, NH₄) led authors to conclude on the quantitative importance of heterotrophic carbon fixation (e.g. Miltner et al., 2005; Alonso-Saéz et al., 2010; Yakimov et al., 2014; Šantrůčková et al., 2018; Akinyede et al., 2020). Individual studies succeeded to follow the label of inorganic carbon in environmental samples into amino acids of microorganisms (Spohn et al., 2019; Spona-Friedl et al., 2020). In the future, the combined application of metabolomics and isotope tracing may help further develop this field of research.

5.2 Heterotrophic CO₂ fixation in different habitats

Measurements of dark DIC fixation with strong evidence of a significant contribution of heterotrophic assimilation of DIC are scarce. In Table 1, we provide a compilation of studies conducted in soils and marine and limnic ecosystems. Where possible, we compared dark DIC fixation rates with heterotrophic activity. In marine and limnic systems, heterotrophic carbon production as a widely applied activity measurement was used. In soils, we compared dark DIC fixation rates with respiration, i.e., CO₂ production. Dark DIC fixation rates in different marine systems range between 0.1 and 206 µg C L⁻¹ d⁻¹, with the highest values found in a eutrophic lagoon and the lowest values in the deep waters of the Mediterranean Sea (Table 1). Data from limnic systems originate from lake sediments with dark DIC fixation rates between 0.12 and 48 mg C m⁻² d⁻¹ (Table 1). Projecting these numbers to only the top 10 cm of sediment in the different lakes studied (which is a gross simplification), values of 1.2–480 µg C L⁻¹ sediment per day are obtained. When compared to rates of bacterial carbon production, dark DIC fixation rates in these habitats accounted for a considerable fraction of total carbon assimilation, occasionally even exceeding it (Table 1). In soils, the dark DIC fixation rates, which were attributed mainly to the activity of heterotrophs, amounted to 0.04 %–39 % of the overall respiration rate (Table 1). Dark DIC fixation rates range from 36 ng C to 23.6 µg C g⁻¹ d⁻¹ covering 3 orders of magnitude (Table 1). The contribution of heterotrophically fixed DIC to biomass carbon of microbes ranged from 0.2 %–1.1 % in temperate forest soil (Akinyede et al., 2020), 0.2 %–4.6 % in temperate forest and field soils (Šantrůčková et al., 2005), to 7 % in arable soil (Miltner et al., 2004). Šantrůčková et al. (2005) estimated the overall heterotrophic CO₂ fixation to be even higher, i.e., 1.9 %–11.3 % taking into account that the labile fraction of the biodegradable organic carbon re-

sulted from metabolites released by spilling reactions of microorganisms due to a limitation in inorganic nutrients or due to the presence of highly reduced energy-rich carbon sources (e.g. Tempest and Neijssel, 1992). A contribution of heterotrophic CO₂ fixation to biomass carbon of 6.5 ± 2.8 % was found in drinking water biofilms and activated sludge (Roslev et al., 2004).

5.3 Carbon biomass stock originating from heterotrophic CO₂ fixation

While it is difficult to derive global estimations from the few studies that measured heterotrophic CO₂ fixation rates in marine, limnic, and terrestrial ecosystems, we may use a conservative approach assuming that at least 1 %–5 % of carbon biomass of all heterotrophs originates from anaerobic DIC fixation. Earth's total living biomass is estimated to amount to about 499–738 Pg C, of which approx. 451–653 Pg C is photoautotrophic biomass (Bar-On et al., 2018). Heterotrophic biomass thus contributes 47–85 Pg C (Table S1 in the Supplement). The uncertainties of the estimates of heterotrophic biomass of the terrestrial subsurface, however, are high (Whitman et al., 1998; McMahon and Parnell, 2014; Bar-On et al., 2018). Nevertheless, following this line of evidence, anaerobic CO₂ fixation contributes between 0.5–5 Pg C to the living biomass.

5.4 Carbon flux related to heterotrophic CO₂ fixation

In terms of annual global heterotrophic production rates, oceans and the terrestrial subsurface (including soils) are the main habitats of heterotrophic CO₂ fixation (Cole et al., 1988; Magnabosco et al., 2018) (Table S2). Recently, Akinyede et al. (2020) estimated a global dark CO₂ fixation rate of all temperate forest soils of 0.26 ± 0.07 Pg C yr⁻¹. We calculated a global heterotrophic C production of 34–245 Pg C yr⁻¹, which would translate into 0.34–12.3 Pg of DIC bound by heterotrophic CO₂ fixation each year. Interestingly, these numbers are consistent with the recently calculated contribution of CO₂ fixation for the integrated epipelagic ocean of ca. 1.2–11 Pg C yr⁻¹ (Baltar and Herndl 2019). This is a significant carbon flux amounting to 0.3 %–14 % of the global net amount of carbon produced annually by photoautotrophs (90–110 Pg C yr⁻¹; Ciais et al., 2013).

Our estimates, as said, are subject to a high uncertainty, which, on the one hand, results from the dependency of the extent of heterotrophic CO₂ fixation on the organic carbon oxidized and, on the other hand, on the predominant environmental conditions. Moreover, data on terrestrial and marine subsurface environments, although large in dimension, are scarce. For these environments, no detailed information on the abundance, growth (yield), and metabolic activity of microbial communities is available, particularly with increasing depth. Most of the deeper subsurface environments, even when harboring considerable living biomass, do not

participate in the global carbon cycle on short and medium timescales (years to decades), but rather on centennial to geological timescales. Nevertheless, in order to provide a first estimate and to be able to roughly evaluate the relevance of heterotrophic CO₂ fixation for all habitats, we adopted this conservative approach (see also Tables S1 and S2).

6 Conclusions

Current models of carbon cycling and carbon sequestration (Gruber et al., 2004; Le Quéré et al., 2009) do not account for heterotrophic CO₂ fixation. Despite the uncertainties in the data on heterotrophic biomass and production rates for some habitats (e.g. the terrestrial subsurface), the numbers presented here represent the first attempt to quantify the global contribution and relevance of heterotrophic CO₂ fixation to carbon cycling. Our results indicate that heterotrophs significantly contribute to global CO₂ fixation – especially (although not restricted to) in habitats experiencing elevated CO₂ concentrations and/or lacking a sufficient supply of degradable organic carbon. In specific environments, this may explain the mismatch between autotrophic C input, consumption, and sequestration as has been observed in marine systems (Baltar and Herndl, 2019; Burd et al., 2010; Reinthaler et al., 2010; Morán et al., 2007; Hoppe et al., 2002; Tait and Schiel, 2013). Particularly in aphotic habitats (which outnumber the photic habitats in both size and volume) such as the dark ocean, seafloor sediments, soils, and the sediments and rocks of the terrestrial subsurface (Miltner et al., 2004, 2005; Yakimov et al., 2014; Wegener et al., 2012), carbon cycling needs to be re-evaluated taking into account anaerobic CO₂ fixation and other inorganic carbon uptake pathways in heterotrophs. In seafloor sediments, wetlands, and marshes, as well as in other habitats where methane oxidation is a key process, a large fraction (10 %–50 %) of heterotrophic biomass potentially originates from heterotrophic DIC fixation. Recently, a time-series study showed a tendency towards higher ratios of dark to light DIC fixation in the top half of the euphotic layer (0–65 m) in the years 2012–2019 than in the preceding years (data started in 1989), which was linked to oceanographic changes (i.e., a deepening of the mixed zone) (Baltar et al., 2019). Moreover, the metabolic theory of ecology posits that heterotrophic metabolism increases more than gross primary production in the ocean in response to warming (see Baltar et al., 2019, and reference therein), which might also make heterotrophic DIC fixation relatively more important in a warmer ocean. In light of global warming leading to an extensive thawing of permafrost soils and providing new habitats for methanotrophs, these processes are expected to become more important in the future. Hence, the potential contribution of heterotrophic CO₂ fixation under climate change conditions clearly deserves further investigation.

Data availability. All data that have been used for our calculations are available in Tables S1 and S2 in the Supplement. No further data sets were used in this article.

Supplement. The supplement related to this article is available online at: <https://doi.org/10.5194/bg-18-3689-2021-supplement>.

Author contributions. AB, ME, and CG conceived the idea for the manuscript. AB, GJH, and CG wrote the manuscript. MSF, ME, MA, FB, and TR substantially commented on and edited the manuscript. MA, MSF, and CG did the literature search on available global carbon data. CG and MA performed the estimation of heterotrophic CO₂ fixation on a global scale.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. We acknowledge Bo Barker Jørgensen for commenting on an earlier draft of the manuscript. We thank Rudolf K. Thauer and Wolfgang Eisenreich for fruitful discussions on heterotrophic CO₂ fixation.

Financial support. Financial support was provided by the Wittgenstein Prize (Austrian Science Fund, project number Z194-B17), by the European Research Council under the European Community's Seventh Framework Program (FP7/2007–2013)/ERC grant agreement no. 268595 (MEDEA project), and the Austrian Science Fund (P 28781-B21) to Gerhard J. Herndl. Financial support was further provided by the Helmholtz Center Munich to Alexander Braun, Martin Elsner, Marina Spona-Friedl, and Christian Griebler.

Review statement. This paper was edited by Steven Bouillon and reviewed by two anonymous referees.

References

- Akinyede, R., Taubert, M., Schrumpf, M., Trumbore, S., and Küsel, K.: Rates of dark CO₂ fixation are driven by microbial biomass in a temperate forest soil, *Soil Biol. Biochem.*, 150, 107950, <https://doi.org/10.1016/j.soilbio.2020.107950>, 2020.
- Alonso-Sáez, L., Galand, P. E., Casamayor, E. O., Pedrós-Alió, C., and Bertilsson, S.: High bicarbonate assimilation in the dark by Arctic bacteria, *ISME J.*, 4, 1581–1590, 2010.
- Arístegui, J., Gasol, J. M., Duarte, C. M., and Herndl, G. J.: Microbial oceanography of the dark ocean's pelagic realm, *Limnol. Oceanogr.*, 54, 1501–1529, 2009.
- Attwood, P. V.: The structure and the mechanism of action of pyruvate-carboxylase, *Int. J. Biochem. Cell B.*, 27, 231–249, 1995.
- Baltar, F. and Herndl, G. J.: Ideas and perspectives: Is dark carbon fixation relevant for oceanic primary production estimates?,

- Biogeosciences, 16, 3793–3799, <https://doi.org/10.5194/bg-16-3793-2019>, 2019.
- Baltar, F., Arístegui, J., Sintes, E., Gasol, J. M., Reinthaler, T., and Herndl, G. J.: Significance of non-sinking particulate organic carbon and dark CO₂ fixation to heterotrophic carbon demand in the mesopelagic northeast Atlantic, *Geophys. Res. Lett.*, 37, 1–6, 2010.
- Baltar, F., Lundin, D., Palovaara, J., Lekunberri, I., Reinthaler, T., Herndl, G. J., and Pinhassi, J.: Prokaryotic responses to ammonium and organic carbon reveal alternative CO₂ fixation pathways and importance of alkaline phosphatase in the mesopelagic North Atlantic, *Front. Microbiol.*, 7, 1670, <https://doi.org/10.3389/fmicb.2016.01670>, 2016.
- Baltar, F., Bayer, B., Bednarsek, N., Deppeler, S., Escribano, R., Gonzalez, C. E., Hansman, R. L., Mishra, R. K., Moran, M. A., Repeta, D. J., Robinson, C., Sintes, E., Tamburini, C., Valentin, L. E., and Herndl, G. J.: Towards integrating evolution, metabolism, and climate change studies of marine ecosystems, *Trends Ecol. Evol.*, 34, 1022–1033, 2019.
- Bar-Even, A., Noor, E., and Milo, R.: A survey of carbon fixation pathways through a quantitative lens, *J. Exp. Bot.*, 63, 2325–2342, 2012.
- Bar-On, Y. M., Phillips, R., and Milo, R.: The biomass distribution on Earth, *P. Natl. Acad. Sci. USA*, 115, 6506–6511, 2018.
- Battley, E. H.: A theoretical study of the thermodynamics of microbial growth using *Saccharomyces cerevisiae* and a different free energy equation, *Q. Rev. Biol.*, 88, 69–96, 2013.
- Beer, C., Reichstein, M., Tomelleri, E., Ciais, P., Jung, M., Carvalhais, N., Rödenbeck, C., Arain, M. A., Baldocchi, D., Bonan, G. B., Bondeau, A., Cescatti, A., Lasslop, G., Lindroth, A., Lomas, M., Luysaert, S., Margolis, H., Oleson, K. W., Rouspard, O., Veenendaal, E., Viovy, N., Williams, C., Woodward, F. I., and Papale, D.: Terrestrial gross carbon dioxide uptake: global distribution and covariation with climate, *Science*, 329, 834–838, 2010.
- Berg, I. A.: Ecological aspects of the distribution of different autotrophic CO₂ fixation pathways, *Appl. Environ. Microbiol.*, 77, 1925–1936, 2011.
- Berg, I. A., Kockelkorn, D., Buckel, W., and Fuchs, G.: A 3-hydroxypropionate/4-hydroxybutyrate autotrophic carbon dioxide assimilation pathway in Archaea, *Science*, 318, 1782–1786, 2007.
- Beulig, F., Heuer, V. B., Akob, D. M., Viehweger, B., Elvert, M., Herrmann, M., Hinrichs, K.-U., and Küsel, K.: Carbon flow from volcanic CO₂ into soil microbial communities of a wetland mofette, *ISME J.*, 9, 746–759, 2015.
- Bräuer, S. L., Kranzler, K., Goodson, N., Murphy, D., Simon, H. M., Baptista, A. M., and Tebo, B. M.: Dark carbon fixation in the Columbia River's estuarine turbidity maxima: molecular characterization of red-type *cbbl* genes and measurement of DIC uptake rates in response to added electron donors, *Estuar. Coast. Shelf S.*, 36, 1073–1083, 2013.
- Burd, A. B., Hansell, D. A., Steinberg, D. K., Anderson, T. R., Arístegui, J., Baltar, F., Beaupre, S. R., Buesseler, K. O., DeHairs, F., Jackson, G. A., Kadko, D. C., Koppelman, R., Lampitt, R. S., Nagata, T., Reinthaler, T., Robinson, C., Robinson, B. H., Tamburini, C., and Tanaka, T.: Assessing the apparent imbalance between geochemical and biochemical indicators of meso- and bathypelagic biological activity: What the @!\$# is wrong with present calculations of carbon budgets?, *Deep-Sea Res. Pt. II*, 57, 1557–1571, 2010.
- Casamayor, E. O., García-Cantizano, J., Mas, J., and Pedrós-Alió, C.: Primary production in estuarine oxic/anoxic interfaces: contribution of microbial dark CO₂ fixation in the Ebro River Salt Wedge Estuary, *Mar. Ecol.-Prog. Ser.*, 215, 49–56, 2001.
- Ciais, P., Sabine, C., Bala, G., Bopp, L., Brovkin, V., Canadell, J., Chhabra, A., DeFries, R., Galloway, J., Heimann, M., Jones, C., Le Quéré, C., Myneni, R. B., Piao, S., and Thornton, P.: Carbon and other biogeochemical cycles, in: *Climate change 2013: The physical science basis. Contribution of working group I to the fifth assessment report of the Intergovernmental Panel on Climate Change*, edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., 465–570, Cambridge, United Kingdom and New York, NY, USA, Cambridge University Press, 2013.
- Cochrane, V. W.: *Physiology of fungi*, John Wiley, New York, 1958.
- Cole, J. J., Findlay, S. E. G., and Pace, M. L.: Bacterial production in fresh and saltwater ecosystems: a cross-system overview, *Mar. Ecol.-Prog. Ser.*, 43, 1–10, 1988.
- Danielopol, D. L., Griebler, C., Gunatilaka, A., and Notenboom, J.: Present state and future prospects for groundwater ecosystems, *Environ. Conserv.*, 30, 104–130, 2003.
- DeLorenzo, S., Bräuer, S. L., Edgmont, C. A., Herfort, L., Tebo, B. M., and Zuber, P.: Ubiquitous dissolved inorganic carbon assimilation by marine bacteria in the Pacific Northwest Coastal Ocean as determined by stable isotope probing, *PLoS ONE*, 7, e46695, <https://doi.org/10.1371/journal.pone.0046695>, 2012.
- Detmer, A. E., Giesenhausen, H. C., Trenkel, V. M., Auf dem Venne, H., and Jochem, F. J.: Phototrophic and heterotrophic pico- and nanoplankton in anoxic depths of the central Baltic Sea, *Mar. Ecol.-Prog. Ser.*, 99, 197–203, 1993.
- Dijkhuizen, L. and Harder, W.: Current views on the regulation of autotrophic carbon dioxide fixation via the Calvin cycle in bacteria, *Antonie van Leeuwenhoek*, 50, 473–487, 1984.
- Doronina, N. V. and Trotsenko, Y. A.: The levels of carbon dioxide assimilation in bacteria with different pathways of 1-carbon metabolism, *Mikrobiologiya*, 53, 885–889, 1984.
- Ehleringer, J. R., Buchmann, N., and Flanagan, L. B.: Carbon isotope ratios in belowground carbon cycle processes, *Ecol. Appl.*, 10, 412–422, 2000.
- Ekendahl, S. and Pedersen, K.: Carbon transformations by attached bacterial populations in granitic ground water from deep crystalline bed-rock of the Stripa research mine, *Microbiology*, 140, 1565–1573, 1994.
- Ensign, S. A., Small, F. J., Allen, J. R., and Sluis, M. K.: New roles for CO₂ in the metabolism of aliphatic epoxides and ketones, *Arch. Microbiol.*, 169, 179–187, 1998.
- Erb, T. J.: Carboxylases in natural and synthetic microbial pathways, *Appl. Environ. Microb.*, 77, 8466–8477, 2011.
- Erb, T. J., Brecht, V., Fuchs, G., Müller, M., and Alber, B. E.: Carboxylation mechanism and stereochemistry of crotonyl-CoA carboxylase/reductase, a carboxylating enoyl-thioester reductase, *P. Natl. Acad. Sci. USA*, 106, 8871–8876, 2009.
- Evans Jr., E. A. and Slotin, L.: The utilization of carbon dioxide in the synthesis of α -ketoglutaric acid, *J. Biol. Chem.*, 136, 301–302, 1940.

- Faber, K., Fessner, W. D., and Turner, N. J.: Science of synthesis: biocatalysis in organic synthesis Vol. 2, 672 pp., Georg Thieme, Stuttgart, 2015.
- Feisthauer, S., Wick, L. Y., Kastner, M., Kaschabek, S. R., Schloemann, M., and Richnow, H. H.: Differences of heterotrophic ^{13}C CO₂ assimilation by *Pseudomonas knackmussii* strain B13 and *Rhodococcus opacus* 1CP and potential impact on biomarker stable isotope probing, *Environ. Microbiol.*, 10, 1641–1651, 2008.
- Fraga, F., Rios, A., Perez, F., and Figueras, F.: Theoretical limits of oxygen:carbon and oxygen : nitrogen ratios during photosynthesis and mineralisation of organic matter in the sea, *Mar. Chem.*, 62, 161–168, 1998.
- Fuchs, G.: Biosynthesis of building blocks, in: *Biology of the prokaryotes*, edited by: Lengeler, J. W., Drews, G., and Schlegel, H. G., 110–160, Thieme, Stuttgart, New York, 1999.
- Giovannoni, S. J. and Stingl, U.: Molecular diversity and ecology of microbial plankton, *Nature*, 437, 343–348, <https://doi.org/10.1038/nature04158>, 2005.
- González, J. M., Fernández-Gómez, B., Fernández-Guerra, A., Gómez-Consarnau, L., Sánchez, O., Coll-Lladó, M., del Campo, J., Escudero, L., Rodríguez-Martínez, R., Alonso-Sáez, L., Latasa, M., Paulsen, I., Nedashkovskaya, O., Lekumberri, I., Pinhassi, J., and Pedrós-Alió, C.: Genome analysis of the proteorhodopsin-containing marine bacterium *Polaribacter* sp. MED152 (Flavobacteria), *P. Natl. Acad. Sci. USA*, 105, 8724–8729, 2008.
- Gruber, N., Friedlingstein, P., Field, C., Valentini, R., Heimann, M., Richey, J. E., Romero-Lankao, P., Schulze, E. D., and Chen, C.-T. A.: The vulnerability of the carbon cycle in the 21st century: an assessment of carbon-climate-human interactions, in: *The global carbon cycle: integrating humans, climate, and the natural world*, edited by: Field, C. B. and Raupach, M. R., 45–76, Island Press, Washington D.C., London, 2004.
- Han, L., Yang, K., Kulowski, K., Wendt-Plienkowski, E., Hutchinson, C. R., and Vining, L. C.: An acyl-coenzyme A carboxylase encoding gene associated with jadomycin biosynthesis in *Streptomyces venezuelae* ISP5230, *Microbiol. UK*, 146, 903–910, 2000.
- Hartman, R. E. and Keen, N. T.: Enzymes catalysing anaplerotic carbon dioxide fixation in *Verticillium albo-atrum*, *Phytopathol.*, 63, 947–953, 1973.
- Hartman, R. E., Keen, N. T., and Long, M.: Carbon dioxide fixation by *Verticillium albo-atrum*, *J. Gen. Microbiol.*, 73, 29–34, 1972.
- Heijnen, J. J. and Roels, J. A.: A macroscopic model describing yield and maintenance relationship in aerobic fermentation processes, *Biotechnol. Bioeng.*, 23, 739–763, 1981.
- Herdnd, G. J. and Reinthaler, T.: Microbial control of the dark end of the biological pump, *Nat. Geosci.*, 6, 718–724, 2013.
- Hesselsoe, M., Nielsen, J. L., Roslev, P., and Nielsen, P. H.: Isotope labeling and microautoradiography of active heterotrophic bacteria on the basis of assimilation of ^{14}C CO₂, *Appl. Environ. Microb.*, 71, 646–655, 2005.
- Hoppe, H. G., Gocke, K., Koppe, R., and Begler, C.: Bacterial growth and primary production along a north-south transect of the Atlantic Ocean, *Nature*, 416, 168–171, 2002.
- Houghton, R. A.: Balancing the global carbon budget, *Annu. Rev. Earth Planet. Sc.*, 35, 313–347, 2007.
- Ingalls, A. E., Shah, S. R., Hansman, R. L., Aluwihare, L. I., Santos, G. M., Druffel, E. R., and Pearson, A.: Quantifying archaeal community autotrophy in the mesopelagic ocean using natural radiocarbon, *P. Natl. Acad. Sci. USA*, 103, 6442–6447, 2006.
- Jitrapakdee, S. and Wallace, J. C.: Structure, function and regulation of pyruvate carboxylase, *Biochem. J.*, 340, 1–16, 1999.
- Jitrapakdee, S., St. Maurice, M., Rayment, I., Cleland, W. W., Wallace, J. C., and Attwood, P. V.: Structure, mechanism and regulation of pyruvate carboxylase, *Biochem. J.*, 413, 369–387, 2008.
- Kellermann, C., Selesi, D., Lee, N., Hügler, M., Esperschütz, J., Hartmann, A., and Griebler, C.: Microbial CO₂ fixation potential in a tar-oil-contaminated porous aquifer, *FEMS Microbiol. Ecol.*, 81, 172–187, 2012.
- Kleiber, M., Smith, A. H., and Black, A. L.: Carbonate as precursor of milk constituents in the intact dairy cow, *J. Biol. Chem.*, 195, 707–714, 1952.
- Kornberg, H. L.: Anaplerotic sequences in microbial metabolism, *Angew. Chem. Int. Edit.*, 4, 558–565, 1965.
- Kornberg, H. L. and Krebs, E. H.: Synthesis of cell constituents from C₂-units by a modified tricarboxylic acid cycle, *Nature*, 179, 988–991, 1957.
- Kotelnikova, S. and Pedersen, K.: Distribution and activity of methanogens and homoacetogens in deep granitic aquifers at Äspö Hard Rock Laboratory, Sweden, *FEMS Microbiol. Ecol.*, 26, 121–134, 1998.
- Krebs, H. A.: Carbon dioxide assimilation in heterotrophic organisms, *Nature*, 147, 560–563, 1941.
- Lazar, C. S., Stoll, W., Lehmann, R., Herrmann, M., Schwab, V. F., Akob, D. M., Nawaz, A., Wubet, T., Buscot, F., Totsche, K.-U., and Küsel, K.: Archaeal diversity and CO₂ fixers in carbonate-/siliciclastic-rock groundwater ecosystems, *Archaea*, 2136287, 1–13, 2017.
- Le Quéré, C., Raupach, M. R., Canadell, J. G., Marland, G., Bopp, L., Ciais, P., Conway, T. J., Doney, S. C., Feely, R. A., Foster, P., Friedlingstein, P., Gurney, K., Houghton, R. A., House, J. I., Huntingford, C., Levy, P. E., Lomas, M. R., Majkut, J., Metzl, N., Ometto, J. P., Peters, G. P., Prentice, I. C., Randerson, J. T., Running, S. W., Sarmiento, J. L., Schuster, U., Sitch, S., Takahashi, T., Viovy, N., van der Werf, G. R., and Woodward, F. I.: Trends in the sources and sinks of carbon dioxide, *Nat. Geosci.*, 2, 831–836, 2009.
- Le Quéré, C., Andrew, R. M., Canadell, J. G., Sitch, S., Korsbakken, J. I., Peters, G. P., Manning, A. C., Boden, T. A., Tans, P. P., Houghton, R. A., Keeling, R. F., Alin, S., Andrews, O. D., Anthoni, P., Barbero, L., Bopp, L., Chevallier, F., Chini, L. P., Ciais, P., Currie, K., Delire, C., Doney, S. C., Friedlingstein, P., Gkritzalis, T., Harris, I., Hauck, J., Haverd, V., Hoppema, M., Klein Goldewijk, K., Jain, A. K., Kato, E., Körtzinger, A., Landschützer, P., Lefèvre, N., Lenton, A., Lienert, S., Lombardozzi, D., Melton, J. R., Metzl, N., Millero, F., Monteiro, P. M. S., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S., O'Brien, K., Olsen, A., Omar, A. M., Ono, T., Pierrot, D., Poulter, B., Rödenbeck, C., Salisbury, J., Schuster, U., Schwinger, J., Séférian, R., Skjelvan, I., Stocker, B. D., Sutton, A. J., Takahashi, T., Tian, H., Tilbrook, B., van der Laan-Luijkx, I. T., van der Werf, G. R., Viovy, N., Walker, A. P., Wiltshire, A. J., and Zaehle, S.: Global Carbon Budget 2016, *Earth Syst. Sci. Data*, 8, 605–649, <https://doi.org/10.5194/essd-8-605-2016>, 2016.
- Lengger, S. K., Rush, D., Maysner, J. P., Blewett, J., Schwartz-Narbonne, R., Talbot, H. B., Middelburg, J. J., Jetten, M. S. M., Schouten, S., Sinninghe Damsté, J. S., and Pancost, R. D.:

- Dark carbon fixation in the Arabian Sea oxygen minimum zone contributes to sedimentary organic carbon (SOM), *Global Biogeochem. Cy.*, 33, 1715–1732, 2019.
- Lliros, M., Alonso-Saéz, L., Gich, F., Plasencia, A., Auguet, O., Casamayor, E. O., and Borrego, C. M.: Active bacteria and archaea cells fixing bicarbonate in the dark along the water column of a stratified eutrophic lagoon, *FEMS Microbiol. Ecol.*, 77, 370–384, 2011.
- Magnabosco, C., Lin, L. H., Dong, H., Bomberg, M., Ghiorse, W., Stan-Lotter, H., Pedersen, K., Kieft, T. L., van Heerden, E., and Onstott, T. C.: The biomass and biodiversity of the continental subsurface, *Nat. Geosci.*, 11, 707–717, 2018.
- McMahon, S. and Parnell, J.: Weighing the deep continental biosphere, *FEMS Microbiol. Ecol.*, 87, 113–120, 2014.
- Melzer, E. and O’Leary M. H.: Anapleurotic CO₂ fixation by phosphoenolpyruvate carboxylase in C₃ plants, *Plant Physiol.*, 84, 58–60, 1987.
- Merlin, C., Masters, M., McAteer, S., and Coulson, A.: Why is carbonic anhydrase essential to *Escherichia coli*?, *J. Bacteriol.*, 185, 6415–6424, 2003.
- Middelburg, J. J.: Chemoautotrophy in the ocean, *Geophys. Res. Lett.*, 38, 1–4, 2011.
- Miltner, A., Richnow, H.-H., Kopinke, F.-D., and Kästner, M.: Assimilation of CO₂ by soil microorganisms and transformation into soil organic matter, *Org. Geochem.*, 35, 1015–1024, 2004.
- Miltner, A., Kopinke, F.-D., Kindler, R., Selesi, D., Hartmann, A., and Kästner, M.: Non-photosynthetic CO₂ fixation by soil microorganisms, *Plant Soil*, 269, 193–203, 2005.
- Molari, M., Manini, E., and Dell’Anno, A.: Dark inorganic carbon fixation sustains the functioning of benthic deep-sea ecosystems, *Global Biogeochem. Cy.*, 27, 212–221, 2013.
- Morán, X. A. G., Pérez, V., and Fernández, E.: Mismatch between community respiration and the contribution of heterotrophic bacteria in the NE Atlantic open ocean: What causes high respiration in oligotrophic waters?, *J. Mar. Res.*, 65, 545–560, 2007.
- Nel, J. A. and Cramer, M. D.: Soil microbial anapleurotic CO₂ fixation in temperate soils, *Geoderma*, 335, 170–178, 2019.
- Noguerola, I., Picazo, A., Lliros, M., Camacho, A., and Borrego, C. M.: Diversity of freshwater Epsilonproteobacteria and dark inorganic carbon fixation in the sulphidic redoxcline of a meromictic karstic lake, *FEMS Microbiol. Ecol.*, 91, fiv086, <https://doi.org/10.1093/femsec/fiv086>, 2015.
- Overbeck, J.: Dark CO₂ uptake – biochemical background and its relevance to in situ bacterial production, *Arch. Hydrobiol. Beiheft*, 12, 38–47, 1979.
- Palovaara, J., Akram, N., Baltar, F., Bunse, C., Forsberg, J., Pedrós-Alió, C., González, J. M., and Pinhassi, J.: Stimulation of growth by proteorhodopsin phototrophy involves regulation of central metabolic pathways in marine planktonic bacteria, *P. Natl. Acad. Sci. USA*, 111, E3650–E3658, 2014.
- Parkinson, S. M., Killham, K., and Wainwright, M.: Assimilation of ¹⁴CO₂ by *Fusarium oxysporum* grown under oligotrophic conditions, *Mycol. Res.*, 94, 959–964, 1990.
- Parkinson, S. M., Jones, R., Meharg, A. A., Wainwright, M., and Killham, K.: The quantity and fate of carbon assimilated from ¹⁴CO₂ by *Fusarium oxysporum* grown under oligotrophic and near oligotrophic conditions, *Mycol. Res.*, 95, 1345–1349, 1991.
- Paulmier, A., Kriest, I., and Oschlies, A.: Stoichiometries of remineralisation and denitrification in global biogeochemical ocean models, *Biogeosciences*, 6, 923–935, <https://doi.org/10.5194/bg-6-923-2009>, 2009.
- Pedersen, K. and Ekendahl, S.: Assimilation of CO₂ and introduced organic compounds by bacterial communities in groundwater from southeastern Sweden deep crystalline bedrock, *Microb. Ecol.*, 23, 1–14, 1992a.
- Pedersen, K. and Ekendahl, S.: Incorporation of CO₂ and introduced organic compounds by bacterial populations in groundwater from deep crystalline bedrock of Stripa mine, *J. Gen. Microbiol.*, 138, 369–376, 1992b.
- Perez, R. C. and Matin, A.: Carbon dioxide assimilation by *Thiobacillus novellus* under nutrient-limited mixotrophic conditions, *J. Bacteriol.*, 150, 46–51, 1982.
- Reinthal, T., Van Aken, H. M., and Herndl, G. J.: Major contribution of autotrophy to microbial carbon cycling in the deep North Atlantic, Åô interior, *Deep-Sea Res. Pt. II*, 57, 1572–1580, 2010.
- Robinson, C.: Microbial respiration, the engine of ocean deoxygenation, *Front. Mar. Sci.*, 5, 533, <https://doi.org/10.3389/fmars.2018.00533>, 2019.
- Robinson, C. and Williams, P. J.: Respiration and its measurement in surface marine waters, in: *Respiration in aquatic ecosystems*, edited by: del Giorgio, P. A. and Williams, P. J., Oxford University Press, Oxford, 2005.
- Romanenko, V. I.: Heterotrophic CO₂ assimilation by water bacterial flora, *Mikrobiologiya*, 33, 679–683, 1964.
- Romanenko, V. I., Overbeck, J., and Sorokin, Y. I.: Estimation of production of heterotrophic bacteria using ¹⁴C, in: *Techniques for the assessment of microbial production and decomposition in fresh waters*, edited by: Sorokin, Y. I. and Kadota, H., IBP Handbook No. 23, Blackwell, Oxford, 82–85, 1972.
- Roslev, P., Larsen, M. B., Jørgensen, D., and Hesselsoe, M.: Use of heterotrophic CO₂ assimilation as a measure of metabolic activity in planktonic and sessile bacteria, *J. Microbiol. Meth.*, 59, 381–393, 2004.
- Santoro, A. L., Bastviken, D., Gudas, C., Tranvik, L., and Enrich-Prast, A.: Dark carbon fixation: an important process in lake sediments, *PLoS ONE*, 8, e65813, <https://doi.org/10.1371/journal.pone.0065813>, 2013.
- Šantrůčková, H., Bird, M. I., Elhottova, D., Novak, J., Pícek, T., Simek, M., and Tykva, R.: Heterotrophic fixation of CO₂ in soil, *Microb. Ecol.*, 49, 218–225, 2005.
- Šantrůčková, H., Kotas, P., Bárta, J., Ulrich, T., Čapek P., Palmtag J., Eloy Alves, R. J., Biasi, C., Diáková, K., Gentsch, N., Gittel, A., Guggenberger, G., Hugelius, G., Lashchinsky, N., Martikainen, P. J., Mikutta, R., Schleper, C., Schneckner, J., Schwab, C., Shibistova, O., Wild, B., and Richter, A.: Significance of dark CO₂ fixation in arctic soils, *Soil Biol. Biochem.*, 119, 11–21, 2018.
- Sauer, U. and Eikmanns, B. J.: The PEP–pyruvate–oxaloacetate node as the switch point for carbon flux distribution in bacteria, *FEMS Microbiol. Rev.*, 29, 765–794, 2005.
- Schink, B.: An alternative to the glyoxylate shunt, *Mol. Microbiol.*, 73, 975–977, 2009.
- Schinner, F., Concina, R., and Binder, H.: Heterotrophic CO₂ fixation by fungi in dependence on the concentration of the carbon source, *Phyton*, 22, 81–85, 1982.
- Scrutton, M. C.: Assay of enzymes of CO₂ metabolism, *Method. Microbiol.*, 6, Part A, 479–541, 1971.

- Signori, C. N., Valentin, J. L., Pollery, R. C. G., and Enrich-Prast, A.: Temporal variability of dark carbon fixation and bacterial production and their relation with environmental factors in a tropical estuarine system, *Estuar. Coast.*, 41, 1089–1101, 2017.
- Smith, A. R., Kieft, B., Mueller, R., Fisk, M. R., Mason, O. U., Popa, R., and Colwell, F. S.: Carbon fixation and energy metabolisms of a seafloor olivine biofilm, *ISME J.*, 13, 1737–1749, 2019.
- Spohn, M., Müller, K., Höschel, C., Mueller, C. W., and Marhan, S.: Dark microbial CO₂ fixation in temperate forest soils increases with CO₂ concentrations, *Glob. Change Biol.*, 26, 1926–1935, 2019.
- Spona-Friedl, M., Braun, A., Huber, C., Eisenreich, W., Griebler, C., Kappler, A., and Elsner, M.: Substrate-dependent CO₂-fixation in heterotrophic bacteria revealed by stable isotope labelling, *FEMS Microbiol. Ecol.*, 96, fiae080, <https://doi.org/10.1093/femsec/fiae080>, 2020.
- Strong, P. J., Xie, S., and Clarke, W. P. Methane as a resource: can the methanotrophs add value?, *Environ. Sci. Technol.*, 49, 4001–4018, 2015.
- Swan, B. K., Martinez-Garcia, M., Preston, C. M., Sczyrba, A., Woyke, T., Lamy, D., Reinthaler, T., Poulton, N. J., Masland, E. D. P., Lluesma Gomez, M., Sieracki, M. E., DeLong, E. F., Herndl, G. J., and Stepanauskas, R.: Potential for chemolithoautotrophy among ubiquitous bacteria lineages in the dark ocean, *Science*, 333, 1296–1300, 2011.
- Tait, L. W. and Schiel, D. R.: Impacts of temperature on primary productivity and respiration in naturally structured macroalgal assemblages, *PLoS ONE*, 8, e74413, <https://doi.org/10.1371/journal.pone.0074413>, 2013.
- Teiro, E., Fernández, A., Álvarez-Salgado, X. A., García-Martín, E. E., Serret, P., and Sobrino, C.: Response of two marine bacterial isolates to high CO₂ concentration, *Mar. Ecol.-Prog. Ser.*, 453, 27–36, 2012.
- Tempest, D. W. and Neijssel, O. M.: Physiological and energetic aspects of bacterial metabolite overproduction, *FEMS Microbiol. Lett.*, 100, 169–176, 1992.
- Tuttle, J. H. and Jannasch, H. W.: Microbial dark assimilation of CO₂ in the Cariaco Trench, *Limnol. Oceanogr.*, 24, 746–753, 1979.
- Vasquez-Cardenas, D., Meysman, F. J. R., and Boschker, H. T. S.: A cross-system comparison of dark carbon fixation in coastal sediments, *Global Biogeochem. Cy.*, 34, 1–14, 2020.
- Vick-Majors, T. J. and Priscu, J. C.: Inorganic carbon fixation in ice-covered lakes of the McMurdo Dry Valleys, *Antarctic Sci.*, 1–10, <https://doi.org/10.1017/S0954102019000075>, 2019.
- von Stockar, U., Maskow, T., Liu, J., Marison, I. W., and Patiño, R.: Thermodynamics of microbial growth and metabolism: An analysis of the current situation, *J. Biotechnol.*, 121, 517–533, 2006.
- Wegener, G., Bausch, M., Holler, T., Thang, N. M., Mollar, X. P., Kellermann, M. Y., Hinrichs, K. U., and Boetius, A.: Assessing sub-seafloor microbial activity by combined stable isotope probing with deuterated water and ¹³C-bicarbonate, *Environ. Microbiol.*, 14, 1517–1527, 2012.
- Werkman, C. H. and Wood, H. G.: Heterotrophic assimilation of carbon dioxide, in: *Advances in Enzymology and Related Areas of Molecular Biology*, edited by: Nord, F. F., 2, 135–182, Interscience Publishers, Inc., 1942.
- Whitman, W. B., Coleman, D. C., and Wiebe, W. J.: Prokaryotes: The unseen majority, *P. Natl. Acad. Sci.*, 95, 6578–6583, 1998.
- Wood, H. G. and Stjernholm, R. L.: Assimilation of carbon dioxide by heterotrophic organisms, in: *The Bacteria: A Treatise on Structure and Function*, Vol. 3, edited by: Gunsalus, I. C. and Stanier, R. Y., Biosynthesis Academic Press, New York, 41–117, 1962.
- Wood, H. G. and Werkman, C. H.: The utilisation of CO₂ in the dissimilation of glycerol by the propionic acid bacteria, *Biochem. J.*, 30, 48–53, 1936.
- Wood, H. G. and Werkman, C. H.: The utilization of CO₂ by the propionic acid bacteria, *Biochem. J.*, 32, 1262–1271, 1938.
- Wood, H. G. and Werkman, C. H.: The position of carbon dioxide-carbon in succinic acid synthesized by heterotrophic bacteria, *J. Biol. Chem.*, 139, 377–381, 1941.
- Wuchter, C., Schouten, S., Boschker, H. T. S., and Sinninghe Damsté, J. S.: Bicarbonate uptake by marine Crenarchaeota, *FEMS Microbiol. Lett.*, 219, 203–207, 2003.
- Yakimov, M. M., La Cono, V., Smedile, F., Crisafi, F., Arcadi, E., Leonardi, M., Decembrini, F., Catalfamo, M., Bargiela, R., Ferrer, M., Golyshin, P. N., and Giuliano, L.: Heterotrophic bicarbonate assimilation is the main process of de novo organic carbon synthesis in hadal zone of the Hellenic Trench, the deepest part of Mediterranean Sea, *Environ. Microbiol. Rep.*, 6, 709–722, 2014.
- Zhang, Y., Qin, W., Hou, L., Zakem, E. J., Wan, X., Zhao, Z., Liu, L., Hunt, K. A., Jiao, N., Kao, S.-J., Tang, K., Xie, X., Shen, J., Li, Y., Chen, M., Dai, X., Liu, C., Deng, W., Dai, M., Ingalls, A. E., Stahl, D. A., and Herndl, G. J.: Nitrifier adaptation to low energy flux controls inventory of reduced nitrogen in the dark ocean, *P. Natl. Acad. Sci. USA*, 117, 4823–4830, 2020.
- Zhao, Y., Liu, P., Rui, J., Cheng, L., Wang, Q., Liu, X., and Yuan, Q.: Dark carbon fixation and chemolithotrophic microbial community in surface sediments of the cascade reservoirs, Southwest China, *Sci. Total Environ.*, 698, 134316, <https://doi.org/10.1016/j.scitotenv.2019.134316>, 2020.
- Zhou, W., Liao, J., Guo, Y., Yuan, X., Huang, H., Yuan, T., and Liu, S.: High dark carbon fixation in the tropical South China Sea, *Cont. Shelf Res.*, 146, 82–88, 2017.
- Zopfi, J., Ferdelman, T. G., Jørgensen, B. B., Teske, A., and Thamdrup, B.: Influence of water column dynamics on sulfide oxidation and other major biogeochemical process in the chemocline of Mariager Fjord (Denmark), *Mar. Chem.*, 74, 29–51, 2001.