


RESEARCH ARTICLE

Haematological and biochemical changes associated with male rats deprived of feed and water

B.S. Okediran*, S.A. Amid, F. Sanusi and K.O. Oladesu



Highlights

- Feed and water deprivation in rats for 6 days led to haemato-biochemical alterations.
- RBC, PCV, WBC, HbC and MCV increased significantly when compared to the control.
- Neutrophils, eosinophils, monocytes and lymphocytes counts rose more than normal.
- MCH, MCHC, glucose, triglyceride and cholesterol dropped below basal parameters.
- TP, albumin, creatinine, BUN, AST and ALT levels were increased above normal values.

RESEARCH ARTICLE

Haematological and biochemical changes associated with male rats deprived of feed and water

B.S. Okediran^{1*}, S.A. Amid², F. Sanusi¹ and K.O. Oladesu¹

¹Department of Veterinary Physiology and Biochemistry, Faculty of Veterinary Medicine, University of Ilorin, Ilorin, Nigeria.

²Department of Veterinary Surgery and Radiology, Faculty of Veterinary Medicine, University of Ilorin, Ilorin, Nigeria.

Received: 20/06/2020; Accepted: 10/01/2021

Abstract: Overnight food and water restriction protocols are common in animal research, however information about the haematological and biochemical changes due to food and water deprivation over longer periods can provide information about the stress experience by these animals. This study was conducted using ten (n = 10) male albino rats housed in metallic cage. The rats were deprived of feed and water consecutively for six days. On the third and sixth days blood samples were obtained for haematological studies while serum was used for biochemical studies. There were significant increases in red cell counts, packed cell volume, haemoglobin concentration and mean corpuscular volume while significant decreases were observed in the mean corpuscular haemoglobin and mean corpuscular haemoglobin concentration as compared to the basal parameters. The white blood cells, neutrophils, eosinophils, monocytes and lymphocytes were significantly increased as compared with the basal parameters. Activities of aspartate and alanine aminotransferase in post feed and water deprivation condition are elevated compared to basal parameters. There were significant decreases in serum glucose, triglyceride, cholesterol concentrations while the total protein, albumin, creatinine, blood urea were significantly elevated. It can be concluded that rats deprived of feed and water predisposes to starvation, dehydration and stress with perturbation of haematological and biochemical parameters.

Keywords: biochemical; deprivation; feed; haematology; water.

INTRODUCTION


Feed and water deprivation is a common procedure performed in association with many different types of experiments mainly in order to reduce variability in investigatory parameters or to facilitate surgical procedures. An animal satisfies its needs for food and water, not just because it experiences an acute feeling of thirst or hunger, but also because it has learned to realize its needs and knows how to meet them within the provisions of the environment in which it lives (Tucci *et al.*, 2006). If the absence of food or water is not dramatic or long lasting, the temporary effects this has on the animal's metabolism or hydration status will have little effect on the animal's welfare. The critical factor is whether the animal learns to change its behaviour to adapt to the availability of food and water over time (Toth and Gardiner, 2000). This situation

is common in the wild and also exists in laboratory experiments when food is rationed. These adjustments are often well tolerated as long as they are within limits to which the animal can adjust. A totally different situation arises if the animal is subjected to sudden, unexpected and chronic food deprivation (Bauer *et al.*, 2004).

Feed and water deprivation leads to some changes in one's eating habits, the amount of received energy and physiological parameters in the blood, however the body continually makes innumerable large and small adjustments to internal and external changes in its environment, to offset changes that threaten to create an abnormal state (Kanizsai *et al.*, 2004). Many of these adjustments are easily visible and relatively dramatic, but still well within what may be defined as 'normal'. The regulation of cellular nutrition during and between meals is an example of these adaptations. As long as the animal is given sufficient food as often as it is accustomed to, these changes are physiological (Stricker, 1984). If however, the supply of food deteriorates significantly (in quality or quantity), the body will no longer be able to regulate its functions within normal limits and pathological conditions will arise. An animal will notice changes in its supply of food rapidly, and this may lead to psychological stress. The degree of stress, in relation to feed and water deprivation, will depend upon the species and also for animals in captivity (Hughes *et al.*, 1994).

Fasting can be expected to cause, among other things, stress, aggressive behaviour, as well as reductions in body weight, body temperature and plasma glucose levels. A reduction in basal metabolic rate, which persists after the period of fasting, has been reported (Penicaud and Le Magnen, 1980). The effect of food deprivation is greatest when it occurs in the dark phase, when the animals are most active, significant reductions in liver weight and glycogen content, as well as increases in levels of glycerol, free fatty acids and acetoacetate have been measured after three hours (Palou *et al.*, 1981). Claassen (1994) reported increases in plasma levels of glucose, urea, lactate and amino acids, a range of hormones including insulin, glucagon and corticosterone, as well as increased activity in the sympathetic nervous system.

*Corresponding Author's Email: okediranbabatunde@gmail.com

 <https://orcid.org/0000-0002-5523-2699>



Food and water deprivation has been shown to cause some changes in hematological and biochemical parameters in blood, however its effect on biochemical parameters is still a matter of debate, it was reported that energy intake decreases during fasting (Sweileh *et al.*, 1992). The commonest haematological and biochemical data available were rats or animals subjected to overnight or maximum of five to six hour feed and water deprivation in which varying physiological changes were observed. This study is to provide a more complete picture of haematological and biochemical changes associated with male rats totally deprived of food and water for a consecutive period of six days.

MATERIALS AND METHODS

Animal ethics

All experimental protocols carried out on the animals were in accordance with the international accepted principles for laboratory animal use and were approved by the Ethics Committee (UIL/FVERC/001/2020) on Laboratory animal use of the Faculty of Veterinary Medicine, University of Ilorin, Nigeria.

Experimental animals

A total of ten ($n = 10$) male albino rats were used for this investigation. The average weight of the rats was 152 ± 3.50 g. These rats were housed in metallic cage under standard room temperature ($20.0\text{ }^{\circ}\text{C}$) and pressure (760 mmHg). They were provided with laboratory animal feed (Fat/oil 6%, crude fibre 5%, calcium 1%, Available phosphorus 0.4%, Lysine 0.85%, Methionine 0.35%, Salt 0.3%, Crude protein 18%, Metabolizable Energy 2900 Kcal/kg manufactured by TOPFEEDS® (Lagos, Nigeria) and water provided. Experimental animals were acclimatized to their environment before the start of the experiment.

Haematological and biochemical parameters

Basal haematological and biochemical parameters were determined before the animals were deprived of feed and water. The animals were deprived of feed and water for six consecutive days. Blood samples were collected on the third and sixth day for haematological and biochemical analyses.

Haematology

Aliquot of blood sample was collected into EDTA bottle for determination of packed cell volume, red blood cell and white blood cell counts as described by Schalm *et al.*, (1975).

Serum preparation

Aliquots of blood samples were collected into plain tubes and centrifuged at 4000 rpm for 10 minutes to separate the sera from the cellular components. The sera were then decanted and stored in Eppendorf tubes for further analyses as described by Schalm *et al.*, (1975).

Biochemical analysis

Blood glucose concentrations were checked by caudal

vein blood sampling and using Bionime glucometer. Total serum protein and albumin concentrations were estimated using Randox® commercial biochemical kits (Randox®, Spain) as described by Tietz (1995) and Grant (1987) respectively. Blood urea nitrogen and creatinine concentrations were estimated using Randox® commercial biochemical kits as described by Henry (1974). The aspartate and alanine aminotransferases activities were assayed as described by Reitman and Frankel (1957), while serum triglyceride and cholesterol concentrations were determined as described by McGowan *et al.*, (1983) and Wybenga *et al.*, (1970) respectively.

Statistical analysis

Results were expressed as mean \pm standard error of mean. Analysis of the data was done using one-way analysis of variance followed by the Duncan multiple range test. A $p < 0.05$ was considered significant. All analyses were done using Graph Pad Prism Version 5.

RESULTS

Table 1 shows the red blood cells and its indices in male rats deprived of feed and water for six consecutive days. There were significant increases ($p < 0.05$) in red cell counts, packed cell volume and haemoglobin concentration compared to the basal parameters. However there was significant increase in mean corpuscular volume, but significant decreases were observed in the mean corpuscular haemoglobin and mean corpuscular haemoglobin concentration compared to the basal parameters.

Table 2 shows the total and differential leucocyte counts white cells counts of rats deprived of feed and water for six consecutive days. Total and differential leucocyte counts are increased.

Table 3 shows the biochemical parameters of male rats deprived of feed and water for six consecutive days. There was a marked decrease in serum glucose concentration especially on the sixth day post feed and water deprivation compared to the basal parameter. However, the lipid parameters, triglyceride and cholesterol were significantly ($p < 0.05$) decreased compared to the basal parameters.

The total protein and albumin are mildly but significantly increased compared to the basal parameters, while creatinine and blood urea showed a mild increase post feed and water deprivation compared to the basal parameters.

There was a significant ($p < 0.05$) increase in the activities of aspartate and alanine aminotransferases post feed and water deprivation compared to the basal parameters.

Table 1: Red blood cells and it indices in male rats deprived of feed and water for six consecutive days

Parameters	Basal	Three days postfeed and water deprivation	Six days post feed and water deprivation
RBC (x 10 ¹² L)	7.38 ± 0.30 ^a	9.31 ± 0.11 ^b	10.20 ± 0.21 ^b
PCV (%)	41.15 ± 0.85 ^a	46.10 ± 0.62 ^b	48.22 ± 0.51 ^b
Hb (g/dl)	14.95 ± 0.15 ^a	16.01 ± 0.11 ^b	18.15 ± 0.12 ^b
MCV (fL)	51.85 ± 2.05 ^a	53.04 ± 1.06 ^b	54.11 ± 1.21 ^b
MCH (Pg)	18.25 ± 1.25 ^a	16.11 ± 1.20 ^b	13.15 ± 1.21 ^c
MCHC (g/dl)	38.25 ± 1.45 ^a	36.12 ± 1.33 ^b	34.12 ± 1.27 ^b

Values within the same rows with different superscripts are significantly different at $p < 0.05$. RBC: Red blood cell; PCV: Packed cell volume; Hb: Haemoglobin; MCV: Mean corpuscular volume; MCH: Mean corpuscular haemoglobin; MCHC: Mean corpuscular haemoglobin concentration.

Table 2: White blood and differentials in male rats deprived of feed and water for six consecutive days

Parameters	Basal	Three days post feed and water deprivation	Six days post feed and water deprivation
TWBC (x 10 ⁹ /L)	7.35 ± 2.11 ^a	12.22 ± 1.07 ^b	15.02 ± 2.11 ^b
Neutrophil (%)	6.11 ± 2.01 ^a	9.37 ± 1.03 ^b	11.14 ± 1.20 ^b
Eosinophils (%)	1.22 ± 0.50 ^a	3.06 ± 1.99 ^b	5.09 ± 1.05 ^b
Monocytes (%)	5.03 ± 1.01 ^a	7.96 ± 1.23 ^b	8.05 ± 2.31 ^b
Lymphocytes (%)	66.50 ± 4.50 ^a	72.75 ± 3.11 ^b	87.08 ± 3.54 ^b

Values within the same rows with different superscripts are significantly different at $p < 0.05$. TWBC: Total white blood cell. ($p < 0.05$) over time compared to the basal parameters.

Table 3: biochemical parameters of male rats deprived of feed and water for six consecutive days

Parameters	Basal	Three day days post feed and water deprivation	Six days post feed and water deprivation
Glucose (mg/dL)	105.33 ± 5.77 ^a	98.34 ± 4.01 ^b	90.03 ± 3.03 ^b
Triglyceride (mg/dL)	67.09 ± 1.54 ^a	61.32 ± 1.23 ^b	57.04 ± 1.03 ^b
Cholesterol (mg/dL)	120.20 ± 0.80 ^a	115.12 ± 0.71 ^b	109.65 ± 0.63 ^b
Total Protein (mg/dL)	6.80 ± 0.10 ^a	9.60 ± 0.16 ^b	12.12 ± 0.32 ^b
Albumin (mg/dL)	4.14 ± 0.04 ^a	4.29 ± 0.14 ^b	4.98 ± 0.13 ^b
Creatinine (mg/dL)	1.53 ± 0.01 ^a	1.65 ± 0.12 ^b	1.68 ± 0.11 ^b
Blood Urea Nitrogen (mg/dL)	3.25 ± 0.11 ^a	5.58 ± 0.35 ^b	8.75 ± 0.22 ^b
Aspartate aminotransferase (U/L)	120.80 ± 0.90 ^a	126.34 ± 0.75 ^b	134.24 ± 0.65 ^b
Alanine aminotransferase (U/L)	34.60 ± 0.80 ^a	38.76 ± 0.65 ^b	42.23 ± 0.75 ^b

Values within the same rows with different superscripts are significantly different at $p < 0.05$.

DISCUSSION

Animals are in normal physiological state when given enough food and water, but when withdrawn, the physiological and pathological variations may take place. Food and water restriction protocols are common in animal research; however information about the physiologic and behavioral impacts of food and water restriction and the potential adverse consequences that the animal will experience as a result of the restriction are limited. Food and water deprivation will affect the animals' health, internal cellular biochemistry, behaviour, absorption rate of substances, carbohydrate and lipid metabolism (Matsuzawa and Sakazume, 1994). Studies reported in literature on the effect of feed deprivation on various haematological indices have been conflicting and inconsistent. In this study, red blood cells count, haemoglobin and hematocrit

were elevated which are similar to the studies of El-Hazmi *et al.* (1987) and Keenan *et al.*, (1994) who reported similar haematological increases in male rats deprived of feed and water for twenty four hours. Conversely, Dewanti *et al.* (2006) showed a significant decrease in hematocrit and haemoglobin concentration. The increase in the haematological parameters observed in this study is attributed to non-availability of nutrients and water needed to augment haemopoiesis, bringing about dehydration, stress and haemoconcentration. The stress could induce the spleen to push more red cells into circulation bringing about relative polycythemia which is slightly macrocytic hypochromic. The relative polycythemia and haemoconcentration bring about increase haemoglobin concentration and packed cell volume.

Total white blood cell counts, neutrophils, eosinophils,

monocytes and lymphocyte, counts all increased over time compared to the basal parameters. Taken together, these changes in white blood cells counts are suggestive of a stress response due to feed and water deprivation. Similarly (Champy *et al.*, 2004) reported significant increase in white cell counts in animals subjected to feed deprivation within days which was attributed to means of coping with stress induced by feed and water deprivation. The longer the animals were stressed the greater the leucocytic response (Kioukia *et al.*, 2000).

Decreases in levels of blood glucose during fasting are due to lack of food intake and non-availability of glucose absorption from the intestines. Serum glucose concentrations in the current study were significantly lower when feed and water was deprived compared to the basal glucose levels measured when the animals were in fed state. The observation is similar to Claassen (1994) who reported decrease plasma glucose concentration in animals deprived of feed for three days. Glucose is the main supply of energy and is maintained between meals by glycogenolysis and possibly by gluconeogenesis in the liver. Glucocorticoid, insulin and glucagon also play significant role to prevent drastic fall in the glucose concentration when the animals were deprived of feed and water (Gursoy *et al.*, 2001; Heiderstadt *et al.*, 2000; Genn *et al.*, 2003).

Many reports have been published on the effect of feed deprivation on blood lipids among healthy animals, with inconsistent and even conflicting findings. The discrepancy was attributed to the amount and type of food intake, physical activity and genetic makeup of the animals (Aksungar *et al.*, 2005). We observed a decrease in the cholesterol and triglyceride concentrations in rats deprived of feed and water compared to the basal parameter, it is suggested that hormone sensitive lipase is activated leading to hydrolysis of esterified cholesterol and triglyceride. The fatty acids generated from the hydrolysis is oxidized to generate energy by the rats deprived of feed and water due to metabolic changes as glucose is not readily available as energy source (Mahboob *et al.*, 1999) reported decrease in plasma triglyceride concentration in human with restricted feed intake. Prolonged fasting induces an increase in plasma glucagon, plasma glucocorticoids, plasma epinephrine and a decrease in plasma insulin concentrations. This leads to hydrolysis of triglyceride in adipose tissue, which causes a rise in plasma free fatty acid concentration (Moitra *et al.*, 1998; Kersten *et al.*, 1999).

Protein constitutes the building block and the basic molecule for any biochemical reaction. They are intimately related with all physiological processes which maintain a simple biochemical system in living condition. The serum hyperalbuminaemia and hyperproteinaemia observed in the rats deprived of feed and water is a sharp response to dehydration and haemoconcentration. This is similar to observation by El-Hazmi *et al.*, (1987) and Keenan *et al.*, (1994) who reported hyperproteinaemia in rats subjected to feed restriction.

Kidney is a major organ involve in body homeostasis and excretion of waste from the body. A mild azotaemia was observed in the serum which is an indication of partial

clearance by the kidney. However Mgan *et al.*, (2011) observed no significant change in the azotaemia though there was mild increase in rats subjected to short time fasting.

The liver is the organ of detoxification and involve in metabolic activities of the animals. The activities of alanine aminotransferase and aspartate aminotransferase were observed to be slightly increased in the serum of rats deprived of feed and water, which is an indication of mild hepatocellular injury due to stress on the liver. The stress resulted in the release of cytoplasmic enzymes from the liver into the serum. It is likely that lipid peroxidation could oxidize the membrane lipid leading to escape of hepatocytes content into the serum. In a study on rats subjected to feed deprivation, the hepatocellular response produces no significant change in aspartate aminotransferase (AST), alanine aminotransferase (ALT) and sorbitol dehydrogenase (SDH) after six hours of deprivation, but after 12, 24 and 36 hours of feed deprivation AST, ALT and SDH were significantly increased in deprived animals compared with non-deprived animals (Lin *et al.*, 2005).

CONCLUSIONS

It is concluded from the data generated that, rats deprived of feed and water exhibit pronounced haematological and biochemical changes. The haematological changes ranges from dehydration to haemoconcentration, producing relative polycythaemia and viscous blood. Also there were stress induced metabolic changes in the liver and the kidney associated with the deprivation of feed and water.

ACKNOWLEDGEMENT

The author wishes to express gratitude to the University of Ilorin and the Department of Veterinary Physiology and Biochemistry, University of Ilorin for granting access to the animal house and laboratory facilities. Mr. Rahman S.A. is appreciated for sourcing healthy rats used for the study.

DECLARATION OF CONFLICT OF INTEREST

The authors declare no conflict of interest.

REFERENCES

- Aksungar, F.B., Eren, A., Ure, S., Teskin, O. and Ates, G. (2005). Effects of intermittent fasting on serum lipid levels, coagulation status and plasma homocysteine levels. *Annual Nutrition Metabolism* **49**(2): 77-82. DOI: 10.1159/000084739.
- Bauer, M., Hamm, A.C. and Bonaus, M. (2004). Starvation response in mouse liver shows strong correlation with life-span-prolonging processes. *Physiology of Genomics* **17**: 230-244. DOI: 10.1152/physiolgenomics.00203.200.
- Champy, M.F., Selloum, M. and Piard, L. (2004). Mouse functional genomics requires standardization of mouse handling and housing conditions. *Mammalian Genome* **15**: 768-783. DOI: 10.1007/s00335-004-2393-1.
- Claassen, V. (1994). *Neglected Factors in Pharmacology and Neuroscience Research: Biopharmaceutics, Animal Characteristics, Maintenance, Testing Conditions*.

- Elsevier ISBN 0-444-81907.
- Dewanti, L., Watanabe, C., Sulistiawati, E. and Ohtsuka, R. (2006). Unexpected changes in blood pressure and haematological parameters among fasting and non-fasting workers during Ramadan in Indonesia. *European Journal of Clinical Nutrition* **60**: 877-881. DOI: 10.1038/sj.ejcn.1602393.
- Grant, G.H. (1987). *Amino acids and Proteins; Fundamentals of Clinical Chemistry*. N.W. Tietz (ed), WB Saunders Company Philadelphia USA. Pp 328-329.
- Gursoy, E., Cardounel, A., Hu, Y. and Muhammed, K. (2001). Biological effects of long-term caloric restriction: adaptation with simultaneous administration of caloric stress plus repeated immobilization stress in rats. *Experimental Biological Medicine* **226**: 97-102. DOI: 10.1177/153537020122600206.
- Heiderstadt, K.M., McLaughlin, R., Wright, D., Walker, S. and Gomez-Sanches, C. (2000). The effect of chronic food and water restriction on open-field behaviour and serum corticosterone levels in rats. *Laboratory Animals* **34**: 20-28. DOI: 10.1258/002367700780578028.
- Henry, R.J. (1974). *Colorimetric Determination of Creatinine in Serum: Clinical Chemistry: Principles and Techniques*. 2nd edn., Harper and Row, New York, Pp: 525-529.
- Hughes, J.E., Amyx, H., Howard, J.L., Nanry, K.P. and Pollard, G.T. (1994). Health effects of water restriction to motivate lever-pressing in rats. *Laboratory Animal Science* **44**: 135-140. PMID: 8028275.
- Kanizsai, P., Garami, A., Solymar, M., Szolcsanyi, J. and Szelenyi, Z. (2009). Energetics of fasting heterothermia in TRPV1-KO and wild type mice. *Physiology Behaviour* **96**: 149-154. DOI: 10.1016/j.physbeh.2008.09.023
- Keenan, K.P., Laroque, P. and Dixit, R. (1998). Need for dietary control by caloric restriction in rodent toxicology and carcinogenicity studies. *Journal of Toxicology and Environmental Health Critical Review* **2**: 135-148. DOI: 10.1080/10937409809524548
- Kersten, S., Seydoux, J., Peters, J.M., Gonzalez, F.J., Desvergne, B. and Wraahli, W. (1999). Peroxisome proliferator activated receptor alpha mediates the adaptive response to fasting. *Journal of Clinical Investigation* **103**: 1489-1498. DOI: 10.1172/JCI6223
- Kioukia, N., Bekris, S., Antoniou, K., Papadopoulou-Daifoti, Z. and Christofidis, I. (2000). Effects of chronic mild stress (CMS) on thyroid hormone function in two rat strains. *Psychoneuroendocrinology* **25**: 247-257. DOI: 10.1016/S0306-4530(99)00051-7.
- Lin, X., Yue, P., Chen, Z. and Schonfeld, G. (2005). Hepatic triglyceride contents are genetically determined in mice: results of a strain survey. *American Journal of Physiology* **288**: G1179-G1189. DOI: 10.1152/ajpgi.00411.2004.
- Mahboob, S., Sattarivand, R., Nouri, M. and Arefhosseini, S. (1999). Effect of Ramadan fasting on serum lipids profiles in normal and hyperlipidemic subjects. *Saudi Medical Journal* **20**: 947-950. DOI: 123456789/52164
- Matsuzawa, T. and Sakazume, M. (1994). Effects of fasting on haematology and clinical chemistry values in the rat and dog. *Comparative Haematology International* **4**: 152-156. DOI: 10.1007/BF00798356.
- Megan, H.N., Kelly, M.S., Hugunin, A. and Karen, L.R. (2011). Effects of Short-Term Fasting in Male Sprague-Dawley Rats. *American Association Laboratory Animal Science* **61** (2): 138-147. PMID: PMC3079815.
- McGowan, M.W., Artiss, J.D., Strandbergh, D.R. and Zak, B. (1983). A peroxidase-coupled method for the colorimetric determination of serum triglycerides. *Clinical Chemistry* **29**: 538-542. DOI:10.1093/clinchem/29.3.538.
- Moitra, J., Mason, M.M. and Olive, M. (1998). Life without white fat: a transgenic mouse. *Genes Development* **12**: 3168-3181. DOI: 10.1101/grad.12.20.3168.
- Palou, A., Remesar, X., Arola, L.I., Herrera, E. and Alemany, M. (1981). Metabolic effects of short term food deprivation in the rat. *Hormonal Metabolic Research* **13**: 326-333. DOI: 10.1015/s-2007-1019258.
- Penicaud, L. and Le Magnen, J. (1980). Recovery of body weight following starvation or food restriction in rats. *Neuroscience Biobehavioural Review* **4**(1): 47-52. DOI: 10.1016/0149-7634(80)90048-2.
- Reitman, S. and Frankel, S. (1957). Colorimetric methods for aspartate and alanine aminotransferase. *American Journal of Clinical Pathology* **28**: 55-60. DOI: 10.1093/ajcp/28.1.56.
- Schalm, D.W., Jain, N.C. and Carrot, E.J. (1975). *Veterinary Haematology*. 3rd edn., Lea and Febiger, Philadelphia, ISBN-13: 978-0812104707, Pp:807.
- Strubbe, J.H., Keijser, J.T., Dijkstra, T. and Prins, A.B. (1986). Interaction between circadian and caloric control of feeding behaviour in the rat. *Physiological Behaviour* **36**: 489-493. DOI: 10.1016/0031-9384(86)90320-3.
- Stricker, E.M. (1984). Biological basis of hunger and satiety: therapeutic implications. *Nutritional Review* **42**: 333-340. DOI: 10.1111/j.1753-4887.1984.tb02249.x.
- Sweileh, N., Schnitzler, A., Hunter, G.R. and Davis, B. (1992). Body composition and energy metabolism in resting and exercising muslims during Ramadan fast. *Journal of Sports Medicine and Physical Fitness* **32**:156-163. PMID: 1434584.
- Tietz, N.W. (1995). *Clinical Guide to Laboratory Tests*. 3rd (ed). W.B. Saunders Company, Philadelphia. Pp. 518-519.
- Tucci, V., Hardy, A. and Nolan, P.M. (2006). A comparison of physiological and behavioural parameters in C57BL/6J mice undergoing food or water restriction regimes. *Behavioural Brain Research* **173**: 22-29. DOI: 10.1016/j.bbr.2006.05.031.
- Toth, L.A. and Gardiner, T.W. (2000). Food and water restriction protocols: physiological and behavioral considerations. *Contemporary Topics* **39**: 9-17. PMID: 11487246.
- Vermeulen, J.K., De Vries, A., Schlingmann, F. and Remie, R. (1997). Food deprivation: common sense or nonsense? *Animal Technology* **48**: 45-54.
- Wingfield, J.C. and Kitaysky, A.S. (2002). Endocrine responses to unpredictable environmental events: Stress or anti-stress hormones? *Integrated Comparative Biology* **42**: 600-609. DOI: 10.1093/ICB/42.3.600.

Wybenga, D.R., Pileggi, V.J., Dirstine, P.I. and Di Giorgio, J. (1970). Direct manual determination of serum total cholesterol with a single stable reagent. *Clinical Chemistry* **16**: 980-984. DOI: 10.1093/clinchem/16.12.980.
