

*Ceylon Journal of Science* 49(1) 2020: 49-59  
DOI: <http://doi.org/10.4038/cjs.v49i1.7705>

## RESEARCH ARTICLE

# Diversity of butterflies in different habitat types of Seethawaka wet zone botanic gardens and Indikadamukalana forest reserve of Sri Lanka

M.U.H. Peiris<sup>1</sup>, C.D. Dangalle<sup>1\*</sup>, N. Pallewatta<sup>1</sup> and S. Wijesundara<sup>2</sup>

<sup>1</sup>Department of Zoology and Environment Sciences, Faculty of Science, University of Colombo, Colombo, Sri Lanka.

<sup>2</sup>National Institute of Fundamental Studies, Kandy, Sri Lanka.

Received: 01/07/2019 ; Accepted: 25/01/2020

**Abstract:** Botanic gardens, forest reserves and other protected areas are established to conserve biodiversity. The present study was conducted to assess the butterfly diversity in a recently established botanic garden of Sri Lanka, the Seethawaka Wet Zone Botanic Gardens (SWZBG) and Indikadamukalana Forest Reserve (IMFR) adjacent to the garden. Three different habitat types within the botanic gardens, the undisturbed forest habitat within the forest reserve and the buffer zone between the Botanic Gardens and Forest Reserve were investigated for butterflies. Butterfly species richness, diversity, endemic species, family composition in different habitats and effects of habitat parameters on butterfly distribution were assessed. Butterflies were collected from five habitats – grassland, shrub, disturbed forest, undisturbed forest, buffer zone, of the SWZBG and IMFR using transect line method with hand netting and fruit baited traps. Collected butterflies were identified using published field guides. Climate and soil parameters of each habitat were measured using standard methods and equipment. Differences of habitat parameters between habitats were estimated using One-Way Analysis of Variance and Tukey's Multiple Comparison method and butterfly diversity and evenness in different habitat types were assessed using Shannon-Weiner diversity index. The study revealed seventy-nine butterfly species from SWZBG and IMFR highlighting the importance of the sites for safeguarding and conservation of butterflies. Species richness and diversity of butterflies was highest in the buffer zone and lowest in the grassland habitat. The most common butterfly family in the study sites was family Nymphalidae. Endemic species were high in the forest habitat types. Significant differences were evident in habitat parameters between habitat types indicating their effect on butterfly species richness and diversity. The study revealed SWZBG and IMFR as important areas for butterfly occurrence when compared with similar botanic gardens and forest reserves elsewhere in the world.

**Keywords:** Buffer zone, Grassland habitat, Nymphalidae, Habitat parameters, Endemic butterflies.

## INTRODUCTION

Butterflies are a taxonomically well studied group within the Lepidoptera and has received considerable attention throughout the world (Ghazoul, 2002) and are represented


by 17,280 species of which roughly two-thirds live in the tropics (Shields, 1989).

Sri Lanka is home to 247 species of butterflies including 31 endemic species and 84 endemic sub-species (van der Poorten and van der Poorten, 2016). Comprehensive publications on the butterfly fauna of Sri Lanka by Woodhouse (1949), d'Abbrera (1998), and more recently by van der Poorten and van der Poorten (2016) provides valuable information on the identification, distribution, biology and food plants of individual species.

Several studies have documented the species composition of butterflies in different localities of Sri Lanka. Biodiversity surveys carried out by IUCN Sri Lanka have recorded butterfly species in Bundala National Park (Bambaradeniya *et al.*, 2001) Muthurajawela Sanctuary (Bambaradeniya *et al.*, 2002a), Maduganga Mangrove Estuary (Bambaradeniya *et al.*, 2002b), Sinharaja Rain Forest (Bambaradeniya *et al.*, 2003) and Anawilundawa Sanctuary (Perera *et al.*, 2005). The butterfly fauna of Udawalawa National Park has been documented by Samarasinghe *et al.*, 1998, and that of Tantirimale Archaeological site, Anuradhapura by Asela *et al.*, 2009. Eventhough the occurrence of butterfly species in different areas of the island have been relatively well documented very little attention has been paid to the study of butterfly ecology prior to the work of van der Poorten and van der Poorten (2016).

The association between different habitat types of a tropical rain forest with diversity of butterfly communities has been demonstrated by Vu and Vu (2011) and according to Vu *et al.*, 2015 habitat variation and disturbances within forests can affect butterfly communities and diversity in complex ways. Studies in Uganda revealed that forest remnants and semi-natural habitats are important for conservation of butterflies (Munyuli, 2013). Habitat type affected butterfly species richness and diversity indices in Bumbuna forest, Northern Sierra Leone and were lower in the disturbed habitats compared to the forest reserve (Sundufu and Dumbuya, 2008). However, contradictory observations have been revealed between butterflies and habitat types of Aokigahara primary woodland of Mount

\*Corresponding Author's Email: [cddangalle@gmail.com](mailto:cddangalle@gmail.com)

 <https://orcid.org/0000-0001-9483-389>



Fiji, Central Japan where butterfly species occurring in disturbed habitats were more voltine and had a wider host plant range (Kitahara, 2004).

Climatic and soil data of habitats are important in determining the distribution, diversity and abundance of butterflies and many studies elsewhere in the world have demonstrated this observation. Species richness of butterflies were found to be closely correlated with air temperature in lowland tropical forests of Bolivia, South America and species numbers differed significantly between seasons (Abrahamczyk *et al.*, 2011). The range and occupancy of the monarch butterfly (*Danaus plexippus plexippus*) of the United States were determined using climate variables which affected host plants of the area (Dilts *et al.*, 2019). Potential evapotranspiration measured in mm/year was a strong predictor of butterfly species richness in extensive terrestrial areas of Canada (Kerr, 2001). Soil type, soil moisture and soil acidity in agricultural landscapes of Iowa, USA were known to affect crop diversity and density which in turn influenced butterfly assemblages (Myers *et al.*, 2015).

In Sri Lanka, Nisviya and Wickramasinghe (2012) revealed that the Northern flank of Knuckles Mountain holds a rich and unique butterfly assemblage and that this maybe due to the plant species with nectar bearing flowers in the area. Larval food plants of butterflies have been documented by Jayasinghe *et al.*, 2014. Investigations in different habitat types of Wasgamuwa National Park, Sri Lanka have revealed that butterfly abundance and richness is lowest in primary forests when compared with secondary forests, wetland margins, shrubland and home gardens and species tend to be more specialized to the habitat they occupy (Slater *et al.*, 2019). The diversity of butterflies in different habitat types in Matara have been recorded by Pathiraja *et al.* (2017). However, the diversity of butterfly species in different habitat types of Sri Lanka and their associations with specific habitat character traits have not been studied in detail. Therefore, the present study was conducted to investigate and record the butterfly

fauna of an important and recently established biodiversity park, Seethawaka Wet Zone Botanic Gardens and a wet zone tropical forest, Indikadamukulana Forest Reserve of Sri Lanka. Different habitat types within the locations were surveyed and species richness associated with habitat parameters was determined.

## MATERIALS AND METHODS

### Study Area

The study was conducted in two locations: Seethawaka Wet Zone Botanic Gardens and the adjacent Indikadamukulana Forest Reserve (6°57'11"N, 80°13'06"E), located in Labugama, Colombo District, Western Province of Sri Lanka (Figure 1).

Three habitat types were selected for the study within the Botanic Gardens as follows:

a. Grassland: The grassland habitat consisted mainly of perennial grasses dominated by the species *Axonopus compressus* and small herbaceous plants such as *Zinnia elegans* and *Tagetes erecta*. The area was devoid of trees. In a few areas of the grassland, landscaped areas with flowers (*Zinnia elegans*, *Turnera ulmifolia*, *Hibiscus radiates*, *Angelonia salicariifolia*) were seen which were known to be changed seasonally by the garden management.

b. Shrub: This habitat type consisted of the plant species *Stachytarpheta urticaefolia*, *Lantana camara*, *Dicranopteris linearis* and *Syzgium* species.

c. Disturbed forest: This habitat consisted of a regenerating secondary forest characterized mainly by *Anacardium occidentale*, *Bridelia moonii*, *Caryota urens*, *Terminalia arjuna*, *Cassia fistula* and *Pueraria phaseoloides*.

Two habitats were selected for the study at the Indikadamukulana Forest Reserve.

a. Undisturbed forest: Primary forest with tall trees and close canopy consisting mainly of *Cinnamom umverum*, *Dioscorea spicata*, *Smilax zeylanica*, *Caryota urens*, *Thottea siliquosa* and *Bambusa vulgaris*.





Figure 1: The study area.



**Figure 2:** Habitat types sampled within the study area. (Adapted from Laborde & Corrales-Ferrayola, 2012)

b. Buffer zone: This habitat was the area between the disturbed and undisturbed forest and consisted mainly of *Trema orientalis*, *Desmodium heterocarpon* and invasive plants such as *Lantana camara*, *Dillenia* sp. and *Clidemia hirta*. (Figure 2).

### Butterfly Sampling

Butterflies were sampled for a period of four months from July 2016 to October 2016 using transect line method with hand netting and fruit baited traps. Each habitat type was surveyed weekly. Two 100 m length transects were established for each habitat type, and the butterfly species seen within a 10m x 10m x 10m area were recorded. Transect surveys were carried out between 8.00 to 18.00 hours on each sampling day utilizing one hour for each transect. Additionally, fruit baited traps were placed in two sites of each habitat type and surveyed weekly 24 hours after placing the trap. Captured specimens were released after photographing (Canon Powershot SX60) and identification.

### Identification of Butterfly Species

Butterfly species were identified using field guides (Jayasinghe *et al.*, 2015) and photographs of publications (Woodhouse, 1949; d'Abrera, 1998; van der Poorten and van der Poorten, 2016).

### Measurement of Habitat Parameters

During each survey climate and soil parameters of the habitat types were measured using standard equipment and methods. The climate parameters, environmental temperature (°C), solar radiation (w/m<sup>2</sup>), relative humidity (%) and wind speed (MPH) were measured using a portable Vantage Pro 2 weather station, and light intensity was measured using a Brannan digital luxmeter (0.01Klx). Rainfall measures of the sampling months were taken from the Department of Meteorology. Canopy cover was recorded using a spherical densitometer (1.04%).

Soil temperature (°C), soil pH and soil electric

conductivity (μS/cm) were measured using standard equipment while soil moisture was measured using the gravimetric method.

### Data Analysis

Species diversity of each habitat type was calculated using the Shannon-Weiner index given below.

$$H = \sum [(p_i) \times \ln(p_i)]$$

H = Shannon-Weiner Index

$p_i$  = Proportion of total sample represented by species  $i$ .

Species evenness of habitat types were calculated using the following equation.

$$E = H / H_{\max}$$

E = Evenness

H = Shannon-Weiner Index

$$H_{\max} = \ln S$$

S = Number of Species

Climate and soil parameters of the five different habitats were compared using One-Way ANOVA of Minitab version 17.0. Tukey's pairwise comparison at 95% confidence was used to determine the significant relationships of habitat parameters between habitat types.

Species diversity and evenness of the butterflies of each habitat type were compared with the habitat parameter differences between the habitat types.

## RESULTS

### Species Composition and Abundance of Butterflies

A total of seventy-nine (79) butterfly species representing the families Papilionidae, Pieridae, Nymphalidae, Lycaenidae, Riodinidae and Hesperidae were recorded from the study area (Table 1). Most of the species represented family Nymphalidae. *Ypthima ceylonica* was the most common butterfly encountered in family Nymphalidae. For family Papilionidae, the most common

species was *Papilio polymnestor parinda*; for Pieridae, *Leptosia nina nina*; Lycaenidae, *Jamides celeno tissama*; Hesperidae, *Iambrix salsala luteipalpis*. *Abisara echerius prunosa* the only species of Family Riodinidae in Sri Lanka was also encountered in the study.

39.24% of the species were endemic and the endangered

Ceylon Rose (*Pachliopta jophon*) was recorded from the shrub habitat and disturbed forest of the Botanic Gardens, and buffer zone. Percentage of endemic species was high in the disturbed and undisturbed forest habitats when compared with the other habitat types (Table 2).

**Table 1:** Butterfly species recorded from different habitat types of the Seethawaka Wet Zone Botanic Gardens and Indikadamukalana Forest Reserve.

Species [Family within brackets]	Habitat
1. * <i>Troides darsius</i> Gray, 1853 [Papilionidae]	Grassland habitat
2. <i>Papilio polytes romulus</i> Cramer, 1775 [Papilionidae]	Shrub habitat
3. <i>Leptosia nina nina</i> Fabricius, 1793 [Pieridae]	Disturbed forest
4. <i>Parantica aglea aglea</i> Stoll, 1782 [Nymphalidae]	Undisturbed forest
5. <i>Junonia atlites atlites</i> Linnaeus, 1763 [Nymphalidae]	Buffer zone
6. <i>Neptis hylas varmona</i> Moore, 1872 [Nymphalidae]	
7. ** <i>Parthenos sylvia cyaneus</i> Moore, 1877 [Nymphalidae]	
8. <i>Orsotriaena medus mandata</i> Moore, 1857 [Nymphalidae]	
9. <i>Mycalesis perseus typhlus</i> Fruhstorfer, 1908 [Nymphalidae]	
10. ** <i>Mycalesis patnia patnia</i> Moore, 1857 [Nymphalidae]	
11. <i>Ypthima ceylonica</i> Hewitson, 1864 [Nymphalidae]	
12. ** <i>Jamides celeno tissama</i> Fruhstorfer, 1916 [Lycaenidae]	
13. <i>Phalanta phalantha phalantha</i> Drury, 1773 [Nymphalidae]	Grassland habitat
14. <i>Junonia almana almana</i> Linnaeus, 1758 [Nymphalidae]	
15. <i>Zizina otis indica</i> Murray, 1874 [Lycaenidae]	
16. ** <i>Ampittia dioscorides singa</i> Evans, 1949 [Hesperiidae]	
17. <i>Junonia iphitha iphitha</i> Cramer, 1779 [Nymphalidae]	Grassland habitat Shrub habitat
18. <i>Catopsilia pyranthe pyranthe</i> Linnaeus, 1758 [Pieridae]	Grassland habitat, Shrub habitat Buffer zone
19. <i>Discolampa ethion ethion</i> Westwood, 1851 [Lycaenidae]	Grassland habitat, Disturbed forest
20. <i>Pelopidas conjuncta narooa</i> Moore, 1878 [Hesperiidae]	Grassland habitat, Buffer zone
21. <i>Parnara bada bada</i> (Moore, 1878) [Hesperiidae]	
22. <i>Eurema hecabe hecabe</i> Linnaeus, 1758 [Pieridae]	Grassland habitat, Shrub habitat
23. * <i>Potanthus satra</i> C. & R. Felder, 1862 [Hesperiidae]	Disturbed forest, Buffer zone
24. <i>Pelopidas agna agna</i> Moore, 1866 [Hesperiidae]	
25. <i>Pelopidas mathias mathias</i> Fabricius, 1798 [Hesperiidae]	
26. <i>Hypolimnas misippus</i> Linnaeus, 1764 [Nymphalidae]	Grassland habitat
27. <i>Hypolimnas bolina bolina</i> Linnaeus, 1758 [Nymphalidae]	Undisturbed forest Buffer zone
28. ** <i>Caltoris philippina seriata</i> Moore, 1878 [Hesperiidae]	Grassland habitat
29. <i>Catopsilia pomona pomona</i> Fabricius, 1775 [Pieridae]	Disturbed forest Buffer zone
30. ** <i>Pachliopta aristolochiae ceylonica</i> Moore, 1881 [Papilionidae]	Grassland habitat, Disturbed forest Undisturbed forest, Buffer zone



	<b>Species [Family within brackets]</b>	<b>Habitat</b>
31.	** <i>Papilio polymnestor parinda</i> Moore, 1881 [Papilionidae]	Shrub habitat, Disturbed forest
32.	<i>Graphium sarpedon teredon</i> Felder & Felder, 1865 [Papilionidae]	Undisturbed forest, Buffer zone
33.	<i>Delias eucharis</i> (Drury, 1773) [Pieridae]	
34.	** <i>Elymnias hypermnestra fraterna</i> Butler, 1871 [Nymphalidae]	
35.	<i>Nacaduba hermnus sidoma</i> Fruhstorfer, 1916 [Lycaenidae]	
36.	<i>Abisara echerius prunosa</i> Moore, 1879 [Riodinidae]	
37.	<i>Iambrix salsala luteipalpis</i> Plötz, 1886 [Hesperiidae]	
38.	<i>Taractrocera maevius</i> Fabricius, 1793 [Hesperiidae]	
39.	* <i>Pachliopta jophon</i> Gray, 1853 [Papilionidae]	Shrub habitat
40.	<i>Graphium agamemnon menides</i> Fruhstorfer, 1904 [Papilionidae]	Disturbed forest
41.	<i>Acraea terpsicore</i> Linnaeus, 1758 [Nymphalidae]	Buffer zone
42.	<i>Mycalesis mineus polyducta</i> Cramer, 1777 [Nymphalidae]	
43.	<i>Castalius rosimon rosimon</i> Fabricius, 1775 [Lycaenidae]	
44.	** <i>Cupha erymanthis placida</i> Moore, 1881 [Nymphalidae]	Shrub habitat, Disturbed forest, Undisturbed forest
45.	** <i>Eurema blanda citrina</i> Moore, 1881 [Pieridae]	Shrub habitat
46.	** <i>Cirroochroa thais lanka</i> Moore, 1872 [Nymphalidae]	Disturbed forest
47.	** <i>Melanitis phedima tambra</i> Moore, 1880 [Nymphalidae]	
48.	** <i>Cethosia nietneri nietneri</i> C. & R. Felder, 1867 [Nymphalidae]	Shrub habitat Undisturbed forest
49.	** <i>Caprona ransonnettii ransonnettii</i> R. Felder, 1868 [Hesperiidae]	Shrub habitat Buffer zone
50.	<i>Papilio crino</i> Fabricius, 1793 [Papilionidae]	Disturbed forest, Undisturbed forest, Buffer zone
51.	* <i>Jamides coruscans</i> Moore, 1877 [Lycaenidae]	Disturbed forest Undisturbed forest
52.	<i>Melanitis leda leda</i> Linnaeus, 1758 [Nymphalidae]	Disturbed forest, Buffer zone
53.	<i>Prosotas nora ardates</i> Moore, 1875 [Lycaenidae]	
54.	<i>Acytolepis puspa felderi</i> Toxopeus, 1927 [Lycaenidae]	
55.	** <i>Chilades pandava lanka</i> Evans, 1925 [Lycaenidae]	
56.	<i>Everes lacturnus lacturnus</i> Godart, 1824 [Lycaenidae]	Undisturbed forest, Buffer zone
57.	* <i>Eurema ormistoni</i> Watkins, 1925 [Pieridae]	Disturbed forest
58.	* <i>Idea iasonia</i> Westwood, 1848 [Nymphalidae]	
59.	** <i>Euploea core asela</i> Moore, 1877 [Nymphalidae]	
60.	<i>Neptis jumbah nalanda</i> Fruhstorfer, 1908 [Nymphalidae]	
61.	<i>Zizula hylax hylax</i> Fabricius, 1775 [Lycaenidae]	
62.	<i>Megisba malaya thwaitesi</i> Moore, 1881 [Lycaenidae]	
63.	** <i>Papilio helenus mooreanus</i> Rothschild, 1895 [Papilionidae]	Undisturbed forest
64.	** <i>Moduza procris calidasa</i> Moore, 1858 [Nymphalidae]	
65.	** <i>Discophora lepida ceylonica</i> Fruhstorfer, 1911 [Nymphalidae]	
66.	** <i>Loxura atymnus arcuata</i> Moore, 1881 [Lycaenidae]	
67.	* <i>Nacaduba sinhala</i> Ormiston, 1924 [Lycaenidae]	
68.	** <i>Jamides alecto meilichius</i> Fruhstorfer, 1916 [Lycaenidae]	
69.	<i>Talicauda nyseus nyseus</i> Guérin-Méneville, 1843 [Lycaenidae]	
70.	<i>Neopithecops zalmora dharma</i> Moore, 1881 [Lycaenidae]	
71.	** <i>Tagiades litigiosa ceylonica</i> Evans, 1932 [Hesperiidae]	
72.	<i>Oriens goloides</i> Moore, 1881 [Hesperiidae]	


	Species [Family within brackets]	Habitat
73.	** <i>Papilio clytia lankeswara</i> Moore, 1879 [Papilionidae]	Buffer zone
74.	** <i>Graphium doson doson</i> C.& R. Felder, 1864 [Papilionidae]	
75.	<i>Spalgis epeus epeus</i> Westwood, 1851 [Lycaenidae]	
76.	<i>Surendra quercetorum discalis</i> Moore, 1857 [Lycaenidae]	
77.	<i>Zizeeria karsandra</i> Moore, 1865 [Lycaenidae]	
78.	□□ <i>Suastus gremius subgrisea</i> Moore, 1878 [Hesperiidae]	
79.	<i>Borbo cinnara</i> Wallace, 1866 [Hesperiidae]	

\* Endemic species

\*\* Endemic sub-species

**Table 2:** Number of endemic butterfly species of different habitat types as a percentage of the total number of endemic species.

Habitat Type	Percentage of Endemic Species (%)
Grassland	25.81
Shrub	45.16
Disturbed Forest	61.29
Undisturbed Forest	54.84
Buffer Zone	48.39



**Figure 3:** Species richness of butterflies in different habitat types of the Botanic Garden and Forest Reserve.

**Table 3:** Shannon-Weiner diversity indices and species evenness for butterfly species of the different habitat types.

Habitat Type	Diversity Index (Mean $\pm$ S.E.)	Species Evenness (Mean $\pm$ S.E.)
Grassland	1.59 <sup>A</sup> $\pm$ 0.42	0.71 <sup>A</sup> $\pm$ 0.18
Shrub	1.78 <sup>B</sup> $\pm$ 0.31	0.77 <sup>A</sup> $\pm$ 0.04
Disturbed Forest	2.41 <sup>B</sup> $\pm$ 0.11	0.78 <sup>A</sup> $\pm$ 0.02
Undisturbed Forest	2.07 <sup>B</sup> $\pm$ 0.21	0.90 <sup>A</sup> $\pm$ 0.03
Buffer Zone	2.71 <sup>B</sup> $\pm$ 0.06	0.88 <sup>A</sup> $\pm$ 0.02

Means with letters that differ along the column are significantly different.

### Diversity of Butterflies in Different Habitats

The highest number of butterfly species and a significantly high diversity index was recorded for the buffer zone, the area between the Botanic Gardens and Forest Reserve. Lowest number of species and a significantly low diversity index was recorded for the grassland habitat found within the Botanic Gardens (Figure 3 and Table 3). *Leptosia nina nina* of family Pieridae was the most common butterfly encountered in the buffer zone habitat while *Junonia atlites*

*atlites* of the same family was common in the grassland. *Ypthima ceylonica* of family Nymphalidae was the most common species in all other habitat types.

Species of all six butterfly families occurred in all habitat types with the exception of the grassland habitat. Family Riodinidae was not found in the grassland habitat. The species evenness of the habitat types indicated the absence of dominant butterfly species (Table 3).

**Table 4:** Climatic and soil parameters of the habitat types at Seethawaka Wet Zone Botanic Gardens and Indikadamukalana Forest Reserve.

Habitat Parameters	Grassland	Shrub Habitat	Disturbed Forest	Undisturbed Forest	Buffer Zone
Environmental Temperature (°C)	29.20 <sup>A</sup> ± 0.36	29.43 <sup>A</sup> ± 0.34	29.36 <sup>A</sup> ± 0.44	29.43 <sup>A</sup> ± 0.37	29.86 <sup>A</sup> ± 0.39
Mean ± S.E.					
Solar Radiation (w/m <sup>2</sup> )	307.90 <sup>A</sup> ± 58.58	174.90 <sup>B</sup> ± 36.84	181.60 <sup>B</sup> ± 33.95	56.70 <sup>C</sup> ± 18.45	212.20 <sup>A</sup> ± 32.77
Mean ± S.E.					
Relative Humidity (%)	76.47 <sup>A</sup> ± 1.53	75.73 <sup>A</sup> ± 1.52	79.00 <sup>A</sup> ± 1.80	77.13 <sup>A</sup> ± 2.16	75.60 <sup>A</sup> ± 2.01
Mean ± S.E.					
Wind Speed (MPH)	3.53 <sup>A</sup> ± 0.58	1.76 <sup>B</sup> ± 0.26	1.33 <sup>B</sup> ± 0.22	0.53 <sup>B</sup> ± 0.14	1.20 <sup>B</sup> ± 0.21
Mean ± S.E.					
Light intensity (Klux)	19.91 <sup>A</sup> ± 3.12	14.83 <sup>A</sup> ± 2.75	15.06 <sup>A</sup> ± 2.61	3.07 <sup>B</sup> ± 1.27	17.98 <sup>A</sup> ± 2.44
Mean ± S.E.					
Canopy Cover (%)	10.65 <sup>A</sup> ± 4.54	57.10 <sup>B</sup> ± 3.83	74.00 <sup>C</sup> ± 2.40	86.48 <sup>D</sup> ± 0.72	65.50 <sup>E</sup> ± 2.24
Mean ± S.E.					
Soil Temperature (°C)	28.41 <sup>A</sup> ± 0.33	26.15 <sup>B</sup> ± 0.33	26.36 <sup>B</sup> ± 0.37	25.25 <sup>B</sup> ± 0.26	28.01 <sup>A</sup> ± 0.49
Mean ± S.E.					
Soil pH	4.59 <sup>A</sup> ± 0.08	4.69 <sup>B</sup> ± 0.06	5.19 <sup>C</sup> ± 0.07	4.60 <sup>A</sup> ± 0.09	4.94 <sup>D</sup> ± 0.07
Mean ± S.E.					
Soil Electric Conductivity (µs/cm)	23.01 <sup>A</sup> ± 2.01	20.19 <sup>B</sup> ± 2.01	22.40 <sup>A</sup> ± 1.50	31.06 <sup>C</sup> ± 3.61	27.24 <sup>A</sup> ± 3.23
Mean ± S.E.					
Soil Moisture (%)	16.97 <sup>A</sup> ± 1.30	13.03 <sup>B</sup> ± 0.81	15.00 <sup>C</sup> ± 0.82	14.62 <sup>C</sup> ± 0.89	13.09 <sup>B</sup> ± 0.62
Mean ± S.E.					

Means with letters that differ along the rows are significantly different.

### Associations of Selected Habitat Parameters with Habitat Types

The buffer zone and grassland habitat differed significantly in certain climatic and soil parameters. When considering the climatic parameters, wind speed of the grassland habitat was significantly high than that of the buffer zone and canopy cover was significantly low. Solar radiation did not differ between the two habitat types. When considering the soil parameters grassland habitat had a significantly low soil pH, whereas soil moisture was significantly high. Soil temperature and soil electric conductivity did not differ between the two habitat types. Air temperature and relative humidity was more or less similar for all habitat types with no significant differences. However, significant differences were evident in the other parameters between certain habitat types (Table 4).

### DISCUSSION

The establishment of botanic gardens, forest reserves and other protected areas is a strategy to avoid the loss of habitats and conserve flora and fauna. Many such areas occur in the world as a single habitat or a network of habitats where potential connectivity is possible (Alexandre *et al.*, 2010). The Seethawaka Wet Zone Botanic Gardens and Indikadamukalana Forest Reserve of Sri Lanka are protected areas that consist of many habitat types where both structural and functional connectivity is possible.

Structural connectivity accounting for the area and spatial configuration of habitats and functional connectivity reflecting the movement of individuals or genes among populations prevents fragmentation of habitats and is essential for successful restoration and conservation management of landscapes (Tischendorf and Fahring, 2000). The current study revealed the presence of a rich butterfly fauna from the Seethawaka Wet Zone Botanic Gardens and Indikadamukalana Forest Reserve emphasizing their function towards conservation of biodiversity. Seventy-nine species of butterflies from six families were recorded from both areas which is comparatively high when considering with similar botanic gardens and forest reserves elsewhere in the world. The Experimental Botanic Gardens of Meghalaya, India with an area of 25 acres and at an altitude of 1000 m above sea level, houses sixty-six species of butterflies of five families (Bora *et al.*, 2014). The botanical garden UPI in Bandung, Indonesia harbour forty species of butterflies (Sanjaya *et al.*, 2016), while ninety-one species of butterflies under five major families occur in Royal Manas National Park of Bhutan which has an area of 1057 km<sup>2</sup> (261,190.4 acres) (Nidup *et al.*, 2014). The Seethawaka Wet Zone Botanic Gardens is a new botanic garden declared in 2015. However, it comprises of an array of vegetational habitats in an area of 105 acres, 100 meters above sea level. The vegetation is known to be provided by the seeds of Indikadamukalana Forest Reserve which is just adjacent to the gardens (Ranwala *et al.*, 2017).

The Botanic Garden has an area that is sufficient for the occupancy of butterflies when compared with other such Gardens (Bora *et al.*, 2014; Nidup *et al.*, 2014; Sanjaya *et al.*, 2016) and suitable elevation. A study in Aralam Wildlife Sanctuary, Kerala revealed that butterfly species preferred habitats with low (< 250 m) and middle (251 – 700 m) elevations compared to habitats with high (>700 m) altitudes (Sreekumar & Balakrishnan, 2001). Thus, the garden and forest reserve together offer an ideal refuge to a diverse community of butterfly species of the country.

Majority of butterfly species in the area was of family Nymphalidae. Diversity studies and checklists in forests (Sundufu and Dumbuya, 2008; Asela *et al.*, 2009; Vu and Vu, 2011), botanic gardens (Bora *et al.*, 2014; Sanjaya *et al.*, 2016), and nature reserves (Nidup *et al.*, 2014, Koneri *et al.*, 2017) have revealed Nymphalidae to be the most common type of butterflies found in such areas.

The buffer zone of the area harboured the highest number of butterfly species while the grassland habitat had the lowest number. A buffer zone is an extension of a protected area, or a transition zone between two different areas with limited human interference (Bhusal, 2014). In the present study the buffer zone was a transition zone between the disturbed and undisturbed forest and significantly differed in canopy cover, solar radiation, soil temperature, soil pH and soil moisture from that of the two habitats on either side of it. Canopy cover was low in the buffer zone than in the forest habitats paving the way to a significantly high solar radiation. The soil temperature was high leading to a significantly lower soil moisture than the soils of the forest habitats; and soil pH of the buffer zone had an intermediate value inbetween the disturbed and undisturbed forest soils. These differences in the climate and soil of the buffer zone may have given rise to a unique vegetation and microhabitats favoured by an array of butterfly species. The buffer zone consisted of many invasive plants such as *Lantana camara*, *Dillenia suffruticosa* and *Clidemia hirta* that were not present in the other habitat types. A study conducted in the Northern flank of Knuckle Forest Reserve of Sri Lanka showed that 56% of butterflies in the region heavily utilized *Lantana camara* and other invasive plants (Nisviya and Wickramasinghe, 2012). And according to Liyanagamage (2016) certain butterfly species of families Papilionidae, Nymphalidae, Pieridae and Hesperidae show preference to *Lantana camara* and families Papilionidae and Nymphalidae to the invasive species *Annona glabra*. Thus, it is highly possible that the high species richness of butterflies in the buffer zone is associated with the presence of invasive plants in the region. *Lantana camara* may attract more butterflies than the other invasive species, as the flowers of the species are known to have a high nectar volume and sugar content and are usually pollinated by Lepidoptera (Carrion-Tacuriet *et al.*, 2012). *Dillenia suffruticosa* have flowers that are nectarless but are pollinated by insects attracted to mature fruits (Smisha *et al.*, 2016). *Clidemia hirta* consists of an inflorescence bearing many nectaries and fruits borne in clusters (Rao and Sagar, 2012). Further, the high species richness of butterflies in the buffer zone could be attributed to the area being an ecotone: a zone of transition between adjacent ecological

systems. In such areas increased species richness occur due to representatives of species characteristic of both of the adjacent communities described as the edge effect (Baker *et al.*, 2002).

The grassland habitat within the Seethawaka Wet Zone Botanic Gardens had the lowest number of butterfly species and a significantly low diversity index. Further, Plum Judy (*Abisara echerius*), the only species of family Riodinidae of Sri Lanka found in all other habitat types was not found in the grassland habitat. A study on European butterflies have revealed that grassland area and landscape composition has an effect on the species richness of butterflies, and larger grasslands situated in landscapes consisting of a high proportion of grasslands tend to have higher butterfly species richness. Further relationships were revealed with butterfly species richness and vegetation height and abundance of flowers of grasslands, and high species richness was evident with increase of vegetation height and flower abundance (Ockinger and Smith, 2006). The present grassland situated within the Seethawaka Wet Zone Botanic Gardens was a small area amidst a mosaic of other habitat types. Further, it consisted mainly of grasses with a few grass flowers and was devoid of tall vegetation. These conditions may have led to the decrease in butterfly species richness and the unfavourable climate and soil properties of the habitat may have aggravated the situation. The canopy cover was significantly low in the grassland habitat resulting in a significantly high solar radiation and wind speed. Wind speed is known to affect butterfly behavior and density, and longer flights and moves require calm conditions (Dennis and Sparks, 2006). Increased wind speed decreased butterfly density (Kuussaari *et al.*, 2016). The number of endemic butterfly species are also known to be effected by wind speed and endemic butterfly species significantly decreased in European grassland dominated habitats with high wind speed (Ozden and Hodgson, 2010). In the present study the lowest percentage of endemic butterfly species (25.81%) was evident in the grassland habitat that had a significantly high wind speed when compared with the other habitat types.

In the present study the majority of endemic species of butterflies were found from the forest habitats on either side of the buffer zone. Endemic butterflies are known to be more abundant in forest habitats preferring evergreen forests (Nunez, 2012; Padhye *et al.*, 2012); mature forests (Lewis *et al.*, 1998), rainforests (Nunez, 2012) and cloud forests (Nunez, 2012). The disturbed forest in the Seethawaka Wet Zone Botanic Gardens and undisturbed forest in the Indikadamukalana Forest Reserve of the present study were similar in many climate and soil features but differed significantly in canopy cover, solar radiation, light intensity, soil pH and soil electric conductivity. The undisturbed forest had a high canopy cover leading to low solar radiation and light intensity; and the soil electric conductivity was high and soil pH slightly acidic. High canopy cover favours regeneration of trees but negatively effects shade-intolerant plants reducing habitat heterogeneity (Wagner *et al.*, 2011). The endemic butterfly species may have preferred the more heterogenous disturbed forest with higher solar radiation and light intensity as was evident by the larger number



of endemic species residing in the habitat. Soil electric conductivity is negatively correlated with soil pH (Aini et al., 2014). Soil properties depend on soil pH and affect many plant characteristics such as height, lateral spread, flower size and number and pollen production. Most plants have an optimum for pH ranging from 5.5 to 6.5 (Gentili et al., 2018). The low pH in the undisturbed forest may adversely affect plant characteristics and thus the endemic butterfly species.

The results in this study show that the Seethawaka Wet Zone Botanic Gardens and Indikadamukalana Forest Reserve are important areas of the country supporting a rich diversity of butterflies. Species of family Nymphalidae were common to the area and all habitat types. The buffer zone with many invasive plants had the highest butterfly species richness and diversity index. The grassland habitat with a significantly low canopy cover and a significantly high light intensity had a low butterfly species richness and diversity index. The study highlights the importance of the Seethawaka Wet Zone Botanic Gardens and Indikadamukalana Forest Reserve in safeguarding and conserving diversity of butterflies of the Island at a time when habitat destruction and other such factors are threatening these charismatic insects. The study enabled to highlight spatial variations in butterflies in a managed botanic garden and an adjacent undisturbed forest reserve. A future study should also focus on examining temporal variations of butterfly communities in these two locations, in relation to climatic parameters.

#### ACKNOWLEDGEMENTS

We are thankful to the Department of Zoology and Environment Sciences, Faculty of Science, University of Colombo for providing laboratory facilities and financial assistance for the current study.

#### DECLARATION OF CONFLICT OF INTEREST

The Authors declare that there is no conflict of interest.

#### REFERENCES

- Abrahamczyk, S., Kluge, J., Gareca, Y., Reichle, S. and Kessler, M. (2011). The influence of climatic seasonality on the diversity of different tropical pollinator groups. *PLoS ONE* **6**(11): 1-9.
- Aini, I.N., Ezrin, M.H. and Aimrun, W. (2014). Relationship between soil apparent electrical conductivity and pH value of Jawa series in oil palm plantation. *Agriculture and Agricultural Science Procedia* **2**: 199-206.
- Alexandre, B., Crouzeilles, R. and Grelle, C.E.V. (2010). How can we estimate buffer zones of protected areas? A proposal using biological data. *Brazilian Journal of Nature Conservation* **8**(2): 165-170.
- Asela, M.D.C., Peiris, R.A.K., Priyankara, S.K.I.U., Jayasekara, R.W. and Karunaratna, D.M.S.S. (2009). Some notes on the butterflies (Lepidoptera: Papilionoidea) of Tantirimale Archaeological site, Anuradhapura district, Sri Lanka. *Journal of Threatened Taxa* **1**(7): 392-394.
- Baker, J., French, K. and Whelan, R.J. (2002). The edge effect and ecotonal species: bird communities across a natural edge in Southeastern Australia. *Ecology* **83**(11): 3048-3059.
- Bambaradeniya, C.N.B., Ekanayake, S.P., Fernando, R.H.S.S., Somaweera, R. and Perera, N. (2001). Biodiversity of the Bundala National Park and Ramsar Wetland. *Proceedings of the Seventh Annual Forestry and Environment Symposium 2001 of the Department of Forestry and Environmental Science, University of Sri Jayawardenepura, Sri Lanka* pp. 50.
- Bambaradeniya, C.N.B., Ekanayake, S.P., Kekulandala, L.D.C.B., Samarawickrama, V.A.P., Ratnayake, N.D. and Fernando, R.H.S.S. (2002a). An assessment of the status of biodiversity in the Muthurajawela Wetland Sanctuary. *Occasional Papers of IUCN Sri Lanka* **3**: pp.48.
- Bambaradeniya, C.N.B., Ekanayake, S.P., Kekulandala, L.D.C.B., Fernando, R.H.S.S., Samarawickrama, V.A.P. and Priyadharshana, T.G.M. (2002b). An assessment of the status of biodiversity in the Maduganga Mangrove Estuary. *Occasional Papers of IUCN Sri Lanka* **1**: pp. 49.
- Bambaradeniya, C.N.B., Perera, M.S.J., Perera, W.P.N., Wickramasinghe, L.M.J., Kekulandala, L.D.C.B., Samarawickrama, V.A.P., Fernando, R.H.S.S. and Samarawickrema, V.A.M.P.K. (2003). Composition of faunal species in the Sinharaja World Heritage Site in Sri Lanka. *The Sri Lanka Forester* **26**: 21-40.
- Bhusal, N. (2014). Buffer zone management system in protected areas of Nepal. *The Third Pole: Journal of Geography Education* **11**: 34-44.
- Bora, A., Meitei, L.R. and Deb, M. (2014). Butterfly species richness and diversity in Experimental Botanic Garden, Botanical Survey of India, ERC, Umiam, Meghalaya, India. *Journal of Entomology and Zoology Studies* **2**(5): 212-217.
- d'Abrera, B. (1998). *The butterflies of Ceylon*. Melbourne, London: Hill House Publishers.
- Carrion-Tacuri, J., Berjano, R., Guerreiro, G., Figueroa, M.E., Tye, A. and Castillo, J.M. (2012). Nectar production by invasive *Lantana camara* and endemic *L. peduncularis* in the Galapagos Islands. *Pacific Science* **66**(4): 435-445.
- Dennis, R.L.H. and Sparks, T.H. (2006). When is a habitat not a habitat? Dramatic resource use changes under differing weather conditions for the butterfly *Plebejus argus*. *Biological Conservation* **129**(3): 291-301.
- Dilts, T.E., Steele, M.O., Engler, J.D., Pelton, E.M., Jepsen, S.J., McKnight, S.J., Taylor, A.R., Fallon, C.E., Black, S.H., Cruz, E.E., Craver, D.R. and Forister, M.L. (2019). Host plants and climate structure associations of the Western Monarch Butterfly. *Frontiers in Ecology and Evolution* **7**: 1-17.
- Gentili, R., Ambrosini, R., Montagnani, C., Caronni, S. and Citterio, S. Effect of soil pH on the growth, reproductive investment and pollen allergenicity of *Ambrosia artemisiifolia* L. *Frontiers in Plant Science* **9**: 1-12.
- Ghazoul, J. (2002). Impact of logging on the richness and diversity of forest butterflies in a tropical dry forest in Thailand. *Biodiversity and Conservation* **11**: 521-541.

- Jayasinghe, H.D., De Alwis, C. and Rajapaksha, S.S. (2015). *A pocket guide to the butterflies of Sri Lanka*. 2<sup>nd</sup> Ed. Butterfly Conservation Society of Sri Lanka.
- Jayasinghe, H.D., Rajapaksha, S.S. and de Alwis, C. (2014). A compilation and analysis of food plants utilization of Sri Lankan butterfly larvae (Papilionoidea). *Taprobanica* **6**(2): 110-131.
- Kerr, J.T. (2001). Butterfly species richness patterns in Canada: Energy, heterogeneity, and the potential consequences of climate change. *Conservation Ecology* **5**(1): 1-10.
- Kitahara, M. (2004). Butterfly community composition and conservation in and around a primary woodland of Mount Fuji, central Japan. *Biodiversity & Conservation* **13**(5): 917-942.
- Koneri, R., Saroyo and Tallei, T.E. (2017). Butterfly diversity varies across habitat types in Tangkoko Nature reserve North Sulawesi, Indonesia. *Journal of Biodiversity and Environmental Sciences* **10**(4): 52-61.
- Kuussaari, M., Rytteri, S., Heikkinen, R.K., Heliola, J. and von Bagh, P. (2016). Weather explains high annual variation in butterfly dispersal. *Proceedings of the Royal Society B* **283**: 1-8.
- Lewis, O.T., Wilson, R.J. and Harper, M.C. (1998). Endemic butterflies on Grande Comore: habitat preferences and conservation priorities. *Biological Conservation* **85**(1-2): 113-121.
- Liyanagamage, M. (2016). A study of the insect assemblages of two major invasive alien plant species (*Annonaglabra*L. and *Lantana camara*L.) in the lowland wet zone in Sri Lanka. *A Dissertation submitted for the Degree of Bachelor of Science in Environment Science, Department of Zoology and Environment Sciences, University of Colombo*.
- Munyuli, M.B.T. (2013). Drivers of species richness and abundance of butterflies in coffee-banana agroforests in Uganda. *International Journal of Biodiversity Science, Ecosystem Services & Management* **9**(4): 298-310.
- Myers, M.C., Mason, J.T., Hoksich, B.J., Cambardella, C.A. and Pfrimmer, J.D. (2015). Birds and butterflies respond to soil-induced habitat heterogeneity in experimental plantings of tallgrass prairie species managed as agroenergy crops in Iowa, USA. *Journal of Applied Ecology* **52**: 1176-1187.
- Nidup, T., Dorji, T. and Tshering, U. (2014). Taxon diversity of butterflies in different habitat types in Royal Manas National Park. *Journal of Entomology and Zoology Studies* **2**(6): 292-298.
- Nisviya, S.S. and Wickramasinghe, S. (2012). Diversity and distribution pattern of butterflies at Northern flank of Knuckles region, Sri Lanka. *Proceedings of the International Forestry and Environment Symposium, Department of Forestry and Environmental Science, University of Sri Jayawardenepura, Sri Lanka*, Pp.25.
- Nunez, R. (2012). The butterflies of Turquino National Park, Sierra Maestra, Cuba (Lepidoptera, Papilionoidea). *Arxius de Miscel-laniaZoologica* **10**: 29-49.
- Ockinger, E. and Smith, H.G. (2006). Landscape composition and habitat area affects butterfly species richness in semi-natural grasslands. *Oecologia* **149**(3): 526-234.
- Ozden, O. and Hodgson, D.J. (2010). Butterflies highlight the conservation value of shrubland and grassland mosaics in cyprriotgariigieue ecosystems. *Top Biodiversity 2010, Conference Proceedings, Intercollege Larnaca, Cyprus* 67-79.
- Padhye, A., Shelke, S. and Dahanukar, N. (2012). Distribution and composition of butterfly species along the latitudinal and habitat gradients of the Western Ghats of India. *Checklist* **8**(6): 1196-1215.
- Pathiraja, G.P.K.S., Perera, H.A.S.D. and Wegiriya, H.C.E. (2017). Diversity of butterflies in different habitat types in Matara district, Sri Lanka. *Sri Lanka-Taiwan Joint Symposium on Advances in Entomological Research*, pp.4.
- Perera, M.S.J., Perera, W.P.N., Rodrigo, R.K., Ekanayake, S.P., Bambaradeniya, C.N.B., Samarawickrama, V.A.P. and Wickramasinghe, L.J.M. (2005). A biodiversity status profile of Anawilundawa Sanctuary – A Ramsar wetland in the western dry zone of Sri Lanka. *Occasional Papers of IUCN Sri Lanka* **9**: pp. 48.
- Ranwala, S.M.W., Dilrukshi, I.A.D.N., Wijesundera, D.S.A. and Attanayake, A.M.A.S. (2017). The first record on biodiversity of the natural arboretum of Seethawaka Wet Zone Botanic Gardens, Sri Lanka. *The Sri Lanka Forester* **38**: 25-45.
- Rao, R.R. and Sagar, K. (2012). Invasive alien weeds of the Western Ghats: Taxonomy and distribution. In: J.R. Bhatt (Eds), *Invasive alien plants: An ecological appraisal for the Indian Subcontinent*, Oxfordshire, UK: 139-161.
- Samarasinghe, M.D.P., Paranagama, P. and Veediyaabandara, S. (1998). Survey of the butterfly fauna of Udawalawa National Park. *Proceedings of the Second Annual Forestry Symposium, University of Sri Jayawardenepura, Sri Lanka*, pp.263-272.
- Sanjaya, Y., Suhara and Rochmayanti, Y. (2016). Role of plant diversity to existence of butterfly in botanical garden UPI Bandung Indonesia. *Journal of Entomology and Zoology Studies* **4**(4): 331-335.
- Shields, O. (1989). World numbers of butterflies. *Journal of the Lepidopterists' Society* **43**(3): 178-183.
- Slater, C., Tolley, C., Fernando, C. and Weston, M.A. (2019). A unique Lepidoptera assemblage in primary forest and understory of central Sri Lanka. *Journal of Asia-Pacific Biodiversity* **12**: 324-327.
- Smisha, K.P., Aswani, K. and Sabu, M. (2016). Reproductive biology of *Dillenia affriticosa* (Griffith) Martelli with emphasis on protandry. *Bangladesh Journal of Botany* **45**(3): 605-611.
- Sreekumar, P.G. and Balakrishnan, M. (2001). Habitat and altitude preferences of butterflies in Aralam Wildlife Sanctuary, Kerala. *Tropical Ecology* **42**(2): 277-281.
- Sundufu, A.J. and Dumbuya, R. (2008). Habitat preferences of butterflies in the Bumbuna forest, Northern Sierra Leone. *Journal of Insect Science* **8**(64): 1-17.
- Tischendorf, L. and Fahring, L. (2000). On the usage and measurement of landscape connectivity. *Oikos* **90**: 7-19.
- Van der Poorten, G.M. and Van der Poorten, N.E. (2016). The butterfly fauna of Sri Lanka. *Distribution Atlas of the Butterflies of Sri Lanka and a Field Guide* 1-30.
- Van der Poorten, N. and Van der Poorten G.M. (2016). *The*

- butterfly fauna of Sri Lanka*. Canada: Lepodon Books.
- Vu, L.V. and Vu, C.Q. (2011). Diversity patterns of butterfly communities (Lepidoptera, Papilionoidea) in different habitat types in a tropical rain forest of Southern Vietnam. *International Scholarly Research Network Zoology* 1-8.
- Vu, L.V., Bonebrake, T.C., Vu, M.Q. and Nguyen, N.T. (2015). Butterfly diversity and habitat variation in a disturbed forest in northern Vietnam. *The Pan-Pacific Entomologist* **91**(1): 29-38.
- Wagner, S., Fischer, H. and Huth, F. (2011). Canopy effects on vegetation caused by harvesting and regeneration treatments. *European Journal of Forest Research* **130** (1): 17-40.
- Woodhouse, L.G.O. (1949). *The butterfly fauna of Ceylon*. 2<sup>nd</sup> Ed. Colombo, Sri Lanka: The Colombo Apothecaries' Company Limited.
-