

Journal of Bioresource Management

Volume 8 | Issue 1

Article 13

Epidemiology and Risk Factor Analysis of Fasciolosis in Buffaloes in District Bagh, Azad Kashmir, Pakistan

Sadaf Fayyaz

Department of Zoology, Woman University Bagh, Azad Jammu and Kashmir.

Muhammad Shahbaz

Department of Zoology, Woman University Bagh, Azad Jammu and Kashmir,
shahbazchattha_pk@yahoo.com

Irfan Baboo

Department of Zoology, Cholistan University of Veterinary and Animal Sciences, Bahawalpur, Pakistan.

Shazia Nazir

Department of Live Stock, Bagh, Azad Jammu and Kashmir.

Misbah Shaukat

Department of Zoology, Woman University Bagh, Azad Jammu and Kashmir.

Follow this and additional works at: <https://corescholar.libraries.wright.edu/jbm>

 Part of the [Zooology Commons](#)

Recommended Citation

Fayyaz, S., Shahbaz, M., Baboo, I., Nazir, S., & Shaukat, M. (2021). Epidemiology and Risk Factor Analysis of Fasciolosis in Buffaloes in District Bagh, Azad Kashmir, Pakistan, *Journal of Bioresource Management*, 8(1).

DOI: <https://doi.org/10.35691/JBM.1202.0172>

ISSN: 2309-3854 online

(Received: Jan 1, 2021; Accepted: Jan 18, 2021; Published: Mar 15, 2021)

This Article is brought to you for free and open access by CORE Scholar. It has been accepted for inclusion in *Journal of Bioresource Management* by an authorized editor of CORE Scholar. For more information, please contact library-corescholar@wright.edu.

Epidemiology and Risk Factor Analysis of Fasciolosis in Buffaloes in District Bagh, Azad Kashmir, Pakistan

© Copyrights of all the papers published in Journal of Bioresource Management are with its publisher, Center for Bioresource Research (CBR) Islamabad, Pakistan. This permits anyone to copy, redistribute, remix, transmit and adapt the work for non-commercial purposes provided the original work and source is appropriately cited. Journal of Bioresource Management does not grant you any other rights in relation to this website or the material on this website. In other words, all other rights are reserved. For the avoidance of doubt, you must not adapt, edit, change, transform, publish, republish, distribute, redistribute, broadcast, rebroadcast or show or play in public this website or the material on this website (in any form or media) without appropriately and conspicuously citing the original work and source or Journal of Bioresource Management's prior written permission.

EPIDEMIOLOGY AND RISK FACTOR ANALYSIS OF FASCIOSIS IN BUFFALOES IN DISTRICT BAGH, AZAD KASHMIR, PAKISTAN

SADAF FAYYAZ¹, MUHAMMAD SHAHBAZ¹, IRFAN BABOO², SHAZIA NAZIR³, MISBAH SHAUKAT¹

¹Department of Zoology, Woman University Bagh, Azad Jammu and Kashmir.

²Department of Zoology, Cholistan University of Veterinary and Animal Sciences, Bahawalpur, Pakistan

³Department of Live Stock, Bagh, Azad Jammu and Kashmir.

⁴Department of Wildlife and Ecology, University of Veterinary and Animal Sciences Lahore, Pakistan.

Corresponding author's email: shahbazchattha_pk@yahoo.com

ABSTRACT

Fasciolosis is a cosmopolitan parasitic disorder cause great economic decline in cattle and buffaloes in terms of poor carcasses quality decrease in meat and milk production, malfunctioning of livers and weight loss. This research work was carried out in district Bagh, Azad Jammu and Kashmir, Pakistan, extending from December 2016 to May 2017. During this study 200 fecal sample from Buffaloes (*Babylus bubalis*) of the study area were collected, it was found that 40 percent fecal samples contain eggs of *Fasciola hepatica* and 60 percent fecal samples were negative. Liver samples of slaughtered buffaloes were also examined during study period and it was found that 74 samples have faciola, out of 200 samples, so 37 per cent samples contain *F. hepatica* and 63 per cent buffaloes liver samples were negative. It was observed that 19.25% animals of the area have parasites in their body, which are causing economic losses to the owners, in the form of less production of milk and meat and more mortality among the livestock and damages to farmers.

Keywords: Epidemiology, buffaloes, faciola, livestock, fasciolosis.

INTRODUCTION

Fasciolosis, a cosmopolitan parasitic disorder responsible for great economic decline in cattle and buffaloes in terms of poor carcasses quality, decrease in meat and milk production, malfunctioning of livers and weight loss (Maqbool et al., 2002). This disease cause major losses in terms of mortality and morbidity in fluky regions. Data collected from some countries of Asia has displayed that buffaloes suffer more frequently from Fasciolosis, than any other parasitic disease (Balasingam, 1962). In the world among all parasitic disorders of ruminants fascioliasis is one of the most common (Massoud et al., 2012). In advanced regions of the world data on epidemiology of several helminthiasis has been

circulated and remedial measures have been taken in these areas for the control of this disease but in developing countries little research work has been done on Fasciolosis (Maqbool et al., 2002). Two liver flukes commonly reported to causes Fasciolosis in ruminants are *F. hepatica* and *F. gigantica*. The life cycle of these trematodes involves snail as an intermediate host. Infected cattle show poor weight increase and dairy cattle have lesser milk yield (Mihreteab et al., 2010). The geographical distribution of trematode species is dependent on the distribution of suitable species of snails (Rahmeto, 2010). Symptoms of Fasciolosis in buffaloes will be imminent after eighty four days eggs will be found in fecal samples have infection after thirteen to fourteen weeks for *F. hepatica* and *F. gigantica*

respectively by shedding eggs of flukes in excreta and up to this time most part of liver may be damaged.

The disease is responsible for obvious financial drop in livestock by declining milk production, weight gain, liver condemnation, fertility and disease is very commonly found in rainy areas of maximum parts of the world and absent where environmental circumstances are critical for the growth of secondary host (Mazhar, 2005). Fasciolosis is the major cause of economic losses in parasitized groups because of limited yield of meat and milk, mortality, abortions and rejection of livers in slaughterhouses, growth retardation in young animals and expenditure on monitoring of this disease. Alves et al. 2011 described that in Brazil, the growing topographical distribution of *F. hepatica* is due to transfer and trade of animals (parasitized) from enzootic disease areas where the localities are not affected but have satisfactory epidemiological surroundings. The encouraging conditions of area are suitable and ensure occurrence of snails (genus *Lymnae*) which are intermediate hosts for *F. hepatica* (Martins et al., 2012). Thus these situations such as adequate and high humidity temperature and rainfall also include the environmental and climatic features that deliver a favorable habitation for these snails, Feasibility of eggs upcoming from naturally infected buffaloes and cattle liability of *L. columella* collected and studied on the farmhouses, to infection by liver fluke (Dracz and Lima, 2014).

Major source of animal protein is derived from Meat of goats, cattle and sheep. There are various species of these but the economically important ones in tropics are *F. gigantica* and in the temperate region is *F. hepatica* (Usip et al., 2014). Numeral surveys have been done by veterinary research groups concerning the occurrence of the disease and studied in many parts of the world (Ozung et al., 2011). The climatic changes that have

occurred in Bungoma in the recent past such as prolonged season of rains may have had an effect on the prevalence and economic importance of Fasciolosis on the rural population (Usip et al., 2014). For great milk yield among the livestock buffalo is thought to be an important kind. Total contribution of milk in country due to buffalo is seventy two percent (Mahadevan, 1978). Buffaloes are subject to the number of significant parasitic diseases. Field reports revealed that more than 27,000 buffaloes lost every year because of different diseases (Junejo and Qureshi, 1993). Milk is mainly produced by the smallholders in Pakistan (Afzal, 2010). Chaudhry et al. 1984 reported that in the alimentary tract, due to combination of many species of helminthic parasites gastrointestinal helminthiasis ailment is caused. Irfan observed in 1984 that in Pakistan where the sanitary situation is very low, the massive financial damages in livestock industry is because of parasites. In Pakistan, Fasciolosis is one of the major factor that is responsible for reduced livestock development (Kendall, 1954).

Babylus bubalis (Asian buffalo) belongs to the class Mammalia (Hussain et al., 2010; Afzal, 2010) as compared to other domesticated bovines species well thought out to be an advantageous animal, and agonized a huge number of internal and external parasitic bacterial, fungal and viral diseases (Raza et al., 2010; Mahfooz et al., 2008).

Helminthiasis, among all parasitic infections has long been documented and still is a criminal principal to deaths in ruminant growth in all across the globe (Alawa et al., 2010). Fascioliasis caused by genus *Fasciola*, is economically most important helminthes ailments of the farm-ruminants (Santos et al., 2015). It is an emerging parasitic pollution, impacts threateningly on both human health and veterinary worldwide (Lazara et al., 2010). It roots huge economic losses in terms of decrease in meat, milk and higher rate of

deaths in all forms of animals (Saleha, 1991).

Two extremely infectious species are identified as and *F. gigantica* and *F. hepatica* first species is normally prevailing in humid regions of various states of the biosphere while the other one endures in a diversity of climatic circumstances (Urquhart et al., 1988). These parasites mainly cause infection in the liver, where they exist in bile duct and hepatic parenchyma and lesion on mucosa and subsequent huge tissue damage (Shaikh et al., 2004, 2005). Proofs indicated that most deaths in bovine is because of fascioliasis (Losos, 1986).

B. bubalis, meat and the prime dairy producing animal not filling the demand and supply gap of milk and meat of the Pakistani people is due to variety of diseases being faced by domesticated animals (Bhutto et al., 2012). Universally, above seven hundred million household ruminants are at threat and economic loss go above 3 billion US dollar per year. Peoples do not gave due concentration to the Fasciolosis in the human population in tropical regions of the world where more than 50 million of the population has been affected. The prevalence rates in Bangladesh in live animals has been reported to vary from 19 to 51% in 8.4 to 31% in sheep (Sangma et al., 2012), buffaloes (Roy, 2016), 10 to 32% in goats (Sammadar et al., 2015) and 21 to 53% in cattle (Islam et al., 2015). In human fascioliasis was traditionally considered as a secondary disease. The explanation of many human endemic regions and the emergence of human reports in many countries, including both prevalence and intensity increases and topographical expansion, have drastically changed the global scenario from the middle of 1990s. Fascioliasis is at present known to be the vector-borne disease presenting the altitudinal, widest latitudinal, longitudinal and distribution known (McAllister and Sultana, 2011).

The global estimation of seventeen million people infected may be an underestimation due to the situation in many countries was previously not known (Mas-Coma et al., 2012). In temperate climatic zones, *F. hepatica* is tremendous significant parasitic of cattle (Bennema et al., 2011). Heavy infection was found in buffaloes as compared to cow and goat (Swarnaka and Sanger, 2014). Production of bovines yield about the earth is majorly affected by trematode invasion (Vercruyssen and Claerebout, 2001). But Fasciolosis increases public to fear not only due to its zoonotic feature. But also its frequency and economic implication to live stock in many regions of the world Ruminants' Fasciolosis is an inhibition in beneficial bovine-farming and for slaughterers and customers too. Parasite of genus *Fasciola* i.e. *F. gigantica* and *F. hepatica* is the causal agent of disease (Fasciolosis) which occur in a wide range of ultimate hosts. It is encouraged on by both environmental changes (warmer, wetter climate) and man-made changes such as an increase in animal travels and growth of livestock farming (Gul, 2017).

MATERIALS AND METHOD

Geographical area of Study

The State Azad Jammu and Kashmir consist of 10 districts and Bagh is one of them. Climate of Bagh has been categorized as warm and temperate. In Bagh the normal annual temperature is 17.3°C. My study area was Bagh.

Fecal Samples

200 stool samples of buffaloes were collected from different localities of city Bagh (During December, 2016 to May, 2017) for the determination of liver fluke eggs, stool samples were taken with the help of forceps. These samples were packed into labelled polythene bags and transferred to the lab. All the collected

samples were examined within 12 hours (Yildirim et al., 2007).

The presence of *Fasciola* was determined in these collected samples by following the sedimentation and floating technique (Urquhart et al., 1988). One gram of stool sample was taken and mixed with water (15 to 20ml) to make solution in cylinder and stir well with glass rod and strained with the help of strainer then put a drop of strained solution on the slide with the help of dropper and place the cover slip on the glass slide. This slide was examined under microscope for the presence of *F. hepatica*. The eggs were counted in each slide. Mask and gloves were used as a hygienic measure.

Liver Samples

A total of 200 samples of liver were collected of slaughtered buffaloes from different slaughter houses of the study area located in Bagh. The liver of freshly slaughtered domestic buffaloes were collected for the study. Firstly the liver was collected and then it was observed keenly, in the beginning, the bile ducts were opened for obtaining flukes. For generalized liver fluke infections incisions were made in different parts of

the bile duct to find the fluke in the liver, gall bladder was also incised and then bile duct opened, first common bile was observed and then smaller ones. Flukes were with care taken up with the help of forceps and needles and thoroughly washed with tap water and then these flukes transferred in saline solution and observed with the help of hand lens.

RESULTS AND DISCUSSION

During the present study of Fasciolosis in buffaloes of district Bagh, extending from December 2016 to May 2017. The presence of *Fasciola* was determined in these collected samples by following the sedimentation and floating technique (Urquhart et al., 1988). Among the collected samples 120 were found negative and 80 were positive having eggs.

During the present study liver samples were also collected from freshly slaughtered animals and these were examined for the presence or absence of *Fasciola* species. Among these 200 samples *F. hepatica* was found in seventy four samples. All the collected samples were examined within 12 hours (Yildirim et al., 2007). Overall prevalence of both liver and fecal samples is 19.25 percent.

Table 1: Data collected through fecal examination.

Sr. No.	Place	Total	Male	Female	Hypher	Non Hypher	Fasoiola Present	Fasiola Absent
1	Arja	25	1	24	2	23	8	17
2	Dhuli	30	2	28	3	25	13	17
3	Baglor	10	2	8	3	7	3	7
4	Chatter	13	0	13	1	12	5	8
5	Hular	20	1	19	1	18	8	12
6	Neela butt	20	0	20	0	20	8	12
7	Dheerkot	25	0	25	0	25	9	16
8	Chamyati	20	1	19	0	20	9	11
9	Jhala	7	0	7	0	7	3	4
10	Mallot	17	0	17	4	13	9	8
11	Bypass	13	2	11	3	10	5	8

Table 2: The percentage of *F. hepatica* in fecal samples collected from Bagh.

Place	Fasciola present	Percentage
Arja	8	32%
Dhuli	13	43%
Baglor	3	30%
Chatter	5	38.46%
Hular	8	40%
Neela butt	8	40%
Dheerkot	9	36%
Chamyati	9	45%
Jhlala	3	42.85%
Mallot	9	53%
Bypass	5	38.46%

Table 3: The percentage of *F. hepatica* collected from slaughter houses.

Month	Total	Male	Female	Hypher	Non Hypher	Fasciola Present	Fasciola Absent
December	30	2	28	2	28	13	17
January	40	4	36	6	34	19	21
February	32	2	30	2	30	12	20
March	35	3	32	3	32	11	24
April	33	2	31	3	30	9	24
May	30	1	29	3	27	10	20

Table 4: The percentage collected through monthly basis.

Month	Fasciola present	Percentage
December	13	56.66%
January	19	47.5%
February	12	37.5%
March	11	31.42%
April	9	27.7%
May	10	33.33%
Total	74	100




Figure 1 (A, B): *F. hepatica* collected from liver of slaughtered animals.

During the present study period, 200 samples of liver were collected from the freshly slaughtered animals (figures 2&3) and 200 stool samples from house hold buffaloes from December 2016 to May 2017. Among these 200 fecal samples, 80 buffaloes fecal samples were found positive for liver fluke, prevalence rate of Fasciolosis is 40%. It is found that Fasciolosis is caused by *F. hepatica* in the study area. This study shows that domestic ruminants are highly infected with *F. hepatica*.




Figure 2: The percentage of fecal sample collected from Bagh.




Figure 3: The percentage of *f. hepatica* from liver of slaughtered animals collected on monthly basis.

Table (1) reveals that *F. hepatica* is most prevalent in buffaloes of the study area. Fecal samples were collected from different places of Bagh like Arja, Dhuli, Baglor, Chatter, Hular, Neela Butt, Dheerkot, Chamtyati, Jhala, Mallot and Bypass. From Arja 25 fecal samples (non hypher 23, hypher 2) during the examination of these fecal samples it was found that Fasciola is present only in eight samples; 30 fecal samples were collected from area of Dhulli and it was found that 13 buffaloes were infected with liver fluke while 17 were not affected; 3/10 buffaloes

of Baglor were found positive, having *Fasciola* eggs in their excreta; From Chatter 13 fecal samples were examined and in 5 samples *Fasciola* was found and 8 samples were free of *Fasciola*; From Hular 20 fecal samples collected, *Fasciola* was present in 8 samples and absent in 12; twenty fecal samples were collected from Neelabutt and it was found that eight samples were positive for liver fluke and twelve were negative; From Chamtyati 20 samples were examined and only nine found positive; from Jhala 7 fecal samples were collected and found three positive for *Fasciola*; From Mallot 17 fecal samples of buffaloes were collected and 9 positive. Total 13 fecal samples from Bypass, 5 were positive for *Fasciola* and 8 were free. Out of 200 buffaloes 80 were found infected by *F. hepatica* and prevalence rate is 40%, eggs of liver flukes were found in only 80 samples out of 200 and it is close to the results of Pfukenyi and Mukaratirwa, who reported 37.1% in Zimbabwe and Abraham and Jude found the prevalence rate 44.8% in Nigeria. So prevalence of liver fluke was thus higher in domestic ruminants. (Maqbool et al., 2002; Aliyu et al., 2014). 200 liver samples were also examined for the determination of rate Fasciolosis in buffaloes, only 74 samples found positive during the present study, showing overall 37%. For the determination of Fasciolosis prevalence in the animals of the study area, liver samples of slaughtered animals were collected from December to May, 2017. Month wise Prevalence rates from December to May 2017 are as 43.33%, 47.55%, 37.5%, 31.42%, 27.27% and 33.33% respectively. The highest prevalence rate is in the month of January, 2017 and lowest in the month of April, 2017. This variation may be due to deworming of animals in different months; highest rate in the month of January, 2017 may be due to lowest cold temperature and less availability of food, weakness of animals and immune system. May be due to non-drenching of in the month of

December and January. These are assumptions for the determination of accurate reason more detailed studied is needed. Carneiro et al., in 2010 reported natural infection by *F. hepatica* in buffaloes for the first time in the south of the state of Espirito Santo, with a positivity rate of 46.67 % among 15 samples. Higher incidences of fascioliasis have been recorded by Pfukenyi and Mukaratirwa, who reported 37.1 % in Zimbabwe and Abraham and Jude recorded 44.8 % in Nigeria.

Marques and Scroferneker conducted a study in 2003 in the state of Rio Grande do Sul and prevalence rate recorded by them was 10.34 % among 377 livers samples of cattle and 20 % in 105 livers samples of buffaloes. The result of the fecal samples are 37 and 40 % liver samples percent which is consistent with the result of Pfukenyi and Mukaratirwa 37.1 % in Zimbabwe, Abraham and Jude 44.8 % in Nigeria.

Alemu et al., (2015) described that Southern Ethiopia using post-mortem examination of liver of each slaughtered animal shows that A total of 500 cattle were examined for the presence of Fasciolosis using both coprological and post mortem examination .The prevalence under coprological and post mortem examinations were found to be 19.4% and 28.6% respectively. This result is lower than my results. Carneiro et al., studied in 2010 that the Fasciolosis in buffaloes of south state of Espirito Santo and found the infection rate 46.6% which is slightly higher to my results. Marques and Scroferneker 2003 carried out in study in the state of Rio Grand Sul and found Fasciolosis rate in fecal and liver samples 10.34% and 20% respectively which is not consistent to my results and manifold lower than my results. Rahman and Mondol in 1983 collected the 762 fecal samples in Bangladesh and found infection rate 66.14% which is higher than my results.

Karim, et al., (2015) in Bangladesh examined 762 fecal samples and found 504 (66.14%) positive with *Fasciola* species. Rahman and Mondol in 1983 and Gupta et al. in 2002 recorded 53 and 70% Fascioliasis in cattle in Bangladesh and India, respectively. Gupta et al. 2002 collected samples and found the prevalence rate 53% which is slightly higher than my results and 70% is too much higher than my findings. Epidemiological studies were undertaken by Maqbool et al., 2002 explained that in Punjab Lahore at slaughter houses and on household buffaloes under the different climatic conditions existing in Punjab province. Infection rate was 25.59 and 10.5 percent, respectively in slaughtered buffaloes and house hold buffaloes. These results are lower than my results.

CONCLUSION

A study was conducted in districts Poonch and Mirpur, Azad Jammu and Kashmir by Ahmad et al., in 2017. They study the prevalence of Fasciolosis in buffaloes from 852 samples, 106 (12.44%) were found positive. The findings of this study are much lower than my results. The variations in the prevalence of fascioliasis may be attributed to the meteorological factors and grazing habits of the host (Shaikh et al., 2004). The inconsistency of my results with the neighboring areas studies may be due to difference of topographical variations.

REFERANCES

- Afzal M (2010). Re-designing smallholder dairy production in Pakistan. *Pak Vet J.*, 30(3): 187-190.
- Ahmad I, Durrani AZ, Ashraf K, Hameed K, Khan O, Abbasi AA, Safdar A (2017). Risk factors analysis of Fasciolosis in buffaloes in Azad Jammu and Kashmir. *Pak J Sci.*, 69(4): 351-354.
- Ahmed S, Nawaz M, Gul R, Zakir RM, Razzaq A (2005). Diversity and prevalence of trematodes in livers of sheep and goat in Quetta, Pakistan. *Pak J Zoo.*, 37(3): 205.
- Alawa CBI, Adamu AM, Gefu JO, Ajanusi OJ, Abdu PA, Chiezey NP (2010). In vivo efficacy of *Vernonia amygdalina* (compositae) against natural helminth infection in Bunaji 360 section in Bunaji (*Bos indicus*) calves. *Pak Vet J.*, 30(4): 215–218.
- Alemu, Abebe B (2015). Fasciolosis: Prevalence, Evaluation of Floatation and Simple Sedimentation Diagnostic Techniques and Monetary Loss due to Liver Condemnation in Cattle Slaughtered at Wolaita Soddo Municipal Abattoir, Southern Ethiopia. *Food Sci Qualit Manag.*, 43(1): 84-98.
- Alves DP, Carneiro MB, Martins IVF, Bernardo CC, Donatele DM, Pereira OS, Almeida BR, Avelar BR, Leao AGC (2011). Distribution and factors associated with *F. hepatica* infection in cattle in the south of Espírito Santo State, Brzil *J Venom Anim Txins.*, 17(3): 271-276.
- Balasingam E (1962). Studies on fascioliasis of cattle and buffaloes in Singapore due to *Fasciola gigantica* Cobbold. *Ceylon Vet J.*, 10(1): 10-29.
- Carneiro MB, Bernardo CC, Jr. Calais A, Alves DP, Jr Pereira OS, Martins IVF (2010). *F. hepatica* em búfalos (*Bubalus bubalis*) no sul do Espírito Santo. *Rev Bras Med.*, 32(2): 89-91.
- Chaudhry NI, Durani MS, Aziz T (1984). Incidence of gastrointestinal parasite in buffalo and cattle of Azad Kashmir. *Pak Vet J.*, 4(1): 60-61.
- Furst T, Keiser J, Utzinger J (2012). Global burden of human food-borne trematodiasis: a

- systematic review and meta-analysis. *Lancet Infect Dis.*, 12(3): 210-221.
- Gul N, Tak H, Fazilli KM, Abdullah I, Sofi TA (2017). Prevalence of Fasciola Infection in Slaughtered Animals in Kashmir. *Glob J Med Res Herid.*, 72(3): 183-188.
- Gupta SC, Singh BP (2002). Fasciolosis in cattle and buffaloes in India. *J of parasitol.*, 16(2): 139-145.
- Islam MH, RN (2015). Ripa. Prevalence of fascioliasis in slaughtered goat in Bengal meat abattoir house and its economic impact on business. *J Chem Bio Phys Sci.*, 5(3): 2684.
- Jabbar M, Green DAG (1982). The status and potential of livestock within the context of agricultural development policy in Bangladesh (No. 183890). Bangladesh Agricultural University, Mymensingh, Department of Agricultural Economics.
- Kakar MN, Kakar JK, Sleman K (2008). Prevalence of endo (trematodes) and ecto-parasites in cows and buffaloes of Quetta, Pakistan. *Pak Vet J.*, 28(1): 34-36.
- Karim MR, Mahmud MS, Giasuddin M (2015). Epidemiological study of bovine Fasciolosis: prevalence and risk factor assessment at Shahjadpur upazila of Bangladesh. *Immunol Infect Dis.*, 3(3): 25-29.
- Kendall SB (1954). Fascioliasis in Pakistan, *Ann Trop Med Parasitol.*, 48(3): 307-313.
- Lazara RA, Vazquez I, Domenech, Robertson LJ (2010). Fascioliasis: can Cuba conquer this emerging parasitosis? *Trends Parasitol.*, 26(1): 26-34.
- Losos GJ (1986). Fascioliasis (*Fasciola Gigantic*): in *Infectious Tropical Diseases of Domestic Animals*, 1st edition. Churchill Livingstone Inc New York USA. 1st ed., 882-902.
- Mahadevan P (1978). Water buffalo research: possible future trends. *World Animal Review*.
- Mahfooz A, Masood MZ, Yousaf A, Akhtar N, Zafar MA (2008). Prevalence and anthelmintic efficacy of Abamectin against gastrointestinal parasites in horses. *Pak Vet J.*, 28(2): 76-78.
- Marques SMT, Scroferneker ML (2003). *F. hepatica* infection in cattle and buffaloes in the State of Rio Grande do Sul, Brazil *Parasitol Latin.*, 58(3): 169-172.
- Mas-Coma MS, Esteban JG, Bargues MD (1999). Epidemiology of human fascioliasis: a review and proposed new classification. *Bull WHO.*, 77(4): 340.
- Mas-Coma S, Bargues MD, Valero MA (2005). Fascioliasis and other plant-borne trematode zoonoses. *Int J parasitol.*, 35(11): 1255-1278.
- Massoud AM, Shalaby HA, El Khateeb RM, Mahmoud MS, Kutkat MA (2012). Effects of Mirazid and myrrh volatile oil on adult *Fasciola gigantica* under laboratory conditions. *Asn Pac J trop Biomed.*, 2(11): 875.
- Mazhar D, Gillmore R, Waxman J (2005). *Cox can Qjm.*, 98(10): 711-718.
- Mir MR, Chishti MZ, Rashid M, Dar SA, Katoch R, Mehraj M, Dar MA, Rasool R (2013). The epidemiology of caprine Fascioliasis in Jammu (J and K)-India. *Int J food Agri Vet Sci.*, 3(1): 233-237.
- Ozung PO, Owai PU, Oni KO (2011). Assessment of the prevalence of fascioliasis of ruminants in Ikom abattoir of cross river state, Nigeria. *Conti J Vet Sci.*, 5(1): 1-5,
- Rahman, Mondal MMH (1983). Helminth parasites of cattle (*Bos indices*) in Bangladesh. *Ind J Parasitol.*, 7(2): 173-174.

- Rahmeto A, Fufa A, Mulugeta B, Solomon M, Bekele M, Alemayehu R (2010). Fasciolosis: Prevalence, financial losses due to liver condemnation and evaluation of a simple sedimentation diagnostic technique in cattle slaughtered at Hawassa Municipal abattoir, southern Ethiopia Hawassa University, Faculty of Vet. Med., Hawassa, Ethiopia. *Ethio Vet J.*, 14 (1): 39-51.
- Raza MA, Murtaza S, Bachaya HA, Qayyum A, Zaman MA (2010). Point prevalence of *Toxocara vitulorum* in large ruminants slaughtered at Multan abattoir. *Pak Vet J.*, 30: 242-244.
- Roy PP, Begum N, Dey AR, Sarker S, Biswas H, Farjana T (2016). Prevalence of gastrointestinal parasites of buffalo at Mongla, Bagerhat. *Int J Nat Soci Sci.*, 3: 59-66.
- Saleha AA (1991). Liver fluke disease (fascioliasis) epidemiology, economic impact and public health significance. *S east Asi J Trop Med Pub Helth.*, 22(4): 361.
- Santos CS, Scherer PO, Vasconcellos MC, Pile EM, Freire LS, Santos JAA, Freire NMS (2015). Registro de *Fasciola hepatica* em eqüinos (*Equus caballus*), caprinos (*Capra hircus*) e ovinos (*Ovis aries*) no município deltaguaí, Rio de Janeiro. *Bra Rev Bra Cin Vet.*, 7(1): 63-64.
- Shaikh AA, Bilqees FM, Khan MM (2004). Bile duct hyperplasia and associated abnormalities in the buffaloes infected with *Fasciola gigantica*. *Pak J Zool.*, 36(3): 231-238.
- Sujon MA, Mostofa M, Jahan MS, Das DR, Rob S (2008). Studies on medicinal plants against gastrointestinal nematodes of goats. *Bangla J Vet Med.*, 6(2): 179-183.
- Tongson LL, Biggers JV, Dayton GO, Bind JM, Knox BE (1978). Surface analysis of WC-Co composite materials (2) Quantitative Auger electron spectrometry. *J Vac Sci Tec.*, 15(3): 1133-1138.
- Urquhart GM, Armour J, Duncan JL, Dunn AM, Jennings FW (1988). *Veterinary Parasitology*, 1st edition. Longman Scientific and Technical, Longman Group UK Ltd., England.
- Usip LPE, Ibanga ES, Edoho HJ, Amadi EC, Utah E (2014). Prevalence of Fascioliasis and the economic loss of condemned liver due to *Fasciola* infection in Cattle slaughtered at three abattoirs in Eket Urban, Akwa Ibom State of Nigeria. *Global Advanced Resh. J Food Sci Technol.*, 3(2): 54-75.