



# Myeloid-Derived Suppressor Cells as a Potential Biomarker and Therapeutic Target in COVID-19

Marianna Rowlands<sup>1</sup>, Florencia Segal<sup>1</sup> and Dominik Hartl<sup>2,3\*</sup>

<sup>1</sup> Novartis Institutes for BioMedical Research (NIBR) Translational Medicine, Cambridge, MA, United States, <sup>2</sup> Novartis Institutes for BioMedical Research (NIBR), Translational Medicine, Basel, Switzerland, <sup>3</sup> Department of Pediatrics I, University of Tübingen, Tübingen, Germany

Clinical presentations of COVID-19 are highly variable, yet the precise mechanisms that govern the pathophysiology of different disease courses remain poorly defined. Across the spectrum of disease severity, COVID-19 impairs both innate and adaptive host immune responses by activating innate immune cell recruitment, while resulting in low lymphocyte counts. Recently, several reports have shown that patients with severe COVID-19 exhibit a dysregulated myeloid cell compartment, with increased myeloid-derived suppressor cells (MDSCs) correlating with disease severity. MDSCs, in turn, promote virus survival by suppressing T-cell responses and driving a highly pro-inflammatory state through the secretion of various mediators of immune activation. Here, we summarize the evidence on MDSCs and myeloid cell dysregulation in COVID-19 infection and discuss the potential of MDSCs as biomarkers and therapeutic targets in COVID-19 pneumonia and associated disease.

**Keywords:** COVID-19, immunology, immunity, MDSC, biomarkers

## OPEN ACCESS

### Edited by:

Dragana Odobasic,  
Monash University, Australia

### Reviewed by:

Girdhari Lal,  
National Centre for Cell Science, India  
Ishak Ozel Tekin,  
Bülent Ecevit University, Turkey

### \*Correspondence:

Dominik Hartl  
[dominik.hartl@med.uni-tuebingen.de](mailto:dominik.hartl@med.uni-tuebingen.de)

### Specialty section:

This article was submitted to  
Molecular Innate Immunity,  
a section of the journal  
Frontiers in Immunology

**Received:** 19 April 2021

**Accepted:** 07 June 2021

**Published:** 18 June 2021

### Citation:

Rowlands M, Segal F and Hartl D  
(2021) Myeloid-Derived Suppressor  
Cells as a Potential Biomarker and  
Therapeutic Target in COVID-19.  
*Front. Immunol.* 12:697405.  
doi: 10.3389/fimmu.2021.697405

## INTRODUCTION

It has been more than a year since the initial reports of an outbreak of pneumonia in the Hubei province of China, and the subsequent identification of a novel *betacoronavirus* severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection as the cause for the coronavirus disease 2019 (COVID-19) (1). During this time, despite global efforts for containment and the declaration of a pandemic on March 2020, there have been more than 126 million confirmed cases of COVID-19 worldwide, and over 3.5 million deaths reported to the World Health Organization (2).

Patients with SARS-CoV-2 infection can experience a range of clinical manifestations, from no symptoms to severe pneumonia, respiratory and/or multiple organ failure (1, 3, 4). Increasing evidence suggests that the immune response to SARS-CoV2 plays a critical role in the pathogenesis of COVID-19 disease. On one end of the spectrum, SARS-CoV-2 can disrupt normal immune responses, resulting in uncontrolled inflammation in severe and critical patients with COVID-19 (5). Specifically, lung infiltration and activation of pro-inflammatory myeloid cells such as monocytes, macrophages and neutrophils, is thought to play a key role in the cytokine storm syndrome and the hyper-inflammatory response observed in severe cases (6–8). On the other hand,

adaptive immune responses elicited by emerging COVID-19 vaccines have shown to be highly protective against severe disease and mortality (9).

Understanding the immunopathology of SARS-CoV-2 can be harnessed for the identification of novel biomarkers for disease progression, as well as potential therapeutic targets for COVID-19. In this review, we summarize the characteristics of COVID-19 related dysregulation of the myeloid cell compartment, and discuss their potential use as biomarkers and future targets for therapeutic intervention.

## MDSC DEFINITION AND FUNCTIONALITY

Myeloid-derived suppressor cells (MDSCs) are defined as innate bone-marrow-derived immune cells suppressing effector T cell responses (10). MDSCs are a heterogeneous population mainly composed of two distinct subtypes, neutrophilic/granulocytic MDSCs (PMN-/G-MDSCs) and monocytic MDSCs (M-MDSCs) (10, 11). Differences compared to terminally differentiated granulocytes and monocytes respectively have been previously described in detail (11) but also key differences are summarized in **Table 1**. While initial investigations focused on T cells as targets of MDSC-mediated suppression, subsequent studies expanded this concept by showing that MDSCs are able to regulate a broad variety of adaptive (T cells, B cells) and innate (natural killer cells, macrophages, dendritic cells) immune cells (12, 13). Beyond dampening immune cell functionalities, MDSCs were further found to promote the development of regulatory T cells (14) and regulatory B cells (15). The effector mechanisms employed by MDSCs to control immune cell subsets depend on the MDSC subtype with PMN-MDSCs mainly use reactive oxygen species (ROS) and arginase I, whereas M-MDSCs use inducible nitric oxide synthase (iNOS) and arginase I to dampen bystander cells. In addition to these major suppressive mechanisms, other immuno-modulatory MDSC functions have been reported, including secretion of anti-inflammatory mediators such as interleukin-10 (IL-10), transforming growth factor beta (TGF- $\beta$ ) or Prostaglandin E2 (PGE2) or the

tryptophan/kynurenine pathway through Indoleamine 2,3 dioxygenase (IDO) (12, 13, 16).

While MDSCs were discovered in malignant diseases and the majority of studies assessed MDSCs in cancer conditions, emerging evidence shows that MDSCs are more diverse and are involved in inflammation, autoimmunity and infection (17). The factors inducing MDSC accumulation and suppressive function in these disease contexts remain only partially defined, but probably include a variety of microenvironmental factors including hypoxia, Granulocyte- and Granulocyte-Macrophage Colony Stimulating Factor (G-/GM-CSF), IL-6, Tumor necrosis factor alpha (TNF- $\alpha$ ), Vascular Endothelial Growth Factor (VEGF), IL-1 $\beta$  and other cytokines, chemokines, damage-associated molecular patterns (DAMPs) and alarmins, such as High mobility group box protein 1 (HMGB1) or S100A8/9 (Calprotectin), checkpoint regulators, such as PD-1/PD-L1 and pathogen-associated molecular patterns (PAMPs), such as flagellin (12, 13, 16, 18, 19). At the transcriptional level, signaling through the transcription factors signal transducer and activator of transcription 3 (STAT3) and STAT5 are key for MDSC expansion (18) (**Figure 1**) whereas transcriptional regulators such as the Inhibitor of Differentiation 1 (Id1) have also been implicated in MDSC expansion (20).

MDSCs were studied in bacterial, viral, parasitic and fungal infections (21). Most so far studied infectious disease settings provided evidence for the accumulation of MDSCs in the peripheral blood and/or the affected tissue, yet the functional role of MDSCs has been reported as both detrimental by downregulating host defense or beneficial by dampening excessive infection-associated inflammation (or other less-defined mechanisms). In some infection models, the protective *vs.* harmful role of MDSCs is rather complex and depends on the animal species/model system used, the stage of disease, and the ratio/balance between pathogens, T cells and MDSC (21–23). Another layer of complexity in host-pathogen interactions is added by the fact that MDSCs as phagocytes can directly act antimicrobial or can be exploited by intracellular pathogens as survival niche. With regards to viral infections, most evidence for MDSC involvement exists for hepatitis B/C (HBV/HCV), human immunodeficiency virus-1 (HIV-1), herpes simplex virus (HSV) or influenza with indications that chronic, rather than acute viral infections induce MDSC expansion (24–26). Intriguingly, MDSC expansion in chronic HCV infection was shown to favor viral persistence (27), whereas in HBV infection MDSC were linked to a protective role by ameliorating hepatic tissue damage (28). In HIV infections, high numbers of MDSC were reported that correlated positively with viral loads, negatively with CD4<sup>+</sup> T cell numbers and dropped upon antiviral therapy (29).


## IMMUNOLOGICAL ASPECTS OF COVID-19

Despite a rapidly increasing number of publications on COVID immunopathogenesis, the precise mechanisms that govern the pathophysiology of the different disease courses of COVID-19 remain poorly defined owing to the complex multi-organ, co-

**TABLE 1** | Differences between MDSCs and other myeloid cells\*.

|  |   |
|--|---|
| Proposed differences between monocytes and M-MDSC                  | <ul style="list-style-type: none"> <li>• Suppressive function</li> <li>• Reduced HLA-DR expression</li> <li>• Increased iNOS expression</li> <li>• Increased ARG1 expression</li> <li>• Increased pSTAT3 expression</li> <li>• Reduced IRF8 expression</li> <li>• Increased cEBP/b expression</li> <li>• Increased S100A8/9 expression</li> <li>• Increased IL-4R (CD124) expression</li> </ul> |
| Proposed differences between neutrophils/granulocytes and PMN-MDSC | <ul style="list-style-type: none"> <li>• Suppressive function</li> <li>• Density (Ficoll gradients)</li> <li>• Increased ROS expression</li> <li>• Increased ARG1 expression</li> <li>• Increased CD33 &amp; CD66b expression</li> </ul>  |

\*non-exhaustive list.



morbidity-, age- and gender-dependent host and evolving viral nature of this condition. Peripheral blood immune signatures across COVID-19 patients revealed changes in both the innate and adaptive arm of immune responses, particularly in B and myelomonocytic cell composition, profoundly altered T cell phenotypes and selective cytokine/chemokine upregulation and SARS-CoV-2-specific antibodies (30). While most studies focused their attention on T cells, more comprehensive immune profiling approaches found that in severe cases the number of T and B lymphocytes, dendritic cells, natural killer (NK) cells, and HLA-DR<sup>high</sup> expressing cells were found to be substantially decreased in COVID-19 disease (31). High-dimensional flow cytometry-analysis focusing on mononuclear phagocyte (MNP) lineages in SARS-CoV-2-infected patients with moderate and severe COVID-19 identified a redistribution of monocyte subsets toward intermediate monocytes and a general decrease in circulating DCs, which coincided with the appearance of MDSCs and a higher frequency of pre-DC2. Furthermore, the presence of a MNP profile was associated with a cluster of COVID-19 non-survivors (32).

Innate immune sensing serves as the first line of antiviral defense and is essential for immunity to viruses. Coronaviruses (CoVs) have evolved several mechanisms to inhibit IFN-I induction and signaling, e.g. suppression of MAV signaling (33, 34). Patients with severe COVID-19 demonstrate remarkably impaired IFN-I signatures as compared to mild or moderate cases and fail to elicit an early IFN-I response (35–37). Perhaps timing is key, as IFN is protective early in disease but later becomes pathogenic. Furthermore, while pathogenic CoVs block IFN signaling, they may actively promote other inflammatory pathways contributing to disease pathology. Among innate immune cells, particularly neutrophil counts were found to be significantly elevated in patients with COVID-19 and correlated with disease severity (38–40). An elevated neutrophil-to-lymphocyte ratio has been further suggested as clinical marker for predicting fatal complications related to Acute Respiratory Distress Syndrome (ARDS) in patients with COVID-19 (38). Increased production of pro-inflammatory cytokines and MDSCs inversely correlated with perforin-expressing NK and CD3+ T cells during disease

progression (38, 39, 41). An early elevation in cytokine levels was associated with a maladapted immune response profile and worse disease outcomes (42). Elevated levels of cytokines, such as IP-10/CXCL10, interleukin-10 and interleukin-6, were shown to predict subsequent clinical progression (30).

There is increasing evidence to suggest that distinct innate immune responses, specifically, underlie the different clinical trajectories of COVID-19 patients and that the hyperinflammatory syndrome in severe COVID-19 results from a dysregulated host innate immune response (43). Transcriptomic, epigenomic, and proteomic analyses revealed widespread dysfunction of peripheral innate immunity in severe and fatal COVID-19, with the most profound disturbances including a prominent neutrophil hyper-activation signature and monocytes with anti-inflammatory features (7, 44, 45). Several studies have identified emergency myelopoiesis as a hallmark of severe or fatal COVID-19 (7, 45–47). Collectively, COVID-19 dysregulates both the innate and the adaptive immune system with a decrease of adaptive T cells and an increase of innate immune cell populations. However, the inter-connectedness of innate and adaptive immune changes remained largely elusive. Very recent studies provide evidence that MDSCs could represent that missing link as discussed below in detail.

## MDSCs IN COVID-19

As outlined above, MDSCs are innate immune cells regulating (mostly downregulating) adaptive immune responses. MDSC activity can be enhanced by malignant or infectious triggers as described for a variety of viral, bacterial and fungal infections. Several recent reports have shown that patients with severe COVID-19 exhibit a dysregulated myeloid cell compartment, with increased MDSC levels and activity correlating with disease severity. In mild COVID-19 disease course, studies have reported an increase in HLA-DR<sup>hi</sup>CD11c<sup>hi</sup> inflammatory monocytes with an interferon-stimulated gene signature, indicative of terminally differentiated monocytes, whereas severe COVID-19 was characterized by a lack of type I IFNs, high levels of HLA-DR<sup>low</sup> classical monocytes and CD10<sup>low</sup>CD101<sup>+</sup>CXCR4<sup>+/-</sup> neutrophils with an immunosuppressive profile in blood and lungs of severe cases, suggestive of emergency myelopoiesis (7, 47–50).

In severe COVID-19 patients with ARDS an increased ratio of MDSC to CD8 effector memory T cell was observed compared to patients hospitalized for COVID-19 moderate pneumonia, with COVID-19-related MDSC expansion directly correlating with lymphopenia and enhanced arginase activity (51). Marked expansion of MDSCs was observed, up to 90% of total circulating mononuclear cells in patients with severe disease, and up to 25% in the patients with mild disease with frequency decreasing with recovery (52). Granulocytic (neutrophil, eosinophil, and basophil) markers were enriched during COVID-19 and discriminated between patients with mild and severe disease, suggestive of PMN-MDSC activity (53). Increased counts of CD15<sup>+</sup>CD16<sup>+</sup> neutrophils, decreased granulocytic expression of integrin CD11b, and Th2-related CRTH2

downregulation in eosinophils and basophils established a COVID-19 signature. Severity was also associated with emergence of PD-L1 checkpoint expression in basophils and eosinophils (53).

In summary, COVID-19 patients show a shift towards an immature myeloid cell profile in peripheral blood together with mature monocytes and segmented neutrophils, likely the result of emergency myelopoiesis in response to the significantly elevated levels of cytokines and other pro-inflammatory mediators in these patients. As myeloid cells are the main immune cell subsets associated with severe COVID-19, identification of their inflammatory and chemotactic gene signatures could be of potential prognostic as well as therapeutic relevance (54).

## MDSCs AS POTENTIAL BIOMARKER IN COVID-19

There is compelling evidence that primarily PMN-MDSC (55) but also M-MDSC subtypes accumulate in COVID-19 patients and are associated with distinct gene and protein signatures. Given the increase of circulating MDSCs in COVID-19 and a correlation with disease outcome, an obvious clinical implication is to consider their utilization as biomarkers of immune dysregulation in COVID-19.

PMN-MDSC expanded during COVID-19 in patients who required intensive care treatments, and correlated with IL-1beta, IL-6, IL-8, and TNF-alpha plasma levels (55, 56). The expression of lectin-type oxidized LDL receptor 1 (LOX-1) on PMN-MDSC, in particular, has been proposed to identify a subset of MDSCs with the most potent immunosuppressive properties, the elevation of which, was found to be more pronounced in patients with ARDS (57). Furthermore, a marked increase in Hexokinase II+ PMN-MDSC was found exclusively in the acute COVID-19 patients with moderate or severe disease (58). PMN-MDSC inhibited T-cell IFN-gamma production upon SARS-CoV-2 peptides stimulation, through TGF-beta- and iNOS-mediated mechanisms, possibly counter-acting virus elimination (55). An observed MDSC decline at convalescent phase was associated to a reduction in TGF-beta and to an increase of inflammatory cytokines in plasma samples (52). Finally, a multivariate regression analysis found a strong association between PMN-MDSC percentages and fatal disease outcome and PMN-MDSC frequencies were higher in non-survivors than survivors (55).

M-MDSCs were also found to expand in blood of COVID-19 patients, suppress T cells and strongly associate with disease severity. More specifically, a population of M-MDSC expressing high carnitine palmitoyltransferase I (CPT1a) and VDAC, were present in the PBMC of the acute COVID-19 patients and correlated with severity of disease (58). Furthermore, M-MDSC frequencies were elevated in blood but not in nasopharyngeal or endotracheal aspirates of COVID-19 patients compared to controls (59). M-MDSCs isolated from COVID-19 patients suppressed T cell proliferation and IFN-gamma production



partly *via* an arginase-1 (Arg-1) dependent mechanism. Furthermore, these patients showed increased Arg-1 and IL-6 plasma levels. COVID-19 patients had fewer T cells, and displayed downregulated expression of the CD3zeta chain (59). In related studies, T cell proliferative capacity *in vitro* was significantly reduced among COVID-19 patients and could be restored through arginine supplementation (51).

Single-cell RNA sequencing (scRNAseq) data from bronchoalveolar lavage (BAL) also revealed the presence of neutrophils and macrophages as a hallmark of severe COVID-19 (54). Among the identified gene signatures, IFITM2, IFITM1, H3F3B, SAT1, and S100A8 gene signatures were highly associated with neutrophils, while CCL8, CCL3, CCL2, KLF6, and SPP1 were associated with macrophages in severe-COVID-19 patients. These findings are in agreement with high levels of calprotectin (S100A8/S100A9) found in plasma of severe cases (47). Genes associated with the inflammatory response and chemotaxis of myeloid cells, phagocytes, and granulocytes were among the top activated functional categories in BAL from severe COVID-19-affected patients (54). A lack of type I IFNs, reduced HLA-DR in myeloid cells and transient expression of IFN-stimulated genes characterized the transcriptome of patients with severe COVID-19 (50). Similarly, in a meta-analysis of transcriptomic data, the upregulation of the monocytic compartment in severe COVID-19, was dependent on the cytokines IL-6 and IL-10, and was characterized by broadly immunosuppressive properties and decreased responsiveness to stimulation (46). Myeloid cells of severe patients showed higher expression of pro-inflammatory cytokines and chemokines such as CXCL8 (60).

ScRNAseq profiling was further used to characterize the PBMC compartment of uninfected controls and COVID-19 patients and cells in paired broncho-alveolar lavage fluid (BALF) (61). A close association of decreased DCs and increased monocytes resembling MDSCs, correlated with lymphopenia and inflammation in the blood of severe COVID-19 patients. Those MDSC-like monocytes were characterized as 'immune-paralyzed'. In contrast, monocyte-macrophages in BALFs of COVID-19 patients produced massive amounts of cytokines and chemokines, but secreted little interferons (61). In similar meta-analyses, an overall upregulation of immunoinhibitory receptors mRNA during SARS-CoV-2 infection, expressed on both lymphoid and myeloid cells were upregulated in nasopharyngeal swabs and autopsies (e.g. BTLA, LAG3, FCGR2B, PDCD1, CEACAM1, CTLA4, CD72, and SIGLEC7), also directly correlated with viral levels (62). Integration of plasma proteomics with nine published scRNAseq datasets also revealed that disease severity in lung tissue is driven by myeloid cell phenotypes and cell-cell interactions with lung epithelial cells and T cells. Epithelial damage more specifically was associated with neutrophil infiltration (63).

Immature myeloid subsets and bronchoalveolar cells of critically-ill COVID-19 patients have been found to express HIF1alpha, a critical regulator of the differentiation and function of MDSCs, and transcriptional targets related to

inflammation (CXCL8, CXCR1, CXCR2, and CXCR4); virus sensing, (TLRs); and metabolism (SLC2A3, PFKFB3, PGK1, GAPDH and SOD2) (64). The up-regulation and participation of HIF1alpha in events such as inflammation and immunometabolism make it a potential biomarker of COVID-19 severity. HIF1alpha and its transcriptionally regulated genes are also expressed in lung cells from severe COVID-19 patients, which may partially explain the hypoxia related events (64).

Calprotectin (S100A8/9) plasma level and a routine flow cytometry assay detecting decreased frequencies of non-classical monocytes have been shown to discriminate patients who develop a severe form of COVID-19, suggesting a prognostic value that deserves prospective evaluation (47). Elevated S100A-family alarmins in myeloid cells and marked enrichment of serum proteins that map to myeloid cells and pathways including cytokines, complement/coagulation, and fluid shear stress were also identified in pediatric MIS-C patients even in the absence of active infection (65). Soluble triggering receptor also expressed on myeloid cells had the best prognostic accuracy for 30-day intubation/mortality (66).

In summary, various observational retrospective investigations suggest that MDSCs, their subsets or MDSC-related markers and signatures (**Figure 1**) could serve as biomarkers for severe COVID-19 (Summarized in **Table 2**), yet prospective and multi-center biomarker-focused studies are required to (i) standardize MDSC assays used and (ii) define the prognostic and/or, if linked to therapeutic treatments, the predictive biomarker potential of MDSCs in COVID-19.

## MDSCs AS POTENTIAL THERAPEUTIC TARGET IN COVID-19

Given the emerging role of MDSCs in COVID-19, another consequent question is how to therapeutically exploit and target this cell population. Based on insights from other, more established disease areas, such as oncology, therapeutic targeting of MDSCs can be achieved through different routes (16, 67): (i) drugs forcing MDSC differentiation into mature cells (e.g. vitamin D3 or retinoic acid), (ii) drugs inhibiting MDSC maturation from cellular precursors (e.g. bevacizumab, tyrosine kinase inhibitors, STAT3 inhibitors, MMP9 inhibitors), (iii) drugs reducing MDSC accumulation in peripheral organs (e.g. CXCR2/CXCR4 antagonists, 5-Fluorouracil, Gemcitabine) or (iv) drugs affecting MDSC inhibitory functions (ROS scavengers, cyclooxygenase 2 (COX2) or phosphodiesterase type 5 (PDE5) inhibitors). Prostaglandin D2 (PGD2) has been proposed a key mediator of lymphopenia in COVID-19 and is known to upregulate M-MDSCs *via* the DP2 receptor signaling in group 2 innate lymphoid cells (ILC2). Targeting PGD2/DP2 signaling using a receptor antagonist such as ramatroban could be used in immunotherapy for immune dysfunction and lymphopenia in COVID-19 disease (68).

Alarmin S100A8 was robustly induced in SARS-CoV-2-infected animal models as well as in COVID-19 patients.

**TABLE 2** | Candidate myeloid biomarkers associated with COVID-19 disease severity.

| Proposed Biomarker  | Description of COVID-19 related findings                 | References       |
|---|--|------------------|
| Neutrophil-to-lymphocyte ratio  | Increased with severity<br>Predicting ARDS complications | (38–40)          |
| MDSC to T-cell and NK-cell ratios                                       | Increased with severity                                  | (38, 39, 41, 51) |
| IP-10/CXCL10, interleukin-10 and interleukin-6                          | Increased with severity<br>Prognostic markers            | (30)             |
| HLA-DR <sup>high</sup> expressing cells                                 | Decreased with disease severity                          | (31)             |
| IFN-I signature   | Downregulated with severity                              | (35–37)          |
| HLA-DR <sup>low</sup> classical monocytes                               | Increased with severity                                  | (7, 47–50)       |
| CD10 <sup>low</sup> CD101 <sup>+</sup> CXCR4 <sup>+/−</sup> neutrophils | Increased with severity                                  | (7, 47–50)       |
| LOX-1 on PMN-MDSC   | Increased with ARDS                                      | (57)             |
| Hexokinase II+ PMN-MDSC   | Increased with severity                                  | (58)             |
| CD15 <sup>+</sup> CD16 <sup>+</sup> CD11b <sup>low</sup> neutrophils    | Increased with severity                                  | (53)             |
| TGF-beta plasma levels  | Increased with severity                                  | (52)<br>(55)     |
| Arg-1 and IL-6 plasma levels  | Increased with severity                                  | (59)             |
| M-MDSC expressing high CPT1a and VDAC                                   | Increased with severity                                  | (58)             |
| HIF1-alpha expression   | Upregulation with severity                               | (64)             |
| Calprotectin (S100A8/9) plasma level                                    | Prognostic marker for severe disease                     | (47)             |
| Soluble triggering receptor   | Prognostic marker for intubation/mortality               | (66)             |

Paquinimod, a specific inhibitor of S100A8/A9, could rescue the pneumonia with substantial reduction of viral loads in SARS-CoV-2-infected mice (69). A group of neutrophils that contributes to the uncontrolled pathological damage and onset of COVID-19 was induced by coronavirus infection. Paquinimod treatment could reduce these neutrophils and regain anti-viral responses, unveiling key roles of S100A8/A9 and aberrant neutrophils in the pathogenesis of COVID-19 and highlighting new opportunities for therapeutic intervention (69).

Therapeutic strategies targeting the migration/recruitment of myeloid cells from bone marrow as mentioned above could be considered for the treatment of COVID-19 induced hyperinflammation and immune dysregulation (39). Inhibitors of CXCR2 or CCR2 and CCR5 may be able to reduce mobilization and migration of MDSC from the bone marrow to the circulation (70, 71). Following compassionate care treatment with the CCR5 blocking antibody leronlimab, a rapid reduction of plasma IL-6, restoration of the CD4/CD8 ratio, and a significant decrease in SARS-CoV-2 plasma viremia was observed. Consistent with reduction of plasma IL-6, single-cell RNA-sequencing also revealed declines in transcriptomic myeloid cell clusters expressing IL-6 and interferon-related genes (72).

Given the expression of inhibitory receptor upregulation observed in a variety of cell subsets during the progression of COVID-19 (62), targeting immuno-inhibitory receptors could also represent an effective therapeutic approach for the treatment of COVID-19 early and reversal of late immune dysregulation and suppression (73). Finally, reprogramming MDSCs by targeting immunometabolism and epigenetics may also hold promise in resolving lung inflammation associated with COVID-19 (74). As patients with severe COVID-19 have an increased inflammatory response that depletes arginine, and subsequently impairs T cell function, inhibition of arginase-1 and/or replenishment of arginine may be a potential future therapeutic approach in preventing/treating severe COVID-19

(75). Furthermore, the fatty acid transport protein 2 (FATP2), responsible for the uptake of arachidonic acid and for the subsequent synthesis of PGE2 was identified as a regulator of the suppressive functions of PMN-MDSCs (76). IDO dependent tryptophan metabolism is another pathway used by MDSCs to inhibit immune responses (77). Targeting metabolic mediators as FATP2 or enzymes such as IDO may be able to reverse MDSC induced suppression of virus-specific T-cell responses seen in severe COVID-19 cases.

Regarding potential targeting of MDSCs in COVID-19, it is essential to define the disease stage and disease severity level where such a therapeutic approach might have the greatest potential and would be beneficial rather than harmful to disease outcome. This consideration is key, as, in analogy to other viral infections (24–26). MDSCs may play a pathogenic or protective role depending on the time-course, pathogen load and severity of the individual disease condition. Given the COVID-19-associated lymphopenia, MDSCs were proposed as causal culprits to decrease T cells and thereby impair T cell-mediated host defense (Figure 1). On the other hand, MDSCs are capable of dampen overshooting tissue inflammation and might be beneficial at certain stages of disease. Therapeutic targeting would make sense at stages where MDSCs cause more harm than good and it is key to first identify those stages precisely.

## CONCLUSIONS

COVID-19 activates the innate immune system and suppresses adaptive T cell responses. MDSCs are key cellular players connecting innate and adaptive immunity. Both M-MDSCs and PMN-MDSCs accumulate in patients with COVID-19 and reflect disease outcome, but what does this mean for the future of COVID-19 diagnosis, monitoring and treatment? Currently,

inflammation, cell-death- and coagulation-associated serum proteins such as CrP, LDH and IL-6 as well as D-Dimers are used to characterize COVID-19 severity and disease progression clinically; it remains to be assessed how MDSC frequencies in peripheral blood and/or airway fluids relate to these clinical serum markers and whether combined/composite biomarker scores composed of both serum proteins and cells (PMN-MDSCs and/or M-MDSCs) could be superior than clinical serum markers alone to monitor and predict the outcome and treatment response in COVID-19. Targeting MDSCs as future therapeutic approach in COVID-19 is farer away, yet could add substantial value, particularly in combination with other immunomodulatory drugs, such as cytokine blockers.

## REFERENCES

- Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical Course and Risk Factors for Mortality of Adult Inpatients With COVID-19 in Wuhan, China: A Retrospective Cohort Study. *Lancet* (2020) 395(10229):1054–62. doi: 10.1016/S0140-6736(20)30566-3
- W. H. O. *Covid-19: World Health Organization*. (2021). Available at: <https://www.who.int/>.
- Manson JJ, Crooks C, Naja M, Ledlie A, Goulden B, Liddle T, et al. Covid-19-associated Hyperinflammation and Escalation of Patient Care: A Retrospective Longitudinal Cohort Study. *Lancet Rheumatol* (2020) 2(10):e594–602. doi: 10.1016/S2665-9913(20)30275-7
- Wiersinga WJ, Rhodes A, Cheng AC, Peacock SJ, Prescott HC. Pathophysiology, Transmission, Diagnosis, and Treatment of Coronavirus Disease 2019 (Covid-19): A Review. *JAMA* (2020) 324(8):782–93. doi: 10.1001/jama.2020.12839
- Berlin DA, Gulick RM, Martinez FJ. Severe Covid-19. *N Engl J Med* (2020) 383(25):2451–60. doi: 10.1056/NEJMcp2009575
- Gomez-Rial J, Rivero-Calle I, Salas A, Martinon-Torres F. Role of Monocytes/Macrophages in Covid-19 Pathogenesis: Implications for Therapy. *Infect Drug Resist* (2020) 13:2485–93. doi: 10.2147/IDR.S258639
- Schulte-Schrepping J, Reusch N, Paclik D, Bassler K, Schlickeiser S, Zhang B, et al. Severe COVID-19 Is Marked by a Dysregulated Myeloid Cell Compartment. *Cell* (2020) 182(6):1419–40.e23. doi: 10.1016/j.cell.2020.08.001
- Merad M, Martin JC. Pathological Inflammation in Patients With COVID-19: A Key Role for Monocytes and Macrophages. *Nat Rev Immunol* (2020) 20(6):355–62. doi: 10.1038/s41577-020-0331-4
- Forni G, Mantovani AR. Covid-19 Commission of Accademia Nazionale dei Lincei. Covid-19 Vaccines: Where We Stand and Challenges Ahead. *Cell Death Differ* (2021) 28(2):626–39. doi: 10.1038/s41418-020-00720-9
- Gabrilovich DI, Nagaraj S. Myeloid-Derived Suppressor Cells as Regulators of the Immune System. *Nat Rev Immunol* (2009) 9(3):162–74. doi: 10.1038/nri2506
- Bronte V, Brandau S, Chen SH, Colombo MP, Frey AB, Greten TF, et al. Recommendations for Myeloid-Derived Suppressor Cell Nomenclature and Characterization Standards. *Nat Commun* (2016) 7:12150. doi: 10.1038/ncomms12150
- Ostrand-Rosenberg S, Fenselau C. Myeloid-Derived Suppressor Cells: Immune-Suppressive Cells That Impair Antitumor Immunity and Are Sculpted by Their Environment. *J Immunol* (2018) 200(2):422–31. doi: 10.4049/jimmunol.1701019
- Veglia F, Perego M, Gabrilovich D. Myeloid-Derived Suppressor Cells Coming of Age. *Nat Immunol* (2018) 19(2):108–19. doi: 10.1038/s41590-017-0022-x
- Serafini P, Mgebroff S, Noonan K, Borrello I. Myeloid-Derived Suppressor Cells Promote Cross-Tolerance in B-Cell Lymphoma by Expanding Regulatory T Cells. *Cancer Res* (2008) 68(13):5439–49. doi: 10.1158/0008-5472.CAN-07-6621
- Sarvaria A, Madrigal JA, Saudemont A. B Cell Regulation in Cancer and Anti-Tumor Immunity. *Cell Mol Immunol* (2017) 14(8):662–74. doi: 10.1038/cmi.2017.35

## AUTHOR CONTRIBUTIONS

MR contributed to manuscript research and writing. FS contributed to manuscript research and writing. DH contributed to manuscript research and writing. All authors contributed to the article and approved the submitted version.

## ACKNOWLEDGMENTS

We wish to thank Alan Abrams, Novartis Institutes for BioMedical Research (Cambridge, USA) for contributing the scientific artwork for **Figure 1**.

- Veglia F, Sanseviero E, Gabrilovich DI. Myeloid-Derived Suppressor Cells in the Era of Increasing Myeloid Cell Diversity. *Nat Rev Immunol* (2021) 2:1–14. doi: 10.1038/s41577-020-00490-y
- Pawelec G, Verschoor CP, Ostrand-Rosenberg S. Myeloid-Derived Suppressor Cells: Not Only in Tumor Immunity. *Front Immunol* (2019) 10:1099. doi: 10.3389/fimmu.2019.01099
- Condamine T, Gabrilovich DI. Molecular Mechanisms Regulating Myeloid-Derived Suppressor Cell Differentiation and Function. *Trends Immunol* (2011) 32(1):19–25. doi: 10.1016/j.it.2010.10.002
- Rieber N, Brand A, Hector A, Graepler-Mainka U, Ost M, Schafer I, et al. Flagellin Induces Myeloid-Derived Suppressor Cells: Implications for Pseudomonas Aeruginosa Infection in Cystic Fibrosis Lung Disease. *J Immunol* (2013) 190(3):1276–84. doi: 10.4049/jimmunol.1202144
- Papaspyridonos M, Matei I, Huang Y, do Rosario Andre M, Brazier-Mitouart H, Waite JC, et al. Id1 Suppresses Anti-Tumour Immune Responses and Promotes Tumour Progression by Impairing Myeloid Cell Maturation. *Nat Commun* (2015) 6:6840. doi: 10.1038/ncomms7840
- Medina E, Hartl D. Myeloid-Derived Suppressor Cells in Infection: A General Overview. *J Innate Immun* (2018) 10(5-6):407–13. doi: 10.1159/000489830
- Dorhoi A, Du Plessis N. Monocytic Myeloid-Derived Suppressor Cells in Chronic Infections. *Front Immunol* (2017) 8:1895. doi: 10.3389/fimmu.2017.01895
- Ost M, Singh A, Peschel A, Mehling R, Rieber N, Hartl D. Myeloid-Derived Suppressor Cells in Bacterial Infections. *Front Cell Infect Microbiol* (2016) 6:37. doi: 10.3389/fcimb.2016.00037
- Goh C, Narayanan S, Hahn YS. Myeloid-Derived Suppressor Cells: The Dark Knight or the Joker in Viral Infections? *Immunol Rev* (2013) 255(1):210–21. doi: 10.1111/imr.12084
- Norris BA, Uebelhoefer LS, Nakaya HI, Price AA, Grakoui A, Pulendran B. Chronic But Not Acute Virus Infection Induces Sustained Expansion of Myeloid Suppressor Cell Numbers That Inhibit Viral-Specific T Cell Immunity. *Immunity* (2013) 38(2):309–21. doi: 10.1016/j.immuni.2012.10.022
- O'Connor MA, Rastad JL, Green WR. The Role of Myeloid-Derived Suppressor Cells in Viral Infection. *Viral Immunol* (2017) 30(2):82–97. doi: 10.1089/vim.2016.0125
- Tacke RS, Lee HC, Goh C, Courtney J, Polyak SJ, Rosen HR, et al. Myeloid Suppressor Cells Induced by Hepatitis C Virus Suppress T-Cell Responses Through the Production of Reactive Oxygen Species. *Hepatology* (2012) 55(2):343–53. doi: 10.1002/hep.24700
- Pallett LJ, Gill US, Quaglia A, Sinclair LV, Jover-Cobos M, Schurich A, et al. Metabolic Regulation of Hepatitis B Immunopathology by Myeloid-Derived Suppressor Cells. *Nat Med* (2015) 21(6):591–600. doi: 10.1038/nm.3856
- Vollbrecht T, Stirner R, Tufman A, Roeder J, Huber RM, Bogner JR, et al. Chronic Progressive HIV-1 Infection Is Associated With Elevated Levels of Myeloid-Derived Suppressor Cells. *AIDS* (2012) 26(12):F31–7. doi: 10.1097/QAD.0b013e328354b43f
- Laing AG, Lorenc A, Del Molino Del Barrio I, Das A, Fish M, Monin L, et al. A Dynamic COVID-19 Immune Signature Includes Associations With Poor Prognosis. *Nat Med* (2020) 26(10):1623–35. doi: 10.1038/s41591-020-1038-6

31. Matic S, Popovic S, Djurdjevic P, Todorovic D, Djordjevic N, Mijailovic Z, et al. Sars-CoV-2 Infection Induces Mixed M1/M2 Phenotype in Circulating Monocytes and Alterations in Both Dendritic Cell and Monocyte Subsets. *PLoS One* (2020) 15(12):e0241097. doi: 10.1371/journal.pone.0241097
32. Kvedaraitė E, Hertwig L, Sinha I, Ponzetta A, Hed Myrberg I, Lourda M, et al. Major Alterations in the Mononuclear Phagocyte Landscape Associated With COVID-19 Severity. *Proc Natl Acad Sci USA* (2021) 118(6):1–12. doi: 10.1073/pnas.2018587118
33. Shi CS, Qi HY, Boularan C, Huang NN, Abu-Asab M, Shelhamer JH, et al. SARS-Coronavirus Open Reading Frame-9b Suppresses Innate Immunity by Targeting Mitochondria and the MAVS/TRAF3/TRAF6 Signalingosome. *J Immunol* (2014) 193(6):3080–9. doi: 10.4049/jimmunol.1303196
34. Gordon DE, Jang GM, Bouhaddou M, Xu J, Obernier K, O'Meara MJ, et al. A SARS-CoV-2-Human Protein-Protein Interaction Map Reveals Drug Targets and Potential Drug-Repurposing. *bioRxiv* (2020). doi: 10.1101/2020.03.22.002386
35. Bastard P, Rosen LB, Zhang Q, Michailidis E, Hoffmann HH, Zhang Y, et al. Autoantibodies Against Type I IFNs in Patients With Life-Threatening COVID-19. *Science* (2020) 370(6515):1–13. doi: 10.1126/science.abd4585
36. Hadjadj J, Yatim N, Barnabei L, Corneau A, Boussier J, Smith N, et al. Impaired Type I Interferon Activity and Inflammatory Responses in Severe COVID-19 Patients. *Science* (2020) 369(6504):718–24. doi: 10.1126/science.abc6027
37. Channappanavar R, Fehr AR, Vijay R, Mack M, Zhao J, Meyerholz DK, et al. Dysregulated Type I Interferon and Inflammatory Monocyte-Macrophage Responses Cause Lethal Pneumonia in SARS-CoV-Infected Mice. *Cell Host Microbe* (2016) 19(2):181–93. doi: 10.1016/j.chom.2016.01.007
38. Chiang CC, Korinek M, Cheng WJ, Hwang TL. Targeting Neutrophils to Treat Acute Respiratory Distress Syndrome in Coronavirus Disease. *Front Pharmacol* (2020) 11:572009. doi: 10.3389/fphar.2020.572009
39. Mann ER, Menon M, Knight SB, Konkel JE, Jagger C, Shaw TN, et al. Longitudinal Immune Profiling Reveals Key Myeloid Signatures Associated With COVID-19. *Sci Immunol* (2020) 5(51):1–11. doi: 10.1126/sciimmunol.abd6197
40. Godkin A, Humphreys IR. Elevated interleukin-6, interleukin-10 and Neutrophil : Lymphocyte Ratio as Identifiers of Severe Coronavirus Disease 2019. *Immunology* (2020) 160(3):221–2. doi: 10.1111/imm.13225
41. Tomic S, Dokic J, Stevanovic D, Ilic N, Gruden-Movsesijan A, Dinic M, et al. Reduced Expression of Autophagy Markers and Expansion of Myeloid-Derived Suppressor Cells Correlate With Poor T Cell Response in Severe Covid-19 Patients. *Front Immunol* (2021) 12:614599. doi: 10.3389/fimmu.2021.614599
42. Lucas C, Wong P, Klein J, Castro TBR, Silva J, Sundaram M, et al. Longitudinal Analyses Reveal Immunological Misfiring in Severe COVID-19. *Nature* (2020) 584(7821):463–9. doi: 10.1038/s41586-020-2588-y
43. Gustine JN, Jones D. Immunopathology of Hyperinflammation in COVID-19. *Am J Pathol* (2021) 191(1):4–17. doi: 10.1016/j.ajpath.2020.08.009
44. Cicco S, Cicco G, Racanelli V, Vacca A. Neutrophil Extracellular Traps (NETs) and Damage-Associated Molecular Patterns (DAMPs): Two Potential Targets for COVID-19 Treatment. *Mediators Inflamm* (2020) 2020:7527953. doi: 10.1155/2020/7527953
45. Wilk AJ, Lee MJ, Wei B, Parks B, Pi R, Martínez-Colón GJ, et al. Multi-Omic Profiling Reveals Widespread Dysregulation of Innate Immunity and Hematopoiesis in COVID-19. *bioRxiv* (2020) 2020.12.18.423363. doi: 10.1101/2020.12.18.423363
46. Reyes M, Filbin MR, Bhattacharyya RP, Sonny A, Mehta A, Billman K, et al. Induction of a Regulatory Myeloid Program in Bacterial Sepsis and Severe COVID-19. *bioRxiv* (2020). doi: 10.1101/2020.09.02.280180
47. Silvin A, Chapuis N, Dunsmore G, Goubet AG, Dubuisson A, Derosa L, et al. Elevated Calprotectin and Abnormal Myeloid Cell Subsets Discriminate Severe From Mild Covid-19. *Cell* (2020) 182(6):1401–18.e18. doi: 10.1016/j.cell.2020.08.002
48. Peruzzi B, Bencini S, Capone M, Mazzoni A, Maggi L, Salvati L, et al. Quantitative and Qualitative Alterations of Circulating Myeloid Cells and Plasmacytoid DC in SARS-CoV-2 Infection. *Immunology* (2020) 161(4):345–53. doi: 10.1111/imm.13254
49. Benlyamani I, Venet F, Coudereau R, Gossez M, Monneret G. Monocyte HLA-DR Measurement by Flow Cytometry in COVID-19 Patients: An Interim Review. *Cytometry A* (2020) 97(12):1217–21. doi: 10.1002/cyto.a.24249
50. Arunachalam PS, Wimmers F, Mok CKP, Perera R, Scott M, Hagan T, et al. Systems Biological Assessment of Immunity to Mild Versus Severe COVID-19 Infection in Humans. *Science* (2020) 369(6508):1210–20. doi: 10.1126/science.abc6261
51. Reizine F, Lesouhaitier M, Gregoire M, Pinceaux K, Gacouin A, Maamar A, et al. SARS-Cov-2-Induced ARDS Associates With MDSC Expansion, Lymphocyte Dysfunction, and Arginine Shortage. *J Clin Immunol* (2021) 41:515–25. doi: 10.1007/s10875-020-00920-5
52. Agrati C, Sacchi A, Bordoni V, Cimini E, Notari S, Grassi G, et al. Expansion of Myeloid-Derived Suppressor Cells in Patients With Severe Coronavirus Disease (COVID-19). *Cell Death Differ* (2020) 27(11):3196–207. doi: 10.1038/s41418-020-0572-6
53. Vitte J, Diallo AB, Boumaza A, Lopez A, Michel M, Allardet-Servent J, et al. A Granulocytic Signature Identifies Covid-19 and Its Severity. *J Infect Dis* (2020) 222(12):1985–96. doi: 10.1093/infdis/jiaa591
54. Shaath H, Vishnubalaji R, Elkord E, Alajez NM. Single-Cell Transcriptome Analysis Highlights a Role for Neutrophils and Inflammatory Macrophages in the Pathogenesis of Severe Covid-19. *Cells* (2020) 9(11):1–19. doi: 10.3390/cells9112374
55. Sacchi A, Grassi G, Bordoni V, Lorenzini P, Cimini E, Casetti R, et al. Early Expansion of Myeloid-Derived Suppressor Cells Inhibits SARS-CoV-2 Specific T-cell Response and may Predict Fatal COVID-19 Outcome. *Cell Death Dis* (2020) 11(10):921. doi: 10.1038/s41419-020-03125-1
56. Takano T, Matsumura T, Adachi Y, Terahara K, Moriyama S, Onodera T, et al. Myeloid Cell Dynamics Correlating With Clinical Outcomes of Severe COVID-19 in Japan. *Int Immunol* (2021) 33(4):241–7. doi: 10.1093/intimm/dxab005
57. Coudereau R, Waeckel L, Cour M, Rimmle T, Pescarmona R, Fabri A, et al. Emergence of Immunosuppressive LOX-1+ Pmn-MDSC in Septic Shock and Severe COVID-19 Patients With Acute Respiratory Distress Syndrome. *J Leukoc Biol* (2021) 28:1–8. doi: 10.1002/JLB.4COVBCR0321-129R
58. Thompson E, Cascino K, Ordonez A, Zhou W, Vaghiasa A, Hamacher-Brady A, et al. Mitochondrial Induced T Cell Apoptosis and Aberrant Myeloid Metabolic Programs Define Distinct Immune Cell Subsets During Acute and Recovered SARS-CoV-2 Infection. *medRxiv* (2020). doi: 10.1101/2020.09.10.20186064
59. Falck-Jones S, Vangeti S, Yu M, Falck-Jones R, Cagigi A, Badolati I, et al. Functional Monocytic Myeloid-Derived Suppressor Cells Increase in Blood But Not Airways and Predict COVID-19 Severity. *J Clin Invest* (2021) 131:1–15. doi: 10.1172/JCI144734
60. Park JH, Lee HK. Re-Analysis of Single Cell Transcriptome Reveals That the NR3C1-CXCL8-Neutrophil Axis Determines the Severity of COVID-19. *Front Immunol* (2020) 11:2145. doi: 10.3389/fimmu.2020.02145
61. Xu G, Qi F, Li H, Yang Q, Wang H, Wang X, et al. The Differential Immune Responses to COVID-19 in Peripheral and Lung Revealed by Single-Cell RNA Sequencing. *Cell Discovery* (2020) 6:73. doi: 10.1038/s41421-020-00225-2
62. Sharif-Askari NS, Sharif-Askari FS, Mdkhana B, Al Heialy S, Alsafar HS, Hamoudi R, et al. Enhanced Expression of Immune Checkpoint Receptors During SARS-CoV-2 Viral Infection. *Mol Ther Methods Clin Dev* (2020) 20:109–21. doi: 10.1016/j.omtm.2020.11.002
63. Filbin MR, Mehta A, Schneider AM, Kays KR, Guess JR, Gentili M, et al. Plasma Proteomics Reveals Tissue-Specific Cell Death and Mediators of Cell-Cell Interactions in Severe COVID-19 Patients. *bioRxiv* (2020). doi: 10.1101/2020.11.02.365536
64. Taniguchi-Ponciano K, Vadillo E, Mayani H, Gonzalez-Bonilla CR, Torres J, Majluf A, et al. Increased Expression of Hypoxia-Induced Factor 1alpha mRNA and its Related Genes in Myeloid Blood Cells From Critically Ill COVID-19 Patients. *Ann Med* (2021) 53(1):197–207. doi: 10.1080/07853890.2020.1858234
65. Ramaswamy A, Brodsky NN, Sumida TS, Comi M, Asashima H, Hoehn KB, et al. Post-Infectious Inflammatory Disease in MIS-C Features Elevated Cytotoxicity Signatures and Autoreactivity That Correlates With Severity. *medRxiv* (2020). doi: 10.1101/2020.12.01.20241364
66. Van Singer M, Brahier T, Ngai M, Wright J, Weckman AM, Erice C, et al. Covid-19 Risk Stratification Algorithms Based on sTREM-1 and IL-6 in



- Emergency Department. *J Allergy Clin Immunol* (2020). doi: 10.1016/j.jaci.2020.10.001
67. Ugel S, Delpozzi F, Desantis G, Papalini F, Simonato F, Sonda N, et al. Therapeutic Targeting of Myeloid-Derived Suppressor Cells. *Curr Opin Pharmacol* (2009) 9(4):470–81. doi: 10.1016/j.coph.2009.06.014
68. Gupta A, Chander Chiang K. Prostaglandin D2 as a Mediator of Lymphopenia and a Therapeutic Target in COVID-19 Disease. *Med Hypotheses* (2020) 143:110122. doi: 10.1016/j.mehy.2020.110122
69. Guo Q, Zhao Y, Li J, Liu J, Yang X, Guo X, et al. Induction of Alarmin S100A8/A9 Mediates Activation of Aberrant Neutrophils in the Pathogenesis of COVID-19. *Cell Host Microbe* (2020) 29(2):222–35.E4. doi: 10.1016/j.chom.2020.12.016
70. Stadtmann A, Zarbock A. Cxcr2: From Bench to Bedside. *Front Immunol* (2012) 3:263. doi: 10.3389/fimmu.2012.00263
71. Lu Z, Zou J, Li S, Topper MJ, Tao Y, Zhang H, et al. Epigenetic Therapy Inhibits Metastases by Disrupting Premetastatic Niches. *Nature* (2020) 579(7798):284–90. doi: 10.1038/s41586-020-2054-x
72. Patterson BK, Seethamraju H, Dhody K, Corley MJ, Kazempour K, Lalezari JP, et al. Disruption of the CCL5/RANTES-CCR5 Pathway Restores Immune Homeostasis and Reduces Plasma Viral Load in Critical Covid-19. *medRxiv* (2020). doi: 10.1101/2020.05.02.20084673
73. Gruber CN, Patel RS, Trachtman R, Lepow L, Amanat F, Krammer F, et al. Mapping Systemic Inflammation and Antibody Responses in Multisystem Inflammatory Syndrome in Children (Mis-C). *Cell* (2020) 183(4):982–95.e14. doi: 10.1016/j.cell.2020.09.034
74. D'Alessio FR, Heller NM. Covid-19 and Myeloid Cells: Complex Interplay Correlates With Lung Severity. *J Clin Invest* (2020) 130(12):6214–7. doi: 10.1172/JCI143361
75. Dean MJ, Ochoa JB, Sanchez-Pino M, Zabaleta J, Garai J, Del Valle L, et al. Transcriptome and Functions of Granulocytic Myeloid-Derived Suppressor Cells Determine Their Association With Disease Severity of COVID-19. *medRxiv* (2021). doi: 10.1101/2021.03.26.21254441
76. Veglia F, Tyurin VA, Blasi M, De Leo A, Kossenkov AV, Donthireddy L, et al. Fatty Acid Transport Protein 2 Reprograms Neutrophils in Cancer. *Nature* (2019) 569(7754):73–8. doi: 10.1038/s41586-019-1118-2
77. Platten M, Nollen EAA, Rohrig UF, Fallarino F, Opitz CA. Tryptophan Metabolism as a Common Therapeutic Target in Cancer, Neurodegeneration and Beyond. *Nat Rev Drug Discovery* (2019) 18(5):379–401. doi: 10.1038/s41573-019-0016-5

**Conflict of Interest:** Authors MR, FS and DH were employed by the company Novartis.

Copyright © 2021 Rowlands, Segal and Hartl. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.