



Mechanobiological Principles Influence the Immune Response in Regeneration: Implications for Bone Healing

Raphael S. Knecht¹, Christian H. Bucher^{1,2}, Sophie Van Linthout^{2,3}, Carsten Tschöpe^{2,3,4}, Katharina Schmidt-Bleek^{1,2†} and Georg N. Duda^{1,2*†}

¹ Julius Wolff Institute and Center for Musculoskeletal Surgery, Charité—Universitätsmedizin Berlin, Berlin, Germany, ² Berlin Institute of Health Center for Regenerative Therapies, Charité—Universitätsmedizin Berlin, Berlin, Germany, ³ German Center for Cardiovascular Research (DZHK), Partner Site Berlin, Berlin, Germany, ⁴ Department of Cardiology, Charité—Universitätsmedizin Berlin, Berlin, Germany

OPEN ACCESS

Edited by:

Elisabetta Ada Cavalcanti-Adam,
Max-Planck-Gesellschaft
(MPG), Germany

Reviewed by:

Jae-Won Shin,
University of Illinois at Chicago,
United States
Mariaceleste Aragona,
University of Copenhagen, Denmark

*Correspondence:

Georg N. Duda
georg.duda@charite.de

†These authors share last authorship

Specialty section:

This article was submitted to
Biomaterials,
a section of the journal
Frontiers in Bioengineering and
Biotechnology

Received: 06 October 2020

Accepted: 22 January 2021

Published: 12 February 2021

Citation:

Knecht RS, Bucher CH, Van
Linthout S, Tschöpe C,
Schmidt-Bleek K and Duda GN (2021)
Mechanobiological Principles
Influence the Immune Response in
Regeneration: Implications for Bone
Healing.
Front. Bioeng. Biotechnol. 9:614508.
doi: 10.3389/fbioe.2021.614508

A misdirected or imbalanced local immune composition is often one of the reasons for unsuccessful regeneration resulting in scarring or fibrosis. Successful healing requires a balanced initiation and a timely down-regulation of the inflammation for the re-establishment of a biologically and mechanically homeostasis. While biomaterial-based approaches to control local immune responses are emerging as potential new treatment options, the extent to which biophysical material properties themselves play a role in modulating a local immune niche response has so far been considered only occasionally. The communication loop between extracellular matrix, non-hematopoietic cells, and immune cells seems to be specifically sensitive to mechanical cues and appears to play a role in the initiation and promotion of a local inflammatory setting. In this review, we focus on the crosstalk between ECM and its mechanical triggers and how they impact immune cells and non-hematopoietic cells and their crosstalk during tissue regeneration. We realized that especially mechanosensitive receptors such as TRPV4 and PIEZO1 and the mechanosensitive transcription factor YAP/TAZ are essential to regeneration in various organ settings. This indicates novel opportunities for therapeutic approaches to improve tissue regeneration, based on the immune-mechanical principles found in bone but also lung, heart, and skin.

Keywords: regeneration, inflammation, mechanobiology, mechano-transduction, immune-mechanics, YAP/TAZ, TRPV4, PIEZO1

INTRODUCTION

The immune response is essential for successful healing after an injury. Even under sterile conditions, the immune system reacts to injury with a pro-inflammatory reaction as a first step (Midwood and Piccinini, 2010; Schmidt-Bleek et al., 2012, 2015). However, a misguided immune cell response leading to a too pronounced pro-inflammatory response or a prolonged anti-inflammatory response ends in an unsuccessful or delayed healing outcome (Wynn, 2004; Braga et al., 2015; Schmidt-Bleek et al., 2015; Dellacherie et al., 2019). Until now, research has mainly

focused on how different cytokines and other small soluble molecules influence the local immune response and subsequently the healing outcome. According to Wolff's law, tissue adaptations in bone in particular are due to physical or mechanical forces during its dynamic homeostatic remodeling (Wolff, 1986; Rho et al., 1998). This force-sensitivity is also remarkably relevant for healing and regeneration such as the healing of bone fractures (Lienau et al., 2005; Schell et al., 2005; Epari et al., 2007; Ghiasi et al., 2017). For example osteocytes show extraordinary mechano-sensitivity and are engaged in regulating other cells in their activity (e.g., osteoblasts and osteoclasts) according to their mechanical sensation they perceive (Bonewald and Johnson, 2008; Prideaux et al., 2016; Qin et al., 2020; Sato et al., 2020). It is becoming increasingly apparent how relevant the crosstalk between immune cells and bone cells is in both homeostasis and regeneration (Tsukasaki and Takayanagi, 2019). However, how the cell niche and its surrounding, notably its mechanical properties, geometry and constitution, are involved in the crosstalk between non-hematopoietic and immune cells has long been overlooked or neglected.

Apparently, the control of cellular responses and cell behavior via mechanical cues is essential for the homeostasis and regeneration of tissue. Therefore, mechanical signals hold great potential for new therapeutic approaches, and they have inherent properties that circumvent some challenging and often application-limiting problems associated with small molecules or proteins. For example, mechanical cues only act locally, and thus, there is no diffusion of the therapeutic stimuli. As a result, mechanical stimuli can, if correctly designed and as needed, remain within the "therapeutic window" for days, weeks, months, or even years and without the need for constant re-application of the therapeutic agent. Moreover, when biomaterials are implanted *in vivo*, they are not mechanically isolated but come into contact with the surrounding tissue and cells. Therefore, forces, pressure, deformations and fluid flow should be considered as potential features for any therapeutical use of mechano-biological cues in addition to the inherent mechanical properties of the biomaterial such as stiffness, (visco-)elasticity, pore size, patterns, shape, and geometrical organization. Interestingly, matching for example the stiffness of the biomaterial to the surrounding tissue could already minimize the foreign body response (Carnicer-Lombarte et al., 2019). Deciphering such mechanical interfaces between biomaterials, surrounding tissue and immune cells will enable new modulation and intervention strategies in regenerative medicine.

So far, the exact mechanism and receptors involved in the perception of the mechanical niche around the cells and how this alters cell behavior have not yet been fully elucidated. This review addresses the question how relevant the mechanical niche is for immune cells in tissue regeneration. We aim to review what is known for different tissue regeneration processes as they are known in bone, lung, heart and skin. With such an approach we intend to decipher the immunomodulatory mechanical features that are either directly perceived by the immune cells themselves or indirectly by non-hematopoietic cells. In the latter case, non-hematopoietic cells act as carriers of mechanical properties to immune cells. Understanding critical mechanical properties that

either induce pathological processes in the body or initiate a successful healing response will help to exploit the full potential of mechanical regulation of cell behavior *in vivo* for future therapeutic approaches.

This review highlights the close crosstalk between the immune system and non-hematopoietic cells during pathological and regenerative processes in bone tissue, with a focus on mechanical constraints. We suggest that the mechanical niche influences ECM composition, non-hematopoietic and immune cells and thereby changes their crosstalk (**Figure 1**). Bone is of particular interest because it is able to regenerate without scarring, which represents the "ideal" healing capacity. At present, studies focusing on mechanical properties that influence crosstalk between immune cells, non-hematopoietic cells and the ECM in the development of pathologies or regeneration in bone are relatively rare. Therefore, we have looked at other tissues such as the lung and the heart, where more studies have already been conducted to decipher the crosstalk between immune cells, non-hematopoietic cells and the ECM from a mechanical point of view. We also looked at skin tissue, which is often used as a model to study wound healing. Finally, we will look back at the bone to identify common principles of interaction between the immune system and the mechanical niche and will point to possible new research opportunities.


BONE HEALING IS MODULATED VIA IMMUNE-MECHANICS

Bone Cells Are Influencing Immune Cell Development

The bone marrow harbors the hematopoietic stem cell (HSC) niche, which gives rise to the myeloid and lymphoid compartment of the immune system (Okamoto et al., 2017). Besides this spatial proximity between the HSC niche and bone, they are also developmentally and functionally connected. Osteoclasts are derived from monocytes in the myeloid compartment of the immune system, while mesenchymal stem cells (MSCs) within the bone marrow give rise to osteoblasts, and also contribute to the HSC niche (Loi et al., 2016). Even after their development from common progenitors, bone and immune cells stay in close contact with each other and more and more mechanisms have now been discovered about how bone cells influence immune regulation and homeostasis (Takayanagi, 2007).

Osteoblasts, osteocytes and osteoclasts have all been described to influence immune cells in physiological and pathological conditions (Tsukasaki and Takayanagi, 2019). HSCs must remain in the bone marrow niche to maintain their stem cell status and to allow the continuous formation of new blood and immune cells (Suárez-Álvarez et al., 2012; Morrison and Scadden, 2014). On the other hand, however, the recruitment of immature and maturing hematopoietic cells into the periphery has been suggested to contribute to host defense and tissue repair (Cottler-Fox et al., 2003; Kollet et al., 2003).

Osteoblasts progenitors, more specifically CXC-chemokine ligand 12 (CXCL12) abundant reticular (CAR) cells, were



shown to contribute to the HSC niche by expressing CXCL12, a chemokine with important functions in the homing and retention of hematopoietic stem/progenitor cells in the bone marrow (Lévesque et al., 2003; Omatsu et al., 2010; Greenbaum et al., 2013). Moreover, CXCL12 from osteoblast progenitors plays an important role for normal B- and T cell development in the bone marrow (Ara et al., 2003; Greenbaum et al., 2013; Tsukasaki and Takayanagi, 2019). The dysregulation of osteoblasts via the overexpression of jagged 1 has been associated with acute myeloid leukemia in mice (Kode et al., 2014). Moreover, sepsis-induced osteoblast ablation has been shown to lower interleukin (IL)-7 levels in the hematopoietic stem cell niche, resulting in lymphopenia and immunodeficiency (Terashima et al., 2016).

Osteocytes were shown to regulate hematopoiesis and myeloid expansion via the secretion of the proliferative factor granulocyte colony-stimulating factor (G-CSF) (Fulzele et al., 2013). Moreover, the ablation of osteocytes in mice was associated with defects in bone marrow, thymus and spleen and subsequently showed severe lymphopenia (Sato et al., 2013). In bone, osteocytes are the main producer of sclerostin, a potent inhibitor of osteoblastic bone formation (Li et al., 2008). Sclerostin-knockout mice however, showed also a depletion of the B cell lineage within the bone marrow and significant reduction of CXCL12 (Cain et al., 2012).

Osteoclasts have been shown to secrete matrix metalloproteinase 9 (MMP9) and cathepsin K, both playing a role in the degradation and inactivation of CXCL12 (Kollet et al., 2006). In addition, the administration of osteoclast stimulator receptor activator of nuclear factor (NF)- κ B (RANK) ligand (RANKL) led to a decrease in CXCL12 and promoted the mobilization of hematopoietic progenitor cells, thereby impairing hematopoiesis (Kollet et al., 2006). Moreover, high calcium concentration produced during bone resorption via osteoclasts has been described to be essential for HSCs niche maintenance (Adams et al., 2006).

Thus, bone progenitor cells and mature bone cells are important for the regulation and maintenance of the HSC niche. Consequently, dysfunctional bone (progenitor) cells in the bone marrow have been found to be associated with immune system pathologies.

Immune Cells Are Regulating the Activity of Bone Cells

Conversely, both lymphoid and myeloid immune cells influence the behavior of bone cells in many bone pathologies by altering the balance of bone formation and resorption (Tsukasaki and Takayanagi, 2019).

Bone resorption is tightly regulated via the RANKL-RANK system (Boyce and Xing, 2008). Signaling via RANKL-RANK together with macrophage colony-stimulating factor (M-CSF) is needed for the maturation and activation of osteoclast precursors and thus promotes bone resorption (Kong et al., 1999; Okamoto et al., 2017). Osteoprotegerin (OPG) is produced by B cells, dendritic cells, MSCs and osteoblasts and can act as decoy receptor for RANKL, blocking RANK mediated signaling in osteoclast precursors and reduce their maturation and activation, leading indirectly to more bone formation (Khosla, 2001). Moreover, T cells also express RANKL and may thereby promote osteoclast maturation and increase bone loss (Theill et al., 2002; Takayanagi, 2007; Weitzmann, 2013). However, the effect of T cells expressing RANKL is often overridden by the secretion of interferon (IFN)- γ by T helper (TH) 1 cells (Takayanagi et al., 2000). IFN- γ interferes with the RANKL-RANK signaling pathway and thus reduces osteoclastogenesis (Takayanagi et al., 2000). While IL-6 secreted by TH2 was found to increase RANKL expression on MSCs and osteoblasts, IL-4 secreted by TH2 has been shown to interfere with the RANKL-RANK signaling pathway and reduce osteoclastogenesis (Takayanagi, 2007; Claes et al., 2012). TH17 cells have been shown to promote osteoclastogenesis in rheumatoid arthritis mainly by promoting RANKL expression in synovial fibroblasts via IL-17 secretion (Sato et al., 2006; Danks et al., 2016).

In addition to the lymphoid cells of the immune system, the myeloid compartment has also been described to regulate bone homeostasis (Loi et al., 2016). Pro-inflammatory cytokines including tumor necrosis factor (TNF), IL-1 and IL-6 secreted by macrophages promoted osteoclastogenesis and increased bone resorption (Boyle et al., 2003; Takayanagi, 2007). However, monocytes and macrophages also support osteoblast differentiation and proliferation through the secretion of bone morphogenetic protein (BMP)-2, BMP-4 and transforming growth factor (TGF)- β 1 (Fromigué et al., 1998; Champagne et al., 2002; Blom et al., 2004). Furthermore, macrophages promote mineralization and osteogenic differentiation of MSCs via b-tricalcium phosphate and oncostatin m (Chen et al., 2014; Guihard et al., 2015). In addition, the polarization of macrophages toward an M2 phenotype via the administration of IL-4 and IL-13 has been shown to promote the mineralization of the soft callus after bone fracture (Schlundt et al., 2018).

These findings suggest that the number and activity of osteoclasts and osteoblasts can be regulated by lymphoid and

myeloid immune cells and therefore shift the balance between bone resorption and bone formation.

Immune Cells Are Crucial Players for Successful Bone Fracture Healing

The close interaction of bone and immune cells already suggests that the immune system also plays a decisive role in bone fracture healing. Indeed, timing of the initial pro-inflammatory and subsequent anti-inflammatory phase during bone healing is a necessity for successful healing (Mountziaris and Mikos, 2008; Schmidt-Bleek et al., 2012, 2014, 2015; Spiller et al., 2015; Maruyama et al., 2020).

The healing cascade after bone fracture can be divided into consecutive but partly overlapping phases (Kolar et al., 2010; Schmidt-Bleek et al., 2015). During the fracture event, blood vessels are disrupted, resulting in hematoma formation. This early fibrin network marks the start of the inflammatory phase and creates a first provisional matrix, into which mainly pro-inflammatory polymorphonuclear leukocyte (PMNs) migrate in and secrete pro-inflammatory cytokines, such as C-C motif chemokine 2 (CCL2) and IL-6 (Yang et al., 2007; Xing et al., 2010; Bastian et al., 2011, 2016). The immigrated PMNs secrete CCL2 and IL-6 recruit, which attracts macrophages and monocytes from the surrounding tissue and vessels into the hematoma (Hurst et al., 2001; Xing et al., 2010; Claes et al., 2012). The recruited macrophages start to phagocytose necrotic cells and debris and secrete further pro-inflammatory cytokines such as TNF, IL-1 β , IL-6, and CCL2, which recruits fibroblasts, MSCs and osteoprogenitor cells (Kon et al., 2001; Bielby et al., 2007; Wu et al., 2013).

In addition, lymphoid cells have also been found to immigrate during the pro-inflammatory phase into the fracture hematoma (Glynn Andrew et al., 1994; Könnecke et al., 2014). Pro-inflammatory factors such as IL-1, IL-6, and TNF have been found to increase RANKL expression on activated T cells, resulting increased osteoclastogenesis (Clowes et al., 2005; Hardy and Cooper, 2009). Even though, activated T cells also secrete interferon (INF)- γ , that interferes with RANKL signaling through downregulating tumor-necrosis factor-receptor-associated factor 6 (TRAF6), it has been suggested, that RANKL mediated promotion of osteoclastogenesis is dominant over the INF- γ mediated downregulation of osteoclastogenesis (Takayanagi, 2007; Greenblatt and Shim, 2013). Moreover, INF- γ secreted by cytotoxic T cells also led to a higher pro-inflammatory M1 macrophage polarization (Schmidt-Bleek et al., 2012). Thus, INF- γ seems to have more pro-inflammatory and osteoclast promoting functions.

This first pro-inflammatory phase, characterized by the formation of the hematoma and recruitment of pro-inflammatory cells, has been found to be critical for a successful healing outcome (Gerstenfeld et al., 2003; Yang et al., 2007; Kolar et al., 2010; Glass et al., 2011). However, within the inflammatory phase, an anti-inflammatory response is needed to resolve inflammation and allow for the transition to the subsequent soft callus and hard callus phase (Serhan and Savill, 2005; Schmidt-Bleek et al., 2015). IL-4, IL-10, and TGF- β secreted

by regulatory T cells (Tregs) interfere with RANKL signaling and reduce osteoclastogenesis and thus the presence of Tregs were associated with a shift from pro- to an anti-inflammatory phase (Arron and Choi, 2000; Takayanagi et al., 2000). In the soft callus phase, connective tissue is formed first, and later replaced by the formation of cartilage via the differentiation of MSCs into chondrocytes (Gerstenfeld et al., 2003; Claes et al., 2012). This soft callus gives initial mechanical stability and serves as a scaffold for new bone formation via endochondral ossification during the hard callus phase, where MSCs and osteoprogenitors are recruited, differentiate into osteoblasts, and start to mineralize the ECM, which can be remodeled to regain form and function of the bone tissue (Bielby et al., 2007; Loi et al., 2016).

Immuno-Modulatory Interventions to Promote Bone Healing

Both the pro- and anti-inflammatory phases are necessary for successful healing and excessive downregulation of either phase can inhibit the healing cascade (Schmidt-Bleek et al., 2015; Rapp et al., 2016). Thus, factors influencing either the pro-inflammatory or anti-inflammatory phase have been shown to largely affect the fracture healing outcome. Most often, the aim of the intervention lays on re-establishing the balance between pro- and anti-inflammatory response, with a focus on preventing a prolonged pro-inflammatory phase.

In mice for example, the transplantation of MSCs to mediate bone repair, was found to be considerably improved with additional downregulation of TNF- α and INF- γ via the administration of aspirin or systemic infusion of Tregs (Liu et al., 2011). The authors propose that INF- γ , secreted by the recipients T-helper cells, downregulate runt-related transcription factor 2 (Runx-2) in the transplanted MSCs and thus inhibits the osteogenic differentiation (Liu et al., 2011).

The administration of a prostacyclin (PGI₂) analog has been shown to improve fracture healing by downregulating pro-inflammatory cytokine release from CD8+ and CD4+ T cells while promoting the anti-inflammatory response via an increased numbers of M2 macrophages in the fracture gap (Wendler et al., 2019). Similarly, Tregs are usually associated with anti-inflammatory functions and have been shown to inhibit osteoclastogenesis by cell-cell contact and the expression of transforming growth factor-beta (TGF- β), IL-4, and IL-10 (Zaiss et al., 2007). Consequently, the adoptive transfer of Tregs improved fracture healing, when they overcome a certain CD8+ effector T (T_{eff}) cell/Treg ratio (Schlundt et al., 2019). Additionally, an elevated concentration of terminally differentiated CD8+ effector memory T cells (CD8+CD11a++CD57+CD28-T cells), a subpopulation known for high pro-inflammatory properties, at the fracture gap significantly delayed the fracture healing cascade in patients. While the depletion of CD8+ T cells by specific antibody therapy was shown to enhance the endogenous fracture regeneration in a murine model, an adoptive transfer of CD8+ T cells however impaired the fracture repair (Reinke et al., 2013). Higher concentration of effector memory T cells can be found in a

more experienced immune system, which usually has a higher propensity for pro-inflammatory effects (Bucher et al., 2019).

Therefore, gaining control over the immune system may lead to promising therapeutic approaches in bone pathologies and regeneration.

Mechanical Properties Are Regulating the Immune Response Following Fracture

Until now, therapeutic approaches have mostly focused on biological factors that influence the immune response. Thus, it has mostly been overlooked so far, how different biophysical and mechanical properties during the healing cascade might directly or indirectly influence the immune response and subsequently suppress or promote bone healing.

At the tissue level, the stability of the fracture gap seems to be closely related to the healing outcome. A stable fracture fixation (minimizing interfragmentary movement) most likely leads to direct bone formation, a moderately fixed fracture may heal by endochondral ossification and an unstable fixed fracture gap almost suppresses the bone healing cascade (Epari et al., 2006, 2007; Claes et al., 2012). How are these mechanically different situations related to the healing process? At the cellular level, cyclic tensile strain applied to osteoblasts was found to induce higher OPG and lower RANKL expression (Tang et al., 2006). Conversely, dynamic compression of osteoblasts *in vitro* showed higher expression of pro-inflammatory IL-6, IL-8, RANKL, and lower expression of OPG, possibly delaying a bone healing response by increased osteoclast activity (Sanchez et al., 2009, 2012). Moreover, compression of osteoblasts stimulated expression of the IL-17 gene. IL-17 has been shown to increase the expression of IL-1, IL-6, IL-8, IL-11 in osteoblast cells in an autocrine manner (Zhang et al., 2010). A function of IL-17 is to stimulate the expression of RANKL on osteoblasts, thereby increasing osteoclastogenesis and osteoclast activity, which may eventually lead to bone erosion (Miossec and Kolls, 2012). Furthermore, systemically elevated levels of IL-8 have been found after bone fracture, and it has been shown that mechanically stimulated osteoblasts release IL-8, which may be the source of the systemically elevated IL-8 levels (Perl et al., 2003; Reinke et al., 2013). IL-8 appears to be involved in several regenerative processes, such as the regulation of revascularization of newly formed tissue after a fracture (Koch et al., 1992; Hou et al., 2014). A lack of sufficient revascularization has been associated with poorer healing outcome (Schmidt-Bleek et al., 2015). In addition, IL-8 is also a chemokine, that attracts polymorphonuclear leukocytes (PMNs) to the site of inflammation (Kolar et al., 2011; Vasconcelos et al., 2016). The attraction of PMNs to the fracture, does not *per se* negatively affect the healing outcome, since an initial pro-inflammatory response is required for a good healing outcome (Schmidt-Bleek et al., 2015). Moreover, IL-8 has been shown to be important for stem cell recruitment and increased chondrogenic gene expression and bone morphogenetic protein (BMP) sensitivity, thereby positively influencing endochondral bone healing (Lin et al., 2019). It appears, that at least the osteoblasts are actively involved in the perception of cyclic stretch and respond to it by secreting pro-inflammatory cytokines.

Moreover, osteoblast proliferation has been found to occur mostly during the pro-inflammatory phase (Gerstenfeld et al., 2003; Watanabe et al., 2009)

Beyond the above-mentioned aspects, little is known about the regulation of the immune-mechanical interplay in bone healing. However, in other organ systems a more profound knowledge on mechano-immunological coupling has been gathered, which we would like to compare with the bone-related knowledge in order to gain a more comprehensive understanding on how immune-mechanics influences tissue regeneration. Although lung, heart, and skin tissue seem at the first very different, the universal feature that we think allows for comparison and extrapolation of information to bone tissue is their remarkable mechanosensitivity, constant exposure to high forces and stress, and of course all tissues are infiltrated by the immune system in health and disease.


LUNG INFLAMMATION FOLLOWING INJURY IS MODULATED VIA IMMUNE-MECHANICS

Positive pressure mechanical ventilation is a life-saving therapy for patients with acute lung injury (ALI), or its more severe form, acute respiratory distress syndrome (ARDS) (Kuipers et al., 2011). However, it can also lead to ventilator-induced lung injury (VILI), which exacerbates pre-existing lung damage and inflammation (Tremblay and Slutsky, 2006; Kuipers et al., 2012) (**Figure 2A**). To reduce the release of inflammatory mediators, it has been suggested that mechanical ventilation with lower tidal volumes should be preferred over high tidal volumes, even though this may cause acidosis and reduced arterial oxygenation (Blanch et al., 1994; Brower et al., 2000). A deeper understanding of how mechanical overdistension of the lung causes inflammatory processes is therefore highly desirable to minimize the side effects of mechanical ventilation in patients with already severe conditions. Moreover, the mechanisms we may learn from understanding how mechanical ventilation induces pro-inflammatory reactions, is an excellent tool to decipher immune-mechanical principles.

ECM-Components as Mechano-Transducer in Inflammation

One possible way how mechanical stimuli can alter the immune response is by changing the ECM-composition. It has been shown, that excessive mechanical ventilation increased the amount of proteoglycan and biglycan in the lung (Al-Jamal and Ludwig, 2001). Biglycan was found to bind to the Toll-like receptor (TLR) 4 on macrophages and to increase the expression of pro-inflammatory TNF- α and macrophage inflammatory protein (MIP)-2 (also known as CXCL2, a chemokine/cytokine sharing the same receptor with human IL-8), thereby triggering or promoting an already ongoing inflammatory response (Schaefer et al., 2005) (**Figure 2B**).

Another ECM component often associated with trauma is hyaluronic acid (HA). HA exists as a more immunological



inert high molecular weight form (HMW-LA), and as pro-inflammatory low molecular weight hyaluronic acid (LMW-HA) form. Mechanical ventilation of rat lungs with high tidal volumes but not low tidal volumes led to an accumulation of LMW-HA in lungs (Mascarenhas et al., 2004). On a cellular level, stretching of pulmonary fibroblasts or alveolar epithelial type II cells (AEIIs) induced *in vitro* the expression of hyaluronan synthase 3 (HAS3); the enzyme that produces LMW-HA (Itano et al., 1999; Mascarenhas et al., 2004; Heise et al., 2011). LMW-HA might then bind to TLR2 or TLR4 on macrophages and increase the expression of IL-8 and IFN- β and finally induces a pro-inflammatory response after mechanical ventilation with high tidal volumes (Jiang et al., 2005; Black et al., 2013) (**Figure 2C**). Moreover, LMW-HA has been shown to be sufficient for inducing epithelial to mesenchymal transition (EMT) of AEIIs, often associated with aberrant wound healing and fibrosis (Heise et al., 2011). Interestingly, stretched fibronectin conformation has been found to bind to TLR4 on lung cancer cells and increased IL-8 expression and release (Cho et al., 2020).

This suggests that not only the presence of fibronectin, but also the mechanical environment and consequently the conformation of fibronectin may have important implications. A stretched fibronectin conformation could be an indicator of mechanically

stressed tissue and could be involved in triggering inflammation by increasing the levels of IL-8.

Non-hematopoietic Cells as Mechano-Transducer in Inflammation and Regeneration

The yes-associated protein 1 (YAP)/ WW domain-containing transcription regulator 1 (WWTR1, also known and more commonly referred to as TAZ) pathway integrates mechanical stimuli, soluble signals, and metabolic information to regulate cell proliferation, cell plasticity and stemness (Totaro et al., 2018; Pocaterra et al., 2020). Although not exclusively regulated by mechanical stimuli, the YAP/TAZ pathway has been shown to be highly mechanosensitive (Dupont et al., 2011). YAP/TAZ can translocate into the cell nucleus by cell-cell contact signaling and/or ECM-integrin-focal adhesion kinase (FAK) signaling (Dupont et al., 2011; Pocaterra et al., 2020) or forces can directly trigger the entry of YAP into the cell nucleus by regulating transport through nuclear pores (Elosegui-Artola et al., 2017). Additionally, cyclic stretch has been shown to increase YAP nuclear localization in AEIIs and subsequently their proliferation rate (Codelia et al., 2014) (**Figure 2D**). This could have a significant impact on lung regeneration, as AEIIs

are considered stem/progenitor cells for the alveolar epithelium in homeostasis and regeneration (Hogan et al., 2014). Hence, an increased proliferation of ACEIIs via YAP/TAZ triggered by cyclic stretching could also promote the regeneration process in other organs. Accordingly, YAP/TAZ inhibition in AEIIs mice showed increased IL-1 β levels and prolonged fibrotic lesions and delayed regeneration compared to normal YAP/TAZ mice (Liu et al., 2016; LaCanna et al., 2019).

In addition to AEIIs, pulmonary fibroblasts also proved to be mechano-sensitive and YAP/TAZ signaling could trigger myofibroblast differentiation and activation. Pulmonary fibroblasts responded to an increase in matrix stiffness to a pathophysiological level with nuclear localization of YAP/TAZ (Liu et al., 2015) (**Figure 2E**). Overexpression of YAP/TAZ in pulmonary fibroblasts resulted in increased fibrogenic potential via an enhanced proliferation rate and matrix synthesis (Liu et al., 2015). Blocking YAP/TAZ signaling in pulmonary fibroblasts reversed the deposition, contraction, and stiffening of the matrix *in vitro* and was able to reverse bleomycin-induced pulmonary fibrosis *in vivo* (Haak et al., 2019). These findings suggest that YAP/TAZ might be an important player during the anti-inflammatory response leading to ECM formation, but also, if not temporal limited, to fibrosis. Therefore, YAP/TAZ could be an attractive target to control the regeneration process by downregulating fibrotic processes and directing the healing process toward regeneration rather than toward scarring.

Immune Cells as Mechano-Transducer in Inflammation and Regeneration

Alternatively, the immune cells might directly sense mechanical properties themselves and respond with pro-inflammatory reactions. A possible way how cells perceive forces such as shear, stretch, osmotic swelling and shrinkage, stiffening and surface expansion is via the transient receptor potential vanilloid 4 (TRPV4). When activated, TRPV4 causes a Ca²⁺ influx and initiates Ca²⁺ mediated signaling cascades (Strotmann et al., 2000; Vriens et al., 2004; Wu et al., 2007; Shin et al., 2012; Jo et al., 2015; Baratchi et al., 2016; Lee et al., 2019; Rosenbaum et al., 2020). Signaling via TRPV4 in VILI was also shown to activate macrophages, increase phagocytosis and caused secretion of pro-inflammatory mediators such as reactive oxygen species (ROS) and nitrogen species (NOS) (Hamanaka et al., 2010). Overall, this suggests that increased matrix stiffness as a result of inflammation, is sufficient to increase pro-inflammatory macrophage activity (Okamoto et al., 2018; Sridharan et al., 2019). Interestingly, there might be a crosstalk between lipopolysaccharide (LPS) mediated activation of macrophages via TLR4 and TRPV4. Blocking TRPV4 signaling prevented LPS-induced acute lung injury and resulted in lower levels of pro-inflammatory molecules such as TNF- α , IL-6, and ROS (Li et al., 2019). Moreover, TRPV4 may lead to a change in the TLR4 signaling pathway, that promotes phagocytosis and bacterial clearance (Scheraga et al., 2020). This nicely demonstrates that there is a crosstalk between pro-inflammatory pathways (TLR4) and mechanosensitive pathways (TRPV4).


PIEZO1 is another mechanosensitive Ca²⁺ ion channel involved in mechanically triggered inflammation. PIEZO1 is highly expressed on macrophages and other immune cells enabling them to sense omnidirectional pressure (Tolar and Wack, 2019). Cyclic hydrostatic pressure can activate an inflammatory response in macrophages and monocytes via PIEZO1 (Solis et al., 2019). Mechanistically, cyclic pressure activates PIEZO1 Ca²⁺ influx in monocytes, thereby increasing the expression and secretion of endothelin1 (ET1) (Solis et al., 2019). ET1 then signals back in an autocrine manner, stabilizing hypoxia-inducible factor 1-alpha (HIF-1 α), which ultimately increases the expression and secretion of the chemoattractant CXCL2 in monocytes (Solis et al., 2019). Finally, neutrophils migrate from the blood along the CXCL2 gradient into the lung and initiate a pro-inflammatory response (Solis et al., 2019) (**Figure 2F**). In summary, this shows that mechanical stimuli play a role in the activation of innate immune cells and in the case of PIEZO1 and TRPV4 can trigger an inflammatory response.

HEART INFLAMMATION IS MODULATED VIA IMMUNE-MECHANICS

In contrast to most other tissues, heart tissue appears to preferentially form fibrotic tissue rather than new muscle tissue in response to damage (Russo et al., 2019). It is assumed that the fibrotic tissue spatially confines the injured area and thus prevents further damage of healthy tissue under constant mechanical stress (Russo et al., 2019). In contrast to healthy tissue, scar and fibrotic tissue have a different structure in their ECM and are probably also more readily formed (Willyard, 2018). Here, the fastest possible restoration of “functional tissue” seems to be of crucial importance, and a regeneration pause is not an option. Consequently, the preferred strategy of forming fibrotic tissue after injury allows the maintenance of structural support at the expense of an active contribution to the dynamic function of the heart. The use of mechanical unloading strategies via axillary pumps, primarily used as a mechanical circulatory support in cardiogenic patients, have been shown to boost endogenous repair in form of reduction in immune cell presence, modulation of the ECM and cardiac metabolism, which go along with a drop in integrin expression (Spillmann et al., 2019; Tschöpe et al., 2019). This example from cardiac regeneration (or lack thereof) illustrates how local immune (im-)balance regulate tissue maintenance and fibrosis after injury while being under constant mechanical stress (**Figure 3A**).

ECM-Components as Mechano-Transducer in Inflammation

Some ECM components in the heart tissue can serve as mechano-transducers. For example, cells responded to a stiffer ECM with increased expression of heparan sulfate proteoglycan agrin (Chakraborty et al., 2017). Since its discovery in the early 90s, many functions of agrin have been identified. The most prominent and one of the earliest discoveries was the role of agrin as a key regulator of the formation of postsynaptic structures at the neuromuscular junction (Ruegg and Bixby,



1998). Surprisingly, agrin was also found on the surface of T cells and contributes as co-stimulatory signal, as well as to the strengthening of the immunological synapse between T cells and antigen-presenting cells (APCs) (Bezakova and Ruegg, 2003). Agrin was also found in the growth plate of bone and was shown to be highly expressed by chondrocytes (Hausser et al., 2007). It was later suggested that agrin is also an important extracellular component of the ECM-integrin-FAK axis and therefore helps to connect the cellular cytoskeleton to the ECM to ensure proper force transmission and matrix stiffness transfer (Haak et al., 2019). Moreover, agrin appears to be essential for the regenerative abilities of the neonatal mouse heart by promoting the division of cardiomyocytes via YAP- and ERK-mediated signaling (Bassat et al., 2017) (Figure 3B). *In vivo*, a single administration of agrin in adult mice after myocardial infarction promoted cardiac regeneration by pleiotropic effects such as cardiomyocyte proliferation, immune modulation and angiogenesis (Bassat et al., 2017). For example, agrin mediated signaling has been shown to cause increased actin cytoskeletal rearrangements, with the result that agrin deficiency impaired phagocytosis in macrophages and ultimately the clearance of tissue debris after injury (Mazzon et al., 2012). Adult mammalian hearts generally have a very low regenerative capacity, likely due to the inability of fully differentiated cardiomyocytes to proliferate (Kikuchi and Poss, 2012; Bigotti et al., 2020). Therefore, therapies that promote

the proliferation of cardiomyocytes via YAP/TAZ-signaling such as the administration of the ECM-component agrin, are an exciting new approach, whose full potential, including its effect on immune cells, could be the focus of further research.

Non-hematopoietic Cells as Mechano-Transducer in Inflammation and Regeneration

Current findings suggest that inflammation is likely to have both negative and positive effects on regeneration, with an initial pro-inflammatory phase being necessary to start the regeneration process, but an uncontrolled and prolonged inflammatory response is detrimental (Coggins and Rosenzweig, 2012). It is therefore not surprising that macrophages and monocytes are essential for a good healing outcome (Morimoto et al., 2006; Van Amerongen et al., 2007; Leblond et al., 2015). One way to promote the local accumulation of monocytes is through cyclic stretching. For example cyclic stretching of rat neonatal cardiomyocytes resulted in increased TLR4 expression and consequently to more TLR4 mediated adhesion of monocytes to cardiomyocytes (Shyu et al., 2010). Additionally, it has been shown that cyclic stretching of rabbit cardiomyocytes increased IL-18 expression (Yoshida et al., 2014). IL-18 is a potent growth factor that induces cardiomyocyte hypertrophy

in vitro (Chandrasekar et al., 2005) and myocardial hypertrophy *in vivo* (Reddy et al., 2010). However, it is believed that the initiation and control of inflammation is regulated by cells in the epicardium such as fibroblast and not by myocytes (Ramjee et al., 2017). For example, *in vitro* mechanical stretching, which mimics cardiac dilatation during heart failure, induces fibroblast activation and stimulates not only the production of ECM but also the production of chemokines, such as the monocyte chemoattractant protein (MCP)-1 and MCP-3, and triggers typical inflammatory pathways (Lindner et al., 2014).

Although an improved healing outcome by YAP/TAZ signaling in cardiomyocytes is often attributed to increased myocyte proliferation (Lin et al., 2014; Leach et al., 2017; Wang et al., 2018), YAP/TAZ may also play an immunomodulatory role in regeneration. It has been found that the loss of YAP/TAZ in the epicardium led to an enhanced fibrotic response and was associated with a decrease in IFN- γ expression, which is a direct transcriptional target of YAP/TAZ (Ramjee et al., 2017) (Figure 3C). Interestingly, IFN- γ has also been described to downregulate fibrotic processes. For example, IFN- γ was shown to increase the number of regulatory T cells after myocardial infarction and lead to less fibrosis (Ramjee et al., 2017) (Figure 3C). Furthermore, IFN- γ counteracts TGF- β induced activation of myofibroblasts and leads particularly to the expression of chemokines attracting pro-inflammatory monocytes (Pappritz et al., 2018). Thus, the role of IFN- γ seems to be ambivalent and might also depend on the presence of other cytokines and cells.

In a pressure overload model by transversal aortic constriction (TAC) the concentrations of TNF- α and IL-1 β were significantly elevated after 3 days and increased levels of the chemokines MCP-1, MIP-2, and TGF- β were found after 7 days (Xia et al., 2009). In contrast to sudden trauma-induced tissue damage, TAC induced injury takes time to develop, which could explain the long span needed for upregulation of cytokines and chemokines. Nevertheless, it has been shown that mechanically induced secretion and activation of TGF- β plays an important role in the activation and transformation of fibroblasts into myofibroblast (Chen and Frangogiannis, 2013; Van Linthout et al., 2014; Pappritz et al., 2018; Steffens et al., 2020). Although myofibroblasts are required for the initial healing response, excessive and prolonged activity of myofibroblasts drives scar formation and fibrosis (Gabbiani, 2003; Darby et al., 2014). In the heart, this increased and prolonged activity of myofibroblasts might be the intrinsically preferred mode of healing after injury, to re-establish mechanical stability as quickly as possible. Recently, it has been found that the increased TGF- β levels after TAC, reduced MMP-8 expression in fibroblasts, which resulted in matrix preservation rather than matrix degradation as well as decreased macrophage-driven inflammation and attenuated cardiomyocyte apoptosis (Russo et al., 2019) (Figure 3D). MMP-8 is known to generate pro-inflammatory matrix fragment proline-glycine-proline (PGP) that promote recruitment and activation of neutrophil cells and polarize macrophages into a pro-inflammatory phenotype (Lin et al., 2008).

Whether this healing mechanism of the heart is something cell- and tissue-inherent, or whether the constant mechanical stress contributes to the preferred fibrotic healing outcome, is still to a large part elusive. Further investigations of the mechanisms involved could help to better understand fibrotic processes and lead to new therapeutic approaches that make use of the mechanical manipulation of the immune system.

Immune Cells as Mechano-Transducer in Inflammation and Regeneration

Surprisingly, either the direct influence of mechanical stimuli on immune cells in the heart has so far been neglected, or it does not play an important role in the development of cardiac pathologies and cardiac regeneration. However, a very recently published study indicates that in aortic valve stenosis causing high shear stress, PIEZO1 can be activated on **monocytes**, resulting in calcium influx and pro-inflammatory responses, such as increased expression of IFN- β 1, IL-1 β , IL-1ra, IL-6, and IL-12 (Baratchi et al., 2020) (Figure 3E).

Recently, YAP/TAZ signaling in immune cells has been described to reduce fibrosis in the heart (Geng et al., 2017; McGeachy, 2017). TAZ has been found to play a regulatory role in the balance of Th17-Treg by promoting acetylation and degradation of the Treg transcription factor Foxp3 (Geng et al., 2017; McGeachy, 2017). Although Tregs produce TGF- β that promotes fibrotic processes, it is thought that Tregs have a more dominant anti-fibrotic function by down-regulating the activity of Th2 cells; one of the major cellular contributors to fibrosis (Gieseck et al., 2018; Bal and Stadhouders, 2019).

This suggests that immune cells are also able to sense mechanical stimuli directly in heart tissue, similar to findings in lung tissue. To what extent this plays a role in initiating an inflammatory response in other organs could be a promising focus of further research.

SKIN INFLAMMATION IS MODULATED VIA IMMUNE-MECHANICS IN WOUND HEALING

The skin is surprisingly mechano-sensitive (Kwon et al., 2018). Probably the most prominent example is the thickening of the epidermis and callus formation after repeated loading at the same site (Goldstein and Sanders, 1998; Silver et al., 2003; Carlson, 2006; Kim et al., 2010). Thickening of the epidermis is suggested to protect the underlying tissue from the excessive stress and to contribute to a better distribution of forces (Thomas et al., 1985; Kim et al., 2010). Impressively, Karl Langer recognized as early as 1861 from the observation that circular hole punches in the skin became elliptical, that skin had oriented pre-tension lines (Langer, 1978a,b,c). Additionally, in the skin, collagen bundles were found to be oriented along the main tensile load-lines/Langer lines (Wilhelmi et al., 1999; Silver et al., 2003). As a result, surgical incisions are often performed parallel to the Langer lines instead of perpendicular to reduce scarring (Wilhelmi et al., 1999). Besides the passive pre-tension of the skin, active stress generation and contraction, mainly produced by

(myo-)fibroblasts, are important for wound closure and reduced scarring. It has been proposed to link mechano-transduction to various skin pathologies such as keloids and hypertrophic scars, and subsequently targeting mechanoreceptors has already been suggested as novel treatment options (Ogawa and Hsu, 2013). For example, pressure has been found to positively influence the wound healing phase by upregulating the apoptosis of myofibroblasts (Costa et al., 1999; Renò et al., 2003). At the beginning of the wound-healing phase, myofibroblasts are thought to have a positive effect by increasing ECM production and its contractile function bringing the two wound edges back together again (Hinz et al., 2012; Humphrey et al., 2014). However, it has been found that excessive matrix production and stiffening of the wound could initiate a vicious cycle leading to prolonged and excessive myofibroblast activity, which ends in fibrosis and hypertrophic scarring (Hinz, 2007; Darby et al., 2014). Interestingly, the mechanical tension applied to a healing wound was already sufficient to produce hypertrophic scarring in mice (Aarabi et al., 2007). Cyclic tensioning of keratinocytes was shown to increase proliferation, while cyclic compression downregulated cell division of keratinocytes (Silver et al., 2003). Thus, it has been known for a relatively long time that the mechanical environment and properties are playing a crucial role for skin wound healing (Tomasek et al., 2002; Silver et al., 2003) (Figure 4A). However, the relationship between the mechanical environment and the immune system, which regulates the healing response, has only recently become the focus of research.

ECM-Components as Mechano-Transducer in Inflammation

The compression of hypertrophic scars was found to release significantly higher levels of IL-1 β compared to normal scars (Renò et al., 2003). This could indicate that the cells within the hypertrophic scar are already pre-sensitized to mechanical stimuli or that more immune cells could be found in the hypertrophic scar. It has been found that the ECM component fibulin-5 plays an important role in inflammation-associated skin pathologies (Nakasaki et al., 2015). The loss of fibulin-5 not only led to a decrease in tissue stiffness, but also reduced the inflammatory response (Nakasaki et al., 2015) (Figure 4B). The increase in stiffness from 2 to 5 kPa was sufficient to enhance the pro-inflammatory response of dermal fibroblasts with increased expression of the chemoattractants CXCL5, CXCL10, and CXCL16, and resulted in increased numbers of macrophages, T cells and mast cells (Nakasaki et al., 2015). Targeting fibulin-5 could prevent the stiffening of skin ECM and could be an attractive target to stop the fibrotic vicious cycle. Another important ECM component that converts mechanical stimuli into an inflammatory response is fibronectin. Dermal fibroblasts have been shown to recognize stretched fibronectin as damage associated molecular patterns (DAMPs), which leads via the TLR4/NF- κ B pathway to increased IL-8 expression (Zheng et al., 2020) (Figure 4C). Moreover, stretched fibronectin showed a higher binding affinity for IL-7 and local accumulation of IL-7 may play an important role in the regulation of the immune response (Ortiz Franyuti et al., 2018) (Figure 4C). IL-7 was found

to be essential for the maintenance of memory Tregs (mTregs) in the skin (Gratz et al., 2013).

Non-hematopoietic Cells as Mechano-Transducer in Inflammation and Regeneration


It is suggested that the focal adhesion kinase (FAK) connects mechanical forces with skin fibrosis via inflammatory signals. In fibroblasts, FAK became activated after a skin injury and its activity can be potentiated by mechanical stretching (Wong et al., 2012). An elevated FAK signaling in fibroblasts then amplified MCP-1/CCL2 signaling and subsequently led to the migration of monocytes toward the wound and the secretion of inflammatory cytokines such as TGF- β ; which possibly triggers a vicious cycle that ends in a hypertrophic scar (Wong et al., 2012). FAK signaling in dermal fibroblasts also proved to be an important player in pathological angiogenesis in hypertrophic scarring (Gao et al., 2019). Recently, TRPV4 was discovered as an important mediator in skin pathologies. In concert with TGF- β , TRPV4 was found to be essential for matrix-stiffness mediated differentiation of dermal fibroblasts into myofibroblasts (Sharma et al., 2017) (Figure 4B) and epithelial-mesenchymal transition (EMT) of epidermal keratinocytes (Sharma et al., 2019). Higher levels of TGF- β could also be triggered by increased ECM stiffness: In human dermal fibroblasts, YAP/TAZ works in conjunction with the transcription factor activator protein (AP)-1 to regulate TGF- β signaling (Qin et al., 2018). Accordingly, YAP/TAZ has already been proposed as an attractive new therapeutic target for reducing fibrosis (Duscher et al., 2014).

In summary, higher ECM stiffness, possibly due to higher concentrations of fibulin-5, increases TGF- β levels and monocyte counts, which promotes the differentiation of dermal fibroblasts into myofibroblasts and EMT of keratinocytes, and ultimately leads to skin fibrosis or hypertrophic scarring. This shows that the increased ECM-stiffness in fibrosis might not only be a consequence of the inflammatory process, but rather plays an active role in initiating the inflammatory process.

DISCUSSION

The Described Immune-Mechanical Principles in the Lung, Heart, and Skin May Also Play a Role in Bone Regeneration

Forces acting on the fracture hematoma/callus (Figure 5A) likely influences the healing outcome by regulating the immune response via: ECM components (Figures 5B,C), indirectly via the sensation of forces by osteoblasts and subsequent cytokine release (Figure 5D), or forces are directly sensed by immune cells (Figure 5D). Fibronectin is an important and ubiquitous component of the ECM with structural and functional roles in homeostasis and regeneration (To and Midwood, 2011). So far, fibronectin has also been found in the growth plate of bone and during new bone formation, which may promote osteoblast differentiation and the mineralization process (Yang et al., 2020). Although higher levels of fibronectin were found in osteoporotic




patients, the downregulation of a fibronectin receptor (β_1 -integrin) on the cell surface reduced fibronectin mediated signaling in osteoblast and might explain the loss of bone despite higher fibronectin concentrations (Yang et al., 2020). Whether these higher fibronectin levels contribute to the osteoporotic pathology via immunomodulatory mechanisms, has not yet been fully elucidated. However, lung cells and dermal fibroblasts have been shown to bind the stretched fibronectin conformation via TLR4, which led to increased IL-8 levels (Cho et al., 2020; Zheng et al., 2020). Since IL-8 also seems to play an important role in bone healing (Lin et al., 2019), mechanical manipulation of the fibronectin conformation may also be an attractive therapeutic target (**Figure 5B**). Furthermore, as in the skin, the mechanical niche during fracture healing could influence the healing outcome via the concentration of IL-7 by the conformational regulation of fibronectin. Subsequent, as described in the skin, this could promote the anti-inflammatory phase via the stabilization of memory Tregs (Gratz et al., 2013) (**Figure 5B**).

Another ECM component that is found in bone and is involved in the regulation of the immune system is agrin (Bezakova and Ruegg, 2003; Hausser et al., 2007). In the heart, the administration of agrin was able to increase the proliferation rate of cardiomyocytes via YAP/TAZ signaling, resulting in improved healing response (Bassat et al., 2017). Potentially, this therapeutic approach could also be translated to enhance bone healing. It has already been found that agrin plays an important role in osteochondral regeneration (Eldridge et al.,

2020) (**Figure 5C**). Since critical-sized bone defects heal via endochondral ossification (Dennis et al., 2015), which involves the formation of cartilage and its subsequent transformation into bone, promoting cartilage formation to heal critical-sized bone defects could be an appropriate approach and is already a focus of current research (Petersen et al., 2018; Lin et al., 2019).

Moreover, the mechanosensitive ion channels PIEZO1 and TRPV4 might also play a role in bone healing. Similar to the findings in lung tissue, IL-8 also attracts PMNs in bone tissue, which have been shown to be a critical player in bone fracture healing (Kovtun et al., 2016; Lin et al., 2019). It could be hypothesized that not only the lung monocytes perceive cyclic pressure via PIEZO1, but also the monocytes in bone healing, which would lead to a contribution to the IL-8 levels found in the fracture hematoma (Kolar et al., 2011) (**Figure 5E**).

In addition, signaling via TRPV4 in lung macrophages has been found to increase ROS levels (Hamanaka et al., 2010; Li et al., 2019) (**Figure 5E**). It could be hypothesized that similar processes may also occur during bone healing. ROS has been shown to interfere with the osteogenic potential of MSCs and the administration of antioxidant resulted in improved bone regeneration *in vivo* (Geißler et al., 2013). However, the situation might be more complex. Recently, it was found that MSCs embedded in a 3D collagen hydrogel might sense nanovibrations via integrins and mechanosensitive ion channels, that increased their osteogenic differentiation (Tsimbouri et al., 2017). Later, the same authors suggested



that MSCs stimulated with nanovibrations produce ROS during osteogenic differentiation, but the ROS themselves do not appear to be a driver of osteogenesis (Orapiriyakul et al., 2020). Whether ROS is only a byproduct during bone healing or whether it takes an active role in this process is still elusive and further studies are required. If ROS does actively influence bone healing, it would be interesting to investigate, whether macrophages in the bone fracture hematoma also sense mechanical stimuli via mechanosensitive ion channels, such as TRPV4, and also produce ROS as they do in lung injuries.

During bone healing, the local mechanical niche changes a lot. The fracture hematoma is soft and relatively dense. While in the subsequent healing phases the stiffness increases steadily. This might have different implications on the cellular behavior of non-hematopoietic and immune cells. On soft and confined substrates, cells tend to show spherical morphology and cytoplasmic localization of YAP/TAZ, while more open and stiff substrates allows for cytoskeletal re-arrangements, cell spreading and nuclear localization of YAP/TAZ (Pancieri et al., 2017). Interestingly, mechanical confinement was not only found to restrict cell spreading, but also reduced the osteogenic differentiation potential of MSCs via a lack of TRPV4 activation (Lee et al., 2019). Also osteoclasts activity is dependent on cytoskeletal re-arrangements (Takayanagi, 2007).

Recent evidence also points out, that T cells are dependent on cytoskeletal re-organization for an efficient immunological synapse formation and T-cell activation (Colin-York et al., 2019; Blumenthal and Burkhardt, 2020). Whether this has also implications in a non-antigen specific, sterile inflammation process, as found during regeneration, could be the scope of future research.

These examples encourage that also during bone regeneration the immune response is modulated by the ECM composition and the mechanical niche. In particular, the ECM components fibronectin and agrin could also have immunomodulatory functions in bone, and TRPV4 and PIEZO1 on macrophages could be important mechano-transducers and thereby influence the outcome of bone healing.

Summary

The understanding of regeneration has come a long way. Today, both the adaptive and the innate immune system are now considered to play a central role in healing and regeneration (Medzhitov, 2008; Julier et al., 2017). Although Julius Wolff and Karl Langer have already in the 19th century put forward the hypothesis that mechanical forces contribute significantly to the biological functions of tissue adaptation and healing (Langer, 1978a,b,c; Wolff, 1986), it is only recently that the




FIGURE 6 | Mechanically or ECM triggered crosstalk between non-hematopoietic cells—immune cells—ECM regulates cell behavior. Immune cells can respond either directly or indirectly to mechanical forces. When immune cells sense the mechanical niche directly (1), they can transmit this information by the cytokine secretion, which either acts in an autocrine manner or on non-hematopoietic cells (2). This connects immune cells within a wound environment not only with each other, but also to non-hematopoietic cells such as smooth muscle cells, fibroblasts, epithelial cells and endothelial cells. Non-hematopoietic cells might then start to secrete ECM-components and change the ECM-composition (3) and subsequently its biophysical properties (4). This could close a feedback loop, and immune cells sense the altered ECM-composition/biophysical cues. Depending on the secreted cytokines of the immune cells, this feedback could be the first step into a vicious cycle or alternatively the first step toward a successful restoration of homeostasis. In the indirect case, tissue-resident cells embedded in the ECM first perceive the mechanical stimuli (5), translate them into a cellular response and might then transmit the stimuli to nearby immune cells by secreting cytokines and/or chemokines (6).

mechanical environment and its influence on a central part of the biological component, the local immune response, has been investigated. Immunomodulatory mechano-transduction can either occur directly in immune cells or indirectly in non-hematopoietic cells. In either case, there is extensive crosstalk between immune and non-hematopoietic cells to determine the appropriate response for a given situation (**Figure 6**). The crosstalk and communication cycle could be proposed as follows: Immune cells perceive the mechanical niche at least by means of integrin, PIEZO1 and TRPV4 [**Figure 6**(1)] and then release cytokines and chemoattractants, which act on nearby immune cells or on non-hematopoietic cells [**Figure 6**(2)]. In response, non-hematopoietic cells alter the components and structure of the ECM [**Figure 6**(3)], which is often accompanied by a change in the mechanical niche [**Figure 6**(4)]. Alternatively, non-hematopoietic cells are assumed to perceive the mechanical niche at least by means of integrins, PIEZO1 and TRPV4 [**Figure 6**(5)] and then transmit the information of the mechanical niche to immune cells or non-hematopoietic cells via cytokines and chemoattractants [**Figure 6**(6)]. This might then alter the activation, differentiation, proliferation and migration of the cells.

In most situations, unphysiological mechanical stimuli or mechanical overstimulation resulted in the release of common pro-inflammatory mediators. Specific ECM components are actively involved in immune regulation, either by acting as DAMPs [LMW-HA (Jiang et al., 2005; Black et al., 2013), fibronectin (Cho et al., 2020), and biglycan (Schaefer et al., 2005)], or by stabilizing ECM-cell interactions [fibulin-5 (Nakasaka et al., 2015), and agrin (Haak et al., 2019)].

In all the tissues described, non-hematopoietic cells often respond to abnormal mechanical stimuli by secreting chemoattractants and cytokines that lure immune cells out from the vasculature and in most cases the secreted cytokines create a pro-inflammatory niche. Within the cell, calcium mediated signaling pathways, FAK and YAP/TAZ are often involved in the translation of the perceived mechanical environment into a cellular response. It is already known that YAP/TAZ plays an important role in the regulation of cell proliferation, and a targeted manipulation of this signaling pathway could help to increase the number of cells after injury (Moya and Halder, 2019). In addition, YAP/TAZ signaling in cardiomyocytes and AEIIs has been found to reduce fibrosis (Liu et al., 2016; Ramjee et al., 2017; LaCanna et al., 2019). In contrast, YAP/TAZ signaling in fibroblasts has been shown to promote fibrosis in the lung, and skin (Liu et al., 2015; Qin et al., 2018; Haak et al., 2019). This indicates that YAP/TAZ signaling can promote regeneration as well as cause fibrosis and a cell specific therapy approach may be necessary to exploit YAP/TAZ signaling.

Immune cells most often perceived the mechanical environment via mechanosensitive ion channels such as PIEZO1 (Solis et al., 2019; Baratchi et al., 2020) and TRPV4 (Hamanaka et al., 2010; Li et al., 2019). The signaling via these receptors provoked at least in monocytes and macrophages the secretion of pro-inflammatory mediators. Interestingly, in most cases when immune cells sensed the mechanical stimuli themselves, they reacted with a pro-inflammatory response. This could provide a possibility for intervention where the restoration of a mechanical niche to pre-injury level could be used to downregulate a pro-inflammatory response.

Conclusion and Future Directions

The above findings suggest that the success of using biomaterials for cell delivery or as a cell modulating *in vivo* micro-niche also depends on the mechanical environment. *In vivo*, the biomaterial is not mechanically isolated, but connected to the mechanical environment. It is conceivable that when cells are encapsulated in the biomaterial, not only the intrinsic mechanical properties of the biomaterial determine the cell behavior, but also how much of the external mechanical stimuli are transmitted through the biomaterial. Biomaterials may not only reduce the diffusion of unwanted DAMPs or pro-inflammatory mediators but may also protect cells from pathological mechanical over-stimulation. These protected cells could then help to downregulate a too pronounced pro-inflammatory reaction.

The examples described here, are possibly just the tip of the iceberg. It seems likely that the described concepts are not only found in the lung, heart, and skin, but also in various tissues throughout the body. Since immune cells are the first cells that interact with a biomaterial and often determine the amount of a foreign body response (Anderson et al., 2008), targeting specifically immune cells via mechanical cues of the biomaterial, seems to be especially promising. If we know how to take advantage of this, it could lead to promising new therapeutic approaches. However, if the local mechanical environment is not well-controlled, it could unintentionally induce pro-inflammatory and pro-fibrotic cascades, which would jeopardize the engineered regenerative effect of the biomaterial. A better understanding of mechano-sensitive principles helps to avoid these negative effects and can even be used under certain

conditions to positively influence and enhance regeneration. It is tempting to speculate that the described immune-mechanical principles are an intrinsic and universal function of the immune system that has been overlooked for a long time. Therefore, in the development of the next generation of biomaterials, it might be worthwhile to move from an approach that focuses mainly on biological factors to a more comprehensive approach that also addresses the mechanical niche and its influence on the immune system.

AUTHOR CONTRIBUTIONS

RK compiled the literature, prepared, and wrote the manuscript. CB, SV, CT, KS-B, and GD outlined the manuscript and edited the manuscript. All authors contributed to the article and approved the submitted version.

FUNDING

This work was supported by Berlin-Brandenburg School for Regenerative Therapies (BSRT, GSC 203), German Research Foundation (DFG, FOR 2165 and SFB 1444), Einstein-Center for Regenerative Therapies (ECRT).

ACKNOWLEDGMENTS

We acknowledge support from the German Research Foundation (DFG) and the Open Access Publication Fund of Charité - Universitätsmedizin Berlin.

REFERENCES

- Aarabi, S., Bhatt, K. A., Shi, Y., Paterno, J., Chang, E. I., Loh, S. A., et al. (2007). Mechanical load initiates hypertrophic scar formation through decreased cellular apoptosis. *FASEB J.* 21, 3250–3261. doi: 10.1096/fj.07-8218com
- Adams, G. B., Chabner, K. T., Alley, I. R., Olson, D. P., Szczepiorkowski, Z. M., Poznansky, M. C., et al. (2006). Stem cell engraftment at the endosteal niche is specified by the calcium-sensing receptor. *Nature* 439, 599–603. doi: 10.1038/nature04247
- Al-Jamal, R., and Ludwig, M. S. (2001). Changes in proteoglycans and lung tissue mechanics during excessive mechanical ventilation in rats. *Am. J. Physiol. - Lung Cell. Mol. Physiol.* 281, 1078–1087. doi: 10.1152/ajplung.2001.281.5.11078
- Anderson, J. M., Rodriguez, A., and Chang, D. T. (2008). Foreign body reaction to biomaterials. *Semin. Immunol.* 20, 86–100. doi: 10.1016/j.smim.2007.11.004
- Ara, T., Itoi, M., Kawabata, K., Egawa, T., Tokoyoda, K., Sugiyama, T., et al. (2003). A Role of CXC chemokine ligand 12/stromal cell-derived factor-1/Pre-B cell growth stimulating factor and its receptor CXCR4 in fetal and adult T cell development in vivo. *J. Immunol.* 170, 4649–4655. doi: 10.4049/jimmunol.170.9.4649
- Arron, J. R., and Choi, Y. (2000). Bone versus immune system. *Nature* 408, 535–536. doi: 10.1038/35046196
- Bal, S. M., and Stadhouders, R. (2019). Tregs in fibrosis: To know your enemy, you must become your enemy. *Sci. Immunol.* 4, 6–9. doi: 10.1126/sciimmunol.aay1160
- Baratchi, S., Almaz, J. G., Darby, W., Tovar-Lopez, F. J., Mitchell, A., and McIntyre, P. (2016). Shear stress mediates exocytosis of functional TRPV4 channels in endothelial cells. *Cell. Mol. Life Sci.* 73, 649–666. doi: 10.1007/s00018-015-2018-8
- Baratchi, S., Zaldivia, M. T. K., Wallert, M., Loseff-Silver, J., Al-Aryahi, S., Zamani, J., et al. (2020). TAVI represents an anti-inflammatory therapy via reduction of shear stress induced, piezo-1-mediated monocyte activation. *Circulation* 142:120.045536. doi: 10.1161/CIRCULATIONAHA.120.045536
- Bassat, E., Mutlak, Y. E., Genzelinakh, A., Shadrin, I. Y., Baruch Umansky, K., Yifa, O., et al. (2017). The extracellular matrix protein agrin promotes heart regeneration in mice. *Nature* 547, 179–184. doi: 10.1038/nature22978
- Bastian, O., Pillay, J., Alblas, J., Leenen, L., Koenderman, L., and Blokhuis, T. (2011). Systemic inflammation and fracture healing. *J. Leukoc. Biol.* 89, 669–673. doi: 10.1189/jlb.0810446
- Bastian, O. W., Koenderman, L., Alblas, J., Leenen, L. P. H., and Blokhuis, T. J. (2016). Neutrophils contribute to fracture healing by synthesizing fibronectin+ extracellular matrix rapidly after injury. *Clin. Immunol.* 164, 78–84. doi: 10.1016/j.clim.2016.02.001
- Bezakova, G., and Ruegg, M. A. (2003). New insights into the roles of agrin. *Nat. Rev. Mol. Cell Biol.* 4, 295–308. doi: 10.1038/nrm1074
- Bielby, R., Jones, E., and McGonagle, D. (2007). The role of mesenchymal stem cells in maintenance and repair of bone. *Injury* 38. doi: 10.1016/j.injury.2007.02.007
- Bigotti, M. G., Skeffington, K. L., Jones, F. P., Caputo, M., and Brancaccio, A. (2020). Agrin-mediated cardiac regeneration: some open questions. *Front. Bioeng. Biotechnol.* 8:594. doi: 10.3389/fbioe.2020.00594
- Black, K. E., Collins, S. L., Hagan, R. S., Hamblin, M. J., Chan-Li, Y., Hallowell, R. W., et al. (2013). Hyaluronan fragments induce IFN β via a novel TLR4-TRIF-TBK1-IRF3- dependent pathway. *J. Inflamm.* 10:23. doi: 10.1186/1476-9255-10-23
- Blanch, L., Fernandez, R., Vallés, J., Sole, J., Roussos, C., and Artigas, A. (1994). Effect of two tidal volumes on oxygenation and respiratory system mechanics during the early stage of adult respiratory distress syndrome. *J. Crit. Care* 9, 151–158. doi: 10.1016/0883-9441(94)90011-6
- Blom, A. B., van Lent, P. L. E. M., Holthuysen, A. E. M., van der Kraan, P. M., Roth, J., van Rooijen, N., et al. (2004). Synovial lining macrophages mediate

- osteophyte formation during experimental osteoarthritis. *Osteoarthr. Cartil.* 12, 627–635. doi: 10.1016/j.joca.2004.03.003
- Blumenthal, D., and Burkhardt, J. K. (2020). Multiple actin networks coordinate mechanotransduction at the immunological synapse. *J. Cell Biol.* 219:e201911058. doi: 10.1083/jcb.201911058
- Bonewald, L. F., and Johnson, M. L. (2008). Osteocytes, mechanosensing and Wnt signaling. *Bone* 42, 606–615. doi: 10.1016/j.bone.2007.12.224
- Boyce, B. F., and Xing, L. (2008). Functions of RANKL/RANK/OPG in bone modeling and remodeling. *Arch. Biochem. Biophys.* 473, 139–146. doi: 10.1016/j.abb.2008.03.018
- Boyle, W. J., Simonet, W. S., and Lacey, D. L. (2003). Osteoclast differentiation and activation. *Nature* 423, 337–342. doi: 10.1038/nature01658
- Braga, T. T., Agudelo, J. S. H., and Camara, N. O. S. (2015). Macrophages during the fibrotic process: M2 as friend and foe. *Front. Immunol.* 6:602. doi: 10.3389/fimmu.2015.00602
- Brower, R. G., Matthay, M. A., Morris, A., Schoenfeld, D., Thompson, B. T., and Wheeler, A. (2000). Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. *N. Engl. J. Med.* 342, 1301–1308. doi: 10.1056/NEJM200005043421801
- Bucher, C. H., Schlundt, C., Wulsten, D., Sass, F. A., Wendler, S., Ellinghaus, A., et al. (2019). Experience in the adaptive immunity impacts bone homeostasis, remodeling, and healing. *Front. Immunol.* 10:787. doi: 10.3389/fimmu.2019.00797
- Cain, C. J., Rueda, R., McLelland, B., Collette, N. M., Loots, G. G., and Manilay, J. O. (2012). Absence of sclerostin adversely affects B-cell survival. *J. Bone Miner. Res.* 27, 1451–1461. doi: 10.1002/jbmr.1608
- Carlson, J. M. (2006). Functional limitations from pain caused by repetitive loading on the skin: a review and discussion for practitioners, with new data for limiting friction loads. *J. Prosthetics Orthot.* 18, 93–103. doi: 10.1097/00008526-200610000-00002
- Carnicer-Lombarte, A., Barone, D. G., Dimov, I. B., Hamilton, R. S., Prater, M., Zhao, X., et al. (2019). Mechanical matching of implant to host minimises foreign body reaction. *bioRxiv [Preprint]*, 1–41. doi: 10.1101/829648
- Chakraborty, S., Njah, K., Pobbati, A. V., Lim, Y. B., Raju, A., Lakshmanan, M., et al. (2017). Agrin as a mechanotransduction signal regulating YAP through the hippo pathway. *Cell Rep.* 18, 2464–2479. doi: 10.1016/j.celrep.2017.02.041
- Champagne, C. M., Takebe, J., Offenbacher, S., and Cooper, L. F. (2002). Macrophage cell lines produce osteoinductive signals that include bone morphogenetic protein-2. *Bone* 30, 26–31. doi: 10.1016/S8756-3282(01)00638-X
- Chandrasekar, B., Mummidi, S., Claycomb, W. C., Mestril, R., and Nemer, M. (2005). Interleukin-18 is a pro-hypertrophic cytokine that acts through a phosphatidylinositol 3-kinase-phosphoinositide-dependent kinase-1-Akt-GATA4 signaling pathway in cardiomyocytes. *J. Biol. Chem.* 280, 4553–4567. doi: 10.1074/jbc.M411787200
- Chen, W., and Frangiannis, N. G. (2013). Fibroblasts in post-infarction inflammation and cardiac repair. *Biochim. Biophys. Acta - Mol. Cell Res.* 1833, 945–953. doi: 10.1016/j.bbamcr.2012.08.023
- Chen, Z., Wu, C., Gu, W., Klein, T., Crawford, R., and Xiao, Y. (2014). Osteogenic differentiation of bone marrow MSCs by β -tricalcium phosphate stimulating macrophages via BMP2 signalling pathway. *Biomaterials* 35, 1507–1518. doi: 10.1016/j.biomaterials.2013.11.014
- Cho, C., Horzempa, C., Longo, C. M., Peters, D. M., Jones, D. M., and McKeown-Longo, P. J. (2020). Fibronectin in the tumor microenvironment activates a TLR4-dependent inflammatory response in lung cancer cells. *J. Cancer* 11, 3099–3105. doi: 10.7150/jca.39771
- Claes, L., Recknagel, S., and Ignatius, A. (2012). Fracture healing under healthy and inflammatory conditions. *Nat. Rev. Rheumatol.* 8, 133–143. doi: 10.1038/nrrheum.2012.1
- Clowes, J. A., Riggs, B. L., and Khosla, S. (2005). The role of the immune system in the pathophysiology of osteoporosis. *Immunol. Rev.* 208, 207–227. doi: 10.1111/j.0105-2896.2005.00334.x
- Codelia, V. A., Sun, G., and Irvine, K. D. (2014). Regulation of YAP by mechanical strain through Jnk and Hippo signaling. *Curr. Biol.* 24, 2012–2017. doi: 10.1016/j.cub.2014.07.034
- Coggins, M., and Rosenzweig, A. (2012). The fire within: cardiac inflammatory signaling in health and disease. *Circ. Res.* 110, 116–125. doi: 10.1161/CIRCRESAHA.111.243196
- Colin-York, H., Javanmardi, Y., Skamrahl, M., Kumari, S., Chang, V. T., Khuon, S., et al. (2019). Cytoskeletal control of antigen-dependent T cell activation. *Cell Rep.* 26, 3369–3379.e5. doi: 10.1016/j.celrep.2019.02.074
- Costa, A. M. A., Peyrol, S., Pôrto, L. C., Comparin, J. P., Foyatier, J. L., and Desmoulière, A. (1999). Mechanical forces induce scar remodeling: Study in non-pressure-treated versus pressure-treated hypertrophic scars. *Am. J. Pathol.* 155, 1671–1679. doi: 10.1016/S0002-9440(10)65482-X
- Cottler-Fox, M. H., Lapidot, T., Petit, I., Kollet, O., DiPersio, J. F., Link, D., et al. (2003). Stem cell mobilization. *Hematology Am. Soc. Hematol. Educ. Program* 419–437. doi: 10.1182/asheducation-2003.1.419
- Danks, L., Komatsu, N., Guerrini, M. M., Sawa, S., Armaka, M., Kollias, G., et al. (2016). RANKL expressed on synovial fibroblasts is primarily responsible for bone erosions during joint inflammation. *Ann. Rheum. Dis.* 75, 1187–1195. doi: 10.1136/annrheumdis-2014-207137
- Darby, I. A., Laverdet, B., Bonté, F., and Desmoulière, A. (2014). Fibroblasts and myofibroblasts in wound healing. *Clin. Cosmet. Investig. Dermatol.* 7, 301–311. doi: 10.2147/CCID.S50046
- Dellacherie, M. O., Seo, B. R., and Mooney, D. J. (2019). Macroscale biomaterials strategies for local immunomodulation. *Nat. Rev. Mater.* 4, 379–397. doi: 10.1038/s41578-019-0106-3
- Dennis, S. C., Berkland, C. J., Bonewald, L. F., and Detamore, M. S. (2015). Endochondral ossification for enhancing bone regeneration: converging native extracellular matrix biomaterials and developmental engineering in vivo. *Tissue Eng. - Part B Rev.* 21, 247–266. doi: 10.1089/ten.teb.2014.0419
- Dupont, S., Morsut, L., Aragona, M., Enzo, E., Giulitti, S., Cordenonsi, M., et al. (2011). Role of YAP/TAZ in mechanotransduction. *Nature* 474, 179–184. doi: 10.1038/nature10137
- Duscher, D., Maan, Z. N., Wong, V. W., Rennert, R. C., Januszyk, M., Rodrigues, M., et al. (2014). Mechanotransduction and fibrosis. *J. Biomech.* 47, 1997–2005. doi: 10.1016/j.jbiomech.2014.03.031
- Eldridge, S. E., Barawi, A., Wang, H., Roelofs, A. J., Kaneva, M., Guan, Z., et al. (2020). Agrin induces long-term osteochondral regeneration by supporting repair morphogenesis. *Sci. Transl. Med.* 12:eaa9086. doi: 10.1126/scitranslmed.aax9086
- Elosegui-Artola, A., Andreu, I., Beedle, A. E. M. M., Lezamiz, A., Uroz, M., Kosmalska, A. J., et al. (2017). Force triggers YAP nuclear entry by regulating transport across nuclear pores. *Cell* 171, 1397–1410.e14. doi: 10.1016/j.cell.2017.10.008
- Epari, D. R., Kassi, J. P., Schell, H., and Duda, G. N. (2007). Timely fracture-healing requires optimization of axial fixation stability. *J. Bone Jt. Surg. - Ser. A* 89, 1575–1585. doi: 10.2106/JBJS.F.00247
- Epari, D. R., Schell, H., Bail, H. J., and Duda, G. N. (2006). Instability prolongs the chondral phase during bone healing in sheep. *Bone* 38, 864–870. doi: 10.1016/j.bone.2005.10.023
- Fromigüé, O., Marie, P. J., and Lomri, A. (1998). Bone morphogenetic protein-2 and transforming growth factor- β 2 interact to modulate human bone marrow stromal cell proliferation and differentiation. *J. Cell. Biochem.* 68, 411–426. doi: 10.1002/(SICI)1097-4644(19980315)68:4<411::AID-JCB2>3.0.CO;2-T
- Fulzele, K., Krause, D. S., Panaroni, C., Saini, V., Barry, K. J., Liu, X., et al. (2013). Myelopoiesis is regulated by osteocytes through Gs α -dependent signaling. *Blood* 121, 930–939. doi: 10.1182/blood-2012-06-437160
- Gabbiani, G. (2003). The myofibroblast in wound healing and fibrocontractive diseases. *J. Pathol.* 200, 500–503. doi: 10.1002/path.1427
- Gao, Y., Zhou, J., Xie, Z., Wang, J., kang Ho, C., Zhang, Y., et al. (2019). Mechanical strain promotes skin fibrosis through LRG-1 induction mediated by ELK1 and ERK signalling. *Commun. Biol.* 2:359. doi: 10.1038/s42003-019-0600-6
- Geißler, S., Textor, M., Schmidt-Bleek, K., Klein, O., Thiele, M., Ellinghaus, A., et al. (2013). In serum veritas-in serum sanitas? Cell non-autonomous aging compromises differentiation and survival of mesenchymal stromal cells via the oxidative stress pathway. *Cell Death Dis.* 4, 1–10. doi: 10.1038/cddis.2013.501
- Geng, J., Yu, S., Zhao, H., Sun, X., Li, X., Wang, P., et al. (2017). The transcriptional coactivator TAZ regulates reciprocal differentiation of T H 17 cells and T reg cells. *Nat. Immunol.* 18, 800–812. doi: 10.1038/ni.3748
- Gerstenfeld, L. C., Cho, T. J., Kon, T., Aizawa, T., Tsay, A., Fitch, J., et al. (2003). Impaired fracture healing in the absence of TNF- α signaling: the role of TNF- α in endochondral cartilage resorption. *J. Bone Miner. Res.* 18, 1584–1592. doi: 10.1359/jbmr.2003.18.9.1584
- Ghiasi, M. S., Chen, J., Vaziri, A., Rodriguez, E. K., and Nazarian, A. (2017). Bone fracture healing in mechanobiological modeling: a review of

- principles and methods. *Bone Reports* 6, 87–100. doi: 10.1016/j.bonr.2017.03.002
- Gieseck, R. L., Wilson, M. S., and Wynn, T. A. (2018). Type 2 immunity in tissue repair and fibrosis. *Nat. Rev. Immunol.* 18, 62–76. doi: 10.1038/nri.2017.90
- Glass, G. E., Chan, J. K., Freidin, A., Feldmann, M., Horwood, N. J., and Nanchahal, J. (2011). TNF- α promotes fracture repair by augmenting the recruitment and differentiation of muscle-derived stromal cells. *Proc. Natl. Acad. Sci. U.S.A.* 108, 1585–1590. doi: 10.1073/pnas.1018501108
- Glynn Andrew, J., Andrew, S. M., Freemont, A. J., and Marsh, D. R. (1994). Inflammatory cells in normal human fracture healing. *Acta Orthop.* 65, 462–466. doi: 10.3109/17453679408995493
- Goldstein, B., and Sanders, J. (1998). Skin response to repetitive mechanical stress: a new experimental model in pig. *Arch. Phys. Med. Rehabil.* 79, 265–272. doi: 10.1016/S0003-9993(98)90005-3
- Gratz, I. K., Truong, H.-A., Yang, S. H.-Y., Maurano, M. M., Lee, K., Abbas, A. K., et al. (2013). Cutting edge: memory regulatory T cells require IL-7 and not IL-2 for their maintenance in peripheral tissues. *J. Immunol.* 190, 4483–4487. doi: 10.4049/jimmunol.1300212
- Greenbaum, A., Hsu, Y. M. S., Day, R. B., Schuettelpelz, L. G., Christopher, M. J., Borgerding, J. N., et al. (2013). CXCL12 in early mesenchymal progenitors is required for haematopoietic stem-cell maintenance. *Nature* 495, 227–230. doi: 10.1038/nature11926
- Greenblatt, M. B., and Shim, J. (2013). Osteoimmunology: a brief introduction. *Immune Netw.* 13:111. doi: 10.4110/in.2013.13.4.111
- Guihard, P., Boutet, M. A., Brounais-Le Royer, B., Gamblin, A. L., Amiaud, J., Renaud, A., et al. (2015). Oncostatin M, an inflammatory cytokine produced by macrophages, supports intramembranous bone healing in a mouse model of tibia injury. *Am. J. Pathol.* 185, 765–775. doi: 10.1016/j.ajpath.2014.11.008
- Haak, A. J., Kostallari, E., Sicard, D., Ligresti, G., Choi, K. M., Caporarello, N., et al. (2019). Selective YAP/TAZ inhibition in fibroblasts via dopamine receptor D1 agonism reverses fibrosis. *Sci. Transl. Med.* 11:6296. doi: 10.1126/scitranslmed.aau6296
- Hamanaka, K., Jian, M. Y., Townsley, M. I., King, J. A., Liedtke, W., Weber, D. S., et al. (2010). TRPV4 channels augment macrophage activation and ventilator-induced lung injury. *Am. J. Physiol. - Lung Cell. Mol. Physiol.* 299, 353–362. doi: 10.1152/ajplung.00315.2009
- Hardy, R., and Cooper, M. S. (2009). Bone loss in inflammatory disorders. *J. Endocrinol.* 201, 309–320. doi: 10.1677/JOE-08-0568
- Hausser, H.-J., Ruegg, M. A., Brenner, R. E., and Ksiazek, I. (2007). Agrin is highly expressed by chondrocytes and is required for normal growth. *Histochem Cell Biol* 127, 363–374. doi: 10.1007/s00418-006-0258-2
- Heise, R. L., Stober, V., Cheluvharaju, C., Hollingsworth, J. W., and Garantziotis, S. (2011). Mechanical stretch induces epithelial-mesenchymal transition in alveolar epithelia via hyaluronan activation of innate immunity. *J. Biol. Chem.* 286, 17435–17444. doi: 10.1074/jbc.M110.137273
- Hinz, B. (2007). Formation and function of the myofibroblast during tissue repair. *J. Invest. Dermatol.* 127, 526–537. doi: 10.1038/sj.jid.5700613
- Hinz, B., Phan, S. H., Thannickal, V. J., Prunotto, M., Desmouliere, A., Varga, J., et al. (2012). Recent developments in myofibroblast biology: paradigms for connective tissue remodeling. *Am. J. Pathol.* 180, 1340–1355. doi: 10.1016/j.ajpath.2012.02.004
- Hogan, B. L. M., Barkauskas, C. E., Chapman, H. A., Epstein, J. A., Jain, R., Hsia, C. C. W., et al. (2014). Repair and regeneration of the respiratory system: complexity, plasticity, and mechanisms of lung stem cell function. *Cell Stem Cell* 15, 123–138. doi: 10.1016/j.stem.2014.07.012
- Hou, Y., Ryu, C. H., Jun, J. A., Kim, S. M., Jeong, C. H., and Jeun, S. S. (2014). IL-8 enhances the angiogenic potential of human bone marrow mesenchymal stem cells by increasing vascular endothelial growth factor. *Cell Biol. Int.* 38, 1050–1059. doi: 10.1002/cbin.10294
- Humphrey, J. D., Dufresne, E. R., and Schwartz, M. A. (2014). Mechanotransduction and extracellular matrix homeostasis. *Nat. Rev. Mol. Cell Biol.* 15, 802–812. doi: 10.1038/nrm3896
- Hurst, S. M., Wilkinson, T. S., McLoughlin, R. M., Jones, S., Horiuchi, S., Yamamoto, N., et al. (2001). IL-6 and its soluble receptor orchestrate a temporal switch in the pattern of leukocyte recruitment seen during acute inflammation. *Immunity* 14, 705–714. doi: 10.1016/S1074-7613(01)00151-0
- Itano, N., Sawai, T., Yoshida, M., Lenas, P., Yamada, Y., Imagawa, M., et al. (1999). Three isoforms of mammalian hyaluronan synthases have distinct enzymatic properties. *J. Biol. Chem.* 274, 25085–25092. doi: 10.1074/jbc.274.35.25085
- Jiang, D., Liang, J., Fan, J., Yu, S., Chen, S., Luo, Y., et al. (2005). Regulation of lung injury and repair by Toll-like receptors and hyaluronan. *Nat. Med.* 11, 1173–1179. doi: 10.1038/nm1315
- Jo, A. O., Ryskamp, D. A., Phuong, T. T. T., Verkman, A. S., Yarishkin, O., Macaulay, N., et al. (2015). TRPV4 and AQP4 channels synergistically regulate cell volume and calcium homeostasis in retinal müller glia. *J. Neurosci.* 35, 13525–13537. doi: 10.1523/JNEUROSCI.1987-15.2015
- Julier, Z., Park, A. J., Briquez, P. S., and Martino, M. M. (2017). Promoting tissue regeneration by modulating the immune system. *Acta Biomater.* 53, 13–28. doi: 10.1016/j.actbio.2017.01.056
- Khosla, S. (2001). Minireview: The OPG/RANKL/RANK System. 142, 5050–5055
- Kikuchi, K., and Poss, K. D. (2012). Cardiac regenerative capacity and mechanisms. *Annu. Rev. Cell Dev. Biol.* 28, 719–741. doi: 10.1146/annurev-cellbio-101011-155739
- Kim, S. H., Kim, S., Choi, H. I., Choi, Y. J., Lee, Y. S., Sohn, K. C., et al. (2010). Callus formation is associated with hyperproliferation and incomplete differentiation of keratinocytes, and increased expression of adhesion molecules. *Br. J. Dermatol.* 163, 495–501. doi: 10.1111/j.1365-2133.2010.09842.x
- Koch, A. E., Polverini, P. J., Kunkel, S. L., Harlow, L. A., DiPietro, L. A., Elner, V. M., et al. (1992). Interleukin-8 as a macrophage-derived mediator of angiogenesis. *Science (80-)* 258, 1798–1801. doi: 10.1126/science.1281554
- Kode, A., Manavalan, J. S., Mosialou, I., Bhagat, G., Rathinam, C. V., Luo, N., et al. (2014). Leukaemogenesis induced by an activating β -catenin mutation in osteoblasts. *Nature* 506, 240–244. doi: 10.1038/nature12883
- Kolar, P., Gaber, T., Perka, C., Duda, G. N., and Buttgerit, F. (2011). “Human early fracture hematoma is characterized by inflammation and hypoxia. In *Clinical Orthopaedics and Related Research (Springer New York LLC)*, 3118–3126. doi: 10.1007/s11999-011-1865-3
- Kolar, P., Schmidt-Bleek, K., Schell, H., Gaber, T., Toben, D., Schmidmaier, G., et al. (2010). The early fracture hematoma and its potential role in fracture healing. *Tissue Eng. - Part B Rev.* 16, 427–434. doi: 10.1089/ten.teb.2009.0687
- Kollet, O., Dar, A., Shvitiel, S., Kalinkovich, A., Lapid, K., Sztainberg, Y., et al. (2006). Osteoclasts degrade endosteal components and promote mobilization of hematopoietic progenitor cells. *Nat. Med.* 12, 657–664. doi: 10.1038/nm1417
- Kollet, O., Shvitiel, S., Chen, Y. Q., Suriawinata, J., Thung, S. N., Dabeva, M. D., et al. (2003). HGF, SDF-1, and MMP-9 are involved in stress-induced human CD34+ stem cell recruitment to the liver. *J. Clin. Invest.* 112, 160–169. doi: 10.1172/JCI17902
- Kon, T., Cho, T. J., Aizawa, T., Yamazaki, M., Nooh, N., Graves, D., et al. (2001). Expression of osteoprotegerin, receptor activator of NF- κ B ligand (osteoprotegerin ligand) and related proinflammatory cytokines during fracture healing. *J. Bone Miner. Res.* 16, 1004–1014. doi: 10.1359/jbmr.2001.16.6.1004
- Kong, Y. Y., Yoshida, H., Sarosi, I., Tan, H. L., Timms, E., Capparelli, C., et al. (1999). OPGL is a key regulator of osteoclastogenesis, lymphocyte development and lymph-node organogenesis. *Nature* 397, 315–323. doi: 10.1038/16852
- Könnecke, I., Serra, A., El Khassawna, T., Schlundt, C., Schell, H., Hauser, A., et al. (2014). T and B cells participate in bone repair by infiltrating the fracture callus in a two-wave fashion. *Bone* 64, 155–165. doi: 10.1016/j.bone.2014.03.052
- Kovtun, A., Bergdolt, S., Wiegner, R., Radermacher, P., Huber-Lang, M., and Ignatius, A. (2016). The crucial role of neutrophil granulocytes in bone fracture healing. *Eur. Cells Mater.* 32, 152–162. doi: 10.22203/eCM.v032a10
- Kuipers, M. T., Aslami, H., Janczy, J. R., Van Der Sluijs, K. F., Vlaar, A. P. J., Wolthuis, E. K., et al. (2012). Ventilator-induced lung injury is mediated by the NLRP3 inflammasome. *Anesthesiology* 116, 1104–1115. doi: 10.1097/ALN.0b013e3182518bc0
- Kuipers, M. T., van der Poll, T., Schultz, M. J., and Wieland, C. W. (2011). Bench-to-bedside review: damage-associated molecular patterns in the onset of ventilator-induced lung injury. *Crit. Care* 15, 1–11. doi: 10.1186/cc10437
- Kwon, S. H., Padmanabhan, J., and Gurtner, G. C. (2018). “Mechanobiology of skin diseases and wound healing,” in *Mechanobiology in Health and Disease* (London: Elsevier), 415–448. doi: 10.1016/B978-0-12-812952-4.00014-3
- LaCanna, R., Liccardo, D., Zhang, P., Tragesser, L., Wang, Y., Cao, T., et al. (2019). Yap/Taz regulate alveolar regeneration and resolution of lung inflammation. *J. Clin. Invest.* 129, 2107–2122. doi: 10.1172/JCI125014
- Langer, K. (1978a). On the anatomy and physiology of the skin. I. The cleavability of the cutis. (Translated from Langer, K. (1861). *Zur Anatomie und Physiologie*

- der Haut. I. Über die Spaltbarkeit der Cutis. Sitzungsbericht der Mathematisch-naturwissenschaftlichen Classe der. *Br. J. Plast. Surg.* 31, 3–8
- Langer, K. (1978b). On the anatomy and physiology of the skin. III. The elasticity of the cutis By Professor K. Langer Presented at the meeting of 27th November 1861. *Br. J. Plast. Surg.* 31, 185–199
- Langer, K. (1978c). On the anatomy and physiology of the skin ii skin tension by Professor K. Langer, presented at the meeting of 27th November 1861. *Br. J. Plast. Surg.* 31, 93–106.
- Leach, J. P., Heallen, T., Zhang, M., Rahmani, M., Morikawa, Y., Hill, M. C., et al. (2017). Hippo pathway deficiency reverses systolic heart failure after infarction. *Nature* 550, 260–264. doi: 10.1038/nature24045
- Leblond, A.-L., Klinkert, K., Martin, K., Turner, E. C., Kumar, A. H., Browne, T., et al. (2015). Systemic and cardiac depletion of M2 macrophage through CSF-1R signaling inhibition alters cardiac function post myocardial infarction. *PLoS ONE* 10:e0137515. doi: 10.1371/journal.pone.0137515
- Lee, H., pyo, Stowers, R., and Chaudhuri, O. (2019). Volume expansion and TRPV4 activation regulate stem cell fate in three-dimensional microenvironments. *Nat. Commun.* 10:529. doi: 10.1038/s41467-019-08465-x
- Lévesque, J. P., Hendy, J., Takamatsu, Y., Simmons, P. J., and Bendall, L. J. (2003). Disruption of the CXCR4/CXCL12 chemotactic interaction during hematopoietic stem cell mobilization induced by gcsf or cyclophosphamide. *J. Clin. Invest.* 111, 187–196. doi: 10.1172/JCI15994
- Li, M., Fang, X. Z., Zheng, Y. F., Xie, Y., Bin, M. A., and X. D., Liu, X. T., et al. (2019). Transient receptor potential vanilloid 4 is a critical mediator in LPS mediated inflammation by mediating calcineurin/NFATc3 signaling. *Biochem. Biophys. Res. Commun.* 513, 1005–1012. doi: 10.1016/j.bbrc.2019.04.020
- Li, X., Ominsky, M. S., Niu, Q. T., Sun, N., Daugherty, B., D'Agostin, D., et al. (2008). Targeted deletion of the sclerostin gene in mice results in increased bone formation and bone strength. *J. Bone Miner. Res.* 23, 860–869. doi: 10.1359/jbmr.080216
- Lienau, J., Schell, H., Duda, G. N., Seebeck, P., Muchow, S., and Bail, H. J. (2005). Initial vascularization and tissue differentiation are influenced by fixation stability. *J. Orthop. Res.* 23, 639–645. doi: 10.1016/j.orthres.2004.09.006
- Lin, D., Chai, Y., Ma, Y., Duan, B., Yuan, Y., and Liu, C. (2019). Rapid initiation of guided bone regeneration driven by spatiotemporal delivery of IL-8 and BMP-2 from hierarchical MBG-based scaffold. *Biomaterials* 196, 122–137. doi: 10.1016/j.biomaterials.2017.11.011
- Lin, M., Jackson, P., Tester, A. M., Diaconu, E., Overall, C. M., Blalock, J. E., et al. (2008). Matrix metalloproteinase-8 facilitates neutrophil migration through the corneal stromal matrix by collagen degradation and production of the chemotactic peptide pro-gly-pro. *Am. J. Pathol.* 173, 144–153. doi: 10.2353/ajpath.2008.080081
- Lin, Z., Von Gise, A., Zhou, P., Gu, F., Ma, Q., Jiang, J., et al. (2014). Cardiac-specific YAP activation improves cardiac function and survival in an experimental murine MI model. *Circ. Res.* 115, 354–363. doi: 10.1161/CIRCRESAHA.115.303632
- Lindner, D., Zietsch, C., Tank, J., Sossalla, S., Fluschnik, N., Hinrichs, S., et al. (2014). Cardiac fibroblasts support cardiac inflammation in heart failure. *Basic Res. Cardiol.* 109, 1–16. doi: 10.1007/s00395-014-0428-7
- Liu, F., Lagares, D., Choi, K. M., Stopfer, L., Marinković, A., Vrbanac, V., et al. (2015). Mechanosignaling through YAP and TAZ drives fibroblast activation and fibrosis. *Am. J. Physiol. - Lung Cell. Mol. Physiol.* 308, L344–L357. doi: 10.1152/ajplung.00300.2014
- Liu, Y., Wang, L., Kikuri, T., Akiyama, K., Chen, C., Xu, X., et al. (2011). Mesenchymal stem cell-based tissue regeneration is governed by recipient T lymphocytes via IFN- γ and TNF- α . *Nat. Med.* 17, 1594–1601. doi: 10.1038/nm.2542
- Liu, Z., Wu, H., Jiang, K., Wang, Y., Zhang, W., Chu, Q., et al. (2016). MAPK-mediated YAP activation controls mechanical-tension-induced pulmonary alveolar regeneration. *Cell Rep.* 16, 1810–1819. doi: 10.1016/j.celrep.2016.07.020
- Loi, F., Córdova, L. A., Pajarinen, J., Lin, T., hua, Yao, Z., and Goodman, S. B. (2016). Inflammation, fracture and bone repair. *Bone* 86, 119–130. doi: 10.1016/j.bone.2016.02.020
- Maruyama, M., Rhee, C., Utsunomiya, T., Zhang, N., Ueno, M., Yao, Z., et al. (2020). Modulation of the inflammatory response and bone healing. *Front. Endocrinol. (Lausanne)*. 11:386. doi: 10.3389/fendo.2020.00386
- Mascarenhas, M. M., Day, R. M., Ochoa, C. D., Choi, W., Il, Y.u., and L., Ouyang, B., et al. (2004). Low molecular weight hyaluronan from stretched lung enhances interleukin-8 expression. *Am. J. Respir. Cell Mol. Biol.* 30, 51–60. doi: 10.1165/rcmb.2002-0167OC
- Mazzon, C., Anselmo, A., Soldani, C., Cibella, J., Ploia, C., Moalli, F., et al. (2012). Agrin is required for survival and function of monocytic cells. *Blood* 119, 5502–5511. doi: 10.1182/blood-2011-09-382812
- McGeachy, M. J. (2017). A Hippo in the Fox(p3) house. *Nat. Immunol.* 18, 709–711. doi: 10.1038/ni.3769
- Medzhitov, R. (2008). Origin and physiological roles of inflammation. *Nature* 454, 428–435. doi: 10.1038/nature07201
- Midwood, K. S., and Piccinini, A. M. (2010). DAMPening inflammation by modulating TLR signalling. *Mediators Inflamm.* 2010:672395. doi: 10.1155/2010/672395
- Miossec, P., and Kolls, J. K. (2012). Targeting IL-17 and T H 17 cells in chronic inflammation. *Nat. Rev. Drug Discov.* 11, 763–776. doi: 10.1038/nrd3794
- Morimoto, H., Takahashi, M., Izawa, A., Ise, H., Hongo, M., Kolattukudy, P. E., et al. (2006). Cardiac overexpression of monocyte chemoattractant protein-1 in transgenic mice prevents cardiac dysfunction and remodeling after myocardial infarction. *Circ. Res.* 99, 891–899. doi: 10.1161/01.RES.0000246113.82111.2d
- Morrison, S. J., and Scadden, D. T. (2014). The bone marrow niche for haematopoietic stem cells. *Nature* 505, 327–334. doi: 10.1038/nature12984
- Mountziaris, P. M., and Mikos, A. G. (2008). Modulation of the inflammatory response for enhanced bone tissue regeneration. *Tissue Eng. - Part B Rev.* 14, 179–186. doi: 10.1089/ten.teb.2008.0038
- Moya, I. M., and Halder, G. (2019). Hippo–YAP/TAZ signalling in organ regeneration and regenerative medicine. *Nat. Rev. Mol. Cell Biol.* 20, 211–226. doi: 10.1038/s41580-018-0086-y
- Nakasaki, M., Hwang, Y., Xie, Y., Kataria, S., Gund, R., Hajam, E. Y., et al. (2015). The matrix protein Fibulin-5 is at the interface of tissue stiffness and inflammation in fibrosis. *Nat. Commun.* 6, 1–11. doi: 10.1038/ncomms9574
- Ogawa, R., and Hsu, C.-K. K. (2013). Mechanobiological dysregulation of the epidermis and dermis in skin disorders and in degeneration. *J. Cell. Mol. Med.* 17, 817–822. doi: 10.1111/jcmm.12060
- Okamoto, K., Nakashima, T., Shinohara, M., Negishi-Koga, T., Komatsu, N., Terashima, A., et al. (2017). Osteoimmunology: the conceptual framework unifying the immune and skeletal systems. *Physiol. Rev.* 97, 1295–1349. doi: 10.1152/physrev.00036.2016
- Okamoto, T., Takagi, Y., Kawamoto, E., Park, E. J., Usuda, H., Wada, K., et al. (2018). Reduced substrate stiffness promotes M2-like macrophage activation and enhances peroxisome proliferator-activated receptor γ expression. *Exp. Cell Res.* 367, 264–273. doi: 10.1016/j.yexcr.2018.04.005
- Omatsu, Y., Sugiyama, T., Kohara, H., Kondoh, G., Fujii, N., Kohno, K., et al. (2010). The essential functions of adipo-osteogenic progenitors as the hematopoietic stem and progenitor cell niche. *Immunity* 33, 387–399. doi: 10.1016/j.immuni.2010.08.017
- Orapiriyakul, W., Tsimbouri, M. P., Childs, P., Campsie, P., Wells, J., Fernandez-Yague, M. A., et al. (2020). Nanovibrational stimulation of mesenchymal stem cells induces therapeutic reactive oxygen species and inflammation for three-dimensional bone tissue engineering. *ACS Nano* 14, 10027–10044. doi: 10.1021/acsnano.0c03130
- Ortiz Franyuti, D., Mitsi, M., and Vogel, V. (2018). Mechanical Stretching of Fibronectin Fibers Upregulates Binding of Interleukin-7. *Nano Lett.* 18, 15–25. doi: 10.1021/acs.nanolett.7b01617
- Panciera, T., Azzolin, L., Cordenonsi, M., and Piccolo, S. (2017). Mechanobiology of YAP and TAZ in physiology and disease. *Nat. Rev. Mol. Cell Biol.* 18, 758–770. doi: 10.1038/nrm.2017.87
- Papritz, K., Savvatis, K., Koschel, A., Miteva, K., Tschöpe, C., and Van Linthout, S. (2018). Cardiac (myo)fibroblasts modulate the migration of monocyte subsets. *Sci. Rep.* 8, 1–11. doi: 10.1038/s41598-018-23881-7
- Perl, M., Gebhard, F., Knöferl, M. W., Bachem, M., Gross, H. J., Kinzl, L., et al. (2003). The pattern of preformed cytokines in tissues frequently affected by blunt trauma. *Shock* 19, 299–304. doi: 10.1097/00024382-200304000-00001
- Petersen, A., Princ, A., Korus, G., Ellinghaus, A., Leemhuis, H., Herrera, A., et al. (2018). A biomaterial with a channel-like pore architecture induces endochondral healing of bone defects. *Nat. Commun.* 9:4430. doi: 10.1038/s41467-018-06504-7

- Pocaterra, A., Romani, P., and Dupont, S. (2020). YAP/TAZ functions and their regulation at a glance. *J. Cell Sci.* 133, 1–9. doi: 10.1242/jcs.230425
- Prideaux, M., Findlay, D. M., and Atkins, G. J. (2016). Osteocytes: the master cells in bone remodelling. *Curr. Opin. Pharmacol.* 28, 24–30. doi: 10.1016/j.coph.2016.02.003
- Qin, L., Liu, W., Cao, H., and Xiao, G. (2020). Molecular mechanosensors in osteocytes. *Bone Res.* 8, 1–24. doi: 10.1038/s41413-020-0099-y
- Qin, W., Xia, W., Fisher, G. J., Voorhees, J. J., and Quan, T. (2018). YAP/TAZ regulates TGF- β /Smad3 signaling by induction of Smad7 via AP-1 in human skin dermal fibroblasts. *Cell Commun. Signal.* 16:18. doi: 10.1186/s12964-018-0232-3
- Ramjee, V., Li, D., Manderfield, L. J., Liu, F., Engleka, K. A., Aghajanian, H., et al. (2017). Epicardial YAP/TAZ orchestrate an immunosuppressive response following myocardial infarction. *J. Clin. Invest.* 127, 899–911. doi: 10.1172/JCI88759
- Rapp, A. E., Bindl, R., Recknagel, S., Erbacher, A., Müller, I., Schrezenmeier, H., et al. (2016). Fracture Healing Is Delayed in Immunodeficient NOD/scid- γ 2L2Ry^c null Mice. *PLoS ONE* 11:e0147465. doi: 10.1371/journal.pone.0147465
- Reddy, V. S., Prabhu, S. D., Mummidhi, S., Valente, A. J., Venkatesan, B., Shanmugam, P., et al. (2010). Interleukin-18 induces EMMPRIN expression in primary cardiomyocytes via JNK/Sp1 signaling and MMP-9 in part via EMMPRIN and through AP-1 and NF- κ B activation. *Am. J. Physiol. - Hear. Circ. Physiol.* 299, 1242–1254. doi: 10.1152/ajpheart.00451.2010
- Reinke, S., Geissler, S., Taylor, W. R., Schmidt-Bleek, K., Juelke, K., Schwachmeyer, V., et al. (2013). Terminally differentiated CD8 + T cells negatively affect bone regeneration in humans. *Sci. Transl. Med.* 5:177ra36. doi: 10.1126/scitranslmed.3004754
- Renò, F., Sabbatini, M., Lombardi, F., Stella, M., Pezzuto, C., Magliacani, G., et al. (2003). In vitro mechanical compression induces apoptosis and regulates cytokines release in hypertrophic scars. *Wound Repair Regen.* 11, 331–336. doi: 10.1046/j.1524-475X.2003.11504.x
- Rho, J. Y., Kuhn-Spearing, L., and Zioupos, P. (1998). Mechanical properties and the hierarchical structure of bone. *Med. Eng. Phys.* 20, 92–102. doi: 10.1016/S1350-4533(98)00007-1
- Rosenbaum, T., Benítez-Angeles, M., Sánchez-Hernández, R., Morales-Lázaro, S. L., Hiriart, M., Morales-Buenrostro, L. E., et al. (2020). Trpv4: A physio and pathophysiological significant ion channel. *Int. J. Mol. Sci.* 21. doi: 10.3390/ijms21113837
- Ruegg, M. A., and Bixby, J. L. (1998). Agrin orchestrates synaptic differentiation at the vertebrate neuromuscular junction. *Trends Neurosci.* 21, 22–27. doi: 10.1016/S0166-2236(97)01154-5
- Russo, I., Cavallera, M., Huang, S., Su, Y., Hanna, A., Chen, B., et al. (2019). Protective effects of activated myofibroblasts in the pressure-overloaded myocardium are mediated through smad-dependent activation of a matrix-preserving program. *Circ. Res.* 124, 1214–1227. doi: 10.1161/CIRCRESAHA.118.314438
- Sanchez, C., Gabay, O., Salvat, C., Henrotin, Y. E., and Berenbaum, F. (2009). Mechanical loading highly increases IL-6 production and decreases OPG expression by osteoblasts. *Osteoarthr. Cartil.* 17, 473–481. doi: 10.1016/j.joca.2008.09.007
- Sanchez, C., Pesesse, L., Gabay, O., Delcour, J. P., Msika, P., Baudouin, C., et al. (2012). Regulation of subchondral bone osteoblast metabolism by cyclic compression. *Arthritis Rheum.* 64, 1193–1203. doi: 10.1002/art.33445
- Sato, K., Suematsu, A., Okamoto, K., Yamaguchi, A., Morishita, Y., Kadono, Y., et al. (2006). Th17 functions as an osteoclastogenic helper T cell subset that links T cell activation and bone destruction. *J. Exp. Med.* 203, 2673–2682. doi: 10.1084/jem.20061775
- Sato, M., Asada, N., Kawano, Y., Wakahashi, K., Minagawa, K., Kawano, H., et al. (2013). Osteocytes regulate primary lymphoid organs and fat metabolism. *Cell Metab.* 18, 749–758. doi: 10.1016/j.cmet.2013.09.014
- Sato, T., Verma, S., Andrade, C. D. C., Omeara, M., Campbell, N., Wang, J. S., et al. (2020). A FAK/HDAC5 signaling axis controls osteocyte mechanotransduction. *Nat. Commun.* 11, 1–18. doi: 10.1038/s41467-020-17099-3
- Schaefer, L., Babelova, A., Kiss, E., Hausser, H. J., Baliova, M., Krzyzankova, M., et al. (2005). The matrix component biglycan is proinflammatory and signals through Toll-like receptors 4 and 2 in macrophages. *J. Clin. Invest.* 115, 2223–2233. doi: 10.1172/JCI23755
- Schell, H., Epari, D. R., Kassi, J. P., Bragulla, H., Bail, H. J., and Duda, G. N. (2005). The course of bone healing is influenced by the initial shear fixation stability. *J. Orthop. Res.* 23, 1022–1028. doi: 10.1016/j.orthres.2005.03.005
- Scheraga, R. G., Abraham, S., Grove, L. M., Southern, B. D., Crish, J. F., Perelas, A., et al. (2020). TRPV4 Protects the Lung from Bacterial Pneumonia via MAPK Molecular Pathway Switching. *J. Immunol.* 204, 1310–1321. doi: 10.4049/jimmunol.1901033
- Schlundt, C., El Khassawna, T., Serra, A., Dienelt, A., Wendler, S., Schell, H., et al. (2018). Macrophages in bone fracture healing: their essential role in endochondral ossification. *Bone* 106, 78–89. doi: 10.1016/j.bone.2015.10.019
- Schlundt, C., Reinke, S., Geissler, S., Bucher, C. H., Giannini, C., Märdian, S., et al. (2019). Individual effector/regulator t cell ratios impact bone regeneration. *Front. Immunol.* 10:1954. doi: 10.3389/fimmu.2019.01954
- Schmidt-Bleek, K., Kwee, B. J., Mooney, D. J., and Duda, G. N. (2015). Boon and bane of inflammation in bone tissue regeneration and its link with angiogenesis. *Tissue Eng. - Part B Rev.* 21, 354–364. doi: 10.1089/ten.teb.2014.0677
- Schmidt-Bleek, K., Schell, H., Lienau, J., Schulz, N., Hoff, P., Pfaff, M., et al. (2014). Initial immune reaction and angiogenesis in bone healing. *J. Tissue Eng. Regen. Med.* 8, 120–130. doi: 10.1002/term.1505
- Schmidt-Bleek, K., Schell, H., Schulz, N., Hoff, P., Perka, C., Buttgerit, F., et al. (2012). Inflammatory phase of bone healing initiates the regenerative healing cascade. *Cell Tissue Res.* 347, 567–573. doi: 10.1007/s00441-011-1205-7
- Serhan, C. N., and Savill, J. (2005). Resolution of inflammation: the beginning programs the end. *Nat. Immunol.* 6, 1191–1197. doi: 10.1038/ni1276
- Sharma, S., Goswami, R., Merth, M., Cohen, J., Lei, K. Y., Zhang, D. X., et al. (2017). TRPV4 ion channel is a novel regulator of dermal myofibroblast differentiation. *Am. J. Physiol. - Cell Physiol.* 312, C562–C572. doi: 10.1152/ajpcell.00187.2016
- Sharma, S., Goswami, R., Zhang, D. X., and Rahaman, S. O. (2019). TRPV4 regulates matrix stiffness and TGF β 1-induced epithelial-mesenchymal transition. *J. Cell. Mol. Med.* 23, 761–774. doi: 10.1111/jcmm.13972
- Shin, S. H., Lee, E. J., Hyun, S., Chun, J., Kim, Y., and Kang, S. S. (2012). Phosphorylation on the ser 824 residue of trpv4 prefers to bind with f-actin than with microtubules to expand the cell surface area. *Cell. Signal.* 24, 641–651. doi: 10.1016/j.cellsig.2011.11.002
- Shyu, K. G., Wang, B. W., Lin, C. M., and Chang, H. (2010). Cyclic stretch enhances the expression of Toll-like Receptor 4 gene in cultured cardiomyocytes via p38 MAP kinase and NF- κ B pathway. *J. Biomed. Sci.* 17. doi: 10.1186/1423-0127-17-15
- Silver, F. H., Siperko, L. M., and Seehra, G. P. (2003). Mechanobiology of force transduction in dermal tissue. *Ski. Res. Technol.* 9, 3–23. doi: 10.1034/j.1600-0846.2003.00358.x
- Solis, A. G., Bielecki, P., Steach, H. R., Sharma, L., Harman, C. C. D. D., Yun, S., et al. (2019). Mechanosensation of cyclical force by PIEZO1 is essential for innate immunity. *Nature* 573, 69–74. doi: 10.1038/s41586-019-1485-8
- Spiller, K. L., Nassiri, S., Witherel, C. E., Anfang, R. R., Ng, J., Nakazawa, K. R., et al. (2015). Sequential delivery of immunomodulatory cytokines to facilitate the M1-to-M2 transition of macrophages and enhance vascularization of bone scaffolds. *Biomaterials* 37, 194–207. doi: 10.1016/j.biomaterials.2014.10.017
- Spillmann, F., Van Linthout, S., Schmidt, G., Klein, O., Hamdani, N., Mairinger, T., et al. (2019). Mode-of-action of the PROPELLA concept in fulminant myocarditis. *Eur. Heart J.* 40, 2164–2169. doi: 10.1093/eurheartj/ehz124
- Sridharan, R., Cavanagh, B., Cameron, A. R., Kelly, D. J., and O'Brien, F. J. (2019). Material stiffness influences the polarization state, function and migration mode of macrophages. *Acta Biomater.* 89, 47–59. doi: 10.1016/j.actbio.2019.02.048
- Steffens, S., Van Linthout, S., Sluijter, J. P. G., Tocchetti, C. G., Thum, T., and Madonna, R. (2020). Stimulating pro-reparative immune responses to prevent adverse cardiac remodelling: consensus document from the joint 2019 meeting of the ESC Working Groups of cellular biology of the heart and myocardial function. *Cardiovasc. Res.* 116, 1850–1862. doi: 10.1093/cvr/cvaa137
- Strotmann, R., Harteneck, C., Nunnenmacher, K., Schultz, G., and Plant, T. D. (2000). OTRPC4, a nonselective cation channel that confers sensitivity to extracellular osmolarity. *Nat. Cell Biol.* 2, 695–702. doi: 10.1038/35036318
- Suárez-Álvarez, B., López-Vázquez, A., and López-Larrea, C. (2012). Mobilization and homing of hematopoietic stem cells. *Adv. Exp. Med. Biol.* 741, 152–170. doi: 10.1007/978-1-4614-2098-9_11

- Takayanagi, H. (2007). Osteoimmunology: Shared mechanisms and crosstalk between the immune and bone systems. *Nat. Rev. Immunol.* 7, 292–304. doi: 10.1038/nri2062
- Takayanagi, H., Ogasawara, K., Hida, S., Chiba, T., Murata, S., Sato, K., et al. (2000). T-cell-mediated regulation of osteoclastogenesis by signalling cross-talk between RANKL and IFN- γ . *Nature* 408, 600–605. doi: 10.1038/35046102
- Tang, L., Lin, Z., and Ming Li, Y. (2006). Effects of different magnitudes of mechanical strain on Osteoblasts in vitro. *Biochem. Biophys. Res. Commun.* 344, 122–128. doi: 10.1016/j.bbrc.2006.03.123
- Terashima, A., Okamoto, K., Nakashima, T., Akira, S., Ikuta, K., and Takayanagi, H. (2016). Sepsis-induced osteoblast ablation causes immunodeficiency. *Immunity* 44, 1434–1443. doi: 10.1016/j.immuni.2016.05.012
- Theill, L. E., Boyle, W. J., and Penninger, J. M. (2002). RANK-L and RANK: T cells, bone loss, and mammalian evolution. *Annu. Rev. Immunol.* 20, 795–823. doi: 10.1146/annurev.immunol.20.100301.064753
- Thomas, S. E., Dykes, P. J., and Marks, R. (1985). Plantar hyperkeratosis: a study of callosities and normal plantar skin. *J. Invest. Dermatol.* 85, 394–397. doi: 10.1111/1523-1747.ep12277052
- To, W. S., and Midwood, K. S. (2011). Plasma and cellular fibronectin: distinct and independent functions during tissue repair. *Fibrogenes. Tissue Repair* 4, 1–17. doi: 10.1186/1755-1536-4-21
- Tolar, P., and Wack, A. (2019). Monocytes work harder under pressure. *Nat. Immunol.* 20, 1422–1424. doi: 10.1038/s41590-019-0523-x
- Tomasek, J. J., Gabbiani, G., Hinz, B., Chaponnier, C., and Brown, R. A. (2002). Myofibroblasts and mechano: regulation of connective tissue remodelling. *Nat. Rev. Mol. Cell Biol.* 3, 349–363. doi: 10.1038/nrm809
- Totaro, A., Panciera, T., and Piccolo, S. (2018). YAP/TAZ upstream signals and downstream responses. *Nat. Cell Biol.* 20, 888–899. doi: 10.1038/s41556-018-0142-z
- Tremblay, L. N., and Slutsky, A. S. (2006). Ventilator-induced lung injury: from the bench to the bedside. *Intensive Care Med.* 32, 24–33. doi: 10.1007/s00134-005-2817-8
- Tschöpe, C., Van Linthout, S., Klein, O., Mairinger, T., Krackhardt, F., Potapov, E. V., et al. (2019). Mechanical unloading by fulminant myocarditis: LV-IMPELLA, ECMELLA, BI-PELLA, and PROPELLA concepts. *J. Cardiovasc. Transl. Res.* 12, 116–123. doi: 10.1007/s12265-018-9820-2
- Tsimbouri, P. M., Childs, P. G., Pemberton, G. D., Yang, J., Jayawarna, V., Orapiriyakul, W., et al. (2017). Stimulation of 3D osteogenesis by mesenchymal stem cells using a nanovibrational bioreactor. *Nat. Biomed. Eng.* 1, 758–770. doi: 10.1038/s41551-017-0127-4
- Tsukasaki, M., and Takayanagi, H. (2019). Osteoimmunology: evolving concepts in bone-immune interactions in health and disease. *Nat. Rev. Immunol.* 19, 626–642. doi: 10.1038/s41577-019-0178-8
- Van Amerongen, M. J., Harmsen, M. C., Van Rooijen, N., Petersen, A. H., and Van Luyn, M. J. A. (2007). Macrophage depletion impairs wound healing and increases left ventricular remodeling after myocardial injury in mice. *Am. J. Pathol.* 170, 818–829. doi: 10.2353/ajpath.2007.060547
- Van Linthout, S., Miteva, K., and Tschöpe, C. (2014). Crosstalk between fibroblasts and inflammatory cells. *Cardiovasc. Res.* 102, 258–269. doi: 10.1093/cvr/cvu062
- Vasconcelos, D. M., Gonçalves, R. M., Almeida, C. R., Pereira, I. O., Oliveira, M. I., Neves, N., et al. (2016). Fibrinogen scaffolds with immunomodulatory properties promote in vivo bone regeneration. *Biomaterials* 111, 163–178. doi: 10.1016/j.biomaterials.2016.10.004
- Vriens, J., Watanabe, H., Janssens, A., Droogmans, G., Voets, T., and Nilius, B. (2004). Cell swelling, heat, and chemical agonists use distinct pathways for the activation of the cation channel TRPV4. *Proc. Natl. Acad. Sci. U.S.A.* 101, 396–401. doi: 10.1073/pnas.0303329101
- Wang, J., Liu, S., Heallen, T., and Martin, J. F. (2018). The Hippo pathway in the heart: pivotal roles in development, disease, and regeneration. *Nat. Rev. Cardiol.* 15, 672–684. doi: 10.1038/s41569-018-0063-3
- Watanabe, Y., Namba, A., Honda, K., Aida, Y., Matsumura, H., Shimizu, O., et al. (2009). IL-1 β stimulates the expression of prostaglandin receptor EP4 in human chondrocytes by increasing production of prostaglandin E2. *Connect. Tissue Res.* 50, 186–193. doi: 10.1080/03008200802588451
- Weitzmann, M. N. (2013). The Role of Inflammatory Cytokines, the RANKL/OPG Axis, and the Immunological Interface in Physiological Bone Turnover and Osteoporosis. *Scientifica (Cairo)*. 2013, 1–29. doi: 10.1155/2013/125705
- Wendler, S., Schlundt, C., Bucher, C. H., Birkigt, J., Schipp, C. J., Volk, H.-D. D., et al. (2019). Immune modulation to enhance bone healing—a new concept to induce bone using prostacyclin to locally modulate immunity. *Front. Immunol.* 10, 1–16. doi: 10.3389/fimmu.2019.00713
- Wilhelmi, B. J., Blackwell, S. J., and Phillips, L. G. (1999). Langer's lines: To use or not to use. *Plast. Reconstr. Surg.* 104, 208–214. doi: 10.1097/00006534-199907000-00033
- Wilyard, C. (2018). Unlocking the secrets of scar-free skin healing. *Nature* 563, S86–S88. doi: 10.1038/d41586-018-07430-w
- Wolff, J. (1986). *The Law of Bone Remodelling*. Berlin, Heidelberg: Springer Berlin Heidelberg. doi: 10.1007/978-3-642-71031-5
- Wong, V. W., Rustad, K. C., Akaishi, S., Sorkin, M., Glotzbach, J. P., Januszky, M., et al. (2012). Focal adhesion kinase links mechanical force to skin fibrosis via inflammatory signaling. *Nat. Med.* 18, 148–152. doi: 10.1038/nm.2574
- Wu, A. C., Raggatt, L. J., Alexander, K. A., and Pettit, A. R. (2013). Unraveling macrophage contributions to bone repair. *Bonekey Rep.* 2:373. doi: 10.1038/bonekey.2013.107
- Wu, L., Gao, X., Brown, R. C., Heller, S., and O'Neil, R. G. (2007). Dual role of the TRPV4 channel as a sensor of flow and osmolality in renal epithelial cells. *Am. J. Physiol. - Ren. Physiol.* 293, 1699–1713. doi: 10.1152/ajprenal.00462.2006
- Wynn, T. A. (2004). Fibrotic disease and the TH1/TH2 paradigm. *Nat. Rev. Immunol.* 4, 583–594. doi: 10.1038/nri1412
- Xia, Y., Lee, K., Li, N., Corbett, D., Mendoza, L., and Frangogiannis, N. G. (2009). Characterization of the inflammatory and fibrotic response in a mouse model of cardiac pressure overload. *Histochem. Cell Biol.* 131, 471–481. doi: 10.1007/s00418-008-0541-5
- Xing, Z., Lu, C., Hu, D., Yu, Y. Y., Wang, X., Colnot, C., et al. (2010). Multiple roles for CCR2 during fracture healing. *DMM Dis. Model. Mech.* 3, 451–458. doi: 10.1242/dmm.003186
- Yang, C., Wang, C., Zhou, J., Liang, Q., He, F., Li, F., et al. (2020). Fibronectin 1 activates WNT/ β -catenin signaling to induce osteogenic differentiation via integrin β 1 interaction. *Lab. Investig.* 100, 1494–1502. doi: 10.1038/s41374-020-0451-2
- Yang, X., Ricciardi, B. F., Hernandez-Soria, A., Shi, Y., Pleshko Camacho, N., and Bostrom, M. P. G. (2007). Callus mineralization and maturation are delayed during fracture healing in interleukin-6 knockout mice. *Bone* 41, 928–936. doi: 10.1016/j.bone.2007.07.022
- Yoshida, T., Friehs, I., Mummidi, S., del Nido, P. J., Addunour-Nakhoul, S., Delafontaine, P., et al. (2014). Pressure overload induces IL-18 and IL-18R expression, but markedly suppresses IL-18BP expression in a rabbit model. IL-18 potentiates TNF- α -induced cardiomyocyte death. *J. Mol. Cell. Cardiol.* 75, 141–151. doi: 10.1016/j.yjmcc.2014.07.007
- Zaiss, M. M., Axmann, R., Zwerina, J., Polzer, K., Gu, E., Skapenko, A., et al. (2007). Treg cells suppress osteoclast formation a new link between the immune system and bone. *Arthritis Rheum* 56, 4104–4112. doi: 10.1002/art.23138
- Zhang, F., Koyama, Y., Sanuki, R., Mitsui, N., Suzuki, N., Kimura, A., et al. (2010). IL-17A stimulates the expression of inflammatory cytokines via celecoxib-blocked prostaglandin in MC3T3-E1 cells. *Arch. Oral Biol.* 55, 679–688. doi: 10.1016/j.archoralbio.2010.06.003
- Zheng, M., Ambesi, A., and and, J., McKeown-Longo, P. (2020). Role of TLR4 receptor complex in the regulation of the innate immune response by fibronectin. *Cells* 9:216. doi: 10.3390/cells9010216

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 Knecht, Bucher, Van Linthout, Tschöpe, Schmidt-Bleek and Duda. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.