

TeV to PeV neutrinos and gamma-rays with Mountain SHALON Mirror Cherenkov Telescope.

V. G. Sinitsyna, T. P. Arsov, S. S. Borisov, F. I. Musin,
S. I. Nikolsky, V. Y. Sinitsyna and G. F. Platonov

P. N. Lebedev Physical Institute, Leninsky pr. 53, Moscow, Russia

Presenter: V. G. Sinitsyna (sinits@sci.lebedev.ru), rus-sinitsina-VG-abs4-og25-poster

The detection of Extraterrestrial Very High Energy Neutrinos by Atmospheric Cherenkov Telescopic System SHALON is discussed. The analysis of results of observation of extensive air showers at height of 3338 m above the sea level by gamma-telescope SHALON at the zenith angles 72° , 76° , 84° , 97° are presented. The observation results are compared with the data of detection of showers according to the direction into the zenith. The observation has been carried out at Tien -Shan station with SHALON gamma-telescope. The SHALON telescopic system has mirror with area of 11.2 m^2 and 144 PMT matrix with angular resolution of experimental method of $0.1^\circ - 0.2^\circ$, and the most in the world angular size of $> 8^\circ$. It allows to control the background of cosmic ray particle emission and the atmospheric transparency continuously with observation that means the increasing of observation efficiency. So it is the telescope characteristics that permit to start the search of local neutrino sources with energy $10^{13} - 10^{16} \text{ eV}$ on EAS generating in mountain-range located at some 7 and more kilometers from gamma-telescope (in Russian the SHALON abbreviation means - the Extensive Air Showers from Neutrino).

1. Introduction

The gamma-source search is the first step of the astronomy to the very high energy range. Only neutrino astronomy in the future can complete the search and investigation of galactic and metagalactic objects where the proton and nucleus acceleration processes are realized. These processes accompanied by generation of gamma-quanta and neutrino which are not scattered in Universe magnetic field, that gives an addition possibilities of investigation of the properties of active stellar objects at a very high energy. The observations have been carried out on P. N. Lebedev Physical Institute Tien - Shan high-mountainous station with gamma-telescope SHALON-1 since 1992 and started operating SHALON-2 [1 – 5].

2. The Observation of Cherenkov Bursts at the Zenith Angle 97°

UHE neutrinos may be searched in observations at large zenith angle. Because of mountain valley screens from background showers, the only particles that can survive are neutrinos with energy $> 10^{13} \text{ eV}$, come under the horizon, interact with mountain matter and product of the interaction produce an extensive air showers. In actual conditions the mirror telescope placement (Fig. 1) the distance till the opposite slope of the gorge is $\sim 7 \text{ km}$ or ~ 16.5 radiation units of length, that is quite enough for the development of an electromagnetic cascade till the structure characteristic for the rarefied atmosphere. The purpose of observations was revealing of background conditions when anthropogenic sources of light are absent. During 206 hours of observations 5 events were detected (Fig. 2a) which have expected angular characteristics of a light burst of an electron-photon cascade developing within a telescope observation angle (Fig. 2b). These showers have energy in the range of about $5 - 15 \text{ TeV}$. All other 265 events of detection of short-range light bursts in the atmosphere have not a narrow angle light direction and are chaotically distributed along the whole matrix or its part of a light-receiver (Fig. 1). These events may be interpreted as a reflection of a Cherenkov burst from a snow mountain slope or

the earth surface. Secondary, the observation of Cherenkov bursts from extensive air showers under the large zenith angles, for example using of horizontal extensive air showers for investigation of an energy spectrum of ultra-high energy cosmic rays is complicated by absorption of Cherenkov photons by a large atmosphere thickness.

4. Conclusion

It is supposed to overcome the main difficulty of observation of EAS, generated by neutrino in conditions of high mountainous observations, connected with the small cross section of neutrino-nuclei inelastic collisions. Two facts allow to carry out the search experiments. The hadron cascades with energy $> 10^{13}$ eV are generated by neutrino in the ground of mountain on the thickness < 300 g/cm². The Cherenkov radiation of the hadron cascades will be observed along the direction of neutrino by gamma-telescope placed on the distance about 7 kms from a mountain slope in area of more than 7×10^5 m². These cascades look like the usual extensive air showers generated in atmosphere with narrow light shape. Presently, the fluxes of galactic gamma-quantum sources Cygnus X-3, Tycho's SNR, Geminga of $< 10^{-14}$ cm⁻² s⁻¹ are observed by SHALON. The appearing of one shower per > 100 observation hours is expected if the flux of neutrino from local sources is 10^{-15} cm⁻² s⁻¹ [1 – 12]. During 206 hours of observations 5 events were detected which have expected angular characteristics of a light burst of an electron-photon cascade developing within a telescope observation angle.

References

- [1] Nikolsky S. I., Sinitsyna V. G., VANT Ser. TFE. 1331 (1987) 30.
- [2] Sinitsyna V. G., Toward a Major Atmospheric Cherenkov Detector -II, ed. Lamb R. C. (Iowa State University) (1993) 91.
- [3] Sinitsyna V. G., Nuovo Cimento 19C(6) (1996) 965.
- [4] Sinitsyna V. G., Nikolsky S. I., et.al., Nucl. Phys. B (Proc. Suppl.) 75A (1999) 352; 97, (2001) 215, 219.
- [5] Nikolsky S. I. and Sinitsyna V. G., Physics of Atomic Nuclei 67(10) (2004) 1900.
- [6] Klapdor-Kliengrothaus H.V., Zuber K., in Teilchenastrophysik, ed. Teubner B.G. (GmbH, Stuttgart, 2000).
- [7] Resvanis L.K., Nucl. Phys. B (Proc. Suppl.) **122**, 24 (2003).
- [8] Degrange B., Toward a Major Atmospheric Cherenkov Detector-I, ed. Fleury P., Vacanti G. (Frontieres) (1992) 171.
- [9] Bugaev E.V. and Shlepin Yu. V., Nucl. Phys. B (Proc. Suppl.) 122, 341 (2003).
- [10] Fargion D., et. al., ApJ 613, 1285 (2004).
- [11] Berezinsky V.S., Gzaev A.Z., JETP lett 25(5) (1977) 276.
- [12] Zatsepin G.T. and Berezinsky V.S., Neutrino Astrophysic Problems 2 (INR, Russian Academy of Science, 1980) 172.