
CMS Conference Report

August 29, 2006

Technicolour and other Beyond the Standard Model alternatives in CMS

P. Kreuzer

Univ. of Athens, Greece

for the CMS collaboration

Presented at “PHYSICS at LHC”, Cracow/Poland, July 2006.

Abstract

The expected signal sensitivity of the $\rho_{TC} \rightarrow W + Z$ decay channel is studied for the CMS detector, under the Technicolour “Straw Man” model. It shows a signal discovery potential at integrated luminosities starting from $\int L dt \simeq 4 \text{ fb}^{-1}$. Secondly, the CMS discovery potential of the heavy Majorana neutrino N_e and the right-handed gauge boson W_R is demonstrated, under the minimal LR symmetric model, at an early stage of the low luminosity running.

1 Search for Technicolour in the $\rho_{TC} \rightarrow W + Z$ channel

Technicolour (TC) stands as an alternative to the elementary Higgs mechanism of the Standard Model (SM) and elegantly solves the naturalness, hierarchy and triviality problems [1, 2]. It introduces a new strong interaction with $(N_{TC}^2 - 1)$ technigluons, at an energy scale $\Lambda_{TC} \sim \nu_{weak} \sim 200$ GeV, providing a dynamical nature to Electroweak Symmetry Breaking (EWSB). The original model was developed and scaled from QCD, in particular the non-zero vacuum expectation value of a technifermion condensate, yielding technipions¹⁾. Technicolour spontaneously breaks electroweak interactions down to electromagnetism and the technipions (Goldstone bosons) become the longitudinal components of the SM gauge bosons W^{+-} and Z. The latter acquire their known masses, proportional to the technipion decay constant $F_\pi = 246$ GeV. As a consequence, the arbitrary introduction of any Higgs doublet is avoided in dynamic EWSB.

“Extended technicolour” (ETC) interactions must be introduced to produce the SM fermion masses: they are embedded in a larger gauge group $SU(N_{TC}) \otimes SU(3)_C \otimes SU(2)_L \otimes U(1)_Y$ and are broken down to colour and technicolour at an energy scale $\Lambda_{ETC} = M_{ETC}/g_{ETC}$. ETC interactions generate the masses of SM quarks and of any light technipion. Moreover, they give rise to quark mixing: experimental limits on Flavor Changing Neutral Currents (FCNC) force the scale Λ_{ETC} to lay around 100-1000 TeV. To obtain quark masses that are large enough then requires an enhancement of the technifermion condensate over that obtained by naive scaling from QCD. This occurs if the technicolour gauge coupling runs very slowly or “walks”. Many technifermions N_D are typically needed in “Walking TC” [1], reducing the expected energy scale (< 1 TeV) of the lightest technicolour resonances technirho (ρ_{TC}) and techniomega (ω_{TC}). The model is completed with topcolour-assisted technicolour (TC2) [3], in order to integrate the generation of the top quark mass.

The present analysis [4] is performed under the phenomenology of the lowest-lying technihadrons, commonly referenced as the technicolour “Straw Man” model (TCSM) [5]. The colour-singlet sector includes the pseudo-scalar technimesons π_{TC} and the vector technimesons ρ_{TC} and ω_{TC} . The decay of ρ_{TC} is expressed as an admixture of π_{TC} and the Standard Model Z and W bosons:

$$\rho_{TC} \rightarrow \cos^2 \chi < \pi_{TC} \pi_{TC} > + 2 \cos \chi \sin \chi < \pi_{TC} W_L > + \sin^2 \chi < W_L W_L > \quad (1)$$

where W_L is the longitudinal mode of the Z or W and $\sin \chi \simeq 1/\sqrt{N_D} \sim 1/3$. The branching fraction $BR(\rho_{TC} \rightarrow W + Z)$ is competing with the two first terms in (1), hence with $M(\pi_{TC})$.

From the experimental point of view, the basic element is the search for a resonance decaying into di-bosons. In particular, the decay channel $\rho_{TC} \rightarrow W + Z$ has the advantage of a very clean final state, namely $3\ell + \nu$; the corresponding production diagram is shown in Fig. 1.




Figure 1: Main $\rho_{TC} \rightarrow W + Z$ production mode at LHC.

Other decay modes including jets, like $\rho_{TC} \rightarrow W + \pi_{TC} \rightarrow \ell \nu b\bar{b}$, have better branching fractions but are more difficult to disentangle from the Standard Model processes.

The most relevant background contributions to the signal in Fig. 1 are $WZ \rightarrow 3\ell + \nu$ (labeled “WZ” below), $ZZ \rightarrow 4\ell$ (labeled “ZZ” below), $Zb\bar{b} \rightarrow 2\ell + X$ (labeled “Z $b\bar{b}$ ” below) and $t\bar{t}$.

1.1 Event Reconstruction and Selection path

All signal and background samples used in this analysis are generated with PYTHIA 6.2 [6]²⁾ with the requirement of at least 3 prompt leptons in the CMS fiducial region. A set of 14 different ρ_{TC} samples are generated within the $[M(\rho_{TC}), M(\pi_{TC})]$ phase space.

The CMS fast simulation (FAMOS _1.4_0 [8]) is used for detector simulation and event reconstruction. Event pileup is taken into account, according to the low instantaneous luminosity scenario of $2 \times 10^{33} \text{cm}^{-2}\text{s}^{-1}$, and nominal

¹⁾ Similarly to QCD, where a quark condensate yields pions at $\Lambda_{QCD} \sim 200$ MeV.

²⁾ The $Zb\bar{b}$ background is generated using COMPHEP [7] interfaced to PYTHIA

CMS Level-1 and High-Level Trigger (HLT) requirements are applied [9]. The main reconstructed objects are leptons (muons and electrons) and the Missing Transverse Energy; their reconstruction quality and efficiency have been validated against the detailed GEANT-based CMS detector simulation [10]. The analysis path is summarized as follows:

- (i) Lepton Selection: 3 high- p_T and isolated electrons or muons.
- (ii) Lepton Trigger: Single- or Two-electron or muon mode.
- (iii) Z : same-flavor/opp.-charge ℓ -pair closest to $M(Z)$, $p_T > (30, 10)$ GeV/c
- (iv) W : 3rd lepton with $p_T > 10$ GeV/c + Missing E_T + $M(W)$ constraint
- (v) $|M(\ell^+\ell^-) - M(Z)| \leq 3\sigma_{M_Z} \cong 7.8$ GeV/c²
- (vi) $p_T(Z)$ and $p_T(W) > 30$ GeV/c³⁾
- (vii) $|\Delta[\eta(Z) - \eta(W)]| \leq 1.2$

The Z and W are reconstructed with a purity of $\sim 99\%$, using the 3 highest- p_T leptons in the event. The Missing E_T is obtained as the vector sum of the jets in the event (“Iterative Cone” algorithm), with an energy resolution of 23% for signal events. The $M(W)$ constraint yields a 2 fold ambiguity in the p_Z component of the reconstructed neutrino: it is found that the most efficient choice for the ρ_{TC} signal is the minimum p_Z solution. The kinematic cuts are illustrated in Fig. 2. The main $t\bar{t}$ reduction is obtained via the Z -mass window requirement (v). The irreducible background $WZ \rightarrow 3\ell + \nu$ is most efficiently separated from the signal via the $\eta(Z) - \eta(W)$ correlation requirement (vii). The p_T cut on Z and W further improves the signal to background ratio, however it is kept




Figure 2: (a) $M(\mu^+\mu^-)$ for $\rho_{TC}(300)$ and $t\bar{t}$; (b) $\Delta[\eta(Z) - \eta(W)]$ for $\rho_{TC}(300)$ and WZ ; (c) $p_T(Z)$ for $\rho_{TC}(300)$ and all backgrounds ($p_T(W)$ is similar); (d) Reconstructed $M(3\ell + \nu)$ for $\rho_{TC}(300)$ and all backgrounds. The vertical lines indicate the applied requirements.

modest in order to preserve the exponential background hypothesis of the $3\ell + \nu$ invariant mass spectrum, used to compute the signal sensitivity. The $\rho_{TC}(300)$ signal and background yields are shown in Fig. 2(d) and the corresponding reconstruction efficiencies are listed in Table 1.

1.2 Signal sensitivity and systematic uncertainties

The sensitivity of each ρ_{TC} benchmark point is computed by taking into account realistic statistical fluctuations for a given integrated luminosity. The sensitivity estimator is defined as the likelihood-ratio $S_L = \sqrt{2 \ln(\mathcal{L}_{S+B}/\mathcal{L}_B)}$, where \mathcal{L}_{S+B} and \mathcal{L}_B are the best-fit likelihoods of the signal-plus-background hypothesis and the null hypothesis

³⁾ For benchmark points with $M(\rho_{TC}) = 200$ GeV/c², the minimum $p_T(Z)$ and $p_T(W)$ threshold is 10 GeV/c.

Table 1: $\sigma \times \text{BR}(\ell = e \text{ or } \mu)$, 3-lepton preselection efficiency, total efficiency and final yield within 3σ of the signal region (Nev), for $\mathcal{L} = 5 \text{ fb}^{-1}$. $\rho_{TC}(300)$ and the main background contributions are shown. The simulation is repeated for all ρ_{TC} benchmark points.

Sample	$\sigma \times \text{BR}(\text{pb})$	$\epsilon(3\text{-lept})$	$\epsilon(\text{Reco}) (\%)$	Nev(5 fb^{-1})
$\rho_{TC} \rightarrow W + Z$	0.13	0.635	25.88 +- 0.40	103
$WZ \rightarrow 3\ell + \nu$	0.39	0.471	9.91 +- 0.11	27
$ZZ \rightarrow 4\ell$	0.07	0.719	15.80 +- 0.14	10
$Zb\bar{b} \rightarrow 2\ell + X$	332	0.046	0.23 +- 0.01	12
$t\bar{t}$	489.72	0.065	0.019 +- 0.001	8

(no signal present). The signal probability density function (p.d.f.) is assumed Gaussian (dominated by detector resolution) and the background p.d.f. is Exponential in all ρ_{TC} fit regions. The output of the fitting procedure is shown in the contour plot over the $[M(\rho_{TC}), M(\pi_{TC})]$ phase space in Fig. 3(a), for various integrated luminosities. A signal sensitivity above 5 is expected for $\mathcal{L} = 3 \text{ fb}^{-1}$ (before including systematic uncertainties).




Figure 3: (a) Signal 5σ Sensitivity curves for various integrated luminosities; (b) sensitivity for $\mathcal{L} = 4 \text{ fb}^{-1}$: the dotted (resp. dashed) curve shows the sensitivity (resp. the 90% C.L. signal upper limit) after including systematic uncertainties.

The ρ_{TC} sensitivity has been simulated for the early CMS data taking phase. Expected detector related systematic uncertainties for $\mathcal{L} = 1 \text{ fb}^{-1}$ are taken into account. While no substantial contribution is found from the tracker and muon system misalignment or the calorimeter miscalibration, the accuracy at which the lepton efficiency will be determined from data affects the result: a 2% uncertainty is considered. Moreover, the lepton fake rate has been simulated on $Zb\bar{b}$ and extrapolated to any $Z + jet(s)$ type background⁴⁾, in order to take into account additional contaminations from pion/kaon decays or from wrongly identified lepton candidates: a single lepton fake rate of $O(10^{-3})$ is obtained with FAMOS, affecting the ρ_{TC} sensitivity as shown below. Finally, a 7.5% uncertainty on the Missing E_T (MET) measurement is considered. The above uncertainties result in the relative ρ_{TC} sensitivity drop $\Delta_{SYS}^{tot} = \sqrt{(\Delta_{Eff})^2 + (\Delta_{Fake})^2 + (\Delta_{MET})^2} = \sqrt{(2.7\%)^2 + (8.5\%)^2 + (6.6\%)^2} = 11\%$. Concerning the generated cross section, introducing Next-to-Leading-Order K-factors for signal and background leads to a relative signal sensitivity increase of 6%; however the latter correction is not included in the final result shown in Fig. 3(b).

2 Detection of heavy Majorana neutrinos and right-handed bosons

Left-right (LR) symmetric models represent another interesting extension of the Standard Model, since they naturally explain parity violation of electroweak interactions. In particular, the minimal LR symmetric model [11, 12] built under the gauge symmetry group $SU_C(3) \otimes SU_L(2) \otimes SU_R(2) \otimes U(1)$ embeds the SM at the scale of the order 1 TeV and the Higgs sector consists of a bi-doublet and two triplets. Three additional gauge bosons W_R and Z' necessarily appear, together with the heavy Majorana neutrino states (N_ℓ) [13]. The latter can provide non-zero masses to their lighter partners ν_ℓ via the see-saw mechanism [14]. The relevance of LR symmetric models has increased since the experimental evidence of neutrino oscillations [15]. Existing experimental data have set lower bounds to the Z' and W_R masses of $O(1)$ TeV [16] and 1.6 TeV [17], respectively, with large uncertainties. This analysis [18] is performed under the assumption $M(W_R) > 1 \text{ TeV}$.

Among several production modes of N_ℓ and W_R in pp collisions, the most promising in terms of cross-section

⁴⁾ A production cross-section of 1047pb per lepton flavor is assumed for $Z + n\text{-jets}$.

and the most suitable for heavy neutrino searches is given in Fig. 4. At LHC energies, the electron flavor N_e is expected to dominate heavier flavors, yielding the signature $pp \rightarrow e + N_e \rightarrow e + eW_R \rightarrow 2e + 2 jets$.




Figure 4: Heavy Majorana neutrino N_ℓ production through a W_R boson.

The main background contributions are expected from SM processes with a lepton pair and at least two jets in the final state, namely WZ (leptonic W decays only and no hadronic Z decays), $Z + jets$, $t\bar{t}$ (leptonic W decays only), ZH and WH .

2.1 Event Reconstruction and Selection path

All signal and background events are generated and their cross section computed with PYTHIA 6.2 [6]. The signal uses default CTEQ5L parton distribution functions [19] and the set of parameters listed in [18].

The reconstruction is performed with the GEANT-based full CMS detector simulation [10]. Event pileup is taken into account, according to the low instantaneous luminosity scenario of $2 \times 10^{33} \text{cm}^{-2}\text{s}^{-1}$, and nominal Level-1 (resp. HLT) electron trigger requirements [9] are applied, yielding a signal efficiency of 100 % (resp. 99 %). All reconstructed electron⁵⁾ candidates are required to satisfy $E_T > 20$ GeV and a Tracker isolation flag is set within a cone of radius 0.3 around the electron track. Jets are reconstructed by the Iterative Cone algorithm, with a minimum E_T requirement of 40 GeV.

A primary selection of at least 2 isolated electrons and 2 jets is made. Furthermore, only events with two isolated electrons are kept (e_1, e_2), with the invariant mass requirement $M_{e_1 e_2} > 200$ GeV, and only the two highest- p_T jets are considered (j_1, j_2). A mass window of 110 GeV (optimized on S/B) is required around the reconstructed heavy neutrino invariant mass $M_{N_e}^{cand} = M_{e j_1 j_2}$ ⁶⁾ and a threshold of 1 TeV is required on the combined system $M_{W_R}^{cand} = M_{e_1 e_2 j_1 j_2}$. The event yields throughout the selection path are shown in Table 2, for the signal benchmark point $(M_{N_e}, M_{W_R}) = (500, 2000)$ GeV (called LRRP below) and for all significant background contributions.

Table 2: Event yields throughout the selection path, for signal and background. Due to processing limitations, only a fraction of $Z + jets$ events are fully simulated.

Step	Signal	$t\bar{t}$	$Z+jets$	ZW	WH
Generated	4965	2.64×10^6	6.2×10^7	6×10^4	11000
Primary Selection	2782	1.5×10^5	-	38	728
2 isolated e	2332	152000	-	15	165
$M_{e_1 e_2} > 200$ GeV	2246	17200	3870	0	72
$M_{N_e}^{cand}$ window	970	3430	1000	0	2
+ $M_{W_R}^{cand} > 1$ TeV	938	198	96	0	0

The $Z + jets$ background has the largest production cross section and is reduced by minimum E_T requirements on reconstructed electron and jet objects. The $M_{e_1 e_2}$ cut dramatically improves the S/B ratio of any type of reaction including a Z . The largest background contribution after full selection is $t\bar{t}$. It has been checked that only leptonic W decay modes from $t\bar{t}$ contribute. Finally, backgrounds containing a Higgs are almost negligible, due to their relatively small production cross section.

The heavy Majorana neutrino search will be performed by first selecting events with $M_{W_R}^{cand} > 1$ TeV, followed by

⁵⁾ For simplicity, positrons are called “electrons” in the text.

⁶⁾ Both combinations $e_1 j_1 j_2$ and $e_2 j_1 j_2$ are kept in the final spectra.

a scan over the reconstructed $M_{N_e}^{cand}$ spectrum. This is illustrated in Fig. 5, for an integrated luminosity of $\int Ldt = 30 \text{ fb}^{-1}$: a large S/B ratio is expected for the LRRP benchmark point.




Figure 5: (a) Reconstructed gauge boson W_R invariant mass; (b) reconstructed heavy Majorana neutrino N_e invariant mass, after a 1 TeV threshold has been required on $M_{W_R}^{cand}$. The signal is shown in open white and the total background in shaded style.

2.2 Signal sensitivity and systematic uncertainties

The expected discovery potential of N_e and W_R at CMS is calculated using the significance estimator $S = 2(\sqrt{N_S - N_B} - \sqrt{N_B}) \geq 5$ [20]. The corresponding discovery contours are shown in Fig. 6, for various integrated luminosities. Invariant mass regions up to $(M_{N_e}, M_{W_R}) = (3.5, 2.3)$ TeV are reachable after 3 years of running at low luminosity ($\int Ldt = 30 \text{ fb}^{-1}$). Lower mass regions (e.g. the LRRP benchmark point) are reachable after only a few fb^{-1} .

The expected uncertainty of this prediction related to various systematic background uncertainties is small, since the background itself is small. The discovery region is mainly limited by the fast drop of the signal cross section at high ratios $r = M_{N_e}/M_{W_R}$ or by the fast drop of signal efficiency at small r , and the contours on Fig. 6 are barely affected by systematic uncertainties. As for the generated signal cross sections, various parton density functions sets have been used to take into account theoretical fluctuations [21]: they lead to a 6% uncertainty on the cross section and to a systematic error of 1-3 % on the significance prediction over whole discovery region.




Figure 6: CMS discovery potential of the heavy Majorana neutrino N_e and the right-handed gauge boson W_R for $\int Ldt = 30, 10$ and 1 fb^{-1} (from outer to inner contour, respectively). The horizontal exclusion line was set by the L3 experiment [22].

3 Conclusions

The signature $\rho_{TC} \rightarrow W + Z$ in the context of the Technicolour “Straw Man” model is studied for the CMS detector. A 5 sigma discovery reach is obtained for an integrated luminosity $\mathcal{L} \simeq 4 \text{ fb}^{-1}$.

The discovery potential of the heavy Majorana neutrino N_e and the right-handed gauge boson W_R is demonstrated, under the minimal LR symmetric model, for only a few fb^{-1} of running at CMS.

Both predictions represent a potential handle into Physics Beyond the Standard Model, at an early stage of the LHC era.

References

- [1] K. D. Lane, “Technicolor 2000”, hep-ph/0007304.
- [2] K. Lane, “Two lectures on technicolor”, hep-ph/0202255.
- [3] C. T. Hill, “Topcolor assisted technicolor”, Phys. Lett. **B345**, 483 (1995), [hep-ph/9411426].
- [4] P. Kreuzer, “Search for Technicolor at CMS in the $\rho_{TC} \rightarrow W + Z$ channel”, CMS Note **2006-135** (2006).
- [5] K. Lane and S. Mrenna, “The collider phenomenology of technihadrons in the technicolor Straw Man Model”, Phys. Rev. **D67**, 115011 (2003), [hep-ph/0210299].
- [6] T. Sjostrand, L. Lonnblad and S. Mrenna, “PYTHIA 6.2: Physics and manual”, hep-ph/0108264.
- [7] A. Pukhov *et al.*, “CompHEP: A package for evaluation of Feynman diagrams and integration over multi-particle phase space. User’s manual for version 33”, hep-ph/9908288.
- [8] The CMS collaboration, “CMS Physics TDR Volume 1, Section 2.6: Fast simulation”, CERN/LHCC **2006-001**, 55 (2006), CMS TDR 8.1.
- [9] The CMS Collaboration, “The TriDAS Project Technical Design Report, Volume 2: Data Acquisition and High-Level Trigger”, CERN/LHCC **2002-26** (2002), CMS TDR 6.2.
- [10] S. Wynhoff *et al.*, “ORCA: CMS Reconstruction Package”, Site located at <http://cmsdoc.cern.ch/orca>.
- [11] R. N. Mohapatra and J. C. Pati, “A ‘NATURAL’ LEFT-RIGHT SYMMETRY”, Phys. Rev. **D11**, 2558 (1975).
- [12] G. Senjanovic and R. N. Mohapatra, “EXACT LEFT-RIGHT SYMMETRY AND SPONTANEOUS VIOLATION OF PARITY”, Phys. Rev. **D12**, 1502 (1975).
- [13] M.Gell-Mann *et al.*, “SUPERGRAVITY. PROCEEDINGS, WORKSHOP AT STONY BROOK, 27-29 SEPTEMBER 1979”, Amsterdam, Netherlands: North-holland (1979) 341p.
- [14] R. N. Mohapatra and G. Senjanovic, “Neutrino mass and spontaneous parity nonconservation”, Phys. Rev. Lett. **44**, 912 (1980).
- [15] C. Giunti and M. Laveder, “Neutrino mixing”, hep-ph/0310238.
- [16] Particle Data Group, S. Eidelman *et al.*, “Review of particle physics”, Phys. Lett. **B592**, 1 (2004).
- [17] G. Barenboim, J. Bernabeu, J. Prades and M. Raidal, “Constraints on the W(R) mass and CP-violation in left-right models”, Phys. Rev. **D55**, 4213 (1997), [hep-ph/9611347].
- [18] S. Gninenko, M. Kirsanov, N. Krasnikov and V. Matveev, “Detection of heavy Majorana neutrinos and right handed bosons”, CMS Note **2006-098** (2006).
- [19] CTEQ, J. Botts *et al.*, “CTEQ parton distributions and flavor dependence of sea quarks”, Phys. Lett. **B304**, 159 (1993), [hep-ph/9303255].
- [20] S. I. Bityukov and N. V. Krasnikov, “Uncertainties and discovery potential in planned experiments”, hep-ph/0204326.
- [21] P. Bertalini, R. Chierci and A. De Roeck, “Guidelines for the estimation of theoretical uncertainties at the LHC”, CMS Note **2005-013** (2005).
- [22] L3, P. Achard *et al.*, “Search for heavy isosinglet neutrino in $e^+ e^-$ annihilation at LEP”, Phys. Lett. **B517**, 67 (2001), [hep-ex/0107014].