

Smoking Guns for On-Shell New Physics at the LHC

Christian Arnesen,¹ Ira Z. Rothstein,¹ and Jure Zupan^{2,3,4}

¹*Carnegie Mellon University, Department of Physics, Pittsburgh Pennsylvania 15213, USA*

²*Theory Division, Department of Physics, CERN, CH-1211 Geneva 23, Switzerland*

³*J. Stefan Institute, Jamova 39, 1000 Ljubljana, Slovenia*

⁴*Faculty of Mathematics and Physics, University of Ljubljana, Jadranska 19, 1000 Ljubljana, Slovenia*

(Received 16 September 2008; published 8 October 2009)

Using Tevatron bounds we derive upper limits on the LHC Higgs boson production rate assuming that no beyond the standard model (BSM) particles are being produced near their mass shell. A violation of these limits would constitute a smoking gun for light BSM particles. Furthermore, we demonstrate how R_T , the ratio of the partially integrated Higgs transverse momentum distribution to the inclusive rate, can also be used as a probe of light BSM particles. This ratio is insensitive to heavy virtual effects and is approximately model independent. The perturbative expansion for R_T has reduced renormalization scale dependence, due to a cancellation of Wilson coefficients. A deviation from the SM value implies that light BSM particles are being produced near their mass shell. A model with colored scalars is used to investigate the model independence of R_T .

DOI: [10.1103/PhysRevLett.103.151801](https://doi.org/10.1103/PhysRevLett.103.151801)

PACS numbers: 13.85.Rm, 12.60.Fr, 14.80.Bn

The Higgs sector of the standard model (SM), responsible for electroweak symmetry breaking, has remained hidden from observation. At a 95% confidence level, LEP data constrain $m_H > 114.4$ GeV [1]. Direct searches at the Tevatron have put upper limits on Higgs event rates at 1–7 times the SM value for Higgs boson masses in the range 110–200 GeV [2]. At the Large Hadron Collider (LHC) particles many times more massive than the top quark will be kinematically accessible. We remain hopeful that beyond the standard model (BSM) particle production will manifest itself in resonances or shoulders of distributions, but we may not be that fortunate. New-physics (NP) signals must be separated from large SM backgrounds in the complicated environment of a hadron collider, and NP discovery may remain elusive. Here we will explore a modest strategy in which we ask, “How can we discern if deviations from the SM are due to particles being produced on-shell?” We approach this question through the framework of effective field theory (EFT).

The premise of EFT is that our theoretical description of low energy observables need not include heavier particles as dynamical degrees of freedom. Instead we can approximate virtual exchange of the massive particles as a set of local contact interactions. The approximation is a power series expansion in p^2/Λ^2 where p^2 is a typical kinematic invariant in the process and the EFT “cutoff” Λ is the mass scale of the exchange. For instance, if the interaction is mediated by pair-produced particles of mass m then $\Lambda \sim 2m$. At each order in the expansion, the set of local operators is the most general one consistent with low energy symmetries, whatever the massive dynamics may be. Absent additional model assumptions, the Wilson coefficients of the local operators must be determined phenomenologically. Then the EFT predicts concrete relationships

between collider observables such as production rates or branching ratios in related channels.

In this Letter, we make model independent EFT predictions for new physics in Higgs production through gluon fusion, which starts at one loop in the SM and will be the dominant Higgs production mechanism at the LHC. The generality of the EFT approach ensures that any massive BSM extension can be accommodated. Conversely, deviations from the EFT predictions signify *model independently* that light particles with $m \sim p$ are being produced. We will demonstrate this for the case of Higgs production, where the presence of new particles with masses $m^2 \sim m_H^2 + p_T^2$ can be probed.

A complete basis of dimension-six EFT operators for the standard model was first constructed in [3]. At leading order in the power expansion, there is just one CP even and one CP odd operator that can modify the Higgs-gluon interaction

$$\mathcal{L}_{\text{eff}} = C_G H G_{\mu\nu}^a G^{a\mu\nu} + \tilde{C}_G H \tilde{G}_{\mu\nu}^a G^{a\mu\nu}, \quad (1)$$

where $G_{\mu\nu}^a$ is the gluon field strength and $\tilde{G}_{\mu\nu}^a = \epsilon_{\mu\nu\rho\sigma} G^{\rho\sigma} / 2$. For $m_H \lesssim 200$ GeV the SM Higgs-gluon interaction is pointlike to an excellent approximation due to the large top-quark mass and can be well described by the same effective Lagrangian, Eq. (1). The infinite-top-mass approximation has been used extensively in the literature since it greatly simplifies the calculation of radiative corrections [4–9]. In this limit, a top loop contributes $C_G^{\text{SM}} = \alpha_s / (12\pi v)$ to the effective Higgs-gluon coupling at leading order. Heavy NP will modify the value of $C_G = C_G^{\text{SM}} + C_G^{\text{NP}}$ and, if CP violating, generate \tilde{C}_G . If we assume that the new physics does not get mass from electroweak symmetry breaking, then the magnitude of \tilde{C}_G is bounded by

upper limits on the electric dipole moment of the neutron $d_n < 2.9 \times 10^{-26} e \text{ cm}$ [10]. For the moment we will neglect CP violating effects and return to this possibility in the conclusions.

Searches at the Fermilab CDF and D0 experiments put upper bounds on Higgs production rates [2,11], which we translate into bounds on C_G . The relevant production mechanisms are $p\bar{p} \rightarrow H(\rightarrow \gamma\gamma)$, $p\bar{p} \rightarrow H(\rightarrow WW^{(*)})$ and $p\bar{p} \rightarrow H(\rightarrow \tau\tau) + 2 \text{ jets}$. The first two channels are gluon fusion dominated. Note that Higgs production with an associated weak boson, which dominates the combined Tevatron Higgs bounds for small Higgs masses [2], are not sensitive to glue-Higgs interactions. The resulting 95% confidence level (C.L.) upper limits on $|C_G/C_G^{\text{SM}}|^2$ are shown in Fig. 1, assuming SM decays for the Higgs boson. This model independent result can be used to constrain the model dependent parameter space of SM extensions. Since gluon fusion is the dominant Higgs production mechanism at the LHC, we can also immediately translate Fig. 1 into model independent bounds on $\sigma(pp \rightarrow H + X)$ at the LHC at next-to-next-to-leading order (NNLO) in the expansion in α_s utilizing Fehip [9]. These LHC experimental bounds are plotted in Fig. 2 for a range of Higgs masses. A measurement exceeding these EFT bounds is a smoking gun for light BSM particles.

There are other model independent signatures for BSM particle production that do not rely on Tevatron input and can thus be improved at LHC with increasing statistics. The observable that we will focus on is the ratio of Higgs production cross section at large transverse momenta to the totally inclusive rate,

$$R_T \equiv \frac{\sigma(H: p_T^H > p_T^{\min})}{\sigma(H)}, \quad (2)$$

with a lab-frame rapidity cut $|y_H| < y_H^{\max}$ in both numerator and denominator. The transverse Higgs production rate is interesting in its own right as a Higgs search channel [12]. The ratio in Eq. (2) has several favorable properties:




FIG. 1 (color online). 95% C.L. upper bound on gluon-fusion Higgs production at the LHC relative to the SM value, $\sigma(H)/\sigma(H)_{\text{SM}} = |C_G/C_G^{\text{SM}}|^2$, coming from Tevatron searches [11] $2.0\text{--}3.0 \text{ fb}^{-1}$: $H(\rightarrow \gamma\gamma)$ (red line); $H(\rightarrow \tau\tau) + 2 \text{ jets}$ (blue line); $H \rightarrow WW^{(*)}$ (dotted line); combined analysis (solid line).

(i) independence from Higgs branching ratios, (ii) reduced perturbative uncertainty relative to the individual cross sections, and most importantly (iii) heavy NP cannot change its value from the SM one. These are *model independent* predictions of the EFT. Figure 3 shows the EFT prediction of this ratio calculated at NLO in α_s using Fehip [9] for a range of Higgs masses in the large- m_t limit. To have a clear EFT model independent interpretation of R_T , the vector boson fusion contributions should be subtracted from the measurement. This can be done precisely at NLO using VBFNLO [13].

The error band in Fig. 3 reflects the perturbative uncertainty, estimated from varying the renormalization scale by a factor of 2 in each direction. While the variation in the transverse and inclusive rates individually are of order 20%, the ratio is relatively insensitive with only 5% variation. A simple way to understand this reduction in errors can be seen from an EFT point of view. The theory calculation has two sources of α_s , from the matrix element and from the matching coefficient. In the large-top-mass limit all of the α_s dependence in the ratio stemming from the matching coefficient cancels. Errors due to parton distribution function (PDF) uncertainties also cancel to large extent in the ratio. A calculation of the cross sections using several different PDFs reveals that these errors are negligible compared to the perturbative errors [7]. When $|C_G^{\text{NP}}| \approx |C_G^{\text{SM}}|$, one needs to consider other corrections to the large-top-mass SM prediction beyond perturbative ones. Kinematic power corrections due to finite top-quark mass are small, $\leq 10\%$, for $m_h, p_T^{\min} \leq 200 \text{ GeV}$ [6], as are contributions through a virtual b -quark loop. The two-loop electroweak corrections modify the $gg \rightarrow h$ cross section at the 5%–8% level [14–16], but are likely smaller for the R_T ratio.

What could we conclude from a deviation from Fig. 3? One strong possibility would be that such a deviation would arise as a consequence of light colored particles. We might hope to see these particles in jet production cross sections, but absent knowledge of their decay widths or




FIG. 2 (color online). 95% C.L. upper bounds on the Higgs production cross section at the LHC from Tevatron Higgs searches [11]. The upper bound includes the error in the theoretical cross section at NNLO, which we conservatively approximate to be 20%. The lower lines are the SM prediction with associated errors.




FIG. 3 (color online). Model independent value of R_T from Eq. (2) at NLO in α_S with $p_T^{\min} = 30$ and $y_H^{\max} = 2$ as a function of Higgs mass. Vector boson fusion contributions are excluded. The uncertainty band is approximated by varying the renormalization scale by a factor of 2.

products, this may be difficult. However, we could learn more about their masses by altering the cuts on the Higgs p_T distribution and seeing how R_T changes. As the cut increases we would expect the deviations to grow. With sufficient statistics we should be able to find the value, p_T , at which the distribution begins to deviate from the SM and infer the presence of on-shell particles with mass $\sim \sqrt{m_H^2 + p_T^2}$. Note that in principle one could also get deviations without light BSM particles if some other higher dimension operators became relevant, for example, four-quark operators. PDF suppression is large though for quark-initiated Higgs production, and LEP bounds constrain the Wilson coefficients of these operators. So this possibility is excluded.

Toy Model.—Let us give a concrete example of how R_T can deviate from the EFT prediction when there are new light particles in the theory. As a toy-model example we extend the SM by adding a color-octet weak-singlet scalar S that couples to the Higgs field through a trilinear term

$$\mathcal{L} = -\frac{\kappa}{2} v H S^a S^a + \dots, \quad (3)$$

after electroweak symmetry breaking, where $v \simeq 246$ GeV is the Higgs vacuum expectation value and “+ ...” refers to other terms in the scalar potential that are not relevant to




FIG. 4 (color online). m_S dependence of R_T for $p_T^H > 30$ GeV (left) and $p_T^H > 100$ GeV (right) for the toy model (solid lines) and for the SM (dashed lines) at LO in α_S . The values of Higgs mass are as indicated. Toy-model calculations performed utilizing [18] are leading order in α_S (with full m_t and m_b dependence). No y_H cut is imposed and κ has been set to unity.

our analysis. The colored scalars contribute virtually through coannihilation, $pp \rightarrow 2S \rightarrow H$, and on-shell production, $pp \rightarrow H + 2S$. For the regions of parameter space explored here, the latter is numerically subleading due to gluon PDF suppression at large momentum fractions. The value of R_T in the SM and in the toy-model extension are shown in Fig. 4 for Higgs momentum cuts $p_T^{\min} = 30$ GeV (left) and $p_T^{\min} = 100$ GeV (right) and a range of octet-scalar masses.

If the scalar’s mass is larger than the typical partonic center-of-mass energy (set by the Higgs mass and momentum), then its effect is simply to shift the Wilson coefficient C_G in Eq. (1). At leading order in α_S ,

$$C_G^{\text{NP}} = \frac{\kappa \alpha_S N_c v}{96 \pi m_S^2}. \quad (4)$$

In this scenario, the transverse and inclusive production rates individually differ from their SM values by a factor $|(C_G^{\text{SM}} + C_G^{\text{NP}})/C_G^{\text{SM}}|^2$, but their ratio is unchanged. Qualitatively we can expect that deviations of R_T from the EFT prediction will only arise when the scalar is sufficiently light. This is seen in Fig. 4 where R_T approaches the SM value when m_S is large. Furthermore, as the cut on the Higgs p_T is increased we would expect this deviation to grow since a larger partonic center-of-mass energy leads to enhanced power corrections. This is a generic feature of all NP models that could modify R_T . Note that since the Higgs branching ratios cancel in the ratio, modification can only come from the Higgs production. In more realistic SM extensions, such as supersymmetry or models with extra singlet scalars, R_T will generally differ from its SM value with the magnitude of the deviation depending on the specific model parameters.

A benefit of the R_T ratio is that it does not rely on the decay properties of the new resonances. In our toy model the octet scalars would also be pair produced (without a Higgs boson), modifying the two-jet production cross section and other QCD observables. Searches in these channels, however, would depend on the scalar decay products, and dedicated studies may be needed. It could be that separating these signals from background is prohibitively challenging, for example, if there is some degeneracy with

another final state as in light stop scenarios in the minimal supersymmetric standard model [17].

Violations of observable predictions of the EFT may be searched for in other channels dominated by gluon fusion as well, for instance $H + 2$ jets. With increasing statistics, the EFT approach outlined here can also be used in channels that proceed at tree level in the SM, in which case EFT contributions would be a subleading effect at low energies. With high p_T cuts in TeV range, these effects may become leading, offering another way to search model independently for TeV resonances. We leave these possibilities for future studies.

Conclusions.—We showed that even if the Tevatron does not discover the Higgs boson, the tightening of constraints on its production may facilitate the search for on-shell BSM particles at the LHC. The Tevatron and LHC results are complementary because the collisions occur at different energies. For example, if the EFT description applies only to Tevatron results, the LHC bounds in Fig. 2 may be violated, acting as a smoking gun for on-shell particle production. Other observables can be used in the indirect searches for on-shell new particles. Figure 3 gives a model independent prediction for the ratio of the Higgs production rate at large transverse momenta to the inclusive rate. Should the data disagree significantly with this prediction then one may conclude that new particles have been produced in the collisions and presumably escaped detection either due to backgrounds or the nature of the decay products. The discriminating power of R_T to indicate on-shell new physics will depend on the magnitude of the deviation both in R_T and the overall production rate, the extent to which cuts on the Higgs decay products allow a measurement of R_T , and the magnitude and uncertainty of sizeable backgrounds. As such, a detailed analysis is beyond the scope of this Letter, however, particularly for an enhanced rate and large R_T deviation, preliminary estimates are promising.

This reasoning assumes that the CP violating Wilson coefficient \tilde{C}_G in Eq. (1) is negligible. This is true if the off-shell new physics is perturbative, as \tilde{C}_G can arise only at two loops and is thus much smaller than C_G^{SM} . Furthermore, the presence of \tilde{C}_G can be bounded model independently from experiment by measuring R_T at several Higgs p_T cuts. Namely, $R_T \propto (C_G^2 + \tilde{C}_G^2)/(C_G^2 + 2\tilde{C}_G^2)$, where the proportionality factor depends on the Higgs mass and p_T^{min} .

In the case of smaller deviations, a more refined analysis would be necessary to determine if the cause is light new physics. Electroweak corrections would have to be included along with the b -quark loop effects. When these effects become important R_T is no longer model independent. Nonetheless one could still use the EFT strategy discussed here to search for light new physics, by first extracting C_G from the inclusive rate and then using it to predict the cut rate.

We have presented a toy model with color-octet scalars. It is important to note, however, should the data fail to produce the correct value of R_T , it does not necessarily imply that the new particles are colored. Nonetheless, a natural place to look for these new particles would be in jet production rates, although our ability to detect such effects will be model dependent.

Finally, we would like to emphasize that even if new BSM particles are discovered in channels not involving the Higgs boson, R_T remains an interesting observable in its own right. For one, it is under better theoretical control due to the cancellation of QCD corrections. In addition, discovery of new particles not involved in Higgs production does not imply that R_T will deviate from its SM value. In fact if R_T did not deviate from its SM value in this scenario, this would provide very useful information in discerning models.

The authors would like to thank T. Ferguson, B. McElarh, T. Hahn, S. Jaeger, T. Rejec, J. Russ, J. Wells and K. Yorita. The work of C. A. and I. Z. R. is supported by DOE Contracts No. DOE-ER-40682-143 and No. DEAC02-6CH03000.

-
- [1] R. Barate *et al.* (LEP Working Group for Higgs boson searches), Phys. Lett. B **565**, 61 (2003).
 - [2] T.T.W. Group (CDF Collaboration and D0 Collaboration), arXiv:0804.3423.
 - [3] W. Buchmuller and D. Wyler, Phys. Lett. B **121**, 321 (1983).
 - [4] R.V. Harlander and W.B. Kilgore, Phys. Rev. Lett. **88**, 201801 (2002).
 - [5] C. Anastasiou and K. Melnikov, Nucl. Phys. B **646**, 220 (2002).
 - [6] C.J. Glosser and C.R. Schmidt, J. High Energy Phys. **12** (2002) 016.
 - [7] V. Ravindran, J. Smith, and W.L. van Neerven, Nucl. Phys. B **665**, 325 (2003).
 - [8] G. Bozzi, S. Catani, D. de Florian, and M. Grazzini, Nucl. Phys. B **737**, 73 (2006).
 - [9] C. Anastasiou *et al.*, Nucl. Phys. B **724**, 197 (2005).
 - [10] C.A. Baker *et al.*, Phys. Rev. Lett. **97**, 131801 (2006).
 - [11] G. Bernardi *et al.* (Tevatron New Phenomena Higgs Working Group and CDF Collaboration and D), arXiv:0808.0534.
 - [12] R.K. Ellis *et al.*, Nucl. Phys. B **297**, 221 (1988).
 - [13] V. Hankele *et al.*, Phys. Rev. D **74**, 095001 (2006).
 - [14] G. Degrossi and F. Maltoni, Phys. Lett. B **600**, 255 (2004).
 - [15] U. Aglietti, R. Bonciani, G. Degrossi, and A. Vicini, Phys. Lett. B **595**, 432 (2004).
 - [16] S. Actis, G. Passarino, C. Sturm, and S. Uccirati, Phys. Lett. B **670**, 12 (2008).
 - [17] M. Carena *et al.*, J. High Energy Phys. **10** (2008) 062.
 - [18] T. Hahn, Comput. Phys. Commun. **140**, 418 (2001).