

Spanning from exotic quarks to Higgs boson with ATLAS



N. Gökhan Ünel

University of California, Irvine

On Behalf of the ATLAS Collaboration

International Conference on Particle Physics, 27-31 October, 2008.

in memoriam of Prof. E. Arik and Her Colleagues



Contents

ATLAS experiment

SM and beyond

- Models w/ additional quarks

ATLAS prospects for

- Fourth Family model
 - quark searches
 - impact to Higgs hunt
- E6 GUT model
 - quark searches
 - impact to Higgs hunt

Conclusions

i n m e m o r i a m



Prof. Fatma Şenel Boydağ
CAST



Prof. İskender Hikmet,
CAST



Mustafa Fidan
CAST



Engin Abat
ATLAS



Berkol Doğan
ATLAS



Prof. Engin Arik
14 Oct. 1948 - 30 Nov. 2007
Participated CERN experiments: Charm-2, Chorus, SMC, CAST, ATLAS



SM and beyond

- SM explains all the experiments conducted so far,
 - BUT:
- What are the fundamental constituents of the matter?
 - Why do we have mass? Why do we exist?
- Is there a basic symmetry between the matter particles and the force carriers?
 - Are there really 4 forces governing the universe?
 - Can they be explained by a Unified Theory?
- Are there 3+1 dimensions in the universe we live in?
- What is the dark matter & blackholes we hear about?
 - Can they be produced in the lab?



What are the options?

SUSY looks as a promising candidate for a general theory

What if there is no SUSY ??

Little Higgs with T-parity? ★

*Fourth Family? ★

UED? ☆

Technicolor ? ★

Compositeness ? ☆

*GUT-inspired E_6 ? ★

★ Models predicting additional quarks


☆ Models predicting additional quarks in some cases

*In this talk, we focus on these models

Recent Hints

Tevatron results (CDF @ ICHEP)

- $Q \rightarrow W + \text{jet}$ channel, pair production (no b-tagging)
- 2σ "fluctuation" in direct searches at high mass tail.
- current limit $m_Q > 311 \text{ GeV}$



Belle $K\pi$ puzzle (Nature V452, 03/2008)

- $A_{\text{sym}}(B^+ \rightarrow K^+ \pi^0)$ vs $A_{\text{sym}}(B^0 \rightarrow K^+ \pi^-)$: difference in SM = 0
 - new CP source: needed for BAU!
- explainable with an extra quark in EW penguin?

$$\frac{n_B}{n_\gamma} = (5.1^{+0.3}_{-0.2}) \times 10^{-10}$$


WMAP

$$\text{KM} \sim 10^{-20}$$

Too Small in SM

$$m_B, m_\tau \cong 300 \text{ GeV} \quad 10^{+13}$$

$$\sim 600 \text{ GeV} \quad 10^{+15}$$




$$\Delta A_{K\pi} = A_{\text{cp}}(K^+ \pi^-) - A_{\text{cp}}(K^+ \pi^0)$$


$$= -0.147 \pm 0.028 @ 5.3\sigma$$




LHC: a gluon collider



gluons, s channel




gluons, t channel



quarks, s channel

LHC is mostly a gluon collider

- quarks are pair produced with large σ .
- quarks will be "easy" to discover:
 - small MET, can distinguish from susy
 - can fully reconstruct the event,
- First searches will look for Q




ATLAS currently


For further details see talk by P.Jenni in this conference

is well and taking data

- with cosmic rays...
- LHC beam expected in April 2009



- Inner Tracking ($|\eta| < 2.5$, 2T solenoid) :
 - Silicon pixels and strips
 - Transition Radiation (e/π separation)
- Calorimetry ($|\eta| < 5$) :
 - EM : Pb-LAr, Accordion shape
 - HAD: Fe/scintillator (centr), Cu/W-LAr (fwd)
- Muon Spectrometer ($|\eta| < 2.7$, 4T toroid) :
 - air-core toroids with muon chambers



first beam event seen in ATLAS

ID:	$\sigma/p_T \approx 5 \times 10^{-4} p_T \oplus 0.001$
	$\sigma(d_0) = 15 \mu\text{m}$ at 20 GeV
ECAL:	$\sigma/E \approx 10\% / \sqrt{E(\text{GeV})} \oplus 0.7\%$
HCAL:	$\sigma/E \approx 50\% / \sqrt{E(\text{GeV})} \oplus 3\%$
Muon:	$\sigma/p_T \approx 10\%$ at 1 TeV/c



Fourth Family (FF)

For further details see talk by S.Sultansoy in this conference

FF is the simplest "modification" to SM as we know it today

● predicts 4 new heavy leptons $m > 100\text{GeV}$

-quarks are to be searched for,

-quasi-degenerate u_4 , d_4 masses expected:



$$|m_{d4} - m_{u4}| < m_W/2$$

● heavy FF quark condensate may play the Higgs role and give mass to fermions

● heavy neutrinos may be the DM candidate we are looking for.

● EW precision data equally compatible with 3SM and 4SM cases.

FF is an economical solution to SM problems.



prospects for FF quarks

Search channels (No FCNC at tree level):




- $Q \rightarrow Wj$ depending on 4×4 CKM, j is from 1st, 2nd or 3rd family

- case w/ dominant 3rd family mixing was studied before

- u_4 easy, d_4 not so easy

- 10 year old work, can be improved.

- w/ 1st 2nd family mixing: recent study



- u_4/d_4 pair $\rightarrow WjWj$, light jets
- semi leptonic mode
- both q_4 reconstructed
- MLM matching not done
- pessimistic results
- background estimated from fit (crystal ball function)

discovery limits:


- 100 pb⁻¹ up to 450 GeV
- 1 fb⁻¹ up to 600 GeV

FF and Higgs



Enhancement to Higgs production from FF quarks

- from to gluon fusion loop $\sigma_{4th}(gg \rightarrow h) \approx 9 \sigma_{SM}(gg \rightarrow h)$

-indirect searches at Tevatron & LHC



LHC @ 14TeV




indirect searches limit the mass of Higgs and the mass of q_4 .
 -> DØ analysis being updated!

FF and Higgs ..

What about the Higgs decays to FF members ?

- In particular to ν_4




Sultansoy, Unel, hep-ph/0707.3266 & Turkish J. Phys. 31, (2007) 295-298

new search channel: $h \rightarrow \nu_4 \bar{\nu}_4$

“Silver & platinum modes” of Higgs search

The diagram shows a Higgs boson (h) decaying into a lepton (lep) and an anti-lepton (lep). The decay proceeds through a loop of ν_4 and $\bar{\nu}_4$ neutrinos. The ν_4 neutrino emits a W^- boson, and the $\bar{\nu}_4$ neutrino emits a W^+ boson, which then interact with the lepton and anti-lepton lines.

Cuhadar-Donszelmann, Karagoz Unel, Ozcan, Sultansoy, Unel, arXiv:0806.4003 [hep-ph], JHEP10(2008)074



Define 3 benchmark points with different mass values all in the $2\mu+4j$ final state $BR(\nu_4 \rightarrow \mu W) = 68\%$ (PRD72,2005, 053006)






	$\sigma_{pp \rightarrow Z \rightarrow \nu_4 \bar{\nu}_4}$ (fb)	m_h (GeV)	$\sigma_{gg \rightarrow h}$ (pb)	m_{ν_4} (GeV)	$BR(h \rightarrow \nu_4 \bar{\nu}_4)$	$\sigma_{pp \rightarrow \nu_4 \bar{\nu}_4 \rightarrow WW \mu \mu}$ (fb)
$S1$	782	N/A	N/A	100	N/A	362
$S2$	782	300	30	100	0.088	1583
$S3$	144	500	10	160	0.055	321

Higgs & v_4 .

PGS based study

Initially same, later on separate cuts for Dirac & Majorana cases

Backgrounds:
 $2V+2\mu, V=W, Z = 5\text{fb}$
 $Z/\gamma +4j = 57\text{pb}$
 $t\bar{t} = 755\text{pb}$






Selection cut	S1	S2	S3	$2\mu 4j$	$t\bar{t}$
At least 2μ	63.6	77.9	84.1	93.3	8.1
$p_T(\mu) > 15 \text{ GeV}$	50.7	55.1	95.1	88.8	29.5
At least $4j$	73.6	82.3	82.6	86.0	88.7
$p_T(j) > 15 \text{ GeV}$	53.3	65.6	72.2	70.4	76.0
$ M_{jj} - M_W < 20 \text{ GeV}$	63.1	60.5	60.3	45.9	52.8
$\Delta R_{\mu j} > 0.4$	64.5	65.9	77.4	83.0	17.4
No j_b	93.6	92.0	91.5	93.6	53.4
$M_{E_T} < 30 \text{ GeV}$	74.4	64.9	68.7	79.4	15.4
Common ϵ_{reco}	3.7	5.7	13.4	24.2	0.012



$$\Delta R = \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2}$$

Higgs & ν_4 ..


$$S = \sqrt{2 \times [(s + b) \ln(1 + \frac{s}{b}) - s]}$$




Majorana case shown



h search via ν_4 pair production



ν_4 search via pair production



Higgs

- 1 fb or less for Majorana
- about 2 fb for Dirac
- if no Higgs Dirac ν_4 can stay undetected



prospects for E6 inspired quarks

E6 GUT, initially suggested by F. Gürsey in 70s

- predicts one iso-singlet quark per family (assume D lightest)

$$\begin{pmatrix} u_L \\ d_L \end{pmatrix}, u_R, d_R, \underline{D_L, D_R} \quad \begin{pmatrix} c_L \\ s_L \end{pmatrix}, c_R, s_R, \underline{S_L, S_R} \quad \begin{pmatrix} t_L \\ b_L \end{pmatrix}, t_R, b_R, \underline{B_L, B_R}$$

$$L_D = \frac{\sqrt{4\pi\alpha_{em}}}{2\sqrt{2}\sin\theta_W} [\bar{u}^\theta \gamma_\alpha (1 - \gamma_5) d \cos\phi + \bar{u}^\theta \gamma_\alpha (1 - \gamma_5) D \sin\phi] W^\alpha$$

$$- \frac{\sqrt{4\pi\alpha_{em}}}{4\sin\theta_W} \left[\frac{\sin\phi \cos\phi}{\cos\theta_W} \bar{d} \gamma_\alpha (1 - \gamma_5) D \right] Z^\alpha$$

$$- \frac{\sqrt{4\pi\alpha_{em}}}{12\cos\theta_W \sin\theta_W} [\bar{D} \gamma_\alpha (4\sin^2\theta_W - 3\sin^2\phi(1 - \gamma_5)) D + \bar{d} \gamma_\alpha (4\sin^2\theta_W - 3\cos^2\phi(1 - \gamma_5)) d] Z^\alpha$$

Mehdiyev, Sultansoy, Unel, Yilmaz, EPJC 49 2007.

Higgs boson could also be in the game:

if $m_h < m_D$, $D \rightarrow h d$ is possible:

$$L_h^M = \frac{m_D}{\nu} \sin^2\phi_L \bar{D}^M D^M h$$

$$- \frac{\sin\phi_L \cos\phi_L}{2\nu} \bar{D}^M [(1 - \gamma^5) m_D + (1 + \gamma^5) m_d] d^M h$$

$$- \frac{\sin\phi_L \cos\phi_L}{2\nu} \bar{d}^M [(1 + \gamma^5) m_D + (1 - \gamma^5) m_d] D^M h$$

$$+ \frac{m_d}{\nu} \cos^2\phi_L \bar{d}^M d^M h$$


Branching Fractions:

- w/o Higgs : $D \rightarrow Wq$ (66%) $D \rightarrow Zq$ (33%)
- with Higgs: $D \rightarrow Wq$ (50%) $D \rightarrow Zq$ (25%) $D \rightarrow hq$ (25%)

Sultansoy, Unel, SN-ATLAS-2007-066, PLB, 669, 1,2008.








ATLAS prospects for E6 quarks



$D\bar{D} \rightarrow$	Final State	Expected Signal	Decay B.R.	Total B.R.
$Z Z d\bar{d}$ 0.33×0.33	$Z \rightarrow \bar{l}l \quad Z \rightarrow \bar{l}l$	$4l + 2 jet$	0.07×0.07	0.0005
	$Z \rightarrow \bar{l}l \quad Z \rightarrow \nu\nu$	$2l + 2 jet + P_T$	$2 \times 0.07 \times 0.2$	0.0030
	$Z \rightarrow \bar{l}l \quad Z \rightarrow q\bar{q}$	$2l + 4 jet$	$2 \times 0.07 \times 0.7$	0.0107
$Z W d u$ $2 \times 0.33 \times 0.67$	$Z \rightarrow \bar{l}l \quad W \rightarrow l\bar{\nu}$	$3l + 2 jet + P_T$	0.07×0.21	0.0065
	$Z \rightarrow \bar{l}l \quad W \rightarrow q\bar{q}$	$2l + 4 jet$	0.07×0.68	0.0211

Mehdiyev, Siodmok, Sultansoy, Ünel
SN-ATLAS-2007-067 EPJC54,2008

$m_D(\text{GeV})$	600	800	1000	
$4l + 2j$	Signal	16	3.7	0.74
	background	3.0	1.3	0.4
	$-\ln p$	21.47	4.78	1.44
$2l + 2j + \cancel{E}_T$	Signal	53	19	4
	background	12	13	5
	$-\ln p$	120	15.81	3.32
$3l + 2j + \cancel{E}_T$	signal	97	18.3	1.8
	background	24.9	9.0	1.05
	$-\ln p$	191.4	20.66	1.69
$2l + 4j$	signal	133	18	2.5
	background	9	3	2.8
	$-\ln p$	983	25.3	2.44
$-\Sigma \ln p$	1315.9	66.5	8.9	
Combined Significance (σ)	51.3	11.3	3.8	



Combination of channels allows discovery in early data taking. 100 fb^{-1} is enough to scan up to 1TeV

Mass (Z(l) jet), GeV Mass (W(lν) jet), GeV


E6 and the Higgs

DD→hj Wj channel studied for a light Higgs + m_D scan


- other channels:
- h→bb possible

D_1	D_2	BR	#expected Higgs/100fb ⁻¹	expected final state
$D \rightarrow h j$	$D \rightarrow h j$	0.029 (0.053)	0.58×10^6 (2.65×10^4)	$2j 4j_b$
$D \rightarrow h j$	$D \rightarrow Z j$	0.092 (0.120)	0.92×10^6 (3.01×10^4)	$2j 2j_b 2l$
$D \rightarrow h j$	$D \rightarrow W j$	0.190 (0.235)	1.9×10^6 (6.04×10^4)	$2j 2j_b l E_{T,miss}$

Background: SM $W^\pm b b j j = 520$ pb



Both h and D reconstruction possible



- The hjWj mode only allows 5σ
 - up to 350 GeV D quark with 1fb⁻¹
 - also a 120 GeV Higgs with 2fb⁻¹
- If m_D < 630 GeV, hjWj channel becomes as promising as H→γγ . (i.e. 8σ in 100 fb⁻¹)




Conclusions

- There are several BSM candidates predicting additional quarks.
- We saw the ATLAS discovery prospects for FF and E6 models:
 - FF model
 - quarks up to 450 GeV with 100 pb⁻¹ if 1st or 2nd & 4th generations mix
 - neutrinos up to 160 GeV with 1fb⁻¹ if Higgs is ≥ 300 GeV
 - E6 model
 - If $m_D < 500$ GeV, ATLAS could discover with 1fb⁻¹ by combining different channels.
 - With 100 fb⁻¹, the observation reach increases to $m_D = 1000$ GeV.
- we saw the potential boost to Higgs searches:
 - FF model
 - $\sigma_{4th}(gg \rightarrow h) \approx 9 \sigma_{SM}(gg \rightarrow h) \Rightarrow$ golden mode reachable w/ 1fb⁻¹ for all h masses.
 - $300 < m_h < 500$ GeV w/ 1fb⁻¹ from v_4 (Dirac & Majorana identification possible)
 - E6 model
 - if $m_h \approx 120$ GeV & $m_D < 630$ GeV, $h_j W_j$ channel becomes as promising as $h \rightarrow \gamma\gamma$
- Watch the “ATLAS” channel, interesting things are about to happen.



thank you..



backup



About CDF hint

Too many events, how could this be?

- Little Higgs model predicts 1 T quark, heavy iso-singlet up type

- similar to the model given in CDF paper
- not enough cross section



- E6-GUT predicts 3 down type, heavy iso-singlet quarks (D, S, B)

- Fourth Family model: u_4, d_4 quarks

- $\Delta m(u_4, d_4) \leq m_W/2$

- A susy model with more quarks?



Only multiple degenerate quarks or anomalous production could explain the number of "signal" events.



Light Higgs & E6

$m_h = 120 \dots 135 \text{ GeV}$

-if $\sin\phi$ becomes 10 times smaller, total cross section increases by few percent.



at $m_H = 120 \text{ GeV}$, best significance is from $H \rightarrow \gamma\gamma$ channel, 8σ at 100 fb^{-1}