

Exclusive High Mass Di-Leptons in CDF

Michael Albrow

Fermi National Accelerator Laboratory, Batavia, IL60510, USA

In the Collider Detector at Fermilab, CDF, we have measured central exclusive production, $p + \bar{p} \rightarrow p + X + \bar{p}$, where X is a pair of leptons or photons and nothing else. In this talk I focus on central masses $M(X) > 8 \text{ GeV}/c^2$. We measured QED production $\gamma\gamma \rightarrow e^+e^-, \mu^+\mu^-$ up to $M(X) = 75 \text{ GeV}/c^2$, and candidates for photoproduction of Upsilon's, $\gamma\mathbb{P} \rightarrow Y(1S), Y(2S), Y(3S)$. I report a search for exclusive photoproduction of Z -bosons, and the status of searches for exclusive two-photons: $p + \bar{p} \rightarrow p + \gamma\gamma + \bar{p}$. These measurements constrain the cross section $\sigma(p + p \rightarrow p + H + p)$ at the LHC.

1 Introduction

By central exclusive production CEP (also called central exclusive diffraction, CED) at the Fermilab Tevatron we mean reactions $p + \bar{p} \rightarrow p + X + \bar{p}$, where X is a simple system fully measured, and “+” are large rapidity gaps ($\Delta y \gtrsim 5$ units) with *no* particles. In the Collider Detector at Fermilab, CDF [1], we cannot detect the forward p or, except for some large $M(X)$ events, the \bar{p} . However we installed scintillation counters (beam shower counters, BSC) along the beam pipe, which detected showers from particles out to pseudorapidity $|\eta| = 7.4$. Requiring them to be empty selects events in which the p and \bar{p} did not fragment and went down the beam pipe.

In the Standard Model the only significant coherent t -channel exchanges over such large rapidity gaps are colour singlets with charge $Q = 0$ and spin J , or effective spin $\alpha(t = 0), \geq 1$. These are the photon, γ , pomeron \mathbb{P} ($C = +1$) and odderon O ($C = -1$). Z -boson exchange would be allowed, but the proton would inevitably break up. The odderon has not yet been convincingly observed; in our observation of exclusive J/ψ and $\psi(2S)$ reported in this meeting by Pinfeld [2, 3] we placed a new limit. In this talk I will report on CDF measurements of exclusive lepton pairs, $p + \bar{p} \rightarrow p + l^+l^- + \bar{p}$ above the charmonium region, see Fig. 1. This includes non-resonant (QED) $\gamma\gamma \rightarrow e^+e^-, \mu^+\mu^-$, and photoproduction: $\gamma\mathbb{P} \rightarrow \Upsilon, Z$. We can also search for exclusive $\mathbb{P}\mathbb{P} \rightarrow \chi_b \rightarrow \Upsilon + \gamma$, but the cross section is very small and its observation probably requires no additional collisions (no pile-up). This process, like exclusive χ_c , is a good test of exclusive Higgs boson production, as the QCD part of the Feynman diagrams is identical. While our CDF studies provide good tests of QCD with large rapidity gaps, and hard pomeron behaviour, they are also precursors to processes with $M(X) \gtrsim 100 \text{ GeV}/c^2$ at the LHC [4], where X can be $h, H, W^+W^-, \tilde{l}^+\tilde{l}^-$ etc., including any exotic particles that couple to gluons or photons and have the right quantum numbers. If states such as Higgs bosons are seen this way at the LHC, their mass, width, spin, C-parity and coupling $\Gamma(Xgg)$ can be determined in unique ways. Even a pair of nearby states, e.g. MSSM $h(140) \rightarrow b\bar{b}$ and $H(150) \rightarrow b\bar{b}$ can be resolved, which is impossible by other means.

2 CDF Detectors

The central detector of CDF [1] has layers of silicon tracking and drift chambers surrounded by a time-of-flight scintillation counter barrel, in a solenoidal field. This is surrounded by electromagnetic and hadronic calorimeters, muon scintillators and tracking chambers. The forward region, $\theta < 3^\circ$, has, on each side, a 48-channel Cherenkov luminosity counter hodoscope, a “miniplug” calorimeter, a set of beam shower counters, BSC, and on the outgoing \bar{p} side, scintillating fiber trackers in Roman pots. The BSC were very important as rapidity gap detectors in no-pile-up events, and for triggering on exclusive events. They are relatively simple scintillation counters around the beam pipes covering $5.5 < |\eta| < 7.4$. Only BSC-1 sees primary particles, and it has two radiation lengths in front to convert photons; the others see showers created in the beam pipe. We have proposed them for CMS [5]; all LHC experiments should have them! Another recommendation to all experiments is to record zero-bias, or bunch-crossing, triggers routinely, e.g. at 1 Hz. These were essential in our CDF exclusive studies. We divided those events into two classes: [a] = probably no interaction, e.g. no tracks, and [b] at least one interaction, with tracks from the beam line. Then for each subdetector, e.g. BSC-1 which had 8 PMTs, we plotted the “hottest” PMT as $\text{Log}(\max \text{ADC counts in BSC-1})$ for events in classes [a] and [b]. One can also plot the sum $\sum_i (\text{ADC}_i)$. Repeating for all subdetectors, one finds cuts that select events with all the CDF detectors empty, except for the state X .




Figure 1: Exclusive two-photon production of lepton pairs, and exclusive photoproduction.

3 CDF Exclusive Physics Programme

In March 2001 some of us proposed [6] to add very forward proton tracking detectors to CDF to look for exclusive Higgs boson production. At that time some theorists (cited in Ref. [6]) had suggested that the cross section could be as high as 10-100 fb, and a signal could be seen using the missing mass technique [7]. The Durham group prediction [8] was a factor $\gtrsim 10^2$ lower, at 0.06 fb (for $M(H) = 120$ GeV), impossibly low for the Tevatron. Incidentally in Ref. [6] we suggested for the first time that exclusive $\gamma\gamma$ production is a good test of the theory, and that troublesome pile-up can be highly suppressed with “fast timing Cherenkov counters”. Given the very large theoretical uncertainty on $\sigma(H)$, the proposal could not proceed at that time, but

we embarked on a program to measure related exclusive processes with cross sections accessible even in the Durham model. We have now measured $p + X + \bar{p}$ final states with $X = \gamma\gamma, \chi_c$ and dijets JJ , all of which have very similar QCD diagrams and issues, and all three are consistent with the Durham group predictions (which have a factor “a few” uncertainty). One can now be certain that it is possible to produce a Higgs boson (if it exists) at the LHC with no other particles, and reasonably confident that their prediction $\sigma(SMH(120)) \sim 1 - 10$ fb at $\sqrt{s} = 14$ TeV is not unrealistic. The cross section can be much higher in some Beyond SM scenarios.

The exclusive processes measured in CDF have different strengths and weaknesses. $X = \gamma\gamma$ is the cleanest as the photons, like the Higgs, have no strong interactions. Looking at the Feynman diagrams through a “QCD-only filter” they are identical; simply the q -loop (mostly u and c) becomes a t -loop and $\gamma\gamma$ is replaced by H . But the cross section is very small. The Durham prediction [9] is 36 fb for $|\eta(\gamma)| < 1$ and $E_T(\gamma) > 5$ GeV. This corresponds to $0.8_{-0.5}^{+1.6}$ events in the CDF search [10]; 3 candidates were found, of which 2 were “perfect” with two single narrow electromagnetic showers, while the third had some characteristics of $\pi^0\pi^0$ (broader showers). Note that the two golden candidates were distilled from 10^{12} inelastic collisions; such things can be done! $X = \chi_c$ is also clean and has a much bigger cross section: $\frac{d\sigma}{dy}(y = 0) = 76 \pm 10(stat) \pm 10(syst)$ nb (CDF [2, 3]), compared with 90 nb (with a large uncertainty) predicted by Durham [11]. The weakness is that the charm mass is small so the process is not very perturbative, the χ_c is colourless but it is still a hadron with final state interactions, and resolving $\chi_{c0}(1P), \chi_{c1}(1P)$, and $\chi_{c2}(1P)$ is difficult. The χ_b is more perturbative than the χ_c , so the theory is under better control, but the cross section is expected to be only about 1/500th that of the χ_c and the decay modes are not well known. For the χ_{b0} the Particle Data Group gives only $\chi_{b0}(1P) \rightarrow \gamma Y(1S) < 6\%$, with no other modes well known, so it will not be possible to give a cross section soon. (Some hadronic decays with poorly known branching fractions are impossible to trigger on in a normal collider environment.) Finally CDF also measured [12] exclusive di-jets, produced by the process $gg \rightarrow gg$ with a colour-cancelling gluon exchange. The cross section is quite large, ≈ 100 pb for two jets with $E_T > 15$ GeV and $|\eta| < 2.5$, with the dijet having $> 80\%$ of the total central mass, i.e. $R_{JJ} > 0.8$. In this region there is an excess of dijets compared with inclusive dijet expectations. The data, out to Jet $E_T^{min} = 35$ GeV, agree within a factor ~ 3 with the Durham prediction [13], but are more than an order of magnitude lower than the DPEMC prediction [14].

The main particle states that can be produced exclusively in double pomeron exchange, DPE , are $\sigma(600), f_0(980), \chi_c, \chi_b$ and H . The first two were observed at the ISR [15] and, less cleanly, at the SPS (fixed target), and the χ_c is now established [2, 3]. The other accessible states are $\gamma\gamma$ and JJ . It will be very hard to get a good ($\lesssim 25\%$) measurement of the χ_b , but H appears to be in reach at the LHC, provided the two protons can be measured. (Double proton tagging for $p + \chi_b + p$ at high enough luminosity to get some events is probably not feasible, due to its low mass.)

In CDF we observed $\gamma\gamma \rightarrow e^+e^-$ collisions with $E_T(e) > 5$ GeV for the first time in hadron-hadron collisions. We published [16] 16 events (background ~ 1.9) with $M(e^+e^-) > 10$ GeV/ c^2 in excellent agreement with the LPAIR Monte Carlo. The highest mass event was at 38 GeV/ c^2 , and all pairs are very back-to-back with $(180^\circ - \Delta\phi) < 2.4^\circ$. We followed that with a search [17] for exclusive Z , in the process finding eight e^+e^- or $\mu^+\mu^-$ events with $M(l^+l^-)$ from 40 to 75 GeV/ c^2 . Again “QED rules”, and the lepton pairs all have $(180^\circ - \Delta\phi) < 0.75^\circ$. This mass reach is as high as (or higher than) e^+e^- at LEP (which has no strongly interacting background) and ep at HERA. This demonstrates that exclusive dileptons can be extracted from the huge backgrounds in hadron-hadron collisions, which is good news as they provide

an excellent (probably the best) calibration of the momentum scale and resolution of the high precision proton spectrometers being planned [4] for ATLAS and CMS. One does not need to see both protons to calibrate the spectrometers, as each one is very well known (ultimately limited by the incoming beam momentum spread, $\frac{dp}{p} \sim 10^{-4}$). The use of exclusive QED dileptons, with precisely known cross section, to measure the machine luminosity (integrated over a period, perhaps days), has been suggested. Unfortunately the precision is likely to be limited by unseen proton dissociation, and knowledge of efficiencies, acceptance and non-exclusive background. (One cannot require no pile-up, as the result would then depend on σ_{inel} which is *a priori* unknown).

Exclusive Z production is allowed in the Standard Model: a radiated virtual photon fluctuates to a $q\bar{q}$ pair, which scatters by hard pomeron exchange on the other proton, followed by $q\bar{q} \rightarrow Z$, as in exclusive vector meson photoproduction, see Fig. 1. However the SM cross section is much too small at the Tevatron: 0.3 fb [18] or 0.21 fb [19]. At the LHC (14 TeV) the predictions are 13 fb [18] and 69 fb [19], which may make an observation possible. A signal at the Tevatron, or a significantly higher cross section at the LHC, would be evidence for new particles with strong and electroweak couplings. White's theory [20] of the supercritical pomeron predicts colour-sextet quarks coupling strongly to the pomeron and to the W and Z , and he expects a much enhanced cross section, but without a quantitative prediction. We used a sample of 3.17×10^5 lepton pairs with $M(l^+l^-) > 40 \text{ GeV}/c^2$, of which 1.83×10^5 were in the Z peak. We required exclusivity over the full range $-7.4 < |\eta| < +7.4$, finding 8 events, agreeing with QED expectations. Fig. 2 shows the mass and azimuthal difference distributions of these events. All events were very back-to-back in the transverse plane, with $(\pi - \Delta\phi) < 0.75^\circ$. One event with $M(\mu^+\mu^-) = 66 \text{ GeV}/c^2$ had a \bar{p} track in the Roman pots; for the others the \bar{p} was out of their acceptance or they were not operational. None of the eight exclusive events were Z -candidates, and a limit was placed: $\sigma_{excl}(Z) < 0.96 \text{ pb}$ at 95% C.L.¹ A nice check of the exclusivity analysis comes from $W \rightarrow l^\pm\nu$ events, which cannot be exclusive, but otherwise are very similar to Z , and which are more abundant. It may be possible to improve this limit using a factor 2-3 more data and including pile-up events, using the requirements of no associated tracks on the l^+l^- vertex, small $(\pi - \Delta\phi)$ and $p_T(l^+l^-)$. We are testing this method on the Upsilon region, with $8 < M(l^+l^-) < 40 \text{ GeV}/c^2$. The QED is a good control, and the photoproduced Y states have cross sections that are within reach, although not very well known. (The HERA data [21] do not resolve clearly the $Y(1S)$, $Y(2S)$ and $Y(3S)$ states and have quite large uncertainties.) Predictions for $\frac{d\sigma}{dy}(Y(1S), y=0)$ are [18, 22, 23, 24] are around 5 – 14 pb. Applying the branching fraction to $\mu^+\mu^-$ or e^+e^- , $B = 0.025$, would give a few hundred events in 2 fb^{-1} (\times the acceptance and efficiency). HERA has provided a nice compilation [25] of exclusive cross sections for vector mesons from ρ to $Y(1S)$ vs. $W = \sqrt{s(\gamma p)}$. In CDF with $y = 0$ we have $W(J/\psi) \sim 80 \text{ GeV}$ and $W(Y(1)) \sim 136 \text{ GeV}$. At HERA the ratio of these cross sections is ~ 300 . We are studying this region both in $\mu^+\mu^-$ and e^+e^- events.

For the dimuons we used a trigger with two muons with $p_T(\mu) > 4 \text{ GeV}/c$ and $|\eta| < 0.6$. The inclusive $M(\mu^+\mu^-)$ spectrum shows the states $Y(1S)$, $Y(2S)$ and $Y(3S)$ as well-separated peaks, see e.g. Ref. [26]; the mass resolution is only $\sigma(M) \approx 50 \text{ MeV}/c^2$, less than the mass differences. Separation is important in order to measure exclusive $\chi_b \rightarrow \Upsilon + \gamma$, which feed differently the three states. Inclusively the ratio $\Upsilon(1S)$:continuum is about 8:1. Requiring

¹Intriguingly, an event with $M(e^+e^-) = 92 \text{ GeV}/c^2$, and $(\pi - \Delta\phi) = 1.25^\circ$ (larger than expected for QED) was rejected; it failed the exclusivity requirement only in the BSC counters on one arm. It has the characteristics of a photoproduced Z but with a proton dissociation. However with only one event no claim can be made.




Figure 2: Mass distribution of exclusive lepton pairs and azimuthal angle difference (from 180°), data and normalised LPAIR prediction.

no other tracks on the $\mu^+\mu^-$ vertex, $\pi - \Delta\phi < 0.1$ rad and $p_T(\mu^+\mu^-) < 1.5$ GeV/c retains both Υ and continuum events, but we cannot yet claim that these events are exclusive. These cuts should be efficient for the QED events and for most of the Y 's; the issue is non-exclusive backgrounds in the presence of other interactions. We are now studying the continuum to see if it is exactly as expected for QED, as a control, and can then give Y -photoproduction cross sections, depending on unknown backgrounds from $\chi_b \rightarrow Y + \gamma$. Unfortunately neither the χ_b production rates nor their radiative decays are known. The $p_T(Y)$ distribution is broader for χ_b -daughters, which may help, but one would like to reconstruct the photons, which is probably not possible with pile-up. The same issues will confront us at the LHC.

For the $Y \rightarrow e^+e^-$ channel we chose to veto pile-up, and used a trigger requiring two electromagnetic showers with $E_T > 2$ GeV and $|\eta| < 2$ m, and forward gaps (BSC-1 empty). Our prime motivation for this trigger is to search for additional exclusive $\gamma\gamma$ candidates, and hopefully to make a definitive observation. Compared with our earlier search, we lowered the trigger threshold from 4 GeV to 2 GeV, took more data, will expand the η -coverage and use better background (π^0) rejection. Analysis is underway. The same trigger collects QED e^+e^- pairs with $M(e^+e^-) > 8$ GeV/ c^2 , and $Y \rightarrow e^+e^-$ decays. The QED data is a good control of our exclusivity cuts, and the Y candidates can be compared with our $Y \rightarrow \mu^+\mu^-$ candidates. As these events are without pile-up, we may have some $Y + \gamma$ candidates attributable to χ_b (but not many are expected). The decay photon tends to be soft (namely 391(442) MeV in the Y -frame) for the $\chi_{b0(b2)}$, and is poorly measured. Other exclusive final states are probably not useful, mainly because of trigger limitations.

Can exclusive Y photoproduction be seen with a forward proton tag at 420 m at the LHC? We have $(1 - x_F)_{1,2} = \frac{1}{\sqrt{s}} \sum_{leptons} p_T e^{+(-)\eta}$ ($1 - x_F$ is the fractional momentum loss of the proton), and the most likely kinematics are $p_{T,1} = p_{T,2} \approx 4.75$ GeV/c, $\eta_1 \approx \eta_2$. To have one proton with $1 - x_F$ as large as (say) 0.01 we need a muon pair with $\eta \sim 2.5$, at the limit of the muon coverage in CMS. A more serious problem with using $p + Y(\rightarrow \mu^+\mu^-) + p$ for proton calibration is that many Υ s, perhaps even most, will be decay products of χ_b states and the proton momenta are then not known. For these reasons the QED $\gamma\gamma \rightarrow \mu^+\mu^-$ with

$M(\mu^+\mu^-) \gtrsim 10 \text{ GeV}/c^2$ will probably be the calibration channel. Electron pairs are less favourable due to final state radiation and bremsstrahlung. With $|\eta_{max}| = 2$ and $M_{min} = 20 \text{ GeV}/c^2$ the cross section is 1.6 pb. One proton will usually have much too small ξ , but fortunately both protons are known and one can be used to calibrate the spectrometers.

In conclusion, five years ago the predictions for exclusive Higgs production had more than two orders of magnitude spread. Since then in CDF we have measured three related processes, exclusive $JJ, \gamma\gamma$ and χ_c , all consistent with the predictions of the Durham group within the quoted factor of “a few”. Especially the χ_c observation means that exclusive Higgs production must happen, if there is a Higgs boson. Our observation of exclusive photoproduced vector mesons demonstrates that exclusive Z photoproduction must be possible, albeit with a small cross section (in the Standard Model) at the LHC. Our observations of exclusive $\gamma\gamma \rightarrow e^+e^-, \mu^+\mu^-$ (with a forward proton detected) are encouraging for forward spectrometer calibrations, and mean that $\gamma\gamma \rightarrow W^+W^-$ (and l^+l^- if sleptons exist) could be measured. This bodes well for a rich physics program with high precision forward spectrometers at the LHC.

Acknowledgements

I acknowledge support from the U.S. Dept. of Energy through Fermi National Accelerator Laboratory.

References

- [1] D. Acosta et al. (CDF Collaboration), Phys. Rev. D **71**, 032001 (2005) and references therein.
- [2] T. Aaltonen et al. (CDF Collaboration), *Observation of exclusive charmonium and $\gamma\gamma \rightarrow \mu^+\mu^-$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96 \text{ TeV}$* , Phys. Rev. Lett. **102**, (2009) 242001.
- [3] J. Pinfeld, *Exclusive charmonium production in CDF*, talk at this conference.
- [4] M.G. Albrow et al. (FP420 R&D Collaboration), *Higgs and new physics with forward protons at the LHC*, arXiv:0806.0302[hep-ex], to be published in J.Inst.
- [5] M.G. Albrow et al., *Forward Shower Counters (FSC) in CMS, a proposal*, CMS Internal (Dec. 2007); M.G. Albrow et al., *Forward physics with rapidity gaps at the LHC*, arXiv:0811.0120, submitted to J. Inst.
- [6] M.G. Albrow et al., *A search for the Higgs boson using very forward tracking detectors with CDF* (2001), hep-ex/0511057.
- [7] M.G. Albrow and A. Rostovtsev, *Searching for the Higgs boson at hadron colliders using the missing mass method*, hep-ph/0009336.
- [8] V.A. Khoze, A.D. Martin and M.G. Ryskin, *Can the Higgs be seen in rapidity gap events at the Tevatron or the LHC?*, Eur.Phys J **C14**, 525 (2000).
- [9] V.A. Khoze, A.D. Martin, M.G. Ryskin and W.J. Stirling, *Diffraction $\gamma\gamma$ production at hadron colliders*, Eur. Phys. J. **C 38**, 475 (2005).
- [10] A. Abulencia et al. (CDF Collaboration), *Search for exclusive $\gamma\gamma$ production in hadron-hadron collisions*, Phys. Rev. Lett. **99**, (2007) 242002.
- [11] V.A. Khoze, A.D. Martin, M.G. Ryskin and W.J. Stirling, *Double diffractive χ meson production at hadron colliders*, Eur. Phys. J. **C 35**, 211 (2004).
- [12] T. Aaltonen et al.(CDF Collaboration), *Observation of exclusive dijet production at the Fermilab Tevatron $p\bar{p}$ collider*, Phys. Rev. D **77**, 052004 (2008).
- [13] J. Monk and A. Pilkington (EXHUME), Comput. Phys. Commun. **175**, 232 (2005).
- [14] M. Boonekamp and T. Kucs (DPEMC), Comput. Phys. Commun. **167**, 217 (2005).

- [15] T. Åkesson et al. (AFS Collaboration), *A search for glueballs and a study of double pomeron exchange at the CERN ISR*, Nucl. Phys. B **264**, 154 (1986).
- [16] A. Abulencia et al. (CDF Collaboration), *Observation of exclusive electron-positron production in hadron-hadron collisions*, Phys. Rev. Lett. **98**, (2007) 112001.
- [17] T. Aaltonen et al. (CDF Collaboration), *Search for exclusive Z-boson production and observation of high-mass $p\bar{p} \rightarrow p\gamma\gamma\bar{p} \rightarrow pl^+l^-\bar{p}$ events in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV*, Phys. Rev. Lett. **102**, (2009) 222002.
- [18] L. Motyka and G. Watt, *Exclusive photoproduction at the Fermilab Tevatron and CERN LHC within the dipole picture*, Phys. Rev. D **78**, 014023 (2008).
- [19] V.P. Goncalves and M.V.T. Machado, *Diffractive photoproduction of Z bosons in coherent interactions at the CERN LHC*, Eur. Phys. J. C **56**, 33 (2008); Erratum-ibid. C **61**:351 (2009).
- [20] A.R. White, *Physics of a sextet quark sector*, Phys. Rev. D **72**, 036007 (2005).
- [21] ZEUS Collaboration, S. Chekanov et al., *Exclusive photoproduction of Υ mesons at HERA*, arXiv:0903.4205 (2009).
- [22] S. Klein and J. Nystrand, *Photoproduction of J/ψ and Υ in pp and $p\bar{p}$ collisions*, hep-ph/0310223.
- [23] A. Bzdak, L. Motyka, L. Szymanowski and J.R. Cudell, *Exclusive J/ψ and Υ hadroproduction and the QCD odderon*, Phys. Rev. D **75**, 094023 (2007).
- [24] A. Rybarska, W. Schafer, and A. Szczurek, *Exclusive photoproduction of Υ : from HERA to Tevatron*, Int. J. Mod. Phys. A **24**, 557 (2009); Phys. Lett. B **668**, 126 (2008).
- [25] See e.g. M. Klein and R. Yoshida, *Collider physics at HERA*, Prog.Part.Nucl.Phys. **61**, 343 (2008); arXiv:0805.3334.
- [26] D. Acosta et al. (CDF Collaboration), *Υ production and polarization in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV*, Phys. Rev. Lett. **88**, 161802 (2002).