

MATHEMATICAL MODELLING AND ANALYSIS  
Volume 15 Number 2, 2010, pages 265–274  
Doi:10.3846/1392-6292.2010.15.265-274  
© Vilnius Gediminas Technical University, 2010

ISSN 1392-6292 print  
ISSN 1648-3510 online  
[www.vgtu.lt/mma/](http://www.vgtu.lt/mma/)

# A Strong Convergence of a Modified Krasnoselskii-Mann Method for Non-Expansive Mappings in Hilbert Spaces\*

Yonghong Yao<sup>1</sup> and Yeal Je Cho<sup>2</sup>

<sup>1</sup>*Department of Mathematics, Tianjin Polytechnic University  
Tianjin 300160, China*

E-mail: [yaoyonghong@yahoo.cn](mailto:yaoyonghong@yahoo.cn)

<sup>2</sup>*Department of Mathematics Education and the RINS  
Gyeongsang National University*

Chinju 660-701, Republic of Korea

E-mail(*corresp.*): [yjcho@gnu.ac.kr](mailto:yjcho@gnu.ac.kr)

Received June 16, 2009; revised October 08, 2009; published online April 20, 2010

**Abstract.** In this paper, we introduce a new method based on the well-known Krasnoselskii-Mann's method for non-expansive mappings in Hilbert spaces. We show that the proposed method has strong convergence for non-expansive mappings.

**Keywords:** non-expansive mapping, fixed point, modified Krasnoselskii-Mann's method, strong convergence, Hilbert space.

**AMS Subject Classification:** 47H05; 47H10.

## 1 Introduction

Let  $H$  be a real Hilbert space. Recall that a mapping  $T : H \rightarrow H$  is said to be non-expansive if

$$\|Tx - Ty\| \leq \|x - y\|, \quad \forall x, y \in H.$$

A point  $x \in H$  is a fixed point of  $T$  provided  $Tx = x$ . Let's denote the set of fixed points of  $T$  by  $F(T)$ , that is,  $F(T) = \{x \in H : Tx = x\}$ . Assume, throughout this paper, that  $F(T) \neq \emptyset$ .

Construction of fixed points of non-expansive mappings is an important subject in the theory of non-expansive mappings and its applications in a number of applied areas, in particular, image recovery and signal processing (see [1]–[28]). It is well-known that Picard's iteration

$$x_{n+1} = Tx_n = \dots = T^{n+1}x$$

---

\* The second author was supported by the Korea Research Foundation Grant funded by the Korean Government (KRF-2008-313-C00050).

of the mapping  $T$  at a point  $x \in H$  may not behave well in general. This means that it may not converge even in the weak topology. One way to overcome this difficulty is to use Krasnoselskii-Mann's method that produces a sequence  $\{x_n\}$  via the recursive manner:

$$x_{n+1} = \alpha_n x_n + (1 - \alpha_n) T x_n, \quad \forall n \geq 0. \quad (1.1)$$

This Krasnoselskii-Mann's method is remarkably useful for finding fixed points of a non-expansive mapping and provides a unified framework for some kinds of algorithms from various different fields. In this respect, the following result is basic and important.

**Theorem 1.** ([19]) *Let  $T$  be a non-expansive mapping on  $H$ . Then the sequence  $\{x_n\}$  defined by the iterative method (1.1) converges weakly to a fixed point of  $T$  provided  $\alpha_n \in [0, 1]$  and  $\sum_{n=0}^{\infty} \alpha_n(1 - \alpha_n) = +\infty$  whenever such fixed points exist.*

However, as in Theorem 1, Krasnoselskii-Mann's method for non-expansive mappings has only weak convergence. Thus a natural question rises: could we obtain a strong convergence result by using the well-known Krasnoselskii-Mann method for non-expansive mappings? In this connection, in 1975, Genel and Lindenstrass [7] gave a counterexample. Hence the modification is necessary in order to guarantee the strong convergence of Krasnoselskii-Mann's method.

Some attempts to construct iteration algorithm so that strong convergence is guaranteed have recently been made. For a sequence  $\{\alpha_n\}$  of real numbers in  $[0, 1]$  and fixed  $u \in C$ , let the sequence  $\{x_n\}$  in  $C$  be iteratively defined by  $x_0 \in C$ ,

$$x_{n+1} = \alpha_n u + (1 - \alpha_n) T x_n, \quad \forall n \geq 0. \quad (1.2)$$

The iterative method is now referred to as the Halpern iterative algorithm. The interest and importance of the Halpern iterative algorithm lie in the fact that strong convergence of the sequence  $\{x_n\}$  is achieved under certain mild conditions on parameter  $\{\alpha_n\}$  in a general Banach space. We recall some relevant important results as follows. In 1977, Lions [11] proved the strong convergence of  $\{x_n\}$  generated by (1.2) to a fixed point of  $T$ , where the real sequence  $\{\alpha_n\}$  satisfies the following conditions:

$$(C1) \quad \lim_{n \rightarrow \infty} \alpha_n = 0, \quad (C2) \quad \sum_{n=0}^{\infty} \alpha_n = \infty, \quad (C3) \quad \lim_{n \rightarrow \infty} \frac{\alpha_n - \alpha_{n-1}}{\alpha_n^2} = 0.$$

Based on this result, many authors considered the strong convergence of the Halpern algorithm under some restrictions on the parameters  $\{\alpha_n\}$ . For related works, see, e.g., [21, 25] and the references therein. Recently, Kim and Xu [9] proposed the following simpler modification of Mann iteration method:

Let  $C$  be a closed convex subset of a Banach space and  $T : C \rightarrow C$  a nonexpansive mapping such that  $Fix(T) \neq \emptyset$ . Define  $\{x_n\}$  in the following

way:

$$\begin{cases} x_0 = x \in C, \\ y_n = \alpha_n x_n + (1 - \alpha_n)Tx_n, \\ x_{n+1} = \beta_n u + (1 - \beta_n)y_n, \end{cases} \quad \forall n \geq 0, \tag{1.3}$$

where  $u \in C$  is an arbitrary (but fixed) element in  $C$ , and  $\{\alpha_n\}$  and  $\{\beta_n\}$  are two sequences in  $(0, 1)$ .

*Remark 1.* The modified Mann iteration scheme (1.3) is a convex combination of a fixed point in  $C$  and the Mann iteration method (1.1). There is no additional projection involved in iteration scheme (1.3).

A strong convergence of iteration scheme (1.3) is proved in [9].

**Theorem 2.** *Let  $C$  be a closed convex subset of a uniformly smooth Banach space  $X$  and let  $T : C \rightarrow C$  be a nonexpansive mapping such that  $Fix(T) \neq \emptyset$ . Given a point  $u \in C$  and given sequences  $\{\alpha_n\}_{n=0}^\infty$  and  $\{\beta_n\}_{n=0}^\infty$  in  $(0, 1)$ , the following conditions are satisfied:*

- (a)  $\beta_n \rightarrow 0, \sum_{n=0}^\infty \beta_n = \infty$  and  $\sum_{n=0}^\infty |\beta_{n+1} - \beta_n| < \infty$ ;
- (b)  $\alpha_n \rightarrow 0, \sum_{n=0}^\infty \alpha_n = \infty$  and  $\sum_{n=0}^\infty |\alpha_{n+1} - \alpha_n| < \infty$ .

*Then  $\{x_n\}_{n=0}^\infty$  defined by (1.2) strongly converges to a fixed point of  $T$ .*

Motivated by the result of Kim and Xu [9], Yao, Chen and Yao [23] introduced a modified version of the algorithm (1.3) with the viscosity method and proved the strong convergence of the proposed algorithm under some mild assumptions on the parameters.

## 2 Preliminaries

In the sequel, we use the notation  $\rightharpoonup$  for weak convergence and  $\rightarrow$  for strong convergence. We will need some lemmas to prove our main results.

**Lemma 1.** ([8]) *Let  $H$  be a real Hilbert space. Let  $T : H \rightarrow H$  be a nonexpansive mapping. Then  $I - T$  is demi-closed at 0, i.e., if  $x_n \rightharpoonup x \in H$  and  $x_n - Tx_n \rightarrow 0$ , then  $x = Tx$ .*

**Lemma 2.** ([21]) *Assume that  $\{a_n\}$  is a sequence of nonnegative real numbers such that  $a_{n+1} \leq (1 - \gamma_n)a_n + \gamma_n \delta_n$  for all  $n \geq 0$ , where  $\{\gamma_n\}$  is a sequence in  $(0, 1)$  and  $\{\delta_n\}$  is a sequence in  $R$  such that*

- (a)  $\sum_{n=0}^\infty \gamma_n = \infty,$  (b)  $\limsup_{n \rightarrow \infty} \delta_n \leq 0$  or  $\sum_{n=0}^\infty |\delta_n \gamma_n| < \infty.$  Then  $\lim_{n \rightarrow \infty} a_n = 0.$

### 3 Main Results

Let  $H$  be a real Hilbert space and  $T : H \rightarrow H$  be a non-expansive mapping. Let  $\lambda$  be a constant in  $(0, 1)$ . For each  $t \in (0, 1)$ , define a mapping  $T_t : H \rightarrow H$  by

$$T_t x = t(\lambda x) + (1 - t)Tx, \quad \forall x \in H.$$

For any  $x, y \in H$ , we have

$$\|T_t x - T_t y\| = \|[t(\lambda x) + (1 - t)Tx] - [t(\lambda y) + (1 - t)Ty]\| \leq [1 - (1 - \lambda)t]\|x - y\|,$$

which implies that  $T_t$  is a contraction. Using Banach's contraction principle, we get that there exists a unique fixed point  $x_t$  of  $T_t$  in  $H$ , i.e.,

$$x_t = t(\lambda x_t) + (1 - t)Tx_t. \quad (3.1)$$

Next, we show the convergence of the net  $\{x_t\}$ .

**Lemma 3.** *Let  $H$  be a real Hilbert space and  $T : H \rightarrow H$  be a non-expansive mapping with  $F(T) \neq \emptyset$ . Then, as  $t \rightarrow 0$ , the net  $\{x_t\}$  defined by (3.1) converges strongly to a fixed point of  $T$ .*

*Proof.* First, we prove that  $\{x_t\}$  is bounded. Take  $u \in F(T)$ . From (3.1), we have

$$\begin{aligned} \|x_t - u\| &= \|t(\lambda x_t) + (1 - t)Tx_t - u\| \leq \lambda t\|x_t - u\| + (1 - t)\|Tx_t - u\| + (1 - \lambda)t\|u\| \\ &\leq [1 - (1 - \lambda)t]\|x_t - u\| + (1 - \lambda)t\|u\|, \end{aligned}$$

which implies that  $\|x_t - u\| \leq \|u\|$ . Hence  $\{x_t\}$  is bounded. Again, from (3.1), it follows that

$$\|x_t - Tx_t\| = t\|\lambda x_t - Tx_t\| \rightarrow 0. \quad (3.2)$$

Next, we show that  $\{x_t\}$  is relatively norm compact as  $t \rightarrow 0$ . Let  $\{t_n\} \subset (0, 1)$  be a sequence such that  $t_n \rightarrow 0$  as  $n \rightarrow \infty$ . Put  $x_n := x_{t_n}$ . From (3.2), we have

$$\|x_n - Tx_n\| \rightarrow 0. \quad (3.3)$$

From (3.1), we get

$$\begin{aligned} \|x_t - u\|^2 &= \langle t(\lambda x_t) + (1 - t)Tx_t - u, x_t - u \rangle \\ &= \lambda t\langle x_t - u, x_t - u \rangle + (1 - t)\langle Tx_t - u, x_t - u \rangle - (1 - \lambda)t\langle u, x_t - u \rangle \\ &\leq [1 - (1 - \lambda)t]\|x_t - u\|^2 - (1 - \lambda)t\langle u, x_t - u \rangle, \end{aligned}$$

which implies that  $\|x_t - u\|^2 \leq \langle u, u - x_t \rangle$ . In particular,

$$\|x_n - u\|^2 \leq \langle u, u - x_n \rangle, \quad \forall u \in F(T). \quad (3.4)$$

Since  $\{x_n\}$  is bounded, without loss of generality, we may assume that  $x_n \rightharpoonup x^* \in H$ . Noticing (3.3), we can use Lemma 1 to get  $x^* \in F(T)$ . Therefore, we can substitute  $x^*$  for  $u$  in (3.4) to get

$$\|x_n - x^*\|^2 \leq \langle x^*, x^* - x_n \rangle.$$

Hence  $x_n \rightharpoonup x^*$  implies that  $x_n \rightarrow x^*$ . This has proved the relative norm compactness of the net  $\{x_t\}$  as  $t \rightarrow 0$ .

To show that the entire net  $\{x_t\}$  converges to  $x^*$ , assume  $x_{t_m} \rightarrow \tilde{x} \in F(T)$ , where  $t_m \rightarrow 0$ . Put  $x_m = x_{t_m}$ . Similarly, we have

$$\|x_m - x^*\|^2 \leq \langle x^*, x^* - x_m \rangle$$

and so

$$\|\tilde{x} - x^*\|^2 \leq \langle x^*, x^* - \tilde{x} \rangle. \tag{3.5}$$

Interchanging  $x^*$  and  $\tilde{x}$ , we obtain

$$\|x^* - \tilde{x}\|^2 \leq \langle \tilde{x}, \tilde{x} - x^* \rangle. \tag{3.6}$$

Adding up (3.5) and (3.6) yields

$$2\|x^* - \tilde{x}\|^2 \leq \|x^* - \tilde{x}\|^2,$$

which implies that  $\tilde{x} = x^*$ . This completes the proof.  $\square$

*Remark 2.* It should be pointed out that Lemma 3 is a new result which is very important for proving the following theorem.

**Theorem 3.** *Let  $H$  be a real Hilbert space and  $T : H \rightarrow H$  be a non-expansive mapping with  $F(T) \neq \emptyset$ . Let  $\{\alpha_n\}$  and  $\{\lambda_n\}$  be two real sequences in  $(0, 1)$ . For given  $x_0 \in C$  arbitrarily, let the sequence  $\{x_n\}$  be generated iteratively by*

$$x_{n+1} = \alpha_n(\lambda_n x_n) + (1 - \alpha_n)Tx_n, \quad \forall n \geq 0. \tag{3.7}$$

*Suppose the following conditions are satisfied:*

- (a)  $\lim_{n \rightarrow \infty} \alpha_n = 0$ ;    (b)  $\lim_{n \rightarrow \infty} \lambda_n = 1$ ;
- (c)  $\sum_{n=0}^{\infty} |\alpha_n - \alpha_{n-1}| < \infty$ ,  $\sum_{n=0}^{\infty} |\lambda_n - \lambda_{n-1}| < \infty$  and  $\sum_{n=0}^{\infty} (1 - \lambda_n)\alpha_n = \infty$ .

*Then the sequence  $\{x_n\}$  generated by (3.7) strongly converges to a fixed point of  $T$ .*

*Proof.* First, we prove that the sequence  $\{x_n\}$  is bounded. Take  $u \in F(T)$ . From (3.7), we have

$$\begin{aligned} \|x_{n+1} - u\| &= \|\alpha_n(\lambda_n x_n) + (1 - \alpha_n)Tx_n - u\| \\ &\leq \alpha_n \lambda_n \|x_n - u\| + (1 - \lambda_n)\alpha_n \|u\| + (1 - \alpha_n)\|x_n - u\| \\ &= [1 - (1 - \lambda_n)\alpha_n]\|x_n - u\| + (1 - \lambda_n)\alpha_n \|u\| \leq \max\{\|x_n - u\|, \|u\|\}. \end{aligned}$$

Hence  $\{x_n\}$  is bounded and so is  $\{Tx_n\}$ . From (3.7), it follows that

$$\begin{aligned} \|x_{n+1}-x_n\| &= \|\alpha_n(\lambda_n x_n)+(1-\alpha_n)Tx_n-\alpha_{n-1}(\lambda_{n-1}x_{n-1})-(1-\alpha_{n-1})Tx_{n-1}\| \\ &= \|\alpha_n\lambda_n(x_n-x_{n-1})+\alpha_n(\lambda_n-\lambda_{n-1})x_{n-1}+(\alpha_n-\alpha_{n-1})(\lambda_{n-1}x_{n-1}) \\ &\quad + (1-\alpha_n)(Tx_n-Tx_{n-1})+(\alpha_{n-1}-\alpha_n)Tx_{n-1}\| \\ &\leq \alpha_n\lambda_n\|x_n-x_{n-1}\|+(1-\alpha_n)\|Tx_n-Tx_{n-1}\| \\ &\quad +|\alpha_n-\alpha_{n-1}|(\lambda_{n-1}\|x_{n-1}\|+\|Tx_{n-1}\|)+\alpha_n|\lambda_n-\lambda_{n-1}|\|x_{n-1}\| \\ &\leq [1-(1-\lambda_n)\alpha_n]\|x_n-x_{n-1}\|+(|\alpha_n-\alpha_{n-1}|+\alpha_n|\lambda_n-\lambda_{n-1}|)M_1, \end{aligned} \tag{3.8}$$

where  $M_1$  is a constant such that  $\sup_n\{\|x_{n-1}\|+\|Tx_{n-1}\|\} \leq M_1$ . Hence, from (3.8) and Lemma 2, we deduce

$$\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = 0.$$

At the same time, we note that

$$\|x_{n+1} - Tx_n\| = \alpha_n\|(\lambda_n x_n) - Tx_n\| \rightarrow 0.$$

Therefore, we have

$$\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0.$$

Next, we prove  $\limsup_{n \rightarrow \infty} \langle x^*, x^* - x_n \rangle \leq 0$ , where  $x^* = \lim_{t \rightarrow 0} x_t$  and  $\{x_t\}$  is the net defined by (3.1). As a matter of fact, we have

$$\begin{aligned} \|x_t - x_n\|^2 &= \langle x_t - x_n, x_t - x_n \rangle = t\langle x_t - x_n, x_t - x_n \rangle - (1-\lambda)t\langle x_t, x_t - x_n \rangle \\ &\quad + (1-t)\langle Tx_t - Tx_n, x_t - x_n \rangle + (1-t)\langle Tx_n - x_n, x_t - x_n \rangle \\ &\leq \|x_t - x_n\|^2 - (1-\lambda)t\langle x_t, x_t - x_n \rangle + (1-t)\langle Tx_n - x_n, x_t - x_n \rangle \\ &\leq \|x_t - x_n\|^2 - (1-\lambda)t\langle x_t, x_t - x_n \rangle + M_2\|Tx_n - x_n\|, \end{aligned}$$

where  $M_2 > 0$  such that  $\sup\{\|x_t - x_n\|, t \in (0, 1), n \geq 0\} \leq M_2$ , which implies that

$$\langle x_t, x_t - x_n \rangle \leq \frac{M_2}{(1-\lambda)t}\|Tx_n - x_n\|.$$

Therefore, we have

$$\limsup_{t \rightarrow 0} \limsup_{n \rightarrow \infty} \langle x_t, x_t - x_n \rangle \leq 0. \tag{3.9}$$

We note that

$$\begin{aligned} \langle x^*, x^* - x_n \rangle &= \langle x^*, x^* - x_t \rangle + \langle x^* - x_t, x_t - x_n \rangle + \langle x_t, x_t - x_n \rangle \\ &\leq \langle x^*, x^* - x_t \rangle + \|x^* - x_t\|\|x_t - x_n\| + \langle x_t, x_t - x_n \rangle \\ &\leq \langle x^*, x^* - x_t \rangle + \|x^* - x_t\|M_2 + \langle x_t, x_t - x_n \rangle. \end{aligned}$$

This together with  $x_t \rightarrow x^*$  and (3.9) implies that

$$\limsup_{n \rightarrow \infty} \langle x^*, x^* - x_n \rangle \leq 0.$$

Finally, we show that  $x_n \rightarrow x^*$ . From (3.7), we have

$$\begin{aligned} \|x_{n+1} - x^*\|^2 &= \langle x_{n+1} - x^*, x_{n+1} - x^* \rangle = \alpha_n \lambda_n \langle x_n - x^*, x_{n+1} - x^* \rangle \\ &+ (1 - \lambda_n) \alpha_n \langle x^*, x^* - x_{n+1} \rangle + (1 - \alpha_n) \langle Tx_n - x^*, x_{n+1} - x^* \rangle \\ &\leq [1 - (1 - \lambda_n) \alpha_n] \|x_n - x^*\| \|x_{n+1} - x^*\| + (1 - \lambda_n) \alpha_n \langle x^*, x^* - x_{n+1} \rangle \\ &\leq \frac{1 - (1 - \lambda_n) \alpha_n}{2} (\|x_n - x^*\|^2 + \|x_{n+1} - x^*\|^2) + (1 - \lambda_n) \alpha_n \langle x^*, x^* - x_{n+1} \rangle, \end{aligned}$$

which implies that

$$\|x_{n+1} - x^*\|^2 \leq [1 - (1 - \lambda_n) \alpha_n] \|x_n - x^*\|^2 + 2(1 - \lambda_n) \alpha_n \langle x^*, x^* - x_{n+1} \rangle.$$

We can check that all the assumptions of Lemma 2 are satisfied. Therefore, we deduce  $x_n \rightarrow x^*$ . This completes the proof.  $\square$

*Remark 3.* The prototype for the iteration parameters are as follows:  $\alpha_n = \frac{1}{\sqrt{n}}$  and  $\lambda_n = 1 - \frac{1}{\sqrt{n}}$  for all  $n \geq 0$ . It is easy to check that these choices satisfy all the conditions of Theorem 3.

*Remark 4.* We conclude the paper with the following observations:

(1) Our algorithm (3.7) is similar to Mann’s algorithm (1.1). It is well-known that Mann’s method for non-expansive mappings has only weak convergence. However, our iterative method (3.7) has strong convergence. Our iterative scheme (3.7) may be interesting which is very different from the algorithm of Kim and Xu (1.3). As a matter of fact, the algorithm (1.3) is a convex combination of  $u$  and the Mann iteration method (1.1).

(2) In order to prove the strong convergence of the algorithm (1.3), Kim and Xu [9] used the convergence of the net  $x_t = tu + (1 - t)Tx_t$ . It is worthy to note that we prove the strong convergence of the algorithm (3.7) using the new Lemma 3. In this respect, our methods are independent from those given in the literature.

## 4 Conclusions

The problem of finding fixed points of nonexpansive mappings has attracted much attention because of its extraordinary utility and broad applicability in many branches of mathematical science and engineering.

Mann’s algorithm is a widely used method for solving a fixed point equation of the form  $Tx = x$ , where  $C$  is a nonempty closed convex subset of a Hilbert space  $H$ , and  $T : C \rightarrow C$  is a nonexpansive mapping. This algorithm generates from an arbitrary initial guess  $x_0 \in C$ , a sequence  $\{x_n\}$  by the recursive manner:

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_nTx_n, \quad \forall n \geq 1,$$

where  $\{\alpha_n\}$  is a sequence in the interval  $[0, 1]$ .

In an infinite-dimensional space  $X$ , Mann's algorithm has only weak convergence, in general. In fact, it is known that if the sequence  $\{\alpha_n\}$  is such that  $\sum_{n=0}^{\infty} \alpha_n(1-\alpha_n) = \infty$ , then Mann's algorithm converges weakly to a fixed point of  $T$  provided the underlying space  $X$  is a Hilbert space or more general, a uniformly convex Banach space which has a Frechet differentiable norm or satisfies Opial's property. It is a very interesting topic of constructing some algorithms such that the strong convergence of proposed algorithms are guaranteed. For this purpose, in this article we present a modified Krasnoselskii-Mann method (3.7) for nonexpansive mappings in Hilbert spaces and show that the proposed method (3.7) has strong convergence. However, we note that in order to obtain the main result of Theorem 3, we have imposed some additional conditions  $\sum_{n=0}^{\infty} |\alpha_n - \alpha_{n-1}| < \infty$ ,  $\sum_{n=0}^{\infty} |\lambda_n - \lambda_{n-1}| < \infty$  and  $\sum_{n=0}^{\infty} (1 - \lambda_n)\alpha_n = \infty$ . Hence this brings us a nature problem: could we weaken or drop these additional assumptions?

## Acknowledgement

The authors are very grateful to the referees for their careful reading, comments and suggestions which improved the presentation of this article.

## References

- [1] H.H. Bauschke. The approximation of fixed points of compositions of nonexpansive mappings in Hilbert space. *J. Math. Anal. Appl.*, **202**:150–159, 1996. (Doi:10.1006/jmaa.1996.0308)
- [2] F.E. Browder and W.V. Petryshyn. Construction of fixed points of nonlinear mappings. *J. Math. Anal. Appl.*, **20**:197–228, 1967. (Doi:10.1016/0022-247X(67)90085-6)
- [3] C. Byrne. A unified treatment of some iterative algorithms in signal processing and image reconstruction. *Inverse Problems*, **20**:103–120, 2004. (Doi:10.1088/0266-5611/20/1/006)
- [4] J.P. Chancelier. Iterative schemes for computing fixed points of nonexpansive mappings in Banach spaces. *J. Math. Anal. Appl.*, **353**:141–153, 2009. (Doi:10.1016/j.jmaa.2008.11.041)
- [5] P.L. Combettes. On the numerical robustness of the parallel projection method in signal synthesis. *IEEE signal Process. Lett.*, **8**:45–47, 2001. (Doi:10.1109/97.895371)
- [6] H.W. Engl and A. Leitao. A Mann iterative regularization for elliptic Cauchy problems. *Numer. Funct. Anal. Optim.*, **22**:861–884, 2001. (Doi:10.1081/NFA-100108313)
- [7] A. Genel and J. Lindenstrass. An example concerning fixed points. *Israel J. Math.*, **22**:81–86, 1975. (Doi:10.1007/BF02757276)
- [8] K. Geobel and W.A. Kirk. Topics in metric fixed point theory. *Cambridge Studies in Advanced Mathematics*, **28**, 1990. Cambridge University Press.
- [9] T.H. Kim and H.K. Xu. Strong convergence of modified Mann iterations. *Nonlinear Anal.*, **61**:51–60, 2005. (Doi: 10.1016/j.na.2004.11.011)



- [10] T.H. Kim and H.K. Xu. Robustness of Mann's algorithm for non-expansive mappings. *J. Math. Anal. Appl.*, **327**:1105–1115, 2007. (Doi:10.1016/j.jmaa.2006.05.009)
- [11] P.L. Lions. Approximation de points fixes de contractions. *C.R.Acad.Sci.Ser.A-B Paris*, **284**:1357–1359, 1977.
- [12] A. Moudafi. Krasnoselski-Mann iteration for hierarchical fixed-point problems. *Inverse Problems*, **23**:1635–1640, 2007. (Doi:10.1088/0266-5611/23/4/015)
- [13] J.W. Peng and J.C. Yao. A new hybrid-extragradient method for generalized mixed equilibrium problems and fixed point problems and variational inequality problems. *Taiwanese Journal of Mathematics*, **12**:1401–1433, 2008.
- [14] C.I. Podilchuk and R.J. Mammone. Image recovery by convex projections using a least-squares constraint. *J. Optim. Soc. Amer.*, **7**:517–521, 1990. (Doi:10.1364/JOSAA.7.000517)
- [15] X. Qin, Y.J. Cho, J.I. Kang and S.M. Kang. Strong convergence theorems for an infinite family of nonexpansive mappings in Banach spaces. *J. Comput. Appl. Math.*, **230**:121–127, 2009. (Doi:10.1016/j.cam.2008.10.058)
- [16] S. Reich. Weak convergence theorems for non-expansive mappings in Banach spaces. *J. Math. Anal. Appl.*, **67**:274–276, 1979. (Doi:10.1016/0022-247X(79)90024-6)
- [17] N. Shioji and W. Takahashi. Strong convergence of approximated sequence for non-expansive mappings. *Proc. Amer. Math. Soc.*, **125**:3641–3645, 1997. (Doi:10.1090/S0002-9939-97-04033-1)
- [18] T. Suzuki. Strong convergence theorems for infinite families of nonexpansive mappings in general Banach spaces. *Fixed Point Theory Appl.*, **2005**:103–123, 2005. (Doi:10.1155/FPTA.2005.103)
- [19] R. Wittmann. Approximation of fixed points of non-expansive mappings. *Arch. Math.*, **58**:486–491, 1992. (Doi:10.1007/BF01190119)
- [20] N.C. Wong, D.R. Sahu and J.C. Yao. Solving variational inequalities involving nonexpansive type mappings. *Nonlinear Anal.*, **69**:4732–4753, 2008. (Doi:10.1016/j.na.2007.11.025)
- [21] H.K. Xu. Another control condition in an iterative method for non-expansive mappings. *Bull. Austral. Math. Soc.*, **65**:109–113, 2002. (Doi:10.1017/S0004972700020116)
- [22] H.K. Xu. Viscosity approximation methods for non-expansive mappings. *J. Math. Anal. Appl.*, **298**:279–291, 2004. (Doi:10.1016/j.jmaa.2004.04.059)
- [23] Y. Yao, R. Chen and J.C. Yao. Strong convergence and certain control conditions of modified Mann iteration. *Nonlinear Anal.*, **68**:1687–1693, 2008.
- [24] Y. Yao and Y.-C. Liou. Weak and strong convergence of Krasnoselski-Mann iteration for hierarchical fixed point problems. *Inverse Problems*, **24**(1):15–22, 2008. (Doi:10.1088/0266-5611/24/1/015015)
- [25] Y. Yao and Y.C. Liou. Strong convergence theorems for infinite countable nonexpansive mappings and image recovery problem. *J. Korean Math. Soc.*, **45**:1591–1600, 2008. ((Doi:0.4134/JKMS.2008.45.6.1591)
- [26] Y. Yao, Y.C. Liou and R. Chen. Convergence theorems for fixed point problems and variational inequality problems. *J. Nonlinear Convex Anal.*, **9**:239–248, 2008.

- [27] Y. Yao, Y.C. Liou and S.M. Kang. Strong convergence of an iterative algorithm on an infinite countable family of nonexpansive mappings. *Appl. Math. Comput.*, **208**:211–218, 2009. (Doi:10.1016/j.amc.2008.11.038)
- [28] D. Youla. Mathematical theory of image restoration by the method of convex projection. In H. Stark (Ed.)(Ed.), *Image Recovery Theory and Applications*. Academic Press, Orlando, pp. 29–77, 1987.