

MIT Joint Program on the Science and Policy of Global Change



A Method for Calculating Reference Evapotranspiration on Daily Time Scales

*William Farmer, Kenneth Strzepke, C. Adam Schlosser, Peter Droogers, and
Xiang Gao*

**Report No. 195
February 2011**

The MIT Joint Program on the Science and Policy of Global Change is an organization for research, independent policy analysis, and public education in global environmental change. It seeks to provide leadership in understanding scientific, economic, and ecological aspects of this difficult issue, and combining them into policy assessments that serve the needs of ongoing national and international discussions. To this end, the Program brings together an interdisciplinary group from two established research centers at MIT: the Center for Global Change Science (CGCS) and the Center for Energy and Environmental Policy Research (CEEPR). These two centers bridge many key areas of the needed intellectual work, and additional essential areas are covered by other MIT departments, by collaboration with the Ecosystems Center of the Marine Biology Laboratory (MBL) at Woods Hole, and by short- and long-term visitors to the Program. The Program involves sponsorship and active participation by industry, government, and non-profit organizations.

To inform processes of policy development and implementation, climate change research needs to focus on improving the prediction of those variables that are most relevant to economic, social, and environmental effects. In turn, the greenhouse gas and atmospheric aerosol assumptions underlying climate analysis need to be related to the economic, technological, and political forces that drive emissions, and to the results of international agreements and mitigation. Further, assessments of possible societal and ecosystem impacts, and analysis of mitigation strategies, need to be based on realistic evaluation of the uncertainties of climate science.

This report is one of a series intended to communicate research results and improve public understanding of climate issues, thereby contributing to informed debate about the climate issue, the uncertainties, and the economic and social implications of policy alternatives. Titles in the Report Series to date are listed on the inside back cover.

Ronald G. Prinn and John M. Reilly
Program Co-Directors

For more information, please contact the Joint Program Office

Postal Address: Joint Program on the Science and Policy of Global Change
77 Massachusetts Avenue
MIT E19-411
Cambridge MA 02139-4307 (USA)

Location: 400 Main Street, Cambridge
Building E19, Room 411
Massachusetts Institute of Technology

Access: Phone: +1(617) 253-7492
Fax: +1(617) 253-9845
E-mail: globalchange@mit.edu
Web site: <http://globalchange.mit.edu/>

A Method for Calculating Reference Evapotranspiration on Daily Time Scales

William Farmer^{*}, Kenneth Strzepek^{*}, C. Adam Schlosser^{*}, Peter Droogers[†], and Xiang Gao^{*}

Abstract

Measures of reference evapotranspiration are essential for applications of agricultural management and water resources engineering. Using numerous esoteric variables, one can calculate daily reference evapotranspiration using the Modified Penman-Monteith methods. In 1985, Hargreaves developed a simplified method for estimating reference evapotranspiration. Similarly, Droogers and Allen improved upon Hargreaves' method in 2002. Both methods provide excellent estimates of average daily rates for a given month, based on monthly climatology. The Hargreaves method also estimates daily rates based on daily data, though the Modified Hargreaves approach developed by Droogers and Allen is largely accepted as a stronger metric. Here efforts are made to improve the functionality of Droogers and Allen's approach and to adapt it to provide daily estimates of reference evapotranspiration based on daily weather. The Hargreaves and Modified Hargreaves are used to calculate daily reference evapotranspiration based on daily data. The coefficients in these equations are then optimized to reduce the root mean squared difference between each estimate and the baseline value calculated by the Modified Penman-Monteith approach. The adapted method for daily reference evapotranspiration proves promising; estimating rates near a root mean squared difference of 1.07 mm/day. These results are validated with data from 1976-1980; here the root mean squared difference is 1.06 mm/day. Results are evaluated spatially and temporally. Weaknesses are seen in the estimates around clearly-defined summers. Further weaknesses are seen in pole-ward regions. Still, at the 1% significance level, the daily optimization of the Modified Hargreaves equation is found to be the best replica of the Modified Penman-Monteith method, globally. Finally, specific caveats and further avenues of research are noted. Overall, the daily Modified-Hargreaves method is advocated for general use in global studies where daily data and variation is of the utmost concern.

Contents

1. INTRODUCTION.....	1
2. METHODS	3
3. RESULTS	5
3.1 Global Analyses of Optimization (1971-1975)	5
3.1.1 Annual Results	5
3.1.2 Seasonal Evaluation	6
3.2 Regional Analyses of Optimization (1971-1975).....	7
4. VALIDATION OF OPTIMIZATION (1976-1980).....	11
4.1 Seasonal Evaluation.....	11
4.2 Regional Evaluation	12
5. CLOSING REMARKS	14
6. REFERENCES.....	17
APPENDIX.....	19

1. INTRODUCTION

Measures of reference evapotranspiration, as defined by Allen *et al.* (1998), are essential to modeling and managing agricultural and water resources, from crop selection to irrigation allocation, streamflow and watershed analysis. Depending on the desired application and time-

^{*} MIT Joint Program on the Science and Policy of Global Change, Cambridge, MA. (Corresponding author: E-mail. whfarmer@mit.edu)

[†] FutureWater, Wageningen, The Netherlands.

step, there are numerous ways to calculate reference evapotranspiration. Due to the high data-demand associated with daily methods, a large number of models have focused on the use of monthly methods of calculation (McKenney and Rosenberg, 1992). While monthly estimates are valuable as a baseline for understanding, further applications require the details acquired at the daily scale. For example, the unique nature of crops and their responses to weather and soil moisture are more appropriately represented at a daily time step.

Currently, the Modified Penman-Monteith, as developed in FAO Irrigation & Drainage Paper No. 56 (1998), is a widely-used method for calculating reference evapotranspiration on a daily time step. Unfortunately, the data demands make this method somewhat problematic, especially for applications in data-poor, developing regions of the globe. These intensive data demands also make this method unpalatable for use with global databases, which often contain only a restricted set of variables. Furthermore, when working with climate change within a set of general circulation models (GCMs), the data demands of the Modified Penman-Monteith method are rarely met in all cases.

Among the key inputs of the Modified Penman-Monteith method are measurements of temperature, wind speed, net radiation, and vapor pressure deficit. While these variables are often available within extensive atmospheric models for coupled climate and weather prediction, comprehensive sets of field measurements containing all these terms are uncommon, even at monthly time steps. As such, Droogers and Allen (2002) confronted the issue of inaccurate or incomplete data on a monthly time step by augmenting the method of Hargreaves *et al.* (1985) to calculate average monthly reference evapotranspiration (mm/day) based on a limited number of variables. Specifically, it was shown that reasonable estimates of reference evapotranspiration could be calculated from monthly average temperature, temperature range and precipitation. This approach is advantageous, as these atmospheric variables are generally available with some confidence across the globe. However, the method remains somewhat unpalatable to most agricultural modeling applications because this reference evapotranspiration represents an estimate for an average day of the month, based on monthly climate conditions. Again, this is problematic because so many applications thrive on daily fluctuations of weather and the variation of those fluctuations.

The purpose of this exercise is to extend the work of Droogers and Allen (2002) by introducing a new algorithm that is able to calculate daily reference evapotranspiration from

daily environmental data. The ability of this algorithm to resolve the daily variations - and the extremes - of reference evapotranspiration will substantially benefit natural resource planning, development planning and water resource engineering. This ability allows for a better evaluation of risk-based assessment and the distribution of current and future climate variables at the daily scale. In the next section, we briefly describe the algorithms and data sets employed, as well as the metrics used to evaluate the daily reference evapotranspiration algorithm. Results from the suite of simulations are then presented, and concluding remarks are provided.

2. METHODS

In general, this exercise will use daily climate data from grid cells across the globe to calculate daily reference evapotranspiration using a number of different methods. This approach is largely a replication of the methods used by Droogers and Allen (2002). The Modified Penman-Monteith method (Allen *et al.*, 1998) will be used as the best approximation, and thus referred to as the baseline estimate. The Hargreaves and Modified-Hargreaves methods (Droogers and Allen 2002) will be compared to the baseline and a measure of their error will be calculated. By varying the coefficients of these equations, the error will be minimized. The results of this variation will be a robust daily method for calculating reference evapotranspiration. The experiment is outlined in detail below.

To generate the required daily inputs for the various evapotranspiration calculations, we employed the Community Land Model Version 3.5 (CLM3.5, Oleson *et al.*, 2004) in “stand-alone” mode in which the atmospheric conditions are prescribed. For this study, atmospheric weather conditions were provided by the National Centers for Environmental Prediction (NCEP). This data, called the NCC dataset, consists of reanalysis data that has been bias corrected to match monthly estimates by the Climate Research Unit (CRU) at the University of East Anglia CRU (Ngo-Duc *et al.*, 2005). CLM was run globally at a spatial resolution of $1^\circ \times 1^\circ$ to provide all the near-surface variables needed to calculate reference evapotranspiration via the Modified Penman-Monteith (Allen *et al.*, 1998), Hargreaves (Har) and Modified-Hargreaves (MH) methods (Droogers and Allen, 2002). For this exercise, daily variables were taken for the period of 1971 through 1976. Looking only at one-degree-square, over-land grids, 25, 252, 525 individual estimates were evaluated.

For this evaluation of the various methods of evapotranspiration calculation, the Modified Penman-Monteith estimate served as the baseline reference evapotranspiration rate. Reference

evapotranspiration was then calculated using the Hargreaves equation (Droogers and Allen, 2002):

$$ET_0 = 0.0023 \cdot 0.408RA \cdot (T_{avg} - 17.8) \cdot TD^{0.5} \quad (1)$$

where RA is incoming solar radiation, T_{avg} is mean daily temperature and TD is the daily temperature range. Similarly, reference evaporation was then calculated by forcing the Modified-Hargreaves equation with daily data, where all variables are the same as in (1), except for an additional term for daily precipitation:

$$ET_0 = 0.0013 \cdot 0.408RA \cdot (T_{avg} + 17) (TD - 0.0123P)^{0.76} \quad (2)$$

It is important to note that Droogers and Allen (2002) optimized the Modified-Hargreaves equation for a monthly input of precipitation, yielding ET_0 in mm/day; here we keep the coefficients the same and use daily data. After this calculation, this method will be expanded upon by applying (2) with daily data and re-estimating the parameters represented by certain coefficients.

As mentioned, the outputs of the Hargreaves equation and the Modified-Hargreaves equation were compared with the Modified Penman-Monteith calculations. Their level of consistency was assessed via the Pearson R^2 correlation coefficient and the root mean squared difference from Droogers and Allen (2002):

$$RMSD = \sqrt{\frac{\sum_{i=1}^n (Pen_i - Calc_i)^2}{n}} \quad (3)$$

where Pen is the reference evapotranspiration calculation from Modified Penman-Monteith and $Calc$ is the same as calculated by the Hargreaves and Modified-Hargreaves. The equation coefficients were estimated by optimizing the correlation coefficient and RMSD using a least-squared regression algorithm. This was done by varying the lettered coefficients of the generalized Hargreaves and Modified-Hargreaves equations given as:

$$ET = a \cdot 0.408RA \cdot (T_{avg} + b) TD^{0.5} \quad (4)$$

$$ET = a \cdot 0.408RA \cdot (T_{avg} + b) (TD - c \cdot P)^d \quad (5)$$

In this manner, the values of a, b, c and d were changed until the RMSD was minimized.

For analyses, the RMSDs and correlation coefficients were evaluated for seven latitudinal regions described in **Table 1**, as well as for each season and on an annual basis. The seasons were defined as December-January-February (DJF), March-April-May (MAM), June-July-August (JJA) and September-October-November (SON). It is important to note that the parameters of equations (4) and (5) were calibrated globally, then the results were assessed regionally and seasonally. No attempt was made to recalibrate the parameters for each season or region. Finally, the results were tested for significance to determine which equation was the best tool for each spatio-temporal region.

Table 1. Seven regions for analysis.

Region	Lower Latitude	Upper Latitude
1	45	60
2	30	45
3	15	30
4	-15	15
5	-30	-15
6	-45	-30
7	-60	-45

3. RESULTS

3.1 Global Analyses of Optimization (1971-1975)

3.1.1 Annual Results

Table 2 summarizes the coefficients of each equation, with their root mean square difference (RMSD) and regression coefficients. Calculations were based on the 25, 252, 525 estimates obtained from all the 1°x1° grids from 1971-1975. The equations noted as ‘daily’ are the equations optimized for daily inputs. The R² and RMSD metrics were obtained by pairing each estimate at every grid point against the corresponding Modified Penman-Monteith result.

Table 2. Statistical measures of the strength of different methods for calculating reference evapotranspiration, 1971-1975.

Equation	R ²	RMSD	a	b	c	d
Hargreaves (Har)	0.8320	1.4720	0.0023	17.8		
Daily Hargreaves (dailyHar)	0.8321	1.1390	0.0028	19.1869		
Modified Hargreaves (MH)	0.8591	1.2630	0.0013	17	0.0123	0.76
Daily Modified Hargreaves (dailyMH)	0.8564	1.0650	0.0019	21.0584	0.0874	0.6278

Modifying the original coefficients of the MH and Har methods (4) and (5) allowed for a reduction in the RMSD of the Hargreaves equation from 1.4720 mm/day to 1.1390 mm/day without corrupting the regression coefficient. This represents a reduction in the RMSD metric of 22.62%.

The original Modified-Hargreaves equation yielded a larger RMSD than the fitted Hargreaves equation, although the regression coefficient is slightly stronger.

The daily Modified-Hargreaves equation yields the most accurate result, coupling a similar regression coefficient with the lowest RMSD, 1.0650 mm/day. This is a 15.68% RMSD improvement on the Modified Hargreaves equation and a 6.50% RMSD improvement on the daily Hargreaves equation.

The optimization of the Modified-Hargreaves equation for daily data was able to reduce the RMSD of all the observations by almost 0.25 mm/day without dramatic changes to the regression coefficient. Nevertheless, it is equally important to understand the effects of this optimization on smaller spatial and temporal scales. To this end, the results of the optimization will be evaluated across different seasons and across different latitudes. Again, note that the results are merely evaluated regionally and seasonally; no recalibration is conducted. Finally, the optimized equations will be validated using a second period of data, from 1976 through 1980.

3.1.2 Seasonal Evaluation

Seasons were defined as December-January-February (DJF, cyclical days 335 through 59), March-April-May (MAM, days 60 through 151), June-July-August (JJA, days 152 through 243) and September-October-November (SON, days 244 through 334). The RMSD for the original Hargreaves (Har), optimized Hargreaves (dailyHar), Modified-Hargreaves (MH) and optimized Modified-Hargreaves (dailyMH) are presented in **Table 3**. (R^2 values can be found in Appendix, Table A1.)

Table 3. Global, seasonal RMSD values, 1971-1975.

Equation	DJF	MAM	JJA	SON	All
Har	1.3082	1.4309	1.6191	1.5832	1.4730
dailyHar	1.0214	1.1265	1.2782	1.1642	1.1405
MH	1.1768	1.2515	1.3358	1.3614	1.2684
dailyMH	0.9369	1.0820	1.1947	1.0820	1.0664

As seen for the annual results in Table 2, for all seasons the RMSD was reduced by each optimization while the R^2 was maintained. In each season, the dailyMH equation provided the lowest RMSD, making it the strongest equation for the calculation of reference evapotranspiration.

It is interesting that almost all seasons provided a slightly higher RMSD than the all-inclusive RMSD. The largest inaccuracies are found in the JJA season, or the summer of the northern hemisphere. Noting that some 61% of the observations were in the northern hemisphere, this high RMSD seems to suggest that the estival seasonality complicates the accuracy of reference evapotranspiration calculations, as will be discussed below.

At first glance, a slight decrease in the effectiveness of the dailyMH equation during the northern-hemisphere summer can be seen (Table 3). In the northern-hemisphere winter and autumn, the dailyMH equation improves upon the MH equation some 20.39% and 20.52%, respectively. In the spring and summer the improvement is only 13.55% and 10.56%, respectively.

3.2 Regional Analyses of Optimization (1971-1975)

Schlosser and Gao (2010) found that the consistency among modeled evapotranspiration is less robust for high latitudes as well as in the tropics. It is therefore important to look at the optimized data under these considerations. Given this, the results were further pooled by 15° latitude bands (Table 1). The number of individual estimates for each geographic region and season is given in **Table 4**.

Table 4. Number of estimates evaluated in each region and season.

N	DJF	MAM	JAJ	SON	All
Region 1	1,552,050	1,586,540	1,586,540	1,569,295	6,294,425
Region 2	1,227,150	1,254,420	1,254,420	1,240,785	4,976,775
Region 3	1,007,100	1,029,480	1,029,480	1,018,290	4,084,350
Region 4	1,440,450	1,472,460	1,472,460	1,456,455	5,841,825
Region 5	660,150	674,820	674,820	667,485	2,677,275
Region 6	271,800	277,840	277,840	274,820	1,102,300
Region 7	67,950	69,460	69,460	68,705	275,575
Global	6,226,650	6,365,020	6,365,020	6,295,835	25,252,525

Tables 5-11 display the seasonal and annual RMSD for all seven regions. (R^2 values can be perused in Appendix, Tables A2-8.) Instances where RMSD was increased after the optimization are highlighted in red. It is important to note here that these tabulated results are

based on the *global, annual-based* optimization (Section 3.1.1) of the RMSD results, made on the annual estimates. In this way, we are testing the robustness and the degree of ubiquity in the global, annual optimization, by the extent to which it holds on a regional and seasonal basis. The coefficients were not recalibrated for each region.

Table 5. Seasonal RMSD values for Region 1, 1971-1975. Instances where RMSD was increased after the optimization are highlighted in red.

Equation	DJF	MAM	JJA	SON	All
Har	0.4350	0.6783	0.9871	0.8929	0.7784
dailyHar	0.4083	0.8711	1.2595	0.7183	0.8712
MH	0.5546	0.7387	0.9381	0.8786	0.7917
dailyMH	0.3967	0.8427	1.1734	0.6990	0.8277

Table 6. Seasonal RMSD values for Region 2, 1971-1975. Instances where RMSD was increased after the optimization are highlighted in red.

Equation	DJF	MAM	JJA	SON	All
Har	0.8951	1.1215	1.8394	1.6697	1.4350
dailyHar	0.7501	1.0370	1.3957	1.1885	1.1183
MH	0.9101	0.9785	1.3918	1.4198	1.1976
dailyMH	0.7085	1.0063	1.3336	1.0866	1.0581

Table 7. Seasonal RMSD values for Region 3, 1971-1975.

Equation	DJF	MAM	JJA	SON	All
Har	1.7926	2.2669	2.4282	2.3633	2.2270
dailyHar	1.2133	1.4935	1.6835	1.6187	1.5132
MH	1.4971	1.6927	1.7867	1.8763	1.7185
dailyMH	1.1209	1.4279	1.5322	1.4667	1.3962

Table 8. Seasonal RMSD values for Region 4, 1971-1975.

Equation	DJF	MAM	JJA	SON	All
Har	1.2673	1.2146	1.2023	1.1443	1.2062
dailyHar	1.0843	0.9962	0.8886	0.9898	0.9904
MH	1.2882	1.2893	1.2500	1.2573	1.2702
dailyMH	1.0120	0.9950	0.8702	0.9705	0.9619

Table 9. Seasonal RMSD values for Region 5, 1971-1975.

Equation	DJF	MAM	JJA	SON	All
Har	2.0884	1.9811	1.8715	2.1295	2.0185
dailyHar	1.6202	1.4585	1.3100	1.5081	1.4770
MH	1.6157	1.7058	1.5422	1.5865	1.6132
dailyMH	1.3939	1.3287	1.1788	1.3601	1.3168

Table 10. Seasonal RMSD values for Region 6, 1971-1975.

Equation	DJF	MAM	JJA	SON	All
Har	1.8595	1.4976	1.1447	1.3004	1.4729
dailyHar	1.4380	1.0777	0.8789	1.0860	1.1363
MH	1.6146	1.3753	1.1236	1.1815	1.3363
dailyMH	1.3832	1.0359	0.8667	1.0720	1.1036

Table 11. Seasonal RMSD values for Region 7, 1971-1975. Instances where RMSD was increased after the optimization are highlighted in red.

Equation	DJF	MAM	JJA	SON	All
Har	0.9401	0.7474	0.5226	0.6738	0.7353
dailyHar	1.1954	0.6556	0.4787	0.8936	0.8470
MH	0.9857	0.8124	0.5683	0.7186	0.7849
dailyMH	1.0951	0.6599	0.4903	0.8234	0.7969

Regions one and seven (**Tables 5 and 11**) show an increase in annual RMSD as a result of the daily, global optimization. Nevertheless, it is important to note that RMSD hovers around 0.80 mm/day, which is lower than the RMSD viewed annually, across the globe (Table 1). More importantly, the increased RMSD were seen exclusively in the summer months, especially JJA. This suggests that the optimization performed on a global, annual basis is likely to be weakened at high-latitude estimates, and is qualitatively consistent with the results of Schlosser and Gao (2010).

Regions two and six represent a majority of the midlatitude regions. The results in region two (**Table 6**), spanning 30°N through 45°N, are similar to those seen in region one. Here, we find that the daily-optimized MH estimate produces a slightly higher RMSD – but only in MAM, where the increase is much smaller than the increases seen for MAM and JJA in region one. Further, for the remaining seasons, the RMSDs for the optimized dailyMH show decreases, most notably in DJF and SON, and thus support the use of the global, annual optimization approach. For JJA, the RMSDs remain slightly higher than all other seasons, though the optimization still reduces it by ~4%. For region 6 (30°S to 45°S), in all seasons the global, annual-based optimizations are improvements on previous methods for calculating reference evapotranspiration. Additionally, there is indication of seasonality in these results as the improvements are 24.68% and 22.86% in MAM and JJA, respectively, while only 14.33% and 9.27% in DJF and SON, respectively.

Regions three and five (**Tables 7 and 9**) represent a large portion of the northern and southern subtropics, respectively. These regions show some of the largest RMSDs encountered –

particularly for the original Har and MH methods. In both regions there is only a weak seasonality present in the RMSD metric. In the northern hemisphere, the season of JJA contains the most inaccurate estimates, as seen in other regions, and the RMSDs are larger than the global averages for almost all seasons (Table 3). The same can be said of DJF and SON in region five, strengthening the characterization of larger estival inaccuracies. Nevertheless, the daily optimization results in a notable improvement on the previous methods. Similar to the southern midlatitude region 6, the percent improvement of the RMSDs indicates a marked seasonality. For region three there is a 22.16% and 21.83% improvement in DJF and SON, respectively, compared with only a 15.64% and 14.25% improvement in MAM and JJA, respectively. For region five the improvement is 22.11% and 23.56% in MAM and JJA, respectively, and only 13.73% and 14.27% in DJF and SON, respectively. Overall, we can characterize the percentage improvement in RMSD in the midlatitudes and sub-tropics for the warm seasons as roughly half of that seen in the cold seasons.

Region four, while covering the largest surface area, encompassing the tropics from 15°S to 15°N, is second in the total number of estimates binned in this zonal discretization (Table 4). Here, the global, annual optimization brings sizeable decreases – at least 20% in all seasons – in the RMSDs. All but DJF result in RMSDs below one millimeter per day. There is no indication of seasonality in the optimization results for this region. Seasonality in optimized estimates would not be expected in this region due to minimal seasonal fluctuations in precipitation and temperature around the equator.

In conclusion, evaluating results by season and by region highlights two important considerations regarding the global, annual-based optimization of the Har and MH methods for calculation of reference evapotranspiration. Firstly, these estimates of reference evapotranspiration continue to be more inaccurate in the subtropics and tropics. Secondly, performing a global, annual based optimization procedure will likely result in little improvement, and even slight degradations, to the previous methods for high-latitude, estival estimates. These notes should be taken into consideration when assessing the application of this method.

This observed seasonality could also be the result of bias introduced by using a RMSD calculated in real-space. The real-space error tends to focus on errors in large values. For example, the optimization of RMSD in real-space would tend to focus on reducing a five-percent difference in large values of PET, like those seen in low latitudes, and ignore a 25% difference in

small values of PET, like those seen in high latitudes. This result could be resolved by examining the RMSD in a space similar to the log-space, but this is suggested for further research. Of course, due to the frequency of zero values, log-space itself is not functional here.

4. VALIDATION OF OPTIMIZATION (1976-1980)

The above discussion analyzes spatial and temporal patterns of 25, 252, 525 estimates used to optimize methods of calculating reference evapotranspiration. It is important to show that these results hold for periods outside of those used for the optimization: that is, a split-sample validation is needed. To this end, the same number of samples was taken from the period 1976-1980 and will be evaluated below. These samples break down similar to those of the period of 1971-1975 (Table 4). Showing that the results seen in the 1971-1975 data hold for the period of 1976 through 1980 provides a validation for the optimization developed here. As shown below, the results are indeed quite similar.

For comparison, **Table 12** shows the global results when the optimized equation is applied to the validation period. The results are qualitatively identical to those shown in Table 2.

Table 12. Statistical measures of the strength of different methods for calculating reference evapotranspiration, 1976-1980.

Equation	R ²	RMSD
Har	0.8344	1.4580
dailyHar	0.8345	1.1315
MH	0.8610	1.2538
dailyMH	0.8584	1.0597

4.1 Seasonal Evaluation

Table 13 presents the RMSD values for all seasons at the global scale. Here, as in the 1971-1975 period, the optimized daily methods out-performed the previous methods for calculating reference evapotranspiration. In all cases, the dailyMH equation provided the lowest RMSD. In addition, we see a peak in RMSD around the northern-hemisphere summer (JJA). Weakened estival improvements are seen as well. The corresponding correlation coefficients can be seen in Appendix, Table A9. As similarly noted in the optimization period, the dailyMH improvement on the MH equation in DJF and SON (19.94% and 20.43%, respectively) is almost double the improvement in MAM and JJA (12.89% and 9.94%, respectively).

Table 13. Global, seasonal RMSD values, 1976-1980.

Equation	DJF	MAM	JJA	SON	All
Har	1.3092	1.4278	1.5961	1.5525	1.4580
dailyHar	1.0203	1.1281	1.2687	1.1388	1.1315
MH	1.1676	1.2494	1.3197	1.3287	1.2538
dailyMH	0.9349	1.0883	1.1885	1.0573	1.0597

4.2 Regional Evaluation

Using the same regions defined in Table 4, the results for 1976-1980 have been evaluated by region and season. **Tables 14-20** present the RMSD values of these results (R^2 values can be perused in Appendix, Tables A10-16).

Table 14. Seasonal RMSD values for Region 1, 1976-1980. Instances where RMSD was increased after the optimization are highlighted in red.

Equation	DJF	MAM	JJA	SON	All
Har	0.4355	0.6808	0.9671	0.8960	0.7735
dailyHar	0.4100	0.8724	1.2473	0.7266	0.8690
MH	0.5437	0.7455	0.9375	0.8837	0.7926
dailyMH	0.3970	0.8476	1.1635	0.7065	0.8270

Table 15. Seasonal RMSD values for Region 2, 1976-1980. Instances where RMSD was increased after the optimization are highlighted in red.

Equation	DJF	MAM	JJA	SON	All
Har	0.9171	1.0872	1.7901	1.6498	1.4102
dailyHar	0.7601	1.0196	1.3632	1.1540	1.0967
MH	0.9241	0.9565	1.3463	1.3937	1.1748
dailyMH	0.7194	1.0027	1.3127	1.0568	1.0448

Table 16. Seasonal RMSD values for Region 3, 1976-1980.

Equation	DJF	MAM	JJA	SON	All
Har	1.7756	2.2232	2.3948	2.2380	2.1706
dailyHar	1.1981	1.4826	1.6630	1.5039	1.4718
MH	1.4669	1.6628	1.7312	1.7501	1.6562
dailyMH	1.1039	1.4351	1.5135	1.3618	1.3626

Table 17. Seasonal RMSD values for Region 4, 1976-1980.

Equation	DJF	MAM	JJA	SON	All
Har	1.2486	1.1897	1.1488	1.1111	1.1740
dailyHar	1.0741	0.9801	0.8724	0.9856	0.9789
MH	1.2718	1.2814	1.2376	1.2439	1.2578
dailyMH	0.9996	0.9806	0.8585	0.9631	0.9504

Table 18. Seasonal RMSD values for Region 5, 1976-1980.

Equation	DJF	MAM	JJA	SON	All
Har	2.1208	2.0802	1.9273	2.1731	2.0759
dailyHar	1.6433	1.5191	1.3691	1.5449	1.5210
MH	1.6309	1.7526	1.5971	1.5987	1.6457
dailyMH	1.4176	1.3761	1.2342	1.3873	1.3546

Table 19. Seasonal RMSD values for Region 6, 1976-1980.

Equation	DJF	MAM	JJA	SON	All
Har	1.8590	1.5589	1.1792	1.3493	1.5062
dailyHar	1.4016	1.1128	0.9042	1.1100	1.1444
MH	1.5751	1.4118	1.1470	1.1967	1.3426
dailyMH	1.3495	1.0662	0.8872	1.0779	1.1061

Table 20. Seasonal RMSD values for Region 7, 1976-1980. Instances where RMSD was increased after the optimization are highlighted in red.

Equation	DJF	MAM	JJA	SON	All
Har	0.9752	0.7352	0.5324	0.6702	0.7444
dailyHar	1.2668	0.6630	0.4947	0.8830	0.8732
MH	1.0210	0.7986	0.5720	0.7194	0.7933
dailyMH	1.1670	0.6619	0.5008	0.8115	0.8207

The results from the 1976-1980 period are almost identical to the results of the 1971-1975 period, lending strength to the validity of the optimized dailyMH method. Again we see a slight weakening of the estimates around the summer months for all regions experiencing some seasonality. In addition, we see an increase of RMSD in the pole-ward latitudes overall and especially in the summer months, though the magnitude of the RMSDs remains small (owing, in part, to the fact that the magnitude of reference evapotranspiration will be lower in these regions).

While degradations in the pole-ward estimates are clear (red text in tables) a glance at the percent improvement highlights the estival reduction in improvement between dailyMH and MH. For region two the JJA is only 2.50%, while the DJF improvement is some 22.15%. For region six the improvements are 24.48% and 22.65% in MAM and JJA, respectively, compared with 14.32% and 9.93% in DJF and SON, respectively. Region five shows improvements of 21.48% and 22.65% in MAM and JJA, respectively, with only 13.08% and 13.22% in DJF and SON, respectively. Finally, region three shows a 24.75% improvement in DJF and a 22.18% improvement in SON, matched against 13.70% and 12.57% improvements in MAM and JJA, respectively. In all cases, the estival improvements are much less than the other two seasons.

The similarity of these results to the optimized period lends support to the effectiveness of the methods employed in this study. Overall, the daily Modified-Hargreaves equation has been demonstrated to be the most consistently performing algorithm at reproducing the Modified Penman-Monteith reference evapotranspiration estimate, but with fewer and more readily available input variables required.

5. CLOSING REMARKS

Overall, these results indicate that, of all the methods examined, the daily Modified Hargreaves method is the most accurate reproduction of the Modified Penman-Monteith approach. Two major concerns have been noted: estival and pole-ward inaccuracies, but the significance of these results is a further consideration.

In order to confidently advocate a single method, efforts were made to test the significance of inaccuracies. **Table 21** represents the results of stringent significance testing. Using a one-tailed, paired t-test, the estimates were tested against the Modified Penman-Monteith value at the 1% significance level. Each equation was tested against the other three in that region. In Table 21, only the equation that performed significantly better than all other equations is noted; a single equation was most significant in all cases.

Again, we see that at the 1% significance level the dailyMH equation is not the strongest in pole-ward summers. The dailyMH equation even fails throughout the southernmost region. These results may be of some concern for individuals focusing solely in these regions. On the whole, it is more important that the dailyMH equation succeeds across all temporal regions when looking at the globe. This result, at the 1% significance level, suggests that it is indeed justified to use the dailyMH equation, over the other methods considered here, for the estimation of daily reference evapotranspiration rates. In particular, this method performs as the most effective surrogate to the Modified Penman-Monteith method, while requiring much fewer input requirements.

Table 21. The results of significance testing, noting the most significantly accurate equation.

Best Equation	DJF	MAM	JJA	SON	Annual
Region 1	dailyMH	Har	MH	dailyMH	Har
Region 2	dailyMH	MH	MH	dailyMH	dailyMH
Region 3	dailyMH	dailyMH	dailyMH	dailyMH	dailyMH
Region 4	dailyMH	dailyMH	dailyMH	dailyMH	dailyMH
Region 5	dailyMH	dailyMH	dailyMH	dailyMH	dailyMH
Region 6	dailyMH	dailyMH	dailyMH	dailyMH	dailyMH
Region 7	Har	dailyHar	dailyHar	Har	Har
Global	dailyMH	dailyMH	dailyMH	dailyMH	dailyMH

This new, fitted, daily Modified-Hargreaves (6) is able to predict daily reference evapotranspiration with some measure of increased confidence, as a supplement to the Modified Penman-Monteith approach, across the globe, allowing for previously-noted caveats. RMSDs of one millimeter/day are considered an acceptable level of error, due to the uncertainties of daily data and estimates.

$$ET_0 = 0.0019 \cdot 0.408RA \cdot (T_{avg} + 21.0584) (TD - 0.0874P)^{0.6278} \quad (6)$$

Figure 1 displays the sample of the original Modified-Hargreaves calculations against the Modified Penman-Monteith equations in red. The blue points are the results of the fitted daily Modified-Hargreaves. The upward movement symbolizes the reduction of the RMSD. In addition, it is evident that the equation is stronger, as a surrogate to Modified Penman-Monteith, in the region of lower reference evapotranspiration.




Figure 1. A sampling (0.001% of 25,252,525 estimates) of Modified-Hargreaves (blue) and fitted Modified-Hargreaves (red) against the Modified Penman-Monteith calculations of reference evapotranspiration.

This method allows researchers to confidently assess the daily rates of reference evapotranspiration, without the use of formulae requiring inputs not commonly observed. These calculations can be used to more accurately calibrate and run daily crop and impact models, and be applied to investigations of potential climate changes in all areas of water resources engineering, planning and development.

Following the methods of Droogers and Allen (2002), there may be a way to introduce a new parameter in an effort to strengthen the accuracy of daily calculations. For this exercise, this was considered to complicate the problem. Efforts were made to use a similar form because, as Droogers and Allen (2002) note, other climate variables are not available with much certainty.

The results of this exercise are far from a perfect representation of methods for calculating daily reference evapotranspiration. Further analysis is always advocated. Firstly, the oddities

seen in high latitudes and summer months could be the result of unfair weighting in the RMSD. The low magnitude of evapotranspiration rates in the high latitudes cause this particular error statistic to focus on observations with larger discrepancies in real-space. Future analyses should explore the possibility for optimizing coefficients based on the RMSD calculated in a space similar to log-space. Allowing the presence of zero values prohibits the use of log-space, but a similar transformation may reduce the effects of this unfair weighting.

Also in the realm of further research: it may be of some interest to measure how much the coefficients would change in the daily equations if those parameters were re-evaluated for each region and season. For this experiment, the interest was in developing a global approach to estimating daily evapotranspiration. It was thus decided that a recalibration for each arbitrary region and season would result in too much complexity and too many equations, with marginal rates of return. It may be that recalibration is warranted for highly localized studies.

Finally, it would be of some interest to understand when monthly methods should be used instead of the daily methods. Calculating the average monthly reference evapotranspiration from the Modified-Hargreaves approach by using monthly data and comparing it to strictly daily results from the daily Modified-Hargreaves method with daily data might shed some light on appropriate use of these equations. This understanding of when monthly data is more applicable than daily data may be of extreme importance to future modeling efforts.

Acknowledgments

The authors gratefully acknowledge the financial support for this work provided by the MIT Joint Program on the Science and Policy of Global Change through a consortium of industrial sponsors and Federal grants.

6. REFERENCES

- Allen, R.G., L.S. Pereira, D. Raes & M. Smith, 1998: *Crop evapotranspiration: Guidelines for computing crop requirements*. Irrigation and Drainage Paper No. 56, FAO, Rome, Italy.
- Droogers, P., & R.G. Allen, 2002: Estimating reference evapotranspiration under inaccurate data conditions. *Irrigation and Drainage Systems* 16: 33-45.
- Hargreaves, G.L., G.H. Hargreaves & J.P. Riley, 1985: Agricultural benefits for Senegal River basin. *Journal of Irrigation and Drainage Engineering*, ASCE 111(2): 113-124.
- McKenney, M.S., and N.J. Rosenberg, 1993: Sensitivity of some potential evapotranspiration estimation methods to climate change. *Agricultural and Forest Methodology* 64:81-110.
- Ngo-Duc, T., Polcher, J., & K. Laval, 2005: A 53-year forcing data set for land surface models, *Journal of Geophysical Research*, 110(D06116): 1-13.
- Oleson, K. W., Y. Dai, G. Bonan, M. Bosilovich, R. Dickinson, P. Dirmeyer, F. Hoffman, P. Houser, S. Levis, G.-Y. Niu, P. Thornton, M. Vertenstein, Z.-L. Yang and X. Zeng, 2004:

Technical description of the Community Land Model (CLM), National Center for Atmospheric Research *Tech. Note NCAR/TN-461+STR*, 173 pp.

Schlosser, C.A., and X. Gao, 2010: Assessing Evapotranspiration Estimates from the Global Soil Wetness Project Phase 2 (GSWP-2) Simulations, *Journal of Hydrometeorology*, 11: 880–897.

APPENDIX

This appendix contains tables of all the R^2 values, matching the tables of RMSD in the text.

Table A 1. Global, seasonal R^2 values, 1971-1975.

Equation	DJF	MAM	JJA	SON	All
Har	0.8624	0.8330	0.8046	0.8130	0.8319
dailyHar	0.8634	0.8330	0.8038	0.8140	0.8319
MH	0.8913	0.8445	0.8302	0.8517	0.8590
dailyMH	0.8904	0.8447	0.8259	0.8473	0.8562

Table A 2. Seasonal R^2 values for Region 1, 1971-1975.

Equation	DJF	MAM	JJA	SON	All
Har	0.1353	0.7245	0.6963	0.7425	0.8260
dailyHar	0.1507	0.7276	0.6965	0.7447	0.8256
MH	0.1534	0.7364	0.7280	0.7690	0.8458
dailyMH	0.1870	0.7515	0.7303	0.7739	0.8460

Table A 3. Seasonal R^2 values for Region 2, 1971-1975.

Equation	DJF	MAM	JJA	SON	All
Har	0.4955	0.7838	0.8030	0.7802	0.8359
dailyHar	0.5044	0.7865	0.8028	0.7829	0.8357
MH	0.5413	0.8125	0.8003	0.8217	0.8637
dailyMH	0.5664	0.8149	0.8052	0.8204	0.8592

Table A 4. Seasonal R^2 values for Region 3, 1971-1975.

Equation	DJF	MAM	JJA	SON	All
Har	0.7367	0.7862	0.8432	0.7786	0.8046
dailyHar	0.7389	0.7868	0.8431	0.7803	0.8057
MH	0.7448	0.7789	0.8362	0.8085	0.8238
dailyMH	0.7503	0.7898	0.8440	0.8096	0.8275

Table A 5. Seasonal R^2 values for Region 4, 1971-1975.

Equation	DJF	MAM	JJA	SON	All
Har	0.6350	0.7752	0.7308	0.6688	0.7077
dailyHar	0.6360	0.7749	0.7320	0.6686	0.7081
MH	0.6951	0.7866	0.7637	0.6956	0.7400
dailyMH	0.6853	0.7845	0.7617	0.6916	0.7354

Table A 6. Seasonal R^2 values for Region 5, 1971-1975.

Equation	DJF	MAM	JJA	SON	All
Har	0.8197	0.6068	0.5588	0.6966	0.6961
dailyHar	0.8206	0.6115	0.5658	0.7005	0.6989
MH	0.8301	0.7180	0.6293	0.7381	0.7725
dailyMH	0.8244	0.6976	0.6251	0.7390	0.7610

Table A 7. Seasonal R^2 values for Region 6, 1971-1975.

Equation	DJF	MAM	JJA	SON	All
Har	0.7181	0.7329	0.5983	0.7334	0.7906
dailyHar	0.7184	0.7345	0.6003	0.7339	0.7906
MH	0.7298	0.7751	0.6340	0.7505	0.8142
dailyMH	0.7367	0.7741	0.6403	0.7552	0.8147

Table A 8. Seasonal R^2 values for Region 7, 1971-1975.

Equation	DJF	MAM	JJA	SON	All
Har	0.5746	0.6151	0.0789	0.6216	0.7086
dailyHar	0.5714	0.6141	0.0807	0.6198	0.7066
MH	0.5883	0.6374	0.1039	0.6311	0.7208
dailyMH	0.5869	0.6370	0.1113	0.6347	0.7197

Table A 9. Global, seasonal R^2 values, 1976-1980.

Equation	DJF	MAM	JJA	SON	All
Har	0.8653	0.8323	0.8065	0.8179	0.8344
dailyHar	0.8662	0.8323	0.8058	0.8189	0.8345
MH	0.8940	0.8435	0.8332	0.8557	0.8610
dailyMH	0.8927	0.8438	0.8289	0.8516	0.8584

Table A 10. Seasonal R^2 values for Region 1, 1976-1980.

Equation	DJF	MAM	JJA	SON	All
Har	0.1088	0.7332	0.6827	0.7311	0.8248
dailyHar	0.1244	0.7364	0.6828	0.7334	0.8246
MH	0.1372	0.7433	0.7146	0.7598	0.8425
dailyMH	0.1627	0.7583	0.7175	0.7646	0.8436

Table A 11. Seasonal R^2 values for Region 2, 1976-1980.

Equation	DJF	MAM	JJA	SON	All
Har	0.5364	0.7856	0.8063	0.7921	0.8404
dailyHar	0.5440	0.7884	0.8062	0.7945	0.8402
MH	0.5793	0.8136	0.8065	0.8275	0.8670
dailyMH	0.5986	0.8168	0.8115	0.8271	0.8629

Table A 12. Seasonal R^2 values for Region 3, 1976-1980.

Equation	DJF	MAM	JJA	SON	All
Har	0.7401	0.7864	0.8499	0.7848	0.8109
dailyHar	0.7420	0.7870	0.8497	0.7863	0.8118
MH	0.7502	0.7799	0.8435	0.8133	0.8295
dailyMH	0.7540	0.7903	0.8513	0.8138	0.8329

Table A 13. Seasonal R² values for Region 4, 1976-1980.

Equation	DJF	MAM	JJA	SON	All
Har	0.6467	0.7755	0.7085	0.6503	0.7023
dailyHar	0.6475	0.7752	0.7096	0.6499	0.7026
MH	0.7035	0.7848	0.7415	0.6755	0.7327
dailyMH	0.6945	0.7845	0.7380	0.6710	0.7285

Table A 14. Seasonal R² values for Region 5, 1976-1980.

Equation	DJF	MAM	JJA	SON	All
Har	0.8240	0.6417	0.5303	0.6979	0.6959
dailyHar	0.8247	0.6457	0.5378	0.7020	0.6986
MH	0.8316	0.7387	0.6051	0.7433	0.7726
dailyMH	0.8266	0.7210	0.5995	0.7422	0.7602

Table A 15. Seasonal R² values for Region 6, 1976-1980.

Equation	DJF	MAM	JJA	SON	All
Har	0.7510	0.7440	0.6120	0.7458	0.8004
dailyHar	0.7510	0.7456	0.6142	0.7462	0.8003
MH	0.7590	0.7821	0.6483	0.7665	0.8254
dailyMH	0.7640	0.7821	0.6528	0.7707	0.8247

Table A 16. Seasonal R² values for Region 7, 1976-1980.

Equation	DJF	MAM	JJA	SON	All
Har	0.5482	0.5604	0.0545	0.5875	0.7021
dailyHar	0.5456	0.5605	0.0573	0.5874	0.7009
MH	0.5590	0.5804	0.0761	0.5920	0.7097
dailyMH	0.5606	0.5853	0.0869	0.6023	0.7129

REPORT SERIES of the MIT Joint Program on the Science and Policy of Global Change

1. **Uncertainty in Climate Change Policy Analysis**
Jacoby & Prinn December 1994
2. **Description and Validation of the MIT Version of the GISS 2D Model** *Sokolov & Stone* June 1995
3. **Responses of Primary Production and Carbon Storage to Changes in Climate and Atmospheric CO₂ Concentration** *Xiao et al.* October 1995
4. **Application of the Probabilistic Collocation Method for an Uncertainty Analysis** *Webster et al.* January 1996
5. **World Energy Consumption and CO₂ Emissions: 1950-2050** *Schmalensee et al.* April 1996
6. **The MIT Emission Prediction and Policy Analysis (EPPA) Model** *Yang et al.* May 1996 (*superseded* by No. 125)
7. **Integrated Global System Model for Climate Policy Analysis** *Prinn et al.* June 1996 (*superseded* by No. 124)
8. **Relative Roles of Changes in CO₂ and Climate to Equilibrium Responses of Net Primary Production and Carbon Storage** *Xiao et al.* June 1996
9. **CO₂ Emissions Limits: Economic Adjustments and the Distribution of Burdens** *Jacoby et al.* July 1997
10. **Modeling the Emissions of N₂O and CH₄ from the Terrestrial Biosphere to the Atmosphere** *Liu* Aug. 1996
11. **Global Warming Projections: Sensitivity to Deep Ocean Mixing** *Sokolov & Stone* September 1996
12. **Net Primary Production of Ecosystems in China and its Equilibrium Responses to Climate Changes**
Xiao et al. November 1996
13. **Greenhouse Policy Architectures and Institutions**
Schmalensee November 1996
14. **What Does Stabilizing Greenhouse Gas Concentrations Mean?** *Jacoby et al.* November 1996
15. **Economic Assessment of CO₂ Capture and Disposal**
Eckaus et al. December 1996
16. **What Drives Deforestation in the Brazilian Amazon?**
Pfaff December 1996
17. **A Flexible Climate Model For Use In Integrated Assessments** *Sokolov & Stone* March 1997
18. **Transient Climate Change and Potential Croplands of the World in the 21st Century** *Xiao et al.* May 1997
19. **Joint Implementation: Lessons from Title IV's Voluntary Compliance Programs** *Atkeson* June 1997
20. **Parameterization of Urban Subgrid Scale Processes in Global Atm. Chemistry Models** *Calbo et al.* July 1997
21. **Needed: A Realistic Strategy for Global Warming**
Jacoby, Prinn & Schmalensee August 1997
22. **Same Science, Differing Policies; The Saga of Global Climate Change** *Skolnikoff* August 1997
23. **Uncertainty in the Oceanic Heat and Carbon Uptake and their Impact on Climate Projections**
Sokolov et al. September 1997
24. **A Global Interactive Chemistry and Climate Model**
Wang, Prinn & Sokolov September 1997
25. **Interactions Among Emissions, Atmospheric Chemistry & Climate Change** *Wang & Prinn* Sept. 1997
26. **Necessary Conditions for Stabilization Agreements**
Yang & Jacoby October 1997
27. **Annex I Differentiation Proposals: Implications for Welfare, Equity and Policy** *Reiner & Jacoby* Oct. 1997
28. **Transient Climate Change and Net Ecosystem Production of the Terrestrial Biosphere**
Xiao et al. November 1997
29. **Analysis of CO₂ Emissions from Fossil Fuel in Korea: 1961-1994** *Choi* November 1997
30. **Uncertainty in Future Carbon Emissions: A Preliminary Exploration** *Webster* November 1997
31. **Beyond Emissions Paths: Rethinking the Climate Impacts of Emissions Protocols** *Webster & Reiner* November 1997
32. **Kyoto's Unfinished Business** *Jacoby et al.* June 1998
33. **Economic Development and the Structure of the Demand for Commercial Energy** *Judson et al.* April 1998
34. **Combined Effects of Anthropogenic Emissions and Resultant Climatic Changes on Atmospheric OH**
Wang & Prinn April 1998
35. **Impact of Emissions, Chemistry, and Climate on Atmospheric Carbon Monoxide** *Wang & Prinn* April 1998
36. **Integrated Global System Model for Climate Policy Assessment: Feedbacks and Sensitivity Studies**
Prinn et al. June 1998
37. **Quantifying the Uncertainty in Climate Predictions**
Webster & Sokolov July 1998
38. **Sequential Climate Decisions Under Uncertainty: An Integrated Framework** *Valverde et al.* September 1998
39. **Uncertainty in Atmospheric CO₂ (Ocean Carbon Cycle Model Analysis)** *Holian* Oct. 1998 (*superseded* by No. 80)
40. **Analysis of Post-Kyoto CO₂ Emissions Trading Using Marginal Abatement Curves** *Ellerman & Decaux* Oct. 1998
41. **The Effects on Developing Countries of the Kyoto Protocol and CO₂ Emissions Trading**
Ellerman et al. November 1998
42. **Obstacles to Global CO₂ Trading: A Familiar Problem**
Ellerman November 1998
43. **The Uses and Misuses of Technology Development as a Component of Climate Policy** *Jacoby* November 1998
44. **Primary Aluminum Production: Climate Policy, Emissions and Costs** *Harnisch et al.* December 1998
45. **Multi-Gas Assessment of the Kyoto Protocol**
Reilly et al. January 1999
46. **From Science to Policy: The Science-Related Politics of Climate Change Policy in the U.S.** *Skolnikoff* January 1999
47. **Constraining Uncertainties in Climate Models Using Climate Change Detection Techniques**
Forest et al. April 1999
48. **Adjusting to Policy Expectations in Climate Change Modeling** *Shackley et al.* May 1999
49. **Toward a Useful Architecture for Climate Change Negotiations** *Jacoby et al.* May 1999
50. **A Study of the Effects of Natural Fertility, Weather and Productive Inputs in Chinese Agriculture**
Eckaus & Tso July 1999
51. **Japanese Nuclear Power and the Kyoto Agreement**
Babiker, Reilly & Ellerman August 1999
52. **Interactive Chemistry and Climate Models in Global Change Studies** *Wang & Prinn* September 1999

Contact the Joint Program Office to request a copy. The Report Series is distributed at no charge.

REPORT SERIES of the **MIT Joint Program on the Science and Policy of Global Change**

53. **Developing Country Effects of Kyoto-Type Emissions Restrictions** Babiker & Jacoby October 1999
54. **Model Estimates of the Mass Balance of the Greenland and Antarctic Ice Sheets** Bugnion Oct 1999
55. **Changes in Sea-Level Associated with Modifications of Ice Sheets over 21st Century** Bugnion October 1999
56. **The Kyoto Protocol and Developing Countries** Babiker et al. October 1999
57. **Can EPA Regulate Greenhouse Gases Before the Senate Ratifies the Kyoto Protocol?** Bugnion & Reiner November 1999
58. **Multiple Gas Control Under the Kyoto Agreement** Reilly, Mayer & Harnisch March 2000
59. **Supplementarity: An Invitation for Monopsony?** Ellerman & Sue Wing April 2000
60. **A Coupled Atmosphere-Ocean Model of Intermediate Complexity** Kamenkovich et al. May 2000
61. **Effects of Differentiating Climate Policy by Sector: A U.S. Example** Babiker et al. May 2000
62. **Constraining Climate Model Properties Using Optimal Fingerprint Detection Methods** Forest et al. May 2000
63. **Linking Local Air Pollution to Global Chemistry and Climate** Mayer et al. June 2000
64. **The Effects of Changing Consumption Patterns on the Costs of Emission Restrictions** Lahiri et al. Aug 2000
65. **Rethinking the Kyoto Emissions Targets** Babiker & Eckaus August 2000
66. **Fair Trade and Harmonization of Climate Change Policies in Europe** Viguier September 2000
67. **The Curious Role of "Learning" in Climate Policy: Should We Wait for More Data?** Webster October 2000
68. **How to Think About Human Influence on Climate** Forest, Stone & Jacoby October 2000
69. **Tradable Permits for Greenhouse Gas Emissions: A primer with reference to Europe** Ellerman Nov 2000
70. **Carbon Emissions and The Kyoto Commitment in the European Union** Viguier et al. February 2001
71. **The MIT Emissions Prediction and Policy Analysis Model: Revisions, Sensitivities and Results** Babiker et al. February 2001 (*superseded* by No. 125)
72. **Cap and Trade Policies in the Presence of Monopoly and Distortionary Taxation** Fullerton & Metcalf March '01
73. **Uncertainty Analysis of Global Climate Change Projections** Webster et al. Mar. '01 (*superseded* by No. 95)
74. **The Welfare Costs of Hybrid Carbon Policies in the European Union** Babiker et al. June 2001
75. **Feedbacks Affecting the Response of the Thermohaline Circulation to Increasing CO₂** Kamenkovich et al. July 2001
76. **CO₂ Abatement by Multi-fueled Electric Utilities: An Analysis Based on Japanese Data** Ellerman & Tsukada July 2001
77. **Comparing Greenhouse Gases** Reilly et al. July 2001
78. **Quantifying Uncertainties in Climate System Properties using Recent Climate Observations** Forest et al. July 2001
79. **Uncertainty in Emissions Projections for Climate Models** Webster et al. August 2001
80. **Uncertainty in Atmospheric CO₂ Predictions from a Global Ocean Carbon Cycle Model** Holian et al. September 2001
81. **A Comparison of the Behavior of AO GCMs in Transient Climate Change Experiments** Sokolov et al. December 2001
82. **The Evolution of a Climate Regime: Kyoto to Marrakech** Babiker, Jacoby & Reiner February 2002
83. **The "Safety Valve" and Climate Policy** Jacoby & Ellerman February 2002
84. **A Modeling Study on the Climate Impacts of Black Carbon Aerosols** Wang March 2002
85. **Tax Distortions and Global Climate Policy** Babiker et al. May 2002
86. **Incentive-based Approaches for Mitigating Greenhouse Gas Emissions: Issues and Prospects for India** Gupta June 2002
87. **Deep-Ocean Heat Uptake in an Ocean GCM with Idealized Geometry** Huang, Stone & Hill September 2002
88. **The Deep-Ocean Heat Uptake in Transient Climate Change** Huang et al. September 2002
89. **Representing Energy Technologies in Top-down Economic Models using Bottom-up Information** McFarland et al. October 2002
90. **Ozone Effects on Net Primary Production and Carbon Sequestration in the U.S. Using a Biogeochemistry Model** Felzer et al. November 2002
91. **Exclusionary Manipulation of Carbon Permit Markets: A Laboratory Test** Carlén November 2002
92. **An Issue of Permanence: Assessing the Effectiveness of Temporary Carbon Storage** Herzog et al. December 2002
93. **Is International Emissions Trading Always Beneficial?** Babiker et al. December 2002
94. **Modeling Non-CO₂ Greenhouse Gas Abatement** Hyman et al. December 2002
95. **Uncertainty Analysis of Climate Change and Policy Response** Webster et al. December 2002
96. **Market Power in International Carbon Emissions Trading: A Laboratory Test** Carlén January 2003
97. **Emissions Trading to Reduce Greenhouse Gas Emissions in the United States: The McCain-Lieberman Proposal** Paltsev et al. June 2003
98. **Russia's Role in the Kyoto Protocol** Bernard et al. Jun '03
99. **Thermohaline Circulation Stability: A Box Model Study** Lucarini & Stone June 2003
100. **Absolute vs. Intensity-Based Emissions Caps** Ellerman & Sue Wing July 2003
101. **Technology Detail in a Multi-Sector CGE Model: Transport Under Climate Policy** Schafer & Jacoby July 2003
102. **Induced Technical Change and the Cost of Climate Policy** Sue Wing September 2003
103. **Past and Future Effects of Ozone on Net Primary Production and Carbon Sequestration Using a Global Biogeochemical Model** Felzer et al. (revised) January 2004

Contact the Joint Program Office to request a copy. The Report Series is distributed at no charge.

REPORT SERIES of the MIT Joint Program on the Science and Policy of Global Change

- 104. A Modeling Analysis of Methane Exchanges Between Alaskan Ecosystems and the Atmosphere** Zhuang *et al.* November 2003
- 105. Analysis of Strategies of Companies under Carbon Constraint** Hashimoto January 2004
- 106. Climate Prediction: The Limits of Ocean Models** Stone February 2004
- 107. Informing Climate Policy Given Incommensurable Benefits Estimates** Jacoby February 2004
- 108. Methane Fluxes Between Terrestrial Ecosystems and the Atmosphere at High Latitudes During the Past Century** Zhuang *et al.* March 2004
- 109. Sensitivity of Climate to Diapycnal Diffusivity in the Ocean** Dalan *et al.* May 2004
- 110. Stabilization and Global Climate Policy** Sarofim *et al.* July 2004
- 111. Technology and Technical Change in the MIT EPPA Model** Jacoby *et al.* July 2004
- 112. The Cost of Kyoto Protocol Targets: The Case of Japan** Paltsev *et al.* July 2004
- 113. Economic Benefits of Air Pollution Regulation in the USA: An Integrated Approach** Yang *et al.* (revised) Jan. 2005
- 114. The Role of Non-CO₂ Greenhouse Gases in Climate Policy: Analysis Using the MIT IGSM** Reilly *et al.* Aug. '04
- 115. Future U.S. Energy Security Concerns** Deutch Sep. '04
- 116. Explaining Long-Run Changes in the Energy Intensity of the U.S. Economy** Sue Wing Sept. 2004
- 117. Modeling the Transport Sector: The Role of Existing Fuel Taxes in Climate Policy** Paltsev *et al.* November 2004
- 118. Effects of Air Pollution Control on Climate** Prinn *et al.* January 2005
- 119. Does Model Sensitivity to Changes in CO₂ Provide a Measure of Sensitivity to the Forcing of Different Nature?** Sokolov March 2005
- 120. What Should the Government Do To Encourage Technical Change in the Energy Sector?** Deutch May '05
- 121. Climate Change Taxes and Energy Efficiency in Japan** Kasahara *et al.* May 2005
- 122. A 3D Ocean-Seaice-Carbon Cycle Model and its Coupling to a 2D Atmospheric Model: Uses in Climate Change Studies** Dutkiewicz *et al.* (revised) November 2005
- 123. Simulating the Spatial Distribution of Population and Emissions to 2100** Asadoorian May 2005
- 124. MIT Integrated Global System Model (IGSM) Version 2: Model Description and Baseline Evaluation** Sokolov *et al.* July 2005
- 125. The MIT Emissions Prediction and Policy Analysis (EPPA) Model: Version 4** Paltsev *et al.* August 2005
- 126. Estimated PDFs of Climate System Properties Including Natural and Anthropogenic Forcings** Forest *et al.* September 2005
- 127. An Analysis of the European Emission Trading Scheme** Reilly & Paltsev October 2005
- 128. Evaluating the Use of Ocean Models of Different Complexity in Climate Change Studies** Sokolov *et al.* November 2005
- 129. Future Carbon Regulations and Current Investments in Alternative Coal-Fired Power Plant Designs** Sekar *et al.* December 2005
- 130. Absolute vs. Intensity Limits for CO₂ Emission Control: Performance Under Uncertainty** Sue Wing *et al.* January 2006
- 131. The Economic Impacts of Climate Change: Evidence from Agricultural Profits and Random Fluctuations in Weather** Deschenes & Greenstone January 2006
- 132. The Value of Emissions Trading** Webster *et al.* Feb. 2006
- 133. Estimating Probability Distributions from Complex Models with Bifurcations: The Case of Ocean Circulation Collapse** Webster *et al.* March 2006
- 134. Directed Technical Change and Climate Policy** Otto *et al.* April 2006
- 135. Modeling Climate Feedbacks to Energy Demand: The Case of China** Asadoorian *et al.* June 2006
- 136. Bringing Transportation into a Cap-and-Trade Regime** Ellerman, Jacoby & Zimmerman June 2006
- 137. Unemployment Effects of Climate Policy** Babiker & Eckaus July 2006
- 138. Energy Conservation in the United States: Understanding its Role in Climate Policy** Metcalf Aug. '06
- 139. Directed Technical Change and the Adoption of CO₂ Abatement Technology: The Case of CO₂ Capture and Storage** Otto & Reilly August 2006
- 140. The Allocation of European Union Allowances: Lessons, Unifying Themes and General Principles** Buchner *et al.* October 2006
- 141. Over-Allocation or Abatement? A preliminary analysis of the EU ETS based on the 2006 emissions data** Ellerman & Buchner December 2006
- 142. Federal Tax Policy Towards Energy** Metcalf Jan. 2007
- 143. Technical Change, Investment and Energy Intensity** Kratena March 2007
- 144. Heavier Crude, Changing Demand for Petroleum Fuels, Regional Climate Policy, and the Location of Upgrading Capacity** Reilly *et al.* April 2007
- 145. Biomass Energy and Competition for Land** Reilly & Paltsev April 2007
- 146. Assessment of U.S. Cap-and-Trade Proposals** Paltsev *et al.* April 2007
- 147. A Global Land System Framework for Integrated Climate-Change Assessments** Schlosser *et al.* May 2007
- 148. Relative Roles of Climate Sensitivity and Forcing in Defining the Ocean Circulation Response to Climate Change** Scott *et al.* May 2007
- 149. Global Economic Effects of Changes in Crops, Pasture, and Forests due to Changing Climate, CO₂ and Ozone** Reilly *et al.* May 2007
- 150. U.S. GHG Cap-and-Trade Proposals: Application of a Forward-Looking Computable General Equilibrium Model** Gurgel *et al.* June 2007
- 151. Consequences of Considering Carbon/Nitrogen Interactions on the Feedbacks between Climate and the Terrestrial Carbon Cycle** Sokolov *et al.* June 2007

REPORT SERIES of the MIT Joint Program on the Science and Policy of Global Change

- 152. Energy Scenarios for East Asia: 2005-2025** *Paltsev & Reilly* July 2007
- 153. Climate Change, Mortality, and Adaptation: Evidence from Annual Fluctuations in Weather in the U.S.** *Deschênes & Greenstone* August 2007
- 154. Modeling the Prospects for Hydrogen Powered Transportation Through 2100** *Sandoval et al.* February 2008
- 155. Potential Land Use Implications of a Global Biofuels Industry** *Gurgel et al.* March 2008
- 156. Estimating the Economic Cost of Sea-Level Rise** *Sugiyama et al.* April 2008
- 157. Constraining Climate Model Parameters from Observed 20th Century Changes** *Forest et al.* April 2008
- 158. Analysis of the Coal Sector under Carbon Constraints** *McFarland et al.* April 2008
- 159. Impact of Sulfur and Carbonaceous Emissions from International Shipping on Aerosol Distributions and Direct Radiative Forcing** *Wang & Kim* April 2008
- 160. Analysis of U.S. Greenhouse Gas Tax Proposals** *Metcalf et al.* April 2008
- 161. A Forward Looking Version of the MIT Emissions Prediction and Policy Analysis (EPPA) Model** *Babiker et al.* May 2008
- 162. The European Carbon Market in Action: Lessons from the first trading period** Interim Report *Convery, Ellerman, & de Perthuis* June 2008
- 163. The Influence on Climate Change of Differing Scenarios for Future Development Analyzed Using the MIT Integrated Global System Model** *Prinn et al.* September 2008
- 164. Marginal Abatement Costs and Marginal Welfare Costs for Greenhouse Gas Emissions Reductions: Results from the EPPA Model** *Holak et al.* November 2008
- 165. Uncertainty in Greenhouse Emissions and Costs of Atmospheric Stabilization** *Webster et al.* November 2008
- 166. Sensitivity of Climate Change Projections to Uncertainties in the Estimates of Observed Changes in Deep-Ocean Heat Content** *Sokolov et al.* November 2008
- 167. Sharing the Burden of GHG Reductions** *Jacoby et al.* November 2008
- 168. Unintended Environmental Consequences of a Global Biofuels Program** *Melillo et al.* January 2009
- 169. Probabilistic Forecast for 21st Century Climate Based on Uncertainties in Emissions (without Policy) and Climate Parameters** *Sokolov et al.* January 2009
- 170. The EU's Emissions Trading Scheme: A Proto-type Global System?** *Ellerman* February 2009
- 171. Designing a U.S. Market for CO₂** *Parsons et al.* February 2009
- 172. Prospects for Plug-in Hybrid Electric Vehicles in the United States & Japan: A General Equilibrium Analysis** *Karplus et al.* April 2009
- 173. The Cost of Climate Policy in the United States** *Paltsev et al.* April 2009
- 174. A Semi-Empirical Representation of the Temporal Variation of Total Greenhouse Gas Levels Expressed as Equivalent Levels of Carbon Dioxide** *Huang et al.* June 2009
- 175. Potential Climatic Impacts and Reliability of Very Large Scale Wind Farms** *Wang & Prinn* June 2009
- 176. Biofuels, Climate Policy and the European Vehicle Fleet** *Gitiaux et al.* August 2009
- 177. Global Health and Economic Impacts of Future Ozone Pollution** *Selin et al.* August 2009
- 178. Measuring Welfare Loss Caused by Air Pollution in Europe: A CGE Analysis** *Nam et al.* August 2009
- 179. Assessing Evapotranspiration Estimates from the Global Soil Wetness Project Phase 2 (GSWP-2) Simulations** *Schlosser and Gao* September 2009
- 180. Analysis of Climate Policy Targets under Uncertainty** *Webster et al.* September 2009
- 181. Development of a Fast and Detailed Model of Urban-Scale Chemical and Physical Processing** *Cohen & Prinn* October 2009
- 182. Distributional Impacts of a U.S. Greenhouse Gas Policy: A General Equilibrium Analysis of Carbon Pricing** *Rausch et al.* November 2009
- 183. Canada's Bitumen Industry Under CO₂ Constraints** *Chan et al.* January 2010
- 184. Will Border Carbon Adjustments Work?** *Winchester et al.* February 2010
- 185. Distributional Implications of Alternative U.S. Greenhouse Gas Control Measures** *Rausch et al.* June 2010
- 186. The Future of U.S. Natural Gas Production, Use, and Trade** *Paltsev et al.* June 2010
- 187. Combining a Renewable Portfolio Standard with a Cap-and-Trade Policy: A General Equilibrium Analysis** *Morris et al.* July 2010
- 188. On the Correlation between Forcing and Climate Sensitivity** *Sokolov* August 2010
- 189. Modeling the Global Water Resource System in an Integrated Assessment Modeling Framework: IGSM-WRS** *Strzepek et al.* September 2010
- 190. Climatology and Trends in the Forcing of the Stratospheric Zonal-Mean Flow** *Monier and Weare* January 2011
- 191. Climatology and Trends in the Forcing of the Stratospheric Ozone Transport** *Monier and Weare* January 2011
- 192. The Impact of Border Carbon Adjustments under Alternative Producer Responses** *Winchester* February 2011
- 193. What to Expect from Sectoral Trading: A U.S.-China Example** *Gavard et al.* February 2011
- 194. General Equilibrium, Electricity Generation Technologies and the Cost of Carbon Abatement** *Lanz and Rausch* February 2011
- 195. A Method for Calculating Reference Evapotranspiration on Daily Time Scales** *Farmer et al.* February 2011

Contact the Joint Program Office to request a copy. The Report Series is distributed at no charge.