

Complete Genome Sequence of *Brevibacterium linens* SMQ-1335

Alessandra G. de Melo,^{a,b} Simon J. Labrie,^{a,b} Jeannot Dumaresq,^c Richard J. Roberts,^d Denise M. Tremblay,^{a,b,e} Sylvain Moineau^{a,b,e}

Département de Biochimie, de Microbiologie et de Bio-informatique, Faculté des Sciences et de Génie, Québec, Québec, Canada^a; Groupe de Recherche en Écologie Buccale, Faculté de Médecine Dentaire, Université Laval, Québec, Québec, Canada^b; Département de Microbiologie et d'Infectiologie, Centre Hospitalier Affilié Universitaire Hôtel-Dieu de Lévis, Lévis, Québec, Canada^c; New England Biolabs, Ipswich, Massachusetts, USA^d; Félix d'Hérelle Reference Center for Bacterial Viruses, Faculté de Médecine Dentaire, Université Laval, Québec, Québec, Canada^e

***Brevibacterium linens* is one of the main bacteria found in the smear of surface-ripened cheeses. The genome of the industrial strain SMQ-1335 was sequenced using PacBio. It has 4,209,935 bp, a 62.6% G+C content, 3,848 open reading frames, and 61 structural RNAs. A new type I restriction-modification system was identified.**

Received 9 September 2016 Accepted 21 September 2016 Published 10 November 2016

Citation de Melo AG, Labrie SJ, Dumaresq J, Roberts RJ, Tremblay DM, Moineau S. 2016. Complete genome sequence of *Brevibacterium linens* SMQ-1335. *Genome Announc* 4(6):e01242-16. doi:10.1128/genomeA.01242-16.

Copyright © 2016 de Melo et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Sylvain Moineau, Sylvain.Moineau@bcm.ulaval.ca.

Brevibacterium linens is a Gram-positive bacterium found on the surface of a variety of washed rind cheeses produced globally (1–3). This non-spore-forming, halotolerant, strictly aerobic chemoorganotroph undergoes a rod-cocci cycle during growth and possesses mesodiaminopimelic acid in its cell wall (1, 2, 4). *B. linens* plays key roles during cheese ripening in the breakdown of lipids and proteins (2), the production of volatile sulfur compounds (5, 6), and the development of color due to carotenoid pigment production (1, 2, 4, 7). Furthermore, the production and accumulation of compatible solutes allows *B. linens* to grow in hyperosmotic environments (4). The genomes of three *B. linens* strains are currently available, of which one is complete (GCA_001606005.1) and two are scaffold and draft sequences (GCA_000167575.1 and GCA_000807915.1, respectively).

The genome of *B. linens* SMQ-1335 was sequenced using one SMRT cell in a PacBio RSII sequencer (Génome Québec Innovation Centre, Montréal, QC, Canada), which generated 76,075 raw subreads of an average length of 9,010 bp that provided an average coverage of 162-fold. The genome was assembled into one contig using HGAP (8). BLASR (9) was used to align and preassemble the sequences using the longest reads as seeds to which all the other subreads were recruited and mapped to correct random errors. The Celera assembler (10) was then used to *de novo* assemble these long and corrected reads into contigs. The sequences were refined using Quiver, wherein the raw reads were aligned on the contigs to generate a consensus sequence containing the complete genome. The single contig had redundant ends of 16,018 bp that were removed from one end for the final assembly. The origin of the genome was set upstream of the gene coding for the replication initiator protein DnaA. Gene prediction and annotation was performed using RAST (11) and BLASTp (12). The *B. linens* SMQ-1335 genome has a high G+C content (62.6%) and is composed of 4,209,935 bp, 3,848 genes, and 61 structural RNAs (49 tRNAs and 12 rRNAs).

In vitro tests revealed that *B. linens* SMQ-1335 is sensitive to vancomycin, daptomycin, gentamicin, tetracycline, and rifampin. However, this strain was insensitive to β -lactam antibiotics (penicillin and ceftriaxone), trimethoprim-sulfamethoxazole (TMP-

SMX), and a second-generation fluoroquinolone (ciprofloxacin). A gene likely coding for the lantibiotic linocin was identified using BAGEL3 (13).

No genes coding for known toxins were found in the genome of SMQ-1335 using Virulence Finder (14), Virulence Factor Database (15), and DBETH (16). RAST and PHASTER (17) identified a putative prophage (31,300 bp). Analysis of this prophage sequence revealed many transposases and integrases, suggesting that this may be a nonfunctional prophage.

The methylome (18) of *B. linens* SMQ-1335 was analyzed to identify DNA methyltransferases and restriction endonucleases (19) as well as specificity subunits (20). The new methyltransferase M.Bli1335I, the restriction endonuclease Bli1335IP, and a new type I RM system were assigned, with the recognition site sequence CGGANNNNNNTTC. The SMQ-1335 genome may contain additional RM systems (types II, III, and IV). For the type II, a new restriction endonuclease (Bli1335II) was also assigned with DTGAAT as the recognition sequence.

Accession number(s). The complete genome sequence of *B. linens* SMQ-1335 is available in GenBank under the accession number CP017150.

ACKNOWLEDGMENTS

We thank the Génome Québec Innovation Centre for performing PacBio sequencing and preliminary genome assembly. R.J.R. is a full-time employee of New England Biolabs, a company that sells research reagents such as DNA MTases. S.M. holds a Tier 1 Canada Research Chair in Bacteriophages.

FUNDING INFORMATION

This work, including the efforts of Sylvain Moineau, was funded by Canada Research Chairs (Chaires de recherche du Canada). This work, including the efforts of Sylvain Moineau, was funded by Gouvernement du Canada | Natural Sciences and Engineering Research Council of Canada (NSERC).

This work, including the efforts of Alessandra G. de Melo, was funded by the National Council for Scientific and Technological Development (CNPq-Brazil) in partnership with CALDO (Canada).

REFERENCES

- Motta AS, Brandelli A. 2008. Properties and antimicrobial activity of the smear surface cheese coryneform bacterium *Brevibacterium linens*. *Eur Food Res Technol* 227:1299–1306. <http://dx.doi.org/10.1007/s00217-008-0856-4>.
- Rattray FP, Fox PF. 1999. Aspects of enzymology and biochemical properties of *Brevibacterium linens* relevant to cheese ripening: a review. *J Dairy Sci* 82:891–909. [http://dx.doi.org/10.3168/jds.S0022-0302\(99\)75308-7](http://dx.doi.org/10.3168/jds.S0022-0302(99)75308-7).
- Breed RS. 1957. Family IX. Brevibacteriaceae breed 1953, p 490–505. In Breed RS, Murray EGD, Smith NR (ed), *Bergey's manual of determinative bacteriology*, 7th ed, Williams and Wilkins Co, Baltimore, MD.
- Onraedt A, Soetaert W, Vandamme E. 2005. Industrial importance of the genus *Brevibacterium*. *Biotechnol Lett* 27:527–533. <http://dx.doi.org/10.1007/s10529-005-2878-3>.
- Arfi K, Landaud S, Bonnarme P. 2006. Evidence for distinct L-methionine catabolic pathways in the yeast *Geotrichum candidum* and the bacterium *Brevibacterium linens*. *Appl Environ Microbiol* 72:2155–2162. <http://dx.doi.org/10.1128/AEM.72.3.2155-2162.2006>.
- Bonnarme P, Psoni L, Spinnler HE. 2000. Diversity of L-methionine catabolic pathways in cheese-ripening bacteria. *Appl Environ Microbiol* 66:5514–5517. <http://dx.doi.org/10.1128/AEM.66.12.5514-5517.2000>.
- Kim SH, Park YH, Schmidt-Dannert C, Lee PC. 2010. Redesign, reconstruction, and directed extension of the *Brevibacterium linens* C₄₀ carotenoid pathway in *Escherichia coli*. *Appl Environ Microbiol* 76:5199–5206. <http://dx.doi.org/10.1128/AEM.00263-10>.
- Chin CS, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C, Clum A, Copeland A, Huddleston J, Eichler EE, Turner SW, Korlach J. 2013. Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing data. *Nat Methods* 10:563–569. <http://dx.doi.org/10.1038/nmeth.2474>.
- Chaisson MJ, Tesler G. 2012. Mapping single molecule sequencing reads using basic local alignment with successive refinement (BLASR): application and theory. *BMC Bioinformatics* 13:238. <http://dx.doi.org/10.1186/1471-2105-13-238>.
- Myers EW, Sutton GG, Delcher AL, Dew IM, Fasulo DP, Flanigan MJ, Kravitz SA, Mobarry CM, Reinert KH, Remington KA, Anson EL, Bolanos RA, Chou HH, Jordan CM, Halpern AL, Lonardi S, Beasley EM, Brandon RC, Chen L, Dunn PJ, Lai Z, Liang Y, Nusskern DR, Zhan M, Zhang Q, Zheng X, Rubin GM, Adams MD, Venter JC. 2000. A whole-genome assembly of *Drosophila*. *Science* 287:2196–2204. <http://dx.doi.org/10.1126/science.287.5461.2196>.
- Overbeek R, Olson R, Pusch GD, Olsen GJ, Davis JJ, Disz T, Edwards RA, Gerdes S, Parrello B, Shukla M, Vonstein V, Wattam AR, Xia F, Stevens R. 2014. The SEED and the rapid annotation of microbial genomes using subsystems technology (RAST). *Nucleic Acids Res* 42:D206–D214. <http://dx.doi.org/10.1093/nar/gkt1226>.
- Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ. 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* 25:3389–3402. <http://dx.doi.org/10.1093/nar/25.17.3389>.
- van Heel AJ, de Jong A, Montalbán-López M, Kok J, Kuipers OP. 2013. BAGEL3: automated identification of genes encoding bacteriocins and (non-) bactericidal posttranslationally modified peptides. *Nucleic Acids Res* 41:W448–W453. <http://dx.doi.org/10.1093/nar/gkt391>.
- Joensen KG, Scheutz F, Lund O, Hasman H, Kaas RS, Nielsen EM, Aarestrup FM. 2014. Real-time whole-genome sequencing for routine typing, surveillance, and outbreak detection of verotoxigenic *Escherichia coli*. *J Clin Microbiol* 52:1501–1510. <http://dx.doi.org/10.1128/JCM.03617-13>.
- Chen L, Xiong Z, Sun L, Yang J, Jin Q. 2012. VFDB 2012 update: toward the genetic diversity and molecular evolution of bacterial virulence factors. *Nucleic Acids Res* 40:D641–D645. <http://dx.doi.org/10.1093/nar/gkr989>.
- Chakraborty A, Ghosh S, Chowdhary G, Maulik U, Chakrabarti S. 2012. DBETH: a database of bacterial exotoxins for human. *Nucleic Acids Res* 40:D615–D620. <http://dx.doi.org/10.1093/nar/gkr942>.
- Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, Wishart DS. 2016. PHASTER: a better, faster version of the PHAST phage search tool. *Nucleic Acids Res* 44:W16–W21. <http://dx.doi.org/10.1093/nar/gkw387>.
- Roberts RJ, Carneiro MO, Schatz MC. 2013. The advantages of SMRT sequencing. *Genome Biol* 14:405. <http://dx.doi.org/10.1186/gb-2013-14-6-405>.
- Roberts RJ, Vincze T, Posfai J, Macelis D. 2015. REBASE—a database for DNA restriction and modification: enzymes, genes and genomes. *Nucleic Acids Res* 43:D298–D299. <http://dx.doi.org/10.1093/nar/gku1046>.
- Murray IA, Clark TA, Morgan RD, Boitano M, Anton BP, Luong K, Fomenkov A, Turner SW, Korlach J, Roberts RJ. 2012. The methylomes of six bacteria. *Nucleic Acids Res* 40:11450–11462. <http://dx.doi.org/10.1093/nar/gks891>.