



# Complete Genome Sequence of Ebrios, a Novel T7virus Isolated from the Ebrie Lagoon in Abidjan, Côte d'Ivoire

Solange Ngazoa-Kakou,<sup>a</sup> Cécile Philippe,<sup>b</sup> Denise M. Tremblay,<sup>b,c</sup> Stéphanie Loignon,<sup>b,c</sup> Aristide Koudou,<sup>a</sup> Ahmed Abole,<sup>a</sup> David Ngolo Coulibaly,<sup>a</sup> Stéphane Kan Kouassi,<sup>a</sup> Mireille Kouamé Sina,<sup>a</sup> Serge Aoussi,<sup>a</sup> Mireille Dosso,<sup>a</sup> Sylvain Moineau<sup>b,c</sup>

<sup>a</sup>Plateforme de Biologie Moléculaire, Département Technique & Technologie, Institut Pasteur de Côte d'Ivoire, Abidjan, Côte d'Ivoire

<sup>b</sup>Département de biochimie, de microbiologie et de bio-informatique, Faculté des sciences et de génie, Québec, Canada

<sup>c</sup>Félix d'Hérelle Reference Center for Bacterial Viruses and GREB, Faculté de médecine dentaire, Université Laval, Québec, Canada

**ABSTRACT** The lytic *Escherichia coli* phage Ebrios was isolated from a water sample collected in Ebrie Lagoon on the Adiopodoumé River in Abidjan (Republic of Côte d'Ivoire, West Africa). The linear genome of this *Podoviridae* family member contains 39,752 bp, has a G+C content of 52.9%, is composed of 53 open reading frames, and is related to the *Stenotrophomonas maltophilia* phage IME15.

Lytic phages are currently being reconsidered as alternatives or complements to antibiotics (1) in many countries, including West Africa (2). Here, a new virulent phage was isolated from a water sample collected in April 2016 from Ebrie Lagoon (Abidjan, Republic of Côte d'Ivoire, West Africa). Briefly, the water sample was filtered (0.45  $\mu$ m) and propagated in the *Escherichia coli* C strain using Luria Bertani (LB) broth and incubated overnight at 37°C (3). Several plaque morphologies were visible on LB plates, but, based on genomic DNA restriction profiles, only one distinct phage was isolated. A single plaque was purified three times, and the phage isolated was designated Ebrios. Twenty-seven bacterial strains were screened to establish the host range of phage Ebrios, consisting of *Escherichia coli* (4 strains), *Salmonella* (23 strains), and *Stenotrophomonas* (1 strain). Phage Ebrios did not form plaques on plates of any of the additional tested strains.

Phage Ebrios particles were stained with uranyl acetate (2%) and observed under an electron microscope (4). Electron micrographs revealed an icosahedral capsid (64  $\pm$  2 nm in diameter) with a short tail, suggesting that phage Ebrios belongs to the *Podoviridae* family.

The genomic DNA of phage Ebrios was extracted from a high-titer lysate (10<sup>9</sup> PFU  $\cdot$  ml<sup>-1</sup>) using a plasmid maxikit (Qiagen) with modifications (5). The sequencing library was prepared with the Nextera XT DNA library preparation kit (Illumina), according to the manufacturer's instructions, and sequenced using a MiSeq reagent kit v2 (500 cycles; Illumina) on a MiSeq instrument. *De novo* assembly was performed with the Ray assembler version 2.2.0 (6). The sequences were analyzed using Geneious software version 11.0.5. Gene annotation was performed using the DNA Master (<http://phagesdb.org/DNAMaster/>) workflow with Glimmer (7), GeneMark (8), and ORF Finder. Open reading frames (ORFs) were also manually validated if they contained at least 30 amino acids, had an initiation codon (AUG, UUG, or GUG), and were preceded by a Shine-Dalgarno sequence.

The genome of phage Ebrios has a G+C content of 52.9% and is composed of 39,752 bp and 53 open reading frames. The Ebrios genome is highly related to the genomes of *Stenotrophomonas maltophilia* phage IME15 (9) and *E. coli* podophage T7

Received 5 March 2018 Accepted 21 May 2018 Published 21 June 2018

**Citation** Ngazoa-Kakou S, Philippe C, Tremblay DM, Loignon S, Koudou A, Abole A, Ngolo Coulibaly D, Kan Kouassi S, Kouamé Sina M, Aoussi S, Dosso M, Moineau S. 2018. Complete genome sequence of Ebrios, a novel T7virus isolated from the Ebrie Lagoon in Abidjan, Côte d'Ivoire. *Genome Announc* 6:e00280-18. <https://doi.org/10.1128/genomeA.00280-18>.

**Copyright** © 2018 Ngazoa-Kakou et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Sylvain Moineau, [Sylvain.Moineau@bcm.ulaval.ca](mailto:Sylvain.Moineau@bcm.ulaval.ca).

(10), with average nucleotide identity (ANIb) values, calculated using JSpeciesWS (11), of 95.27% (79.80% query coverage) and 76.85% (66.36% query coverage), respectively. The first nucleotide of the Ebrios genome was defined based on its alignment with the phage IME15 genome.

Functions of gene products were assigned based on results from BLASTP searches (12). Twenty-nine proteins were annotated with a specific function. Bioinformatics analysis indicated that 44 of 53 deduced Ebrios proteins are related to proteins found in the *S. maltophilia* phage IME15. Four of the remaining proteins have homologs in other podophages, namely ORF9 (*Yersinia* phage phiYeO3-12), ORF18 (*Enterobacteria* phage 13a), ORF35 (*E. coli* phage CICC 80001), and ORF42 (*E. coli* phage EG1). ORF10 appears to be unique to phage Ebrios. Other predicted proteins, such as tail fiber (ORF42) or capsid (ORF35) proteins, were similar to those of T7viruses that infect *E. coli*.

Phage Ebrios was deposited in the Félix d'Hérelle Reference Center for Bacterial Viruses ([www.phage.ulaval.ca](http://www.phage.ulaval.ca)) under the catalog number HER552.

**Accession number(s).** The complete genome sequence of phage Ebrios is available in GenBank under the accession number [MG966531](https://doi.org/10.1093/nar/gki487).

## ACKNOWLEDGMENTS

We acknowledge funding from the Pasteur Institute Côte d'Ivoire, the Natural Sciences and Engineering Research Council of Canada, and the Canadian Institutes of Health Research.

S.M. holds the Tier 1 Canada Research Chair in Bacteriophages.

## REFERENCES

1. Abedon ST, Kuhl SJ, Blasdel BG, Kutter EM. 2011. Phage treatment of human infections. *Bacteriophage* 1:66–85. <https://doi.org/10.4161/bact.1.2.15845>.
2. Nagel TE, Chan BK, De Vos D, El-Shibiny A, Kang'ethe EK, Makumi A, Pirnay J-P. 2016. The developing world urgently needs phages to combat pathogenic bacteria. *Front Microbiol* 7:882. <https://doi.org/10.3389/fmicb.2016.00882>.
3. Kakou Ngazoa ES, Kouya D, Koudou AA, Abole A, Guessend KN, Sylla A, Sina Kouamé M, Coulibaly Ngolo D, Kouassi Kan S, Meite S, Aoussi S, Dosso M. 2017. New identification of phages in water and in fishes from Lagoon Ebrié and their application for multidrug strains in Abidjan, West Africa. *World J Pharma Res* 6:84–89.
4. Fortier L-C, Moineau S. 2007. Morphological and genetic diversity of temperate phages in *Clostridium difficile*. *Appl Environ Microbiol* 73:7358–7366. <https://doi.org/10.1128/AEM.00582-07>.
5. Deveau H, Van Calsteren M-R, Moineau S. 2002. Effect of exopolysaccharides on phage-host interactions in *Lactococcus lactis*. *Appl Environ Microbiol* 68:4364–4369. <https://doi.org/10.1128/AEM.68.9.4364-4369.2002>.
6. Boisvert S, Laviolette F, Corbeil J. 2010. Ray: simultaneous assembly of reads from a mix of high-throughput sequencing technologies. *J Comput Biol* 17:1519–1533. <https://doi.org/10.1089/cmb.2009.0238>.
7. Delcher AL, Bratke KA, Powers EC, Salzberg SL. 2007. Identifying bacterial genes and endosymbiont DNA with Glimmer. *Bioinformatics* 23:673–679. <https://doi.org/10.1093/bioinformatics/btm009>.
8. Besemer J, Borodovsky M. 2005. GeneMark: web software for gene finding in prokaryotes, eukaryotes and viruses. *Nucleic Acids Res* 33:W451–W454. <https://doi.org/10.1093/nar/gki487>.
9. Huang Y, Fan H, Pei G, Fan H, Zhang Z, An X, Mi Z, Shi T, Tong Y. 2012. Complete genome sequence of IME15, the first T7-like bacteriophage lytic to pan-antibiotic-resistant *Stenotrophomonas maltophilia*. *J Virol* 86:13839–13840. <https://doi.org/10.1128/JVI.02661-12>.
10. Henn MR, Sullivan MB, Stange-Thomann N, Osburne MS, Berlin AM, Kelly L, Yandava C, Kodira C, Zeng Q, Weiand M, Sparrow T, Saif S, Giannoukos G, Young SK, Nusbaum C, Birren BW, Chisholm SW. 2010. Analysis of high-throughput sequencing and annotation strategies for phage genomes. *PLoS One* 5:e9083. <https://doi.org/10.1371/journal.pone.0009083>.
11. Richter M, Rosselló-Móra R, Oliver Glöckner F, Peplies J. 2016. JSpeciesWS: a Web server for prokaryotic species circumscription based on pairwise genome comparison. *Bioinformatics* 32:929–931. <https://doi.org/10.1093/bioinformatics/btv681>.
12. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *J Mol Biol* 215:403–410. [https://doi.org/10.1016/S0022-2836\(05\)80360-2](https://doi.org/10.1016/S0022-2836(05)80360-2).