

Movement sequencing in normal aging: speech, oro-facial and finger movements

Running title: Speech production in aging

Authors: Bilodeau-Mercure, Mylène¹, Kirouac, Vanessa^{1,2}, Langlois, Nancy^{1,2}, Ouellet, Claudie¹, Gasse, Isabelle, and Tremblay, Pascale^{1,2}

1. Centre de Recherche de l'Institut Universitaire en Santé Mentale de Québec, 2601 de la canardière, Québec City, QC, Canada
2. Département de Réadaptation, Faculté de Médecine, Université Laval, Québec City, QC, Canada

Corresponding author:

Pascale Tremblay, Ph.D.

Département de Réadaptation, Université Laval, 1050 avenue de la Médecine, Québec (QC), CANADA, G1V 0A6

Email: Pascale.Tremblay@fmed.ulaval.ca

Phone: +001 418 663 5000 ext. 4738

Fax: +001 418 656-5476

Introduction

Speech is one of the most distinguishing features of the humankind. The act of speaking is an extremely complex behavior, both cognitively and at the sensorimotor level. It begins with an intention to communicate, continues to the translation of the message into words, which are converted in syllables that in turn need to be ordered serially (*i.e.* sequenced) before articulation can begin. The final output stage of this complex process requires the coordination of multiple sensorimotor components for the production of fluent speech, including the respiratory system, which provides the airflow necessary to set the vocal folds into vibration, the laryngeal muscles that convert the flow of air from the lungs into speech sounds (phonation), and, finally, the supra-laryngeal muscles that change the configuration of the vocal tract to convert the laryngeal output into sequences of vowels and consonants (articulation). In spite of this complexity, the chain of events that leads to the production of speech occurs within several hundreds of milliseconds. Indeed, adult speakers may produce as many as six to nine syllables per second ([Kent 2000](#)). Despite the importance of communication on quality of life, the manner and extent to which speech behaviour, from respiration to articulation, change throughout adulthood, as well as the nature of the cognitive, physiological, and neurobiological mechanisms that underlie these changes is not well understood.

Previous studies have shown age-related changes in voice *fundamental frequency* [F0] (i.e. the acoustical correlate of voice *pitch*, which ranges from low to high) ([Decoster and Debruyne 1997](#); [Honjo and Isshiki 1980](#); [Hunter et al. 2012](#); [Linville 1996](#); [Mueller 1997](#); [Ramig 1983b](#)), which would begin as early as ~50 years ([D'Haeseleer et al. 2011](#)). Older adults also have higher *jitter* – a measure of cycle-to-cycle variation of vocal F0 (i.e. the acoustical correlate of pitch fluctuations) – compared to younger adults ([Wilcox and Horii 1980](#)). In addition to changes in F0 and jitter, voice loudness also changes (decrease) in aging ([Baker et al. 2001](#)), affecting

males more than females ([Goy, Fernandes, Pichora-Fuller, & van Lieshout, 2013](#)). A decline in speech rate has also been reported for the repetition of words or sentences ([Fozo and Watson 1998](#); [Wohlert and Smith 1998](#)) or directed speech ([Duchin and Mysak 1987](#); [Searl et al. 2002](#)), even when pauses between sentences were excluded from the calculation of speech rate, suggesting that the duration of speech sounds becomes longer with age ([Ramig 1983a](#); [Ryan 1972](#)). This is indeed consistent with the results of a few studies that have shown an age-related increase in the duration of individual speech sounds and syllables during repetition of words or sentences ([Morris and Brown 1987](#); [Ryan and Burk 1974](#); [Smith et al. 1987](#)). There is also limited evidence that aging affects speech intelligibility, that is, the capacity to produce speech sounds that can be recognized ([Shuey 1989](#)). In this study, participants were asked to listen and write down to a series of words embedded in a carrier phrase pronounced by young and older adults, and they misunderstood significantly more often the final consonant pronounced by older compared to younger adults, suggesting an age-related decline in speech intelligibility. Consistent with this finding, others researchers have reported that a group of 20 speech-language pathologists rated older adults (67 to 81 years old) as being less intelligible than younger adults (21 to 28 years old) in a diadochokinetic (DDK) task ([Parnell and Amerman 1987](#)).

Though it is clear that the speech system undergoes important changes with age, little is known about the nature and scope of the underlying biological aging mechanisms. One approach to uncover the nature of these mechanisms is to compare aging of speech skills to the aging of other finely controlled movements (such as finger and oro-facial movements). This is particularly relevant given the apparent relationship between speech and finger movements and between speech and oro-facial movements ([Gentilucci 2003](#); [Gentilucci et al. 2008](#); [Tremblay and Gracco 2009](#); [Tremblay and Gracco 2010](#)). Though this is not without some controversy, behavioural studies have shown that when adults manipulate an object at the same time as they produce

syllables, the size of the object manipulated influences the degree of mouth opening, demonstrating a link between hand and speech movements ([Gentilucci 2003](#); [Gentilucci et al. 2008](#)). Recent studies also suggest that several motor preparatory mechanisms, such as motor response selection mechanisms, engage similar neural resources for speech and oro-facial movements ([Tremblay and Gracco 2009](#); [Tremblay and Gracco 2010](#)) as well as for speech and manual movements ([Tremblay et al. 2008](#)). In this context, it is possible that aging of shared motor control mechanisms affects movement control in a general fashion. There is abundant literature documenting a decline of manual motor control with aging ([Aoki and Fukuoka 2010](#); [Cacola et al. 2013](#); [Cousins et al. 1998](#); [Jimenez-Jimenez et al. 2011](#); [Ruiz et al. 2007](#)). For example, it has been shown that older adults are slower in producing different kinds of finger movements, including the production of sequences of 3 to 5 finger taps triggered visually ([Cacola et al. 2013](#)) or tapping a key with one finger as rapidly as possible during 10 seconds ([Cousins et al. 1998](#)). A relationship has also been found between age and response time in a task requiring participants to tap their thumb with their index finger and then with each finger in rapid successions, whereby response time increased with age ([Ruiz et al. 2007](#)). Interestingly, such age effects appear to develop relatively late. For instance, it has been shown that older adults (65-92 years), but not middle-aged adults (40-63 years) were slower than younger adults (18-32 years) in producing rapid multi-finger tapping movements ([Cacola et al. 2013](#)).

Despite evidence of a relationship between speech, oro-facial and manual movements, to our knowledge, no study to date has examined whether aging mechanisms are movement-specific or domain-general. The goal of the present study was therefore to examine the effect of aging on the ability to produce sequences of fine motor actions (speech, oro-facial and manual movements) varying in complexity levels. We hypothesized that all sequences of movements

would be affected by age and by sequence complexity. We also expected to find interactions between age and sequence complexity, with more complex sequences being more affected by aging across all kinds of movements. Because response sequencing is likely a domain-general mechanism, we expected to find similar effects of age on motor sequence complexity across movement types.

1. Methods

1.1. Participants

85 participants were recruited to participate in the study. Of these, nine were excluded (~11%) either due to recording problems during the experiment ($n = 5$), difficulty complying with task demands in a specific condition ($n = 1$), because they did not complete the hearing assessment ($n = 1$), or because they did not meet the inclusion criteria ($n = 2$). The final group therefore contained 76 participants (mean age 52.95 ± 19.01 SD; range: 22-93 years; 50 females). As can be seen in Table 1, this group was divided into 4 subgroups based on age (Group 1: 22-34 years; Group 2: 37-54; Group 3: 55-69; Group 4: 70-93). For Group 4, all but one participant were aged between 70 and 83 years, and there was a 93-year-old participant. All participants were right-handed, as assessed by the Edinburgh Handedness Inventory ([Oldfield 1971](#)), native speakers of Canadian French with a mean of 16.895 ± 4.203 years of education (range: 6-29 years). All participants had normal or corrected-to-normal vision and no self-reported speech, voice, language, psychological, neurological or neurodegenerative disorder at the time of the study and all were non-smokers. Participants were screened for depression using the Geriatric Depression Scale ([Yesavage et al. 1982](#)) and their cognitive level was assessed using the Montreal Cognitive Assessment scale (MOCA) ([Nasreddine et al. 2003](#)). Participants'

characteristics are reported in Table 1. The study was approved by the Institutional Ethical Committee of the Institut Universitaire en Santé Mentale de Québec (#293-2012).

1.2. Hearing assessment

Pure tone audiometry was performed using a clinical audiometer (AC40, Interacoustic) for each ear separately, at the following frequencies: .25, .5, 1, 2, 3, 4, 6, 8, 12 and 16 kHz. For each participant, a standard pure tone average (PTA: average of threshold at .5, 1 and 2 kHz) was computed for the left and right ear and used as a covariate in the statistical analyses. PTAs are used in clinical settings as a measure of hearing loss for speech because most speech sounds fall within this range ([Stach 2010](#)). The result of the hearing assessment is provided in Table 1.

1.3. Procedure

Participants were seated in a quiet room in front of a laptop computer (Thinkpad W510, Lenovo). Following a short practice session, participants were asked to produce sequences of speech, oro-facial and finger movements in separate blocks. The task consisted in the production of meaningless sequences of (i) three French syllables (SPEECH), (ii) three oro-facial movements (MOUTH), and (iii) three finger movements (FINGER) (see Table 2). Trials were randomly interleaved with short inter-trial intervals ranging from 500 to 1250 ms (with a mean of 875 ms). Participants' oral responses (speech and mouth) were recorded using a high quality multidirectional headworn microphone (Shure, Beta 53) connected to a sound card (Fast Track C400, M-audio), which was in turn connected to a laptop computer. All oral responses were recorded with the software Audacity (Open source). Finger movements were recorded using a USB response pad (Cedrus, model models RB-830). Throughout the procedure, participants' fingers rested on the response pad (see Figure 6a).

1.4. Stimuli and motor responses

The syllables used in the SPEECH condition were complex (CCVC) syllables: /krik/, /drad/, /broub/, and /grug/ (see Table 2). Meaningless syllables were used to avoid linguistic top-down effects that can facilitate speech production and because they are useful in the evaluation of maximal performance. Indeed, difficult syllable tasks could reveal a decline in maximal performance differences (“reduced reserve”). This is important because a reduced reserve can impair a person’s flexibility, that is, the ability to adjust speech output to different situations, and can also reveal whether the process of speaking is becoming overall more difficult ([Kent et al. 1987](#)). The oro-facial movements used in the MOUTH condition were (1) a kissing movement with the lips (“kiss”), (2) the production of a popping sound made with the two lips being pressed and opened (“pop”), (3) a sound made by pressing the tip of the tongue against the alveoles and then releasing the tongue (“tic”), and (4) a clapping sound also made with the tip of the tongue (“clap”). All movements produced a distinct sound. For the FINGER condition, participants were asked to position their right hand (thumb, index, middle, or ring finger) on the response pad and to press specific buttons when instructed.

All motor responses (SPEECH, MOUTH and FINGER) were triggered either visually or auditory. In the auditory condition, participants were presented, through high quality headphones (Shure, SRH440), with recordings of (1) the syllables, (2) the sounds of the oro-facial movements, (3) the color of the button to press (“red, blue, green, yellow”). Auditory stimuli were read by native speaker of French Canadian in a soundproof room, and recorded with Sound Studio 3.5.4 software (Felt Tip Software) at a sampling rate of 44 KHz. Stimuli were edited using Wave Pad Sound Editor 4.53 (NHC Software) to standardize their duration to 1200 ms and normalize the root mean square (RMS) intensity of the sound files. In the visual condition,

participants were presented with (1) the syllables written on the computer screen, (2) the name of oro-facial movement (“kiss, pop, tic and clap”), and (3) the name of the color of the button they needed to press (“red, blue, green, yellow”). All stimuli (visual and auditory) were presented using Presentation Software (Neurobehavioral System, CA, USA). The presentation of the stimuli lasted for 1800 ms and was followed, after an average of 1050 ms, by a green-coloured visual response cue (□) that remained on the screen for 3500 ms. At the end of this period, a red-coloured stop cue (□) was presented indicating to participants to stop responding. The stop cue remained on the screen until the beginning of the following trial, which occurred, on average, 875ms later (range 750 ms – 1250 ms).

Two types of sequences were performed. The sequences were either of simple, in which at least two of the three movements were identical and performed one after the other (ex: /krik krik krik/ or /pop pop clac/), or more complex, in which three different movements were produced (ex: /drad krik grug/ or /tic clac kiss/). Examples of sequences are reported in Table 2. The experiment included 24 trials of each condition (3 movements x 2 complexity levels x 2 modalities) for a total of 288 trials. These trials were divided into six experimental runs (2 runs for each movement modality). Within each run, the complexity of the movements was randomized while the other factors (stimuli modality and movement type) were kept constant to avoid task-switching effects not of interest in this experiment. The order of the runs was counter-balanced across participants. For all type of responses (SPEECH, MOUTH and FINGER), the different movements (e.g.: krik/, /drad/, /broub/, and /grug/) were produced a similar number of times (i.e. between 16 and 19 times) across modality and complexity level.

2.4 Behavioural data analysis

Data analyses focused on performance, measured as a percentage of errors per sequences for each condition (number of incorrect movements divided by total number of movement produced). Errors included errors of commission, errors of omission and production of additional movements. Accuracy was calculated based on the number of runs included in the analysis. A run was kept only if at least 50% of sequences were completed. For the analyses of SPEECH and MOUTH mistakes, a research assistant naive to the purpose of the study listened to and transcribed the participants' responses. For FINGER, the responses were recorded directly to disk and verified. The percentages of errors by condition and by age group are reported in Table 3.

2.5 Statistical analyses

First, a 3x2x2x4 ANCOVA was run using SPSS (IBM, version 22) to analyze performance (percentage of error by sequence), with three within-subject factors (Movement [SPEECH, MOUTH and FINGER], Complexity [simple, complex] and Modality [visual, auditory]), and one between-subject factor (Group [group 1, group 2, group 3 and group 4]). Two covariates were included in the statistical model (sex and the right PTA) to control for potential sex and hearing differences. Since a strong correlation was found between the right and the left PTA ($n = 80.876$, $p = 0.000$), only the right PTA was included in the analyses to control for hearing sensitivity while avoiding over fitting the data. Significant effects revealed by the ANCOVA were explored using FDR-corrected post hoc tests ([Benjamini and Hochberg 1995](#); [Genovese et al. 2002](#)) ($q = 0.05$, $i = 25$ tests).

2. Results

2.1. Number of errors by sequence

The four-way ANCOVA conducted on the percentage of errors revealed significant main effects of Movement ($F_{(2,140)} = 6.288, p = 0.002$), Complexity ($F_{(1,70)} = 21.578, p < 0.001$) and Group ($F_{(3,70)} = 9.648, p < 0.001$). Interactions between Group and Movement ($F_{(6,140)} = 5.835, p < 0.001$), between Complexity and Group ($F_{(3,70)} = 7.675, p < 0.001$), between Movement and Modality ($F_{(2,140)} = 5.889, p = 0.004$), between Movement, Modality and Group ($F_{(6,140)} = 4.236, p = 0.001$), between Movement and Complexity ($F_{(2,140)} = 8.241, p < 0.001$), between Movement, Complexity and Group ($F_{(6,140)} = 3.184, p = 0.006$) and between Modality and Complexity ($F_{(1,70)} = 6.635, p = 0.012$) were also found. No effects of PTA or Sex were found.

As can be seen in Figure 1a, in general, participants were more accurate in producing simple than complex sequences ($t_{(75)} = -20.465, p < 0.001$). As shown in Figure 1b, with age, there was an overall increase in error rate. Post hoc tests revealed that the eldest participants (group 4) made more mistakes than all other groups. Performance in this group differed significantly from group 1 ($t_{(36)} = -6.584, p = 0.002$), group 2 ($t_{(33)} = -4.753, p < 0.001$) and group 3 ($t_{(37)} = -2.908, p = 0.009$). Group 1 also made significantly fewer mistakes than group 3 ($t_{(39)} = -3.732, p = 0.001$). Group 2 and group 3 did not differ from each other. The main effect of Movement is illustrated in Figure 2, which shows that participants made less mistakes during FINGER compared to SPEECH ($t_{(75)} = 12.855, p < 0.001$) and MOUTH ($t_{(75)} = 14.233, p < 0.001$).

As shown in Figure 3, for the Modality by Movement interaction, post-hoc tests revealed that in the MOUTH condition, participants made more mistakes in the auditory modality than in the visual modality ($t_{(75)} = 12.500, p < 0.001$). In contrast, in the SPEECH and FINGER conditions, they made more mistakes in the visual than in the auditory modality (SPEECH: $t_{(75)} = -4.022, p < 0.001$; FINGER: $t_{(75)} = -2.216, p = 0.037$). For the 3-way interaction between Modality, Group and Movement, post-hoc tests revealed that, within each group, the direction of the stimulus

Modality effect was different in the MOUTH compared to the SPEECH condition (group 1: $t_{(19)} = -3.582, p = 0.003$; group 2: $t_{(16)} = -5.796, p < 0.001$; group 3: $t_{(20)} = -6.034, p < 0.001$; group 4: $t_{(17)} = -6.926, p < 0.001$). The effect also differed between MOUTH and FINGER (group 1: $t_{(19)} = 5.551, p < 0.001$; group 2: $t_{(16)} = 8.507, p < 0.001$; group 3: $t_{(20)} = 5.448, p < 0.001$; group 4: $t_{(17)} = 7.076, p < 0.001$). The modality difference between FINGER and SPEECH was significant for group 3 ($t_{(20)} = -2.510, p = 0.027$) and group 4 ($t_{(17)} = -2.704, p = 0.021$). For the Modality by Complexity interaction, post-hoc tests revealed that for the complex sequences, participants made more mistakes in auditory than visual modality. There was no effect of modality for the simple sequences.

To decompose the 3-way interaction between Complexity, Group and Movement, a series of 3 additional 2x4 ANCOVAs were run, one for each type of movement, to examine age and sequence complexity effects within each movement type, with Complexity as a within subject factor and Group as a between subject factor. Two covariates were included in the statistical model (sex and the right PTA) to control for potential sex and hearing differences. As can be seen in Figure 4, these analyses revealed that Complexity effects were present in the SPEECH and MOUTH conditions but not in the FINGER condition. As show in Figure 5a, for SPEECH, the ANCOVA revealed a main effect of Complexity ($F_{(1,70)} = 18.596, p < 0.001$) and Group ($F_{(3,70)} = 5.928, p = 0.001$) and an interaction between Complexity and Group ($F_{(3,70)} = 6.049, p = 0.001$). Analysis revealed no effect of PTA or sex. FDR-corrected post hoc t-tests ([Benjamini and Hochberg 1995](#); [Genovese et al. 2002](#)) ($q = 0.05; i = 10$) revealed that the difference between simple and complex sequences was significant for all groups (group 1: $t_{(19)} = -7.005, p < 0.001$; group 2: $t_{(16)} = -11.737, p < 0.001$; group 3: $t_{(20)} = -12.354, p < 0.001$; group 4: $t_{(17)} = -8.668, p < 0.001$). Moreover, the difference between simple and complex sequences was larger for group 2

($t_{(35)} = 3.548, p = 0.002$), group 3 ($t_{(39)} = 4.508, p < 0.001$) and group 4 ($t_{(36)} = 3.711, p = 0.001$) compared to group 1.

As show in 5b, for MOUTH, the ANCOVA revealed a main effect of Complexity ($F_{(1,70)} = 19.075, p < 0.001$), and Group ($F_{(3,70)} = 11.764, p < 0.001$), and an interaction between Complexity and Group ($F_{(3,70)} = 5.868, p = 0.001$). No effects of PTA or sex were found. FDR-corrected post-hoc tests ($q = 0.05, i = 10$) ([Benjamini and Hochberg 1995](#); [Genovese et al. 2002](#)) revealed that the difference between simple and complex sequences was significant for all groups (Group 1: $t_{(20)} = -5.949, p < 0.001$; group 2 : $t_{(16)} = -10.503, p < 0.001$; group 3 : $t_{(20)} = -11.483, p < 0.001$; group 4 : $t_{(17)} = -8.328, p < 0.001$). Moreover, the difference between simple and complex sequences was larger for group 2 ($t_{(35)} = 4.216, p < 0.001$), group 3 ($t_{(39)} = 4.071, p < 0.001$) and group 4 ($t_{(36)} = 3.489, p = 0.002$) compared to group 1.

For FINGER, the ANCOVA revealed an effect of right PTA ($F_{(1,70)} = 8.044, p = 0.006$) but no significant effect of complexity or age. Because we expected to find age effects on finger movements, we decided to explore the manual movements further by conducting additional analyses on reaction time (RT) and sequence duration in order to determine whether age affected finger movements in terms of timing rather than accuracy. For this analysis, we extracted RT and sequence durations only for the correct trials. Sequences that started before the response cue or that finished after the end of the trial were removed from the statistical analysis. Trials containing outliers, defined as values $\pm 2SD$ from the participant's mean, were also removed from the analysis. One participant was excluded from the duration and RT analysis because in the visual condition too many sequences (73%) were incorrect. Two separate $2 \times 2 \times 4$ ANCOVA were conducted on the resulting data, one for the RT and one for duration, with Complexity and

Modality as the within-subject factors, and Group as the between subject factor. For RT, the ANCOVA revealed significant main effects of Modality ($F_{(1,69)} = 4.130, p = 0.046$) and Complexity ($F_{(1,69)} = 6.7552, p = 0.011$) but no interaction. FDR-corrected post-hoc t-tests ([Benjamini and Hochberg 1995](#); [Genovese et al. 2002](#)) ($q = 0.05, i = 2$) revealed that participants were slower in visual than auditory condition ($t_{(74)} = -11.675, p < 0.001$). As shown in Figure 6b, in general, RT were shorter for simple compared to complex sequences ($t_{(74)} = -13.386, p < 0.001$).

For sequence duration, the ANCOVA revealed significant main effects of Complexity ($F_{(1,69)} = 4.945, p = 0.029$) and Group ($F_{(3,69)} = 4.670, p = 0.005$). Interactions between Complexity and Group ($F_{(3,69)} = 5.437, p = 0.002$) and between Complexity and Modality ($F_{(1,69)} = 4.182, p = 0.045$) were also found. FDR-corrected post-hoc t-tests ([Benjamini and Hochberg 1995](#); [Genovese et al. 2002](#)) ($q = 0.05, i = 12$) were conducted to explore these effects. As shown in Figure 6c, in general, simple sequences were shorter than complex sequences ($t_{(74)} = -7.935, p < 0.001$). As show in Figure 6d, Post hoc tests revealed that the youngest participants (group 1) were faster than all other groups. Performance in this group differed significantly from group 2 ($t_{(35)} = -3.070, p = 0.006$), group 3 ($t_{(39)} = -3.255, p = 0.004$) and group 4 ($t_{(35)} = -3.680, p = 0.002$). For the Group by Complexity interaction, post-hoc tests revealed a complexity effect (complex > simple) in all groups except for group 1 (group 2: $t_{(16)} = -4.556, p = 0.001$; group 3: $t_{(20)} = -5.298, p < 0.001$; group 4: $t_{(16)} = -5.548, p < 0.001$). The youngest participants showed no complexity effect on response duration. For the Modality by Complexity interaction, post-hoc tests revealed that the difference between simple sequences and complex sequences was larger in the visual condition than in auditory condition ($t_{(74)} = 5.261, p < 0.001$).

3. Discussion

The goal of the present study was to examine the effect of aging on the ability to produce sequences of fine motor actions (speech, oro-facial and manual movements) varying in complexity levels, while controlling for hearing, in healthy adults. Given the inherently sequential nature of speech, sequencing difficulties can be particularly detrimental to communication efficiency in older ages. Despite the importance of communication in aging, the extent and underlying causes of articulatory and speech sequencing difficulties are still unknown, i.e. whether they are related to peripheral factors such as decreased oral muscle endurance, or to neurobiological factors such as less efficient neural mechanisms or structural damage to the brain regions involved in speech production. Here, we aimed at characterizing the extent of these difficulties using a behavioural approach and by conducting an analysis of errors. As was expected, an overall age-related performance decline was observed. When movements were examined separately, differences in the effect of aging and sequence complexity were found across movement types, with speech and oro-facial movements showing age-related accuracy decline but not finger movements. For finger movements, however, additional analyses revealed an increase in response duration with age. These findings are discussed in the following paragraphs.

3.1. Speech production

In the present study, we examined accuracy during a sequential speech production task in healthy young and older adults and we found a significant decrease in performance with age and, importantly, we found that this decline was stronger for complex sequences. From group 1 to 4, a ~55% decline was observed for the simple sequences and ~52% decline for the complex sequences. Several prior studies have shown a decline in speech rate with age ([Duchin and Mysak 1987](#); [Fozo and Watson 1998](#); [Ramig 1983a](#); [Ryan 1972](#); [Searl et al. 2002](#); [Wohlert and](#)

[Smith 1998](#)) but few study have examined accuracy and only a few have shown that older adults are less intelligible than younger adults ([Parnell and Amerman 1987](#); [Shuey 1989](#)). The present results demonstrate, for the first time, a decline in speech sequencing skills in healthy adults.

The underlying causes of the decline in speech skills with aging remain unknown. In previous studies of speech rate, the stimuli used were usually sentences or words; it is therefore possible that a decline in the efficiency of linguistic processing (e.g. syntax, lexical access, etc.) may account for the observed slowing, or at least for a part of it. In contrast, in the present study, we used meaningless syllable sequences and we still observed an age-related decline in accuracy, which suggests that the decline in efficiency of linguistic processes is not the only factor contributing to the observed decline of speech skills in aging. One possibility is that physiological changes in the oro-facial sphere could be contributing to the observed decline in accuracy. For example, it has been shown that older adults exhibit decreased oral tactile sensitivity ([Calhoun et al. 1992](#); [Wohlert 1996b](#); [Wohlert and Smith 1998](#)), as well as decreased lip strength ([Wohlert and Smith 1998](#)), and decreased maximal tongue strength ([Neel and Palmer 2012](#)). However, one study has shown that tongue maximal strength is a poor predictor of articulation rate ([Neel and Palmer 2012](#)), suggesting limited contribution of these physiological factors. Another, more likely possibility is that decline in speech accuracy is related to a decline in the neural planning and control of speech movements. Consistent with this hypothesis, recent studies from our group have shown important age-related changes in the structure and function of brain areas involved in speech motor control, including the premotor cortex and the supplementary motor area ([Bilodeau-Mercure et al. 2014](#); [Tremblay et al. 2013](#)). Given the importance of communication in aging, and the fact that speaking is intrinsically a sequential behavior, further research is needed to better understand the cause of these sequencing difficulties. Moreover, further research needs to establish the range of these difficulties, and whether they are modulated by factors such as

syllable complexity and syllable frequency. This is necessary to establish a more complete picture of the changes that the speech system undergoes with age and help design new interventions to remediate speech difficulties based on a better understanding of the specific articulatory difficulties faced by elderly adults.

3.2. Oro-facial movements

In the present study, we observed that older adults made significantly more mistakes than younger adults when they produced sequences of oro-facial movements. This effect was significantly stronger for the complex sequences. From group 1 to 4, a ~76% decline was observed for simple sequences and ~66% decline for complex sequences suggesting a decline in oro-facial motor control with aging. Very few studies have explored the production of oro-facial movements in aging. In one study, age-related changes in perioral reflex movement in response to innocuous mechanical stimulation of the lip vermillion have been reported ([Wohlert 1996a](#)). In this study, older women (67-85 years) produced less reflexive responses to stimulation and their response had lower amplitude and longer latency than younger women (20-25 years). These changes in oral reflex movements can be due to physical changes like decreased lip strength ([Wohlert and Smith 1998](#)), or decreased oral tactile sensitivity ([Calhoun et al. 1992](#); [Wohlert 1996b](#); [Wohlert and Smith 1998](#)). In the present study, these factors could be responsible for the performance decline that we observed. However, changes in motor planning and execution are also likely to play a role in the etiology of these age-related changes. Indeed, though performance decreased with an increase in complexity in all groups, we found that the effect of the complexity increased with age consistent with a decline in the neural control of movement planning.

The finding of a similar gradual decline in speech and oro-facial movements suggests a common underlying aging mechanism, not specific to speech movements. This aging process

could be related to the planning of movements involving the face and mouth or to the execution of these movements. Though only a few study have examined the relationship between oro-facial movements and speech in adulthood and particularly in aging, a link between oro-facial movements and speech has been shown, consistent with the idea of shared mechanisms ([Alcock 2006](#); [Alcock et al. 2000](#); [Tremblay and Gracco 2009](#); [Tremblay and Gracco 2010](#)). For example, some studies have shown that motor response selection involves similar neural resources for speech and oro-facial movements ([Tremblay and Gracco 2009](#); [Tremblay and Gracco 2010](#)). Moreover, patient with speech impairments also have difficulty executing oro-facial movements ([Alcock 2006](#); [Alcock et al. 2000](#)) suggesting shared mechanisms. In line with previous results, the present findings support the notion that speech and oro-facial movements engage common motor control mechanisms, including movement sequencing. Additional studies are needed to continue to explore the common and separate etiology of these changes, whether related to peripheral factors such as decreased oro-facial strength or sensibility, or to central factors such as motor planning, in particular motor sequencing.

3.3. Finger movements

In the present study, no effect of age or complexity was found on accuracy of finger movements. This finding was unexpected given that many studies have shown an age-related decline in the accuracy of manual movements ([Chaput and Proteau 1996](#); [Christou and Enoka 2011](#); [Goggin and Meeuwsen 1992](#); [Pohl et al. 1996](#); [Yan et al. 1998](#)). There are, however, several differences between the tasks used in these studies and the one that was used here. First, in the present study, only the fingers were used, whereas, in many others studies, participants performed more complex movements involving the whole arm ([Chaput and Proteau 1996](#); [Goggin and Meeuwsen 1992](#); [Pohl et al. 1996](#); [Yan et al. 1998](#)). Moreover, in our study, accuracy

was measured as the number of correct movements produced (i.e. pressing with the correct finger); it was thus a simple dichotomous dependent variable with only two possible outcomes (correct/incorrect). In others studies, in contrast, accuracy was measured continuously in terms of movement precision ([Chaput and Proteau 1996](#); [Christou and Enoka 2011](#); [Goggin and Meeuwsen 1992](#); [Pohl et al. 1996](#); [Yan et al. 1998](#)). For example, in a recent study, participants were asked to lift and lower light loads with their index finger and to stop their movements at a specific angle from the other fingers ([Christou and Enoka 2011](#)). Older adults had more difficulty stopping their movement at the specified angle compared to the younger adults, though they could still stop the movement. It is possible that, should we have measured movement trajectories, we could have found age-related differences in movement precision in our task. Consistent with this hypothesis, we conducted additional analyses of the manual movements and we found age effects on response duration, consistent with the notion that, though globally accurate (i.e. on target), the finger movements of older adults differed from those produced by younger adults. We also show that more complex sequences of finger movements are particularly affected by age in terms of response duration. These results are consistent with previous studies that found an effect of aging on the time required to produce sequences of movements ([Aoki and Fukuoka 2010](#); [Cacola et al. 2013](#); [Cousins et al. 1998](#); [Ruiz et al. 2007](#)).

Taken together, our results suggest that, in a simple finger movement (button pressing) tasks, age effects affect duration more strongly than global accuracy. Indeed, the percentage of errors was very low across conditions. This may suggest that the orofacial tasks had a higher difficulty level compared to the finger movement task. The other possibility is that response accuracy declines more quickly in the oro-facial sphere than it does in the manual action sphere, perhaps due to the inherent complexity of speech movements, which require the coordination of several different muscles. Further studies are needed to replicate these findings, and determine

the extent to which the aging of speech and oro-facial movements follows a different trajectory from that of finger movements.

3.4. Stimuli modality

In the present study, we observed that stimuli modality affected performance differently depending of the type of movement produced. For SPEECH and FINGER, participants were less accurate in the visual compared to the auditory condition. In contrast, for MOUTH, participants were less accurate in the auditory than in the visual modality. Participants reported having difficulty recognizing the sounds of the oro-facial movements. In day-to-day situations, these sounds usually occur in the presence of visual information (e.g. shape of the mouth, degree of opening) or other forms of contextual information including speech, such as listening to a kissing lip movement when saying goodbye to a loved one the phone. In contrast, people frequently listen to speech without visual information, for example, during phone conversations. This may explain the difficulty related to the auditory MOUTH condition. It is important to note that, since all analyses were corrected for hearing threshold, this effect cannot be attributed to a hearing decline otherwise it would have affected all movements equally. The present results suggest that the modality of the movement trigger differentially affect accuracy of oro-facial and speech movements in normal aging, which may have important implications for rehabilitation.

5. Conclusion

This study provides important new empirical evidence that orofacial motor control declines in cognitively healthy elderly adults (both males and females). These results suggest that the motor speech system undergoes significant decline over time that affect oro-facial movements. Here, we follow the general hypothesis that age-related speech difficulties, both

perceptual and motor, have a multifactorial aetiology that includes central factors (such as speech motor planning), and possibly peripheral factors as well (such as tactile sensibility, decrease muscular endurance). Further studies are needed to continue explore the distinct impact of these different factors on the ability to communicate in aging. This is crucial since communication difficulties are considered to be of great importance by elderly adults ([Jacobs-Condit 1984](#)). Indeed, these difficulties often lead to social participation that is less diverse, more restricted to home settings, and involves fewer relationships ([Law 2002](#)). It is therefore important that we gain a better understand of the extent to which age affects speech production and its underlying mechanisms, to be able to detect and remediate these important health issues.

6. Acknowledgments

This work was supported by grants from the Fonds Québécois de le Recherche – Société et Culture (FRQ-SC) and from the Fonds Québécois de le Recherche – Santé (FRQ-S) to P.T. and by a start-up grant from the Institut Universitaire en Santé Mentale de Québec (IUSMQ) to P.T. These sponsors played no role in the design, execution, analysis and interpretation of data, or writing of the study. We thank Isabelle Deschamps for her comments on previous versions of this manuscript.

7. References

Alcock K (2006) The development of oral motor control and language Down's syndrome, research and practice : the journal of the Sarah Duffen Centre / University of Portsmouth 11:1-8

- Alcock KJ, Passingham RE, Watkins KE, Vargha-Khadem F (2000) Oral dyspraxia in inherited speech and language impairment and acquired dysphasia *Brain and language* 75:17-33
doi:10.1006/brln.2000.2322
- Aoki T, Fukuoka Y (2010) Finger tapping ability in healthy elderly and young adults *Medicine and science in sports and exercise* 42:449-455 doi:10.1249/MSS.0b013e3181b7f3e1
- Baker KK, Ramig LO, Sapir S, Luschei ES, Smith ME (2001) Control of vocal loudness in young and old adults *Journal of speech, language, and hearing research : JSLHR* 44:297-305
- Benjamini Y, Hochberg Y (1995) Controlling the False Discovery Rate: a Practical and Powerful Approach to Multiple Testing *Journal of the Royal Statistical Society Series B (Methodological)* 57:289-300
- Bilodeau-Mercure M, Lortie CL, Sato M, Guitton MJ, Tremblay P (2014) The neurobiology of speech perception decline in aging *Brain structure & function* doi:10.1007/s00429-013-0695-3
- Cacola P, Roberson J, Gabbard C (2013) Aging in movement representations for sequential finger movements: a comparison between young-, middle-aged, and older adults *Brain and cognition* 82:1-5 doi:10.1016/j.bandc.2013.02.003
- Calhoun KH, Gibson B, Hartley L, Minton J, Hokanson JA (1992) Age-related changes in oral sensation *The Laryngoscope* 102:109-116 doi:10.1288/00005537-199202000-00001
- Chaput S, Proteau L (1996) Aging and motor control *The journals of gerontology Series B, Psychological sciences and social sciences* 51:P346-355
- Christou EA, Enoka RM (2011) Aging and movement errors when lifting and lowering light loads *Age (Dordrecht, Netherlands)* 33:393-407 doi:10.1007/s11357-010-9190-4

- Cousins MS, Corrow C, Finn M, Salamone JD (1998) Temporal measures of human finger tapping: effects of age *Pharmacology, biochemistry, and behavior* 59:445-449
- D'Haeseleer E, Depypere H, Claeys S, Wuyts FL, Baudonck N, Van Lierde KM (2011) Vocal characteristics of middle-aged premenopausal women *Journal of voice : official journal of the Voice Foundation* 25:360-366 doi:10.1016/j.jvoice.2009.10.016
- Decoster W, Debruyne F (1997) The ageing voice: changes in fundamental frequency, waveform stability and spectrum *Acta oto-rhino-laryngologica Belgica* 51:105-112
- Duchin SW, Mysak ED (1987) Disfluency and rate characteristics of young adult, middle-aged, and older males *Journal of communication disorders* 20:245-257
- Fozo MS, Watson BC (1998) Task complexity effect on vocal reaction time in aged speakers *Journal of voice : official journal of the Voice Foundation* 12:404-414
- Genovese CR, Lazar NA, Nichols T (2002) Thresholding of Statistical Maps in Functional Neuroimaging Using the False Discovery Rate *NeuroImage* 15:870-878
doi:<http://dx.doi.org/10.1006/nimg.2001.1037>
- Gentilucci M (2003) Grasp observation influences speech production *The European journal of neuroscience* 17:179-184
- Gentilucci M, Dalla Volta R, Gianelli C (2008) When the hands speak *Journal of physiology, Paris* 102:21-30 doi:10.1016/j.jphysparis.2008.03.002
- Goggin NL, Meeuwsen HJ (1992) Age-related differences in the control of spatial aiming movements *Research quarterly for exercise and sport* 63:366-372
doi:10.1080/02701367.1992.10608758
- Honjo I, Isshiki N (1980) Laryngoscopic and voice characteristics of aged persons *Archives of otolaryngology (Chicago, Ill : 1960)* 106:149-150

Hunter EJ, Kapsner-Smith M, Peard P, Engar MZ, Brown WR (2012) Age and speech production: a 50-year longitudinal study *Journal of the American Geriatrics Society* 60:1175-1177
doi:10.1111/j.1532-5415.2012.03983.x

Jacobs-Condit LE (1984) *Gerontology and Communication Disorders*. American Speech-Language-Hearing Association, Rockville

Jimenez-Jimenez FJ et al. (2011) Influence of age and gender in motor performance in healthy subjects *Journal of the neurological sciences* 302:72-80 doi:10.1016/j.jns.2010.11.021

Kent RD (2000) Research on speech motor control and its disorders: a review and prospective *J Commun Disord* 33:391-427; quiz 428 doi:S0021-9924(00)00023-X [pii]

Kent RD, Kent JF, Rosenbek JC (1987) Maximum performance tests of speech production *The Journal of speech and hearing disorders* 52:367-387

Law M (2002) Participation in the occupations of everyday life *The American journal of occupational therapy : official publication of the American Occupational Therapy Association* 56:640-649

Linville SE (1996) The sound of senescence *Journal of voice : official journal of the Voice Foundation* 10:190-200

Morris R, Brown WS (1987) Age-related voice measures among adult women *Journal of voice* 1:43

Mueller PB (1997) The aging voice *Seminars in speech and language* 18:159-168; quiz 168-159
doi:10.1055/s-2008-1064070

Nasreddine ZS, Chertkow H, Phillips N, Bergman H, Whitehead V (2003) Sensitivity and Specificity of The Montreal Cognitive Assessment (MoCA) for Detection of Mild Cognitive Deficits *Can J Neurol Sci* 30

- Neel AT, Palmer PM (2012) Is tongue strength an important influence on rate of articulation in diadochokinetic and reading tasks? *Journal of speech, language, and hearing research* : JSLHR 55:235-246 doi:10.1044/1092-4388(2011/10-0258)
- Oldfield RC (1971) The assessment and analysis of handedness: the Edinburgh inventory *Neuropsychologia* 9:97-113
- Parnell MM, Amerman JD (1987) Perception of oral diadochokinetic performances in elderly adults *Journal of communication disorders* 20:339-351
- Pohl PS, Winstein CJ, Fisher BE (1996) The locus of age-related movement slowing: sensory processing in continuous goal-directed aiming *The journals of gerontology Series B, Psychological sciences and social sciences* 51:P94-102
- Ramig LA (1983a) Effects of physiological aging on speaking and reading rates *Journal of communication disorders* 16:217-226
- Ramig LA (1983b) Effects of physiological aging on vowel spectral noise *Journal of gerontology* 38:223-225
- Ruiz PJ, Bernardos VS, Bartolome M, Torres AG (2007) Caput timed tests quantify age-related motor decline in normal subjects *Journal of the neurological sciences* 260:283-285 doi:10.1016/j.jns.2007.04.034
- Ryan WJ (1972) Acoustic aspects of the aging voice *Journal of gerontology* 27:265-268
- Ryan WJ, Burk KW (1974) Perceptual and acoustic correlates of aging in the speech of males *Journal of communication disorders* 7:181-192
- Searl JP, Gabel RM, Fulks JS (2002) Speech disfluency in centenarians *Journal of communication disorders* 35:383-392
- Shuey EM (1989) Intelligibility of older versus younger adults' CVC productions *Journal of communication disorders* 22:437-444

- Smith BL, Wasowicz J, Preston J (1987) Temporal characteristics of the speech of normal elderly adults *Journal of speech and hearing research* 30:522-529
- Stach BA (2010) *Clinical Audiology: An Introduction*. Delmar, Clifton Park, NY, USA
- Tremblay P, Dick AS, Small SL (2013) Functional and structural aging of the speech sensorimotor neural system: functional magnetic resonance imaging evidence *Neurobiology of aging* 34:1935-1951 doi:10.1016/j.neurobiolaging.2013.02.004
- Tremblay P, Gracco VL (2009) Contribution of the pre-SMA to the production of words and non-speech oral motor gestures, as revealed by repetitive transcranial magnetic stimulation (rTMS) *Brain research* 1268:112-124 doi:10.1016/j.brainres.2009.02.076
- Tremblay P, Gracco VL (2010) On the selection of words and oral motor responses: evidence of a response-independent fronto-parietal network *Cortex* 46:15-28
doi:10.1016/j.cortex.2009.03.003
- Tremblay P, Shiller DM, Gracco VL (2008) On the time-course and frequency selectivity of the EEG for different modes of response selection: evidence from speech production and keyboard pressing *Clinical neurophysiology : official journal of the International Federation of Clinical Neurophysiology* 119:88-99
- Wilcox KA, Horii Y (1980) Age and changes in vocal jitter *Journal of gerontology* 35:194-198
- Wohlert AB (1996a) Reflex responses of lip muscles in young and older women *Journal of speech and hearing research* 39:578-589
- Wohlert AB (1996b) Tactile perception of spatial stimuli on the lip surface by young and older adults *Journal of speech and hearing research* 39:1191-1198
- Wohlert AB, Smith A (1998) Spatiotemporal stability of lip movements in older adult speakers *Journal of speech, language, and hearing research : JSLHR* 41:41-50

Yan JH, Thomas JR, Stelmach GE (1998) Aging and rapid aiming arm movement control
Experimental aging research 24:155-168 doi:10.1080/036107398244292

Yesavage JA, Brink TL, Rose TL, Lum O, Huang V, Adey M, Leirer VO (1982) Development
and validation of a geriatric depression screening scale: a preliminary report Journal of
psychiatric research 17:37-49

FIGURES




Fig. 1 Overall percentage of errors displayed as function of (a) sequence complexity (simple and complex) and (b) age group.




Fig. 2 Percentage of errors displayed as function of age group for each movement.




Fig. 3 Percentage of errors displayed as function of stimulus modality (visual or auditory) for each movement: (a) SPEECH, (b) MOUTH, (c) FINGER. Asterisks indicate significant differences.




Fig. 4 Complexity effect ([percentage of errors complex sequences] – [percentage of errors for simple sequences]) displayed as function of age group, separately for each movement: (a) SPEECH, (b) MOUTH, and (c) FINGER. Asterisks indicate significant differences.




Fig. 5 Percentage of errors for the simple and complex sequences displayed as function age group for: (a) SPEECH, (b) MOUTH.

Finger movements




Fig. 6 (a) Position of the fingers on USB response pad (Cedrus, model models RB-830) during the FINGER task. The bar graphs illustrate the results for the additional analyses conducted on the finger movements. Asterisks indicate significant differences. **(b)** Reaction time as a function of complexity; **(c)** Response duration as function age group **(d)** Response duration as a function of sequence complexity.

TABLES

Table 1: Participants' characteristics by age group

		Age	Years of education	MOCA*	Depression scale**	Laterality Quotient***	Right ear PTA (Hz)	Left ear PTA (Hz)
Group 1 (22-34 years) (n=20, 13 females)	Minimum	22	13	26	0	90	-28.67	-27
	Maximum	34	24	30	8	100	-16	-12
	Mean \pm SD	28.05 \pm 4.17	17.70 \pm 2.64	28.80 \pm 1.20	2.50 \pm 2.28	99.50 \pm 2.24	-20.12 \pm 3.76	-19.28 \pm 3.47
Group 2 age (37-54 years) (n=17, 8 females)	Minimum	37	11	25	0	70	-39.33	-36
	Maximum	54	24	30	9	100	-15.33	-12
	Mean \pm SD	45.94 \pm 5.45	16.65 \pm 4.17	27.64 \pm 1.56	1.65 \pm 2.50	97.06 \pm 8.49	-22.02 \pm 6.25	-19.39 \pm 5.41
Group 3 (55-69 years) (n=21, 17 females)	Minimum	55	12	26	0	90	-42.33	-35.67
	Maximum	69	29	30	8	100	-18.67	-13
	Mean \pm SD	61.81 \pm 5.17	18.14 \pm 4.07	28.19 \pm 1.63	1.48 \pm 2.18	99.52 \pm 2.18	-25.81 \pm 6.17	-23.22 \pm 6.18
Group 4 (70-93 years) (n=18, 12 females)	Minimum	70	6	23	0	78.95	-48.67	-51.33
	Maximum	93	24	30	9	100	-20.33	-17.67
	Mean \pm SD	76.89 \pm 5.71	14.78 \pm 5.19	26.67 \pm 1.68	2.28 \pm 2.59	97.07 \pm 6.06	-32.46 \pm 7.79	-32.07 \pm 7.85

* Max score at the MOCA is 30

** Max score for the depression scale is 30

***Max score for the laterality quotient is 100

Table 2: Examples of sequence for each type of stimulus

Movement type	Stimuli	Simple sequence	Complex sequence
SPEECH	krik drad	krik krik krik	krik grug drad
	broub grug	krik krik grug	drad broub krik
MOUTH	pop bec (kiss)	bec bec bec	bec pop tic
	tic clac	pop tic tic	pop tic bec
	bleu (blue) (thumb)		
FINGER	jaune (yellow) (index)	vert vert vert	jaune bleu rouge
	rouge (red) (middle finger)		
	vert (green) (ring finger)	jaune jaune vert	vert jaune bleu

Table 3: Mean accuracy (percentage of errors) and standard deviation for each condition and each group

Group	Group 1 (22-34 years)		Group 2 (37-54 years)		Group 3 (55-69 years)		Group 4 (70-93 years)	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD
SPEECH auditory simple sequences	12.34	16.65	9.72	9.56	9.80	9.62	18.46	9.72
SPEECH auditory complex sequences	21.63	17.47	24.05	14.44	28.37	15.96	39.82	15.05
SPEECH visual simple sequences	12.46	11.32	10.41	9.61	18.00	18.60	35.97	21.72
SPEECH visual complex sequences	22.44	14.75	29.43	14.90	37.33	21.63	52.71	23.31
MOUTH auditory simple sequences	8.00	8.36	14.69	7.97	23.38	13.88	30.94	18.41
MOUTH auditory complex sequences	22.76	13.49	39.78	13.41	45.40	10.51	51.62	13.44
MOUTH visual simple sequences	3.69	2.65	6.30	8.54	12.34	10.92	17.20	12.67
MOUTH visual complex sequences	3.54	3.97	11.74	9.54	19.28	13.88	25.87	13.33
FINGER auditory simple sequences	0.90	1.17	0.80	1.26	0.93	1.61	1.36	1.81
FINGER auditory complex sequences	1.27	1.91	1.17	2.01	1.04	1.84	1.57	1.53
FINGER visual simple sequences	0.92	1.23	1.59	3.35	1.43	1.32	2.27	3.45
FINGER visual complex sequences	1.79	1.68	1.58	2.06	1.94	3.57	5.52	11.50