

## RESEARCH LETTER

10.1002/2015GL066235

Catherine M. O'Reilly, Sapna Sharma,  
Derek K. Gray, and Stephanie E.  
Hampton joint first authors

## Key Points:

- Lake surface waters are warming rapidly but are spatially heterogeneous
- Ice-covered lakes are typically warming at rates greater than air temperatures
- Both geomorphic and climate factors influence lake warming rates

## Supporting Information:

- Figures S1–S4 and Tables S1–S4

## Correspondence to:

C. M. O'Reilly,  
oreilly@ilstu.edu

## Citation:

O'Reilly, C. M., et al. (2015), Rapid and highly variable warming of lake surface waters around the globe, *Geophys. Res. Lett.*, 42, 10,773–10,781, doi:10.1002/2015GL066235.

Received 16 OCT 2015

Accepted 14 NOV 2015

Published online 16 DEC 2015

©2015. The Authors.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

## Rapid and highly variable warming of lake surface waters around the globe

Catherine M. O'Reilly<sup>1</sup>, Sapna Sharma<sup>2</sup>, Derek K. Gray<sup>3</sup>, Stephanie E. Hampton<sup>4</sup>, Jordan S. Read<sup>5</sup>, Rex J. Rowley<sup>1</sup>, Philipp Schneider<sup>6</sup>, John D. Lenters<sup>7</sup>, Peter B. McIntyre<sup>8</sup>, Benjamin M. Kraemer<sup>8</sup>, Gesa A. Weyhenmeyer<sup>9</sup>, Dietmar Straile<sup>10</sup>, Bo Dong<sup>11</sup>, Rita Adrian<sup>12</sup>, Mathew G. Allan<sup>13</sup>, Orlane Anneville<sup>14</sup>, Lauri Arvola<sup>15</sup>, Jay Austin<sup>16</sup>, John L. Bailey<sup>17</sup>, Jill S. Baron<sup>18</sup>, Justin D. Brookes<sup>19</sup>, Elvira de Eyto<sup>20</sup>, Martin T. Dokulil<sup>21</sup>, David P. Hamilton<sup>22</sup>, Karl Havens<sup>23</sup>, Amy L. Hetherington<sup>24</sup>, Scott N. Higgins<sup>25</sup>, Simon Hook<sup>26</sup>, Lyubov R. Izmet'seva<sup>27</sup>, Klaus D. Joehnk<sup>28</sup>, Kulli Kangur<sup>29</sup>, Peter Kasprzak<sup>30</sup>, Michio Kumagai<sup>31</sup>, Esko Kuusisto<sup>32</sup>, George Leshkevich<sup>33</sup>, David M. Livingstone<sup>34</sup>, Sally MacIntyre<sup>35</sup>, Linda May<sup>36</sup>, John M. Melack<sup>37</sup>, Doerthe C. Mueller-Navarra<sup>38</sup>, Mikhail Naumenko<sup>39</sup>, Peeter Noges<sup>40</sup>, Tiina Noges<sup>40</sup>, Ryan P. North<sup>41</sup>, Pierre-Denis Plisnier<sup>42</sup>, Anna Rigosi<sup>19</sup>, Alon Rimmer<sup>43</sup>, Michela Rogora<sup>44</sup>, Lars G. Rudstam<sup>24</sup>, James A. Rusak<sup>45</sup>, Nico Salmaso<sup>46</sup>, Nihar R. Samal<sup>47</sup>, Daniel E. Schindler<sup>48</sup>, S. Geoffrey Schladow<sup>49</sup>, Martin Schmid<sup>50</sup>, Silke R. Schmidt<sup>12</sup>, Eugene Silow<sup>27</sup>, M. Evren Soylu<sup>51</sup>, Katrin Teubner<sup>52</sup>, Piet Verburg<sup>53</sup>, Ari Voutilainen<sup>54</sup>, Andrew Watkinson<sup>55</sup>, Craig E. Williamson<sup>56</sup>, and Guoqing Zhang<sup>57</sup>

<sup>1</sup>Department of Geography-Geology, Illinois State University, Normal, Illinois, USA, <sup>2</sup>Department of Biology, York University, Toronto, Ontario, Canada, <sup>3</sup>Department of Biological and Environmental Sciences, California University of Pennsylvania, California, Pennsylvania, USA, <sup>4</sup>Center for Environmental Research, Education and Outreach, Washington State University, Pullman, Washington, USA, <sup>5</sup>U.S. Geological Survey-Center for Integrated Data Analytics, Middleton, Wisconsin, USA, <sup>6</sup>Norwegian Institute for Air Research, Kjeller, Norway, <sup>7</sup>LimnoTech, Ann Arbor, Michigan, USA, <sup>8</sup>Center for Limnology, University of Wisconsin-Madison, Madison, Wisconsin, USA, <sup>9</sup>Department of Ecology and Genetics/Limnology, Uppsala University, Uppsala, Sweden, <sup>10</sup>Limnological Institute, University of Konstanz, Konstanz, Germany, <sup>11</sup>Department of Atmospheric and Environmental Sciences, State University of New York at Albany, Albany, New York, USA, <sup>12</sup>Department of Ecosystem Research, Leibniz Institute of Freshwater Ecology and Inland Fisheries, Berlin, Germany, <sup>13</sup>Environmental Research Institute, University of Waikato, Hamilton, New Zealand, <sup>14</sup>National Institute for Agricultural Research, UMR Centre Alpin de Recherche sur les Réseaux Trophiques des Ecosystèmes Limniques, Thonon-Les-Bains, France, <sup>15</sup>Lammi Biological Station, University of Helsinki, Lammi, Finland, <sup>16</sup>Large Lakes Observatory, University of Minnesota, Duluth, Duluth, Minnesota, USA, <sup>17</sup>Ministry of Environment and Climate Change, Cooperative Freshwater Ecology Unit, Laurentian University of Sudbury, Sudbury, Ontario, Canada, <sup>18</sup>U.S. Geological Survey, Fort Collins Science Center, Colorado State University, Fort Collins, Colorado, USA, <sup>19</sup>Water Research Centre, Environment Institute, School of Earth and Environmental Science, University of Adelaide, Adelaide, South Australia, Australia, <sup>20</sup>Fisheries Ecosystems Advisory Services, Marine Institute, Furnace, Newport, Ireland, <sup>21</sup>Research Institute for Limnology, University of Innsbruck, Mondsee, Austria, <sup>22</sup>Environmental Research Institute, University of Waikato, Hamilton, New Zealand, <sup>23</sup>Florida Sea Grant and UF/IFAS, University of Florida, Gainesville, Florida, USA, <sup>24</sup>Department of Natural Resources, Cornell University, Ithaca, New York, USA, <sup>25</sup>International Institute for Sustainable Development-Experimental Lakes Area, Winnipeg, Manitoba, Canada, <sup>26</sup>National Aeronautics and Space Administration, Jet Propulsion Laboratory, California Institute of Technology, Pasadena, California, USA, <sup>27</sup>Institute of Biology, Irkutsk State University, Irkutsk, Russia, <sup>28</sup>Commonwealth Scientific and Industrial Research Organization Land and Water Flagship, Black Mountain, Canberra, Australia, <sup>29</sup>Institute of Agricultural and Environmental Sciences, Estonian University of Life Sciences, Tartu, Estonia, <sup>30</sup>Department of Experimental Limnology, Leibniz Institute of Freshwater Ecology and Inland Fisheries, Berlin, Germany, <sup>31</sup>Ritsumeikan University, Kusatsu, Japan, <sup>32</sup>Finnish Environment Institute, Helsinki, Finland, <sup>33</sup>NOAA/Great Lakes Environmental Research Laboratory, Ann Arbor, Michigan, USA, <sup>34</sup>Department of Water Resources and Drinking Water, Eawag (Swiss Federal Institute of Aquatic Science and Technology), Dübendorf, Switzerland, <sup>35</sup>Department of Ecology, Evolution and Marine Biology, University of California, Santa Barbara, California, USA, <sup>36</sup>Centre for Ecology & Hydrology, Bush Estate, Midlothian, UK, <sup>37</sup>Bren School of Environmental Science and Management, University of California, Santa Barbara, California, USA, <sup>38</sup>Department of Biology, University of Hamburg, Hamburg, Germany, <sup>39</sup>Hydrology Laboratory, Limnology Institute, Russian Academy of Sciences, St. Petersburg, Russian Federation, <sup>40</sup>Centre for Limnology, Institute of Agricultural and Environmental Sciences, Estonian University of Life Sciences, Tartu, Estonia, <sup>41</sup>Institute of Coastal Research, Helmholtz-Zentrum Geesthacht, Geesthacht, Germany, <sup>42</sup>Department of Earth Sciences, Royal Museum for Central Africa, Tervuren, Belgium, <sup>43</sup>Israel Oceanographic and Limnological Research, The Kinneret Limnological Laboratory, Migdal, Israel, <sup>44</sup>National Research Council Institute of Ecosystem Study, Verbania Pallanza, Italy, <sup>45</sup>Dorset Environmental Science Centre, Ontario Ministry of the Environment and Climate Change, Dorset, Ontario, Canada, <sup>46</sup>IASMA Research and Innovation Centre, Istituto Agrario di S. Michele all'Adige-Fondazione E. Mach, Trento, Italy, <sup>47</sup>Department of Natural Resources and Earth Systems Research Center, University of New

Hampshire, Durham, New Hampshire, USA, <sup>48</sup>School of Aquatic and Fishery Sciences, University of Washington, Seattle, Washington, USA, <sup>49</sup>Tahoe Environmental Research Center, Department of Civil and Environmental Engineering, University of California, Davis, California, USA, <sup>50</sup>Department of Surface Waters-Research and Management, Eawag (Swiss Federal Institute of Aquatic Science and Technology), Kastanienbaum, Switzerland, <sup>51</sup>Department of Civil Engineering, Melikah University, Kayseri, Turkey, <sup>52</sup>Department of Limnology and Biological Oceanography, Faculty of Life Sciences, University of Vienna, Vienna, Austria, <sup>53</sup>National Institute of Water and Atmospheric Research, Hamilton, New Zealand, <sup>54</sup>Department of Biology, University of Eastern Finland, Kuopio, Finland, <sup>55</sup>Seqwater, Ipswich, Queensland, Australia, <sup>56</sup>Department of Biology, Miami University, Oxford, Ohio, USA, <sup>57</sup>CAS Center for Excellence in Tibetan Plateau Earth Sciences, Chinese Academy of Sciences, Beijing, China

**Abstract** In this first worldwide synthesis of in situ and satellite-derived lake data, we find that lake summer surface water temperatures rose rapidly (global mean =  $0.34^{\circ}\text{C decade}^{-1}$ ) between 1985 and 2009. Our analyses show that surface water warming rates are dependent on combinations of climate and local characteristics, rather than just lake location, leading to the counterintuitive result that regional consistency in lake warming is the exception, rather than the rule. The most rapidly warming lakes are widely geographically distributed, and their warming is associated with interactions among different climatic factors—from seasonally ice-covered lakes in areas where temperature and solar radiation are increasing while cloud cover is diminishing ( $0.72^{\circ}\text{C decade}^{-1}$ ) to ice-free lakes experiencing increases in air temperature and solar radiation ( $0.53^{\circ}\text{C decade}^{-1}$ ). The pervasive and rapid warming observed here signals the urgent need to incorporate climate impacts into vulnerability assessments and adaptation efforts for lakes.

## 1. Introduction

Lakes hold a large majority of Earth's liquid freshwater, support enormous biodiversity, and provide key provisioning and cultural ecosystem services to people around the world. Climate change is among the greatest threats to lakes [Carpenter *et al.*, 2011], yet empirical knowledge of global lake responses remains fragmented, in need of the syntheses that already have catalyzed major climate change initiatives for marine, terrestrial, and atmospheric systems. Previous analyses have been restricted to temperature trends in either remotely sensed or in situ data, each of which have geographic and morphological biases. For example, satellite-inferred water temperature data are generally restricted to lakes  $>10,000$  ha [Schneider and Hook, 2010; MacCallum and Merchant, 2012] omitting  $>90\%$  of the world's lakes that are small and shallow and may respond differently to climate change [Wetzel, 1990; Winslow *et al.*, 2015], and previous efforts using in situ data tended to be geographically restricted, primarily in north temperate latitudes (e.g., [Livingstone and Dokull, 2001; Austin and Colman, 2007]). As a result, while there is sufficient evidence to indicate the effects of climate change for individual lakes or lake regions, there is limited understanding of large-scale spatial patterns in lake responses or how those patterns are influenced by various climatic and geomorphic factors. By integrating satellite and in situ surface water temperature trends for lakes worldwide, we were able to balance the biases inherent to each data type [Hampton, 2013], capturing broad spatial coverage as well as geomorphic variability across a range of lake sizes (Figure S1 and Table S1 in the supporting information).

Understanding the trajectories of temperature change in inland waters is a foundational step in advancing science on a broad diversity of societally important issues. Even seemingly small changes in lake temperature profoundly affect key physical and biological processes through nonlinear dynamics [Adrian *et al.*, 2009]. The diverse array of lake sizes and shapes on earth suggests that patterns of lake warming should be highly variable, both in space and time. Key drivers of surface water temperature include absorbed solar irradiance and heat exchange with the atmosphere, which is controlled by air temperature, solar radiation, humidity, ice cover, and wind [Edinger *et al.*, 1968], but is also mediated by local factors such as lake surface area and depth [Schmid *et al.*, 2014]. These morphometric factors vary enormously across lakes, and recent rates of change in climate variables are also spatially heterogeneous [Wild, 2012; Eastman and Warren, 2013; Ji *et al.*, 2014]. Accordingly, we examined relationships between lake surface temperatures and climatic and geomorphic drivers in order to better understand and predict global trends of lake warming.

## 2. Data and Methodology

### 2.1. Data Set

We used a database that incorporates lake summer surface water temperatures (LSSWT) and climate variables (air temperatures, radiation, and cloud cover) from 1985 to 2009 [Sharma *et al.*, 2015]. The database includes

LSSWT derived from in situ and/or satellite measurements, providing a global distribution of lake data. LSSWT were calculated as 3 month mean temperatures. Generally, for lakes situated in the Northern Hemisphere, summer was defined as the period 1 July–30 September (JAS), whereas in the Southern Hemisphere summer was 1 January–31 March (JFM). Exceptions were latitudes less than 23.5°, for which the JAS metric was used south of the equator, and the JFM metric was used north of the equator to avoid the cloudy wet season in the tropics allowing for an increased number of cloud-free satellite observations [Schneider and Hook, 2010; Sharma *et al.*, 2015]. The only exception to this time period were the in situ data for Toolik Lake, Alaska, for which June–August were used due to the early onset of winter in September at this high latitude. In situ data are point collected, whereas the satellite data represent an areal mean of at least 9 km<sup>2</sup>. We chose lakes for which there were at least 13 years of data (118 in situ sampled lakes and 128 satellite-sampled lakes; there were 11 lakes sampled by both in situ and satellite methods) (Table S2). These lakes had data relatively evenly distributed across the 25 year time period and were not missing data at the beginning and the end of the time period. The median record length was 22 years; the 75% quartile was 24, and the 25% quartile was 19 years.

Metadata for each lake included latitude, longitude, elevation, surface area, volume, mean depth, and maximum depth for each lake as well as climate variables [Sharma *et al.*, 2015]. Air temperature was gridded data at 0.5° resolution from Climatic Research Unit time series version 3.21. Surface solar radiation data were from the 1° × 1° satellite product from NASA/Global Energy and Water Cycle Experiment Surface Radiation Budget shortwave radiation data set version 3.0 available from 1985 to 2007. Cloud cover were from a 1° regridded version of the NOAA 5-channel Advanced Very High Resolution Radiometer cloud imagery record with percent coverage statistics derived using Pathfinder Atmosphere's Extended processing system [Heidinger *et al.*, 2010; Stubenrauch *et al.*, 2013].

## 2.2. Global Average Rate of Change and Trend Calculations

The global average warming rate for lake summer surface water temperatures was calculated following the approach used by the International Panel on Climate Change [Hartmann *et al.*, 2013a, 2013b]. For each lake, we calculated the temperature anomalies relative to its 1985–2009 mean. We then used linear regression across the annual globally averaged anomalies to determine the global LSSWT warming rate. For lakes with both in situ and satellite temperature data, we used only the in situ values. Calculations were done in R [R Development Core Team, 2014]. For each individual lake, we used Sen slopes to calculate trends in LSSWT, air temperature, cloud cover, and shortwave radiation. To obtain the most robust trends for each variable, we used all available data and did not gap-match across data sets. Sen slopes and significance were calculated in R using the “openair” package [Carslaw and Ropkins, 2012] (Table S2). A variety of sources contribute uncertainty to individual lake trends (Table S2); however, given that our approach is to compare trends across lakes, noise is more likely to obscure patterns rather than to create them.

## 2.3. Proximal Similarity Analysis

A proximal similarity analysis was completed using the Getis-Ord  $G_i^*$  statistic, and resulting maps were generated in ArcGIS 10.2. This analysis identified subcontinent regions on the globe where lake temperatures were trending similarly to surrounding lakes within that area relative to the global trend. The Getis-Ord  $G_i^*$  statistic is a z score based on two characteristics for each lake: its trend value (Sen slope) and its proximity to other lakes with similar values. This is computed by, first, summing one point value and that of a number of proximal observations within an approximately 1300 km radius (a subcontinent regional sum). That regional sum is then compared proportionally to the sum of the global data set. Such a comparison results in a z score for each observation, and observations with a regional sum significantly higher or lower than the global sum are considered to have statistically significant regional similarity above or below the global trend. In other words, statistically significant high or low z scores identify high clustering of point locations with high or low values within the data set [Getis and Ord, 1992; Ord and Getis, 1995; ESRI, 2013]. The distance of 1300 km was determined by computing the average distance that included at least 5% of the total input lake temperature points in the aggregate data set. This means that even in areas where data points are less dense (e.g., South America and Africa), one lake will still be compared in the analysis with a handful of regional neighbors. The Getis-Ord  $G_i^*$  statistic is computed using a false discovery rate correction to resolve multiple testing and spatial dependency concerns in the data set [Caldas de Castro and Singer, 2006; ESRI, 2013]. This process generates a z score and p value for each lake. For visualization purposes, we used the z score of each

lake point to generate a kernel density layer showing regions wherein exist a statistically significant density of lake points with similar temperature trends (high or low).

In our case, the hotspot analysis identified lakes with high or low temperature trends within the global data set that are in close proximity to other lakes with similarly high or low temperature trends. Because this analysis is based on the distribution of the data, an identification of a cool spot does not necessarily mean these are cooling areas. Rather, it means they are areas of concentration of lakes with trends less than the mean of the data set. Areas that are labeled nonsignificant do not have a statistically significant relationship between their trend and proximity to other lakes with similar trends.


#### 2.4. Regression Tree of Lake Temperature Trends

A regression tree analysis was performed on LSSWT trends to identify suites of factors that correspond with the warming trends observed across the widely distributed lakes in our database. Prior to conducting these analyses on LSSWT trends, we used preliminary models of interannual variation in LSSWT to assess whether the available set of environmental variables offers reasonable predictions of surface temperature in any given lake for any given year. Predictors included winter and summer mean air temperature, % cloud cover, and shortwave radiation, as well as geomorphic characteristics of the lakes (elevation, surface area, and maximum depth) and we tested three regression model approaches (Table S3). Subsequently, in the regression tree analyses, we included trends in winter and summer air temperature, % cloud cover, shortwave radiation, geomorphic characteristics, and mean winter air temperatures (as a proxy for ice cover). We included winter climate trends because some regions are experiencing greater climate change during the winter, and winter conditions can strongly influence summer water temperatures. Regression trees iteratively divide data into two homogenous, mutually exclusive groups based on a threshold in an explanatory variable while minimizing the variation (sum of squares) of the response variable within the two groups [Breiman *et al.*, 1984; De'ath and Fabricius, 2000; De'ath, 2002]. Regression trees can perform well with complex ecological data that exhibit high-order interactions, multicollinearity, and nonlinear relationships between predictor variables [De'ath and Fabricius, 2000; De'ath, 2002]. We included both in situ and satellite data for the 11 lakes that had both; and both in situ and satellite data appeared in the same leaf for each lake, improving our confidence in the resulting model. We also included a "datatype" variable that distinguished between in situ and satellite data, and this was not significant. An  $n$ -fold cross-validation procedure was used. The most parsimonious regression tree was selected by pruning the tree to the level where the complexity parameter minimized the cross-validation error [Sharma *et al.*, 2012]. The regression tree was pruned to a depth in which the complexity parameter minimized the cross validation error. The percent variation ( $R^2$ ) explained by the regression tree was calculated as follows:  $R^2 = 1 - \text{Relative Error}$  [Sharma *et al.*, 2012]. All regression trees were developed using the "rpart" package in R [R Development Core Team, 2014] and also using JMP 10 (SAS Institute, Inc.).


### 3. Results and Discussion

Our synthesis shows that lake summer surface water temperatures (LSSWT) are warming significantly, with a mean trend of  $0.34^\circ\text{C decade}^{-1}$  (95% CI: 0.16–0.52), across 235 globally distributed lakes between 1985 and 2009 (Figure 1). This warming rate is consistent with the rapid annual average increase in air temperatures ( $0.25^\circ\text{C decade}^{-1}$ ) and ocean surface temperatures ( $0.12^\circ\text{C decade}^{-1}$ ) over a similar time period (1979–2012) [Hartmann *et al.*, 2013a]. The difference between the overall trend for summer air and lake temperatures was not statistically significant across these sites, indicating broad global coherence in air and lake temperature trends. However, for individual lakes, air and lake temperature trends often diverged (Figure 2), emphasizing the importance of understanding the various factors that control lake heat budgets rather than assuming lake temperatures will respond similarly to air temperatures.


Although warming is widespread, LSSWT trends range from  $-0.7$  to  $1.3^\circ\text{C decade}^{-1}$  and show clear regional variation. Previous studies that have used only satellite data, necessarily constrained by the technology to focus on larger lakes, also reported a range of warming rates, in step with or exceeding that of air temperature [Schneider *et al.*, 2009; Layden *et al.*, 2015]. Our data set allowed exploration of a range of potential drivers across a broader suite of lakes. Within our data set, no single geographic (latitude and elevation) or morphometric factor (depth, volume, and surface area) adequately explained this variation, since correlations between these geomorphic factors and LSSWT were weak ( $<0.2$ ) to moderate ( $<0.4$ ) (Figure S2). Warm-water



**Figure 1.** Map of trends in lake summer surface temperatures from 1985 to 2009. Most lakes are warming, and there is large spatial heterogeneity in lake trends. Note that the magnitudes of cooling and warming are not the same.



**Figure 2.** Lake summer surface water temperature (LSSWT) trends varied widely. Although the slope of the linear regression line between LSSWT trends and air temperature trends was not significantly different from 1, there was wide variation in both air and lake temperature trends. LSSWT trends significant at  $p < 0.1$  are indicated by a black central dot within a data point. Included are the 1:1 line and counts ( $n$ ) and % in each quadrant. Histograms show distribution of data along that axis.



and cool-water lakes showed similar ranges of warming rates (Figure 2). Furthermore, lake warming rates were heterogeneous even within regions; both warming and cooling trends occurred in high-latitude lakes (e.g., Alaska) and in nearby lakes within several regions (e.g., Central Europe and Tibetan Plateau). Proximal similarity analyses indicated that in both the Laurentian Great Lakes region and in Northern Europe, lakes were warming significantly faster than the global average (Figure S3), confirming previous findings [Schneider and Hook, 2010; Hook et al., 2012]. In contrast, lakes in southeastern North America were warming significantly more slowly than the global average (Figure S3).

Our interannual models of LSSWT indicate good predictions of LSSWT from the available climate and geomorphic factors. The multiple regression model suggests that only elevation and winter shortwave radiation were not significant predictors of LSSWT and aggregate predictive power was high (multiple regression:  $R^2 = 0.82$ , RMSE =  $0.42^\circ\text{C}$ , Table S3). Even without taking into account factors such as wind, relative humidity, water transparency, and residence time, the model explained as much variation as interannual studies that have included additional variables [Sharma et al., 2008]. The mixed effect and year-specific multiple regressions provide further evidence that our predictor set is appropriate and powerful, and all three statistical approaches point to summer air temperature as the single most important and consistent predictor of LSSWT (Table S3). These results underscore the fact that LSSWT is under tight physical control by climate drivers and geomorphic characteristics.

This large heterogeneity in LSSWT trends is associated with diverse climate and geomorphic factors. Major climate changes over the last few decades include increases in air temperatures [Karl et al., 2015], shifts in cloud cover and type in many regions [Eastman and Warren, 2013], and increases or decreases in solar radiation at various locations around the globe [Wild, 2012]. The explanatory power of these climate variables is mediated by the morphometric properties of individual lakes that affect the efficiency of heat transfer [Toffolon et al., 2014]. In our data set, % cloud cover, air temperature, and shortwave radiation are only weakly to moderately correlated ( $<0.4$ ) suggesting that multicollinearity is minimal and that each of these variables should be used to model lake surface water temperatures.

Regression tree analysis implied both nonlinear effects and complex interactions among variables ( $R^2 = 64\%$  for full model,  $R^2 = 45\%$  for pruned model; Figure 3a). Winter ice cover appears to be a key factor influencing LSSWT trends (Figure 3a). Mean winter air temperatures of  $-0.4^\circ\text{C}$  marked the first split in the regression tree for predicting lake temperature trends. This mean winter air temperature has previously been associated with ice formation [Weyhenmeyer et al., 2004], and we found that the split accurately reflected the division between lakes in our database that become seasonally covered by ice (hereafter referred to as ice-covered lakes) versus lakes with no seasonal ice cover (ice-free lakes) (Table S4). On average, ice-covered lakes are warming significantly faster than lakes that do not experience ice cover (Wilcoxon  $p < 0.0001$ ; ice-covered median  $0.48^\circ\text{C decade}^{-1}$ , 95% CI 0.45 to 0.55; warm-winter median  $0.25^\circ\text{C decade}^{-1}$ , 95% CI 0.19 to 0.31). Among ice-covered lakes, LSSWT trends were related to both geomorphic characteristics and cloud cover trends. In contrast, LSSWT trends in ice-free lakes were more closely associated with trends in air temperature and solar radiation.

Ice-covered lakes are typically warming faster than ambient air temperatures, and lake morphology affected the strength of this response. The world's deepest ice-covered lakes warmed twice as fast as the overlying air temperatures, consistent with previous single-lake studies (e.g., [Austin and Colman, 2007; Hampton et al., 2008]). For these large, deep lakes, the combination of shorter ice duration [Magnuson, 2000] and rising air temperatures can lead to earlier summer stratification that results in surface waters warming more rapidly than air [Austin and Colman, 2007], whereas in smaller, shallower lakes, surface water temperatures should more closely track changes in air temperature [Toffolon et al., 2014]. In addition, summer shortwave radiation trends were significantly greater for ice-covered lakes (Wilcoxon  $p < 0.0001$ ) (Figure 3a), and the lakes exhibiting the highest warming rates also experienced substantial decreases in summer cloud cover (leaves A and C, Table S4). Thus, we infer that the highest warming rates occur in ice-covered lakes that are subject to a combination of shorter ice duration, decrease in cloud cover, and increase in both summer air temperature and shortwave radiation (Figure S4). A more detailed exploration of the interplay between these various climate drivers, by investigating the impact of temperature-related changes in ice-cover relative to increases in shortwave radiation, for example, would be beneficial for improving our ability to predict lake changes.

Ice-free lakes are warming more slowly, frequently at rates similar to or less than those of air temperature. This pattern accords with theoretical predictions based on the psychrometric properties of air and water, which dictate that long-term rates of temperature change should be lower for lakes than air [Schmid *et al.*, 2014]. Among ice-free lakes in our survey, the main exceptions were observed in Florida and Australia, where surface temperatures of multiple lakes changed faster than air temperatures (cooling in Florida and warming in Australia). Aside from certain small lakes (primarily) in Australia, the highest warming rates among ice-free lakes were cases where both summer air temperature and summer shortwave radiation increased (leaf H, Figure S4). Although 10% of lakes showed cooling trends (Figure 2), underlying reasons for cooling were apparently site specific (Table S2).

#### 4. Conclusion

The high level of spatial heterogeneity in lake warming rates found in this study runs counter to the common assumption of general regional coherence. Lakes for which warming rates were similar in association with particular geomorphic or climatic predictors (i.e., lakes within a “leaf”) showed weak geographic clustering (Figure 3b), contrary to previous inferences of regional-scale spatial coherence in lake warming trends [Palmer *et al.*, 2014; Wagner *et al.*, 2012]. In fact, similarly responding lakes were broadly distributed across the globe, indicating that lake characteristics can strongly mediate climatic effects. The heterogeneity in surface warming rates underscores the importance of considering interactions among climate and geomorphic factors that are driving lake responses and prevents simple statements about surface water trends; one cannot assume that any individual lake has warmed concurrently with air temperature, for example, or that all lakes in a region are warming similarly. Predicting future responses of lake ecosystems to climate change relies upon identifying and understanding the nature of such interactions.

Consequences of this extensive warming are numerous and diverse. The global average lake summer surface water warming rate found here implies a 20% increase in algal blooms and a 5% increase in toxic blooms over the next century [Brookes and Carey, 2011; Rigosi *et al.*, 2015], as well as a 4% increase in methane emissions from lakes during the next decade. Increased evaporation associated with warming can lead to declines in lake water level, with implications for water security [Vorosmarty, 2000; Hanrahan *et al.*, 2010], substantial economic consequences [Gronewold and Stow, 2014], and in some cases, complete ecosystem loss (e.g., [Smol and Douglas, 2007]). Already, changes in thermal structure and mixing have decreased productivity of some lakes, which threaten human communities that depend on fisheries as a nutritional and economic resource [O'Reilly *et al.*, 2003]. Lakes with high rates of surface temperature change may appear more likely to experience major ecosystem changes [Smol *et al.*, 2005; Smol and Douglas, 2007], but we caution that even lakes with low rates of change may be under ecosystem stress if the initial water temperatures are already near physiological maxima [Tewksbury *et al.*, 2008]. The widespread warming reported here suggests that large changes in Earth's freshwater resources and their processes are not only imminent but already under way.

#### References

- Adrian, R., *et al.* (2009), Lakes as sentinels of climate change, *Limnol. Oceanogr.*, 54(6), 2283–2297.
- Austin, J. A., and S. M. Colman (2007), Lake Superior summer water temperatures are increasing more rapidly than regional temperatures: A positive ice-albedo feedback, *Geophys. Res. Lett.*, 34, L06604, doi:10.1029/2006GL029021.
- Breiman, L., J. H. Friedman, R. A. Olshen, and C. J. Stone (1984), *Classification and Regression Trees, The Wadsworth Statistics Probability Series*, CRC Press, Boca Raton, Fla.
- Brookes, J. D., and C. C. Carey (2011), Resilience to blooms, *Science*, 334(6052), 46–47.
- Caldas de Castro, M., and B. H. Singer (2006), Controlling the false discovery rate: A new application to account for multiple and dependent tests in local statistics of spatial association, *Geogr. Anal.*, 38(2), 180–208.
- Carpenter, S. R., E. H. Stanley, and M. J. Vander Zanden (2011), State of the world's freshwater ecosystems: Physical, chemical, and biological changes, *Annu. Rev. Environ. Resour.*, 36(1), 75–99.
- Carlsaw, D. C., and K. Ropkins (2012), Openair—An R package for air quality data analysis, *Environ. Modell. Software*, 27–28, 52–61, doi:10.1016/j.envsoft.2011.09.008.
- De'ath, G. (2002), Multivariate regression trees: A new technique for modeling species-environment relationships, *Ecology*, 83(4), 1105–1117.
- De'ath, G., and K. E. Fabricius (2000), Classification and regression trees: a powerful yet simple technique for the analysis of complex ecological data, *Ecology*, 81, 3178–3192.
- Eastman, R., and S. G. Warren (2013), A 39-yr survey of cloud changes from land stations worldwide 1971–2009: Long-term trends, relation to aerosols, and expansion of the tropical belt, *J. Clim.*, 26(4), 1286–1303.
- Edinger, J. E., D. W. Duttweiler, and J. C. Geyer (1968), The response of water temperatures to meteorological conditions, *Water Resour. Res.*, 4(5), 1137–1143, doi:10.1029/WR004i005p01137.

#### Acknowledgments

Data used in this study are available in Sharma *et al.* [2013]. Funding in support of this work came from the following sources: NASA Earth Science Division ROSES INCA and Science of Terra and Aqua programs and NASA ROSES E.2; NSF awards 1147666, 1136637, 1030242, 1128040, and 1204267; USDA National Institute of Food and Agriculture Hatch 0226747; Estonian Ministry of Education and Research IUT21-2; University of Nebraska Institute of Agriculture and Natural Resources; Inter-American Institute for Global Change Research CRN3038; German Research Foundation DFG STR 499/6-1; Russian Ministry of Education and Science research project GR 01201461929, Russian Science Foundation Project 14-14-00400; and The David and Lucille Packard Foundation. We thank M. Moore and T. Kratz for support in developing this initiative, N. Barabás and D. Lofton for discussion, N. Keuler for help with statistical analyses, and K. Woo for technical assistance. D.K. Gray, S.E. Hampton, C.M. O'Reilly, and S. Sharma conceived the idea for this paper and co-led this project. D.K. Gray, S.E. Hampton, B.M. Kraemer, P.B. McIntyre, C. M. O'Reilly, J.S. Read, R.J. Rowley, S. Sharma, D. Straile, and G.A. Weyhenmeyer conducted analyses incorporated into this paper. D.K. Gray, B.M. Kraemer, C.M. O'Reilly, J.S. Read, R.J. Rowley, and S. Sharma drafted figures, tables, and headings for this paper. B. Dong, D.K. Gray, S.E. Hampton, B.M. Kraemer, P.B. McIntyre, C. M. O'Reilly, J.S. Read, R.J. Rowley, P. Schneider, and S. Sharma wrote sections of the text for this paper. All authors provided critical feedback, edits, and/or commented on drafts of this paper. J.D. Brookes, D.P. Hamilton, S. Hook, J.D. Lenters, D. Livingstone, P.B. McIntyre, C.M. O'Reilly, J.S. Read, and P. Schneider played important roles in the early development of the Global Lake Temperature Collaboration.



- ESRI (2013), ArcGIS Desktop Help 10.2. [Available at <http://resources.arcgis.com/en/help/main/10.2/>]
- Getis, A., and J. K. Ord (1992), The analysis of spatial association by use of distance statistics, *Geogr. Anal.*, *24*, 189–206, doi:10.1111/j.1538-4632.1992.tb00261.x.
- Gronewold, A. D., and C. A. Stow (2014), Water loss from the Great Lakes, *Science*, *343*(6175), 1084–1085.
- Hampton, S. E. (2013), Understanding lakes near and far, *Science*, *342*(6160), 815–816.
- Hampton, S. E., L. R. Izmeteva, M. V. Moore, S. L. Katz, B. Dennis, and E. A. Silow (2008), Sixty years of environmental change in the world's largest freshwater lake—Lake Baikal, Siberia, *Global Change Biol.*, *14*(8), 1947–1958, doi:10.1111/j.1365-2486.2008.01616.x.
- Hanrahan, J. L., S. V. Kravtsov, and P. J. Roebber (2010), Connecting past and present climate variability to the water levels of Lakes Michigan and Huron, *Geophys. Res. Lett.*, *37*, L01701, doi:10.1029/2009GL041707.
- Hartmann, D. L., et al. (2013a), *IPCC Fifth Assessment Report, Climate Change 2013: The Physical Science Basis*, edited by T. F. Stocker et al., Cambridge Univ. Press, Cambridge, U. K.
- Hartmann, D. L., et al. (2013b), *Observations: Atmosphere and Surface Supplementary Material. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by S. T. F. D. Qin et al., Cambridge Univ. Press, Cambridge, U. K.
- Heidinger, A. K., W. C. Straka, C. C. Molling, J. T. Sullivan, and X. Wu (2010), Deriving an inter-sensor consistent calibration for the AVHRR solar reflectance data record, *Int. J. Remote Sens.*, *31*(24), 6493–6517.
- Hook, S. R., R. C. Wilson, S. MacCallum, and C. J. Merchant (2012), [Global climate] Lake surface temperature [in “State of the climate in 2011”], *Bull. Am. Meteorol. Soc.*, *93*(7), S18–S19, doi:10.1175/2012BAMSStateoftheClimate.
- Ji, F., Z. Wu, J. Huang, and E. P. Chassignet (2014), Evolution of land surface air temperature trend, *Nat. Clim. Change*, *4*(6), 462–466.
- Karl, T. R., A. Arguez, B. Huang, J. H. Lawrimore, J. R. McMahon, M. J. Menne, T. C. Peterson, R. S. Vose, and H.-M. Zhang (2015), Possible artifacts of data biases in the recent global surface warming hiatus, *Science*, *348*, 1469–1472.
- Layden, A., C. Merchant, and S. MacCallum (2015), Global climatology of surface water temperatures of large lakes by remote sensing, *Int. J. Climatol.*, doi:10.1002/joc.4299.
- Livingstone, D. M., and M. T. Dokulil (2001), Eighty years of spatially coherent Austrian lake surface temperatures and their relationship to regional air temperature and the North Atlantic Oscillation, *Limnol. Oceanogr.*, *46*(5), 1220–1227.
- MacCallum, S. N., and C. J. Merchant (2012), Surface water temperature observations of large lakes by optimal estimation, *Can. J. Remote Sens.*, *38*(1), 25–45.
- Magnuson, J. J. (2000), Historical trends in lake and river ice cover in the Northern Hemisphere, *Science*, *289*(5485), 1743–1746, doi:10.1126/science.289.5485.1743.
- O'Reilly, C. M., S. R. Alin, P.-D. Plisnier, A. S. Cohen, and B. A. McKee (2003), Climate change decreases aquatic ecosystem productivity of Lake Tanganyika, Africa, *Nature*, *424*(6950), 766–768.
- Ord, J. K., and A. Getis (1995), Local spatial autocorrelation statistics: Distributional issues and an application, *Geogr. Anal.*, *27*(4), 286–306.
- Palmer, M. E., N. D. Yan, and K. M. Somers (2014), Climate change drivers coherent trends in physics and oxygen content in North American lakes, *Clim. Change*, *124*, 285–299, doi:10.1007/s10584-014-1085-4.
- R Development Core Team (2014), R: A language and environment for statistical computing, R Foundation for Statistical Computing, Vienna, Austria. [Available at <http://www.R-project.org/>]
- Rigosi, A., et al. (2015), Determining the probability of cyanobacterial blooms: the application of Bayesian networks in multiple lake systems, *Ecol. Appl.*, *25*(1), 186–199, doi:10.1890/13-1677.1.
- Schmid, M., S. Hunziker, and A. Wüest (2014), Lake surface temperatures in a changing climate: A global sensitivity analysis, *Clim. Change*, *124*(1–2), 301–315.
- Schneider, P., and S. J. Hook (2010), Space observations of inland water bodies show rapid surface warming since 1985, *Geophys. Res. Lett.*, *37*, L22405, doi:10.1029/2010GL045059.
- Schneider, P., S. J. Hook, R. G. Radocinski, G. K. Corlett, G. C. Hulley, S. G. Schladow, and T. E. Steissberg (2009), Satellite observations indicate rapid warming trend for lakes in California and Nevada, *Geophys. Res. Lett.*, *36*, L22402, doi:10.1029/2009GL040846.
- Sharma, S., S. C. Walker, and D. A. Jackson (2008), Empirical modelling of lake water-temperature relationships: A comparison of approaches, *Freshwat. Biol.*, *53*(5), 897–911.
- Sharma, S., P. Legendre, D. Boisclair, S. Gauthier, and S. J. Smith (2012), Effects of spatial scale and choice of statistical model (linear versus tree-based) on determining species–habitat relationships, *Can. J. Fish. Aquat. Sci.*, *69*, 2095–2111.
- Sharma, S., et al. (2015), A global database of lake surface temperatures collected by in situ and satellite methods from 1985–2009, *Sci. Data*, *2*.
- Smol, J. P., and M. S. V. Douglas (2007), Crossing the final ecological threshold in high Arctic ponds, *Proc. Natl. Acad. Sci. U.S.A.*, *104*(30), 12,395–12,397.
- Smol, J. P., et al. (2005), Climate-driven regime shifts in the biological communities of arctic lakes, *Proc. Natl. Acad. Sci. U.S.A.*, *102*(12), 4397–4402.
- Stubenrauch, C. J., et al. (2013), Assessment of global cloud datasets from satellites: Project and database initiated by the GEWEX radiation panel, *Bull. Am. Meteorol. Soc.*, *94*(7), 1031–1049.
- Tewksbury, J. J., R. B. Huey, and C. A. Deutsch (2008), Putting the heat on tropical animals, *Science*, *320*(5881), 1296.
- Toffolon, M., S. Piccolroaz, B. Majone, A. Soja, F. Peeters, M. Schmid, and A. Wüest (2014), Prediction of surface temperature in lakes with different morphology using air temperature, *Limnol. Oceanogr.*, *59*(6), 2185–2202.
- Vorosmarty, C. J. (2000), Global water resources: Vulnerability from climate change and population growth, *Science*, *289*(5477), 284–288, doi:10.1126/science.289.5477.284.
- Wagner, A., S. Hülsmann, L. Paul, R. J. Paul, T. Petzoldt, R. Sachse, T. Schiller, B. Zeis, J. Benndorf, and T. U. Berendonk (2012), A phenomenological approach shows a high coherence of warming patterns in dimictic aquatic systems across latitude, *Mar. Biol.*, *159*(11), 2543–2559.
- Wetzel, R. G. (1990), Land-water interfaces: Metabolic and limnological regulators, *Verh. Internat. Verein Limnol.*, *24*(1), 6–24.
- Weyhenmeyer, G. A., M. Meili, and D. M. Livingstone (2004), Nonlinear temperature response of lake ice breakup, *Geophys. Res. Lett.*, *31*, L07203, doi:10.1029/2004GL019530.
- Wild, M. (2012), Enlightening global dimming and brightening, *Bull. Am. Meteorol. Soc.*, *93*(1), 27–37.
- Winslow, L. A., J. S. Read, G. J. A. Hansen, and P. C. Hanson (2015), Small lakes show muted climate change signal in deepwater temperatures, *Geophys. Res. Lett.*, *42*, 355–361, doi:10.1002/2014GL062325.