



Bazzucchi, Ilenia, De Vito, Giuseppe, Felici, Francesco, Dewhurst, Susan, Sgadari, A. and Sacchetti, Massimo (2015) Effect of exercise training on neuromuscular function of elbow flexors and knee extensors of type 2 diabetic patients. *Journal of Electromyography and Kinesiology*, 25 (5). pp. 815-823.

Downloaded from: <http://insight.cumbria.ac.uk/2048/>

*Usage of any items from the University of Cumbria's institutional repository 'Insight' must conform to the following fair usage guidelines.*

Any item and its associated metadata held in the University of Cumbria's institutional repository Insight (unless stated otherwise on the metadata record) may be copied, displayed or performed, and stored in line with the JISC fair dealing guidelines (available [here](#)) for educational and not-for-profit activities

**provided that**

- the authors, title and full bibliographic details of the item are cited clearly when any part of the work is referred to verbally or in the written form
  - a hyperlink/URL to the original Insight record of that item is included in any citations of the work
- the content is not changed in any way
- all files required for usage of the item are kept together with the main item file.

**You may not**

- sell any part of an item
- refer to any part of an item without citation
- amend any item or contextualise it in a way that will impugn the creator's reputation
- remove or alter the copyright statement on an item.

The full policy can be found [here](#).

Alternatively contact the University of Cumbria Repository Editor by emailing [insight@cumbria.ac.uk](mailto:insight@cumbria.ac.uk).

## Accepted Manuscript

Effect of exercise training on neuromuscular function of elbow flexors and knee extensors of type 2 diabetic patients

I. Bazzucchi, G. De Vito, F. Felici, S. Dewhurst, A. Sgadari, M. Sacchetti

PII: S1050-6411(15)00144-3

DOI: <http://dx.doi.org/10.1016/j.jelekin.2015.06.008>

Reference: JJEK 1869

To appear in: *Journal of Electromyography and Kinesiology*

Received Date: 13 March 2015

Revised Date: 23 June 2015

Accepted Date: 23 June 2015



Please cite this article as: I. Bazzucchi, G. De Vito, F. Felici, S. Dewhurst, A. Sgadari, M. Sacchetti, Effect of exercise training on neuromuscular function of elbow flexors and knee extensors of type 2 diabetic patients, *Journal of Electromyography and Kinesiology* (2015), doi: <http://dx.doi.org/10.1016/j.jelekin.2015.06.008>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1 TITLE

2 **Effect of exercise training on neuromuscular function of elbow flexors and knee**  
3 **extensors of type 2 diabetic patients**

4  
5 AUTHORS

6 Bazzucchi I.<sup>1</sup>, De Vito G.<sup>2</sup>, Felici F.<sup>1</sup>, Dewhurst S.<sup>3</sup>, Sgadari A.<sup>4</sup>, Sacchetti M.<sup>1</sup>

7  
8 AFFILIATIONS

9 1. Department of Movement, Human and Health Sciences, University of Rome "Foro  
10 Italic", Piazza Lauro De Bosis 6, 00135 Rome, Italy

11 2. Institute for Sport and Health, University College Dublin, Belfield, D4, Ireland

12 3. Department of Medical and Sport Sciences, University of Cumbria, Lancaster  
13 LA13JD, United Kingdom

14 4. Department of Geriatrics, Neurosciences and Orthopedics, Catholic University of  
15 Sacred Heart, Largo Francesco Vito 2, 00168 Rome, Italy

16  
17 RUNNING TITLE

18 Diabetes and neuromuscular function

19  
20 CONTACT INFORMATION

21 \*Corresponding Author: Ilenia Bazzucchi

22 Phone: +39 06 36 733 291

23 Fax: +39 06 36 733 203

24 e-mail: [ilenia.bazzucchi@uniroma4.it](mailto:ilenia.bazzucchi@uniroma4.it)

25  
26 FUNDING

27 This work was supported by a grant from the University of Rome "Foro Italic"

28 (grant no. RIC112013).

29  
30 KEY WORDS

31 Type II Diabetes; Electromyography; Isokinetic strength; Muscle fatigue

## 1 ABSTRACT

2

3 PURPOSE: The effects of exercise training on neuromuscular function of arm and  
4 leg muscles in type 2 diabetic patients (T2D) was investigated.

5 METHODS: Eight T2D sedentary male patients ( $61.0 \pm 2.3$  years) and eight sedentary  
6 healthy age matched control subjects (H,  $63.9 \pm 3.8$  years) underwent a 16-week  
7 supervised combined endurance and resistance exercise program. Before and after  
8 training, maximal isometric (MVIC), isokinetic (15, 30, 60, 120, 180,  $240^\circ\text{s}^{-1}$ ) torque  
9 and muscle endurance of the elbow flexors (EF) and knee extensors (KE) were  
10 assessed. Simultaneously, surface electromyographic signals from biceps brachii  
11 (BB) and vastus lateralis (VL) muscles were recorded and muscle fiber conduction  
12 velocity (MFCV) estimated.

13 RESULTS: Following training, maximal torque of the KE increased during MVIC  
14 and isokinetic contractions at 15 and  $30^\circ\text{s}^{-1}$  in the T2D ( $+19.1 \pm 2.7\%$  on average;  
15  $p < 0.05$ ) but not in the H group ( $+7 \pm 0.9\%$   $p > 0.05$ ). MFCV recorded from the VL  
16 during MVIC and during isokinetic contractions at 15 and  $30^\circ\text{s}^{-1}$  increased  
17 ( $+11.2 \pm 1.6\%$  on average;  $p < 0.01$ ), but in the diabetic group only. Muscular  
18 endurance was lower in T2D ( $20.1 \pm 0.7$  s) compared to H ( $26.9 \pm 1.3$  s), with an  
19 associated increase in the MFCV slope after training in the KE muscles only.

20 CONCLUSION: The effect of a combined exercise training on muscle torque appears  
21 to be angular velocity-specific in diabetic individuals, with a more pronounced effect  
22 on KE muscles and at slow contraction velocities, along with an associated increase  
23 in the MFCV. MFCV appears to be a more sensitive marker than torque in detecting  
24 the early signs of neuromuscular function reconditioning.

25

26

27

## 1 INTRODUCTION

2 Deleterious changes in muscle contractile properties ([Oberbach et al. 2006](#)) as well  
3 as degeneration of the motor nerves ([Ijzerman et al., 2011](#)) in type 2 diabetes increase  
4 the risk of physical disability ([Andersen, 2012](#); [Drouin et al., 2009](#)), with a reduction  
5 in muscle strength, power and muscle mass being well documented in this patient  
6 population ([Leenders et al., 2013](#); [Park et al., 2006, 2009](#); [Shah et al., 2011](#); [Volpato](#)  
7 [et al., 2012](#)). Recent findings suggest that these muscular deficiencies are greater in  
8 the lower than in the upper limbs. Further, the extent of diabetes-related muscle  
9 weakness is seen to be dependent on both the muscle contraction type and velocity  
10 with the greater deficits seen at the higher contraction velocities ([Sacchetti et al.,](#)  
11 [2013](#)). This trend is similar to what is observed for the decay of muscle function that  
12 occurs with aging ([Bazzucchi et al. 2004](#)). The combination of diabetes and aging  
13 may accelerate the strength decline especially in the lower limbs and compromise the  
14 quality of life of these individuals. Indeed, older men with type 2 diabetes have been  
15 found to have a two- to three-fold increased risk of developing physical disability, in  
16 which the declines in the functional capacity of the neuromuscular system is a  
17 contributor ([Park et al., 2006](#)).

18 The benefits of exercise training for counteracting the detrimental effect of diabetes  
19 on both glyco-metabolic control and neuromuscular function have been well reported.  
20 Resistance training in particular has been seen to improve both muscle strength and  
21 mass as well as insulin sensitivity ([Brooks et al., 2007](#); [Cauza et al., 2005](#); [Dunstan et](#)  
22 [al., 2002](#); [Holten et al., 2004](#); [LeBrasseur et al., 2011](#); [Mann et al., 2014](#)). For these  
23 reasons, it may be desirable that type 2 diabetes patients would begin an appropriate  
24 exercise training program as soon as possible after diagnosis to prevent, or at least  
25 limit, the decline in neuromuscular function. More information, however, is needed  
26 on the effects of exercise training on the torque-angular velocity relationship, which  
27 reflects the fundamental mechanisms of force generation during different contractile  
28 tasks ([Larsson et al., 1979](#)). Similarly, muscle endurance, another important  
29 determinant of performance during functional task/activities of daily living, has been

1 poorly explored in relation with diabetes, with the few investigations performed  
2 leading to conflicting results ([Almeida et al., 2008](#); [Andersen et al., 2005](#); Andersen,  
3 1998; [Fritschi & Quinn, 2010](#)). The use of surface electromyography (EMG), a non-  
4 invasive measure of myoelectric activity, allows a greater understanding of the  
5 activation patterns of the muscle of interest. In particular, the propagation velocity of  
6 the action potential along the muscle fiber (muscle fiber conduction velocity, MFCV)  
7 is one of the physiological characteristics correlated to the derangement of  
8 neuromuscular function since it is directly related to sarcolemmal  
9 excitability/function (Merletti et al., 2003) with Sacchetti et al. (2013) reporting a  
10 decrease in the MFCV of the vastus lateralis muscle in diabetic patients during  
11 isometric contractions in comparison to healthy matched individuals.

12 The present study, thus, has been designed to understand to what extent qualities such  
13 as muscle strength and endurance are compromised in diabetic patients in the early  
14 phases of the disease, and whether an active lifestyle, combined with a limited  
15 number of training sessions (i.e. sustainable by all patients), is effective in preserving  
16 their neuromuscular function. Specifically, in this study a particular emphasis was  
17 paid to the characterization of the torque-velocity relationship, muscle fatigability and  
18 the myoelectric activity of diabetic individuals both pre and post training when  
19 compared to their age matched healthy counterpart.

## 1 METHODS

2 **Subjects.** Eight type 2 diabetic patients (T2D) and eight healthy control subjects (H)  
3 gave their informed consent to participate in the study, which was approved by the  
4 local ethics committee. All diabetic patients were treated with diet and oral  
5 hypoglycemic agents (metformin) but not insulin or other drugs, were free from  
6 clinical signs of diabetic peripheral neuropathy and had relatively short history of the  
7 diseases (mean diabetes duration of  $5.2 \pm 1.3$  years). They were accepted for study  
8 only if they had total HbA<sub>1c</sub> levels <9% on therapy. Subjects' main characteristics are  
9 shown in table 1. None of the subjects were involved in a regular exercise program  
10 for at least 6 months before entering the study. All the age-matched healthy subjects  
11 had normal glucose tolerance (assessed by a 75-g oral glucose tolerance test), and  
12 none were taking any medication.

13

14 **Exercise training program.** All subjects underwent a 16-week exercise training  
15 program, designed following the ACSMs guidelines for exercise participation in  
16 individuals with type 2 diabetes (ACSM 2010). Exercise was performed 3 times  
17 weekly under the direct supervision of a sport scientist. Each session started with a  
18 warm-up consisting in 10-min low intensity endurance exercises on a cycle ergometer  
19 or a treadmill and 3 sets of 15 abdominal crunches separated by 2-min rest. The core  
20 of the training session incorporated aerobic training followed by the resistance  
21 training. Subjects rested for 4 minutes between the two modalities of training. At the  
22 end of the training, subjects performed also 10 minutes of cool down with stretching  
23 exercises. The aerobic training was performed on a treadmill or a bicycle ergometer.  
24 In order to equally distribute sessions between treadmill and bicycle ergometer, all  
25 subjects carried out 2 sessions on treadmill and 1 on bicycle ergometer in week 1 and  
26 *vice versa* 2 sessions on bicycle ergometer and 1 on treadmill in week 2. This  
27 alternation was repeated for the subsequent weeks. Participants progressed from 20  
28 minutes per session at 40-60% of the heart rate reserve (HRR; weeks 1-8) to 40  
29 minutes per session at 60-80% of the HRR (weeks 9-16). Heart rate monitors (Polar,

1 Finland) were used to adjust the workload in order to achieve the target heart rate. For  
2 the resistance training, 1 repetition maximum (1-RM) was assessed twice on 5  
3 different weight machines (leg press, leg extension, bench press, cable curl, cable pull  
4 down). The training loads were calculated with respect to the highest 1-RM value  
5 obtained at baseline and at week 9. Resistance training consisted of 3 sets of 10  
6 repetitions at a load progressing from 60 to 80% of 1RM.

7

8 **Overview of the experimental protocol.** Each subject visited the laboratory on three  
9 occasions. In the first visit, subjects were familiarized with the experimental  
10 procedures. Participants then returned to the laboratory on two additional days, the  
11 first before the training period (PRE) and the second 5 days after the 16 weeks of  
12 training program (POST). The same experimental protocol was followed both PRE  
13 and POST. The elbow flexion (EF) and knee extension (KE) torques of the dominant  
14 limb were measured with a dynamometer (Kin-Com, Chattanooga, USA).  
15 Participants were seated comfortably on the dynamometer and stabilized by chest,  
16 waist and thigh straps. The elbow angle was fixed at 90° (180°, full extension) with  
17 the upper arm parallel to the trunk and the forearm in a neutral position (halfway  
18 between pronation and supination). The wrist was secured in a padded cuff attached  
19 to the load cell. The rotational center of the lever arm was aligned to the distal lateral  
20 epicondyle of the humerus. The knee joint was set at a 90° angle (180°, full  
21 extension) as well as the hip joint. The lower leg was attached to the lever arm of the  
22 dynamometer with the ankle secured in a resistance pad. The center of rotation of the  
23 lever arm was aligned to the lateral femoral epicondyle of the knee.

24 The surface electromyographic signals (EMG) were recorded with a linear array of  
25 four electrodes (silver bars 5 mm long, 1 mm thick, 10 mm apart; OTBioelettronica,  
26 Turin, Italy) from the biceps brachii (BB) and from the vastus lateralis (VL) muscles.  
27 These two muscles were considered as representative of upper and lower limbs  
28 muscle respectively as previously reported ([Bazzucchi et al. 2004](#); [Harwood et al.](#)  
29 [2008](#); [Theou et al. 2013](#)). After gentle skin abrasion and cleaning with ethyl alcohol,



1 electrodes were attached on the skin over the BB along the line connecting the  
2 acromion to the cubital fossa, and over the VL on the line from the anterior spina  
3 iliaca superior to the lateral side of the patella. The optimal position and orientation  
4 of the electrodes were determined to be conveniently distant from the innervation  
5 zone and the tendon after checking that there was clear propagation in one direction  
6 of the action potentials without change in shape.(Bazzucchi et al. 2005). A reference  
7 electrode was placed around the wrist and ankle of the contralateral limb,  
8 respectively. To ensure the same electrode placement throughout the two  
9 experimental sessions, individual maps of the upper arm were made on transparent  
10 plastic by marking the position of permanent skin blemishes with respect to the  
11 electrodes. Three EMG signals were detected in a single-differential mode. Two  
12 double-differentials were computed off-line and were used for further analysis.  
13 Signals were amplified (x1000), band-pass filtered (10 Hz to 450 Hz), sampled at  
14 2048 Hz (EMG-USB2 amplifier, OTBioelettronica, Turin, Italy), recorded and stored  
15 on a personal computer.

16

17 **Experimental Tests.** During the test trial, the following parameters were evaluated:  
18 (1) maximal voluntary isometric contractions (MVIC); (2) isokinetic concentric  
19 contractions; (3) isometric fatiguing task.

20 1) MVIC. The joint angle was fixed at 90° (180°, full extension) for both the elbow  
21 and the knee. The MVIC task consisted of rapidly increasing the force exerted to a  
22 maximum. Visual feedback was provided to the subjects by setting a target line on  
23 the computer screen at a value 20% higher than the best MVIC. All subjects were  
24 verbally encouraged to exceed the target force, producing a maximal contraction “as  
25 hard as possible” and to maintain it for at least 2–3 s before relaxing. A minimum of  
26 three maximal attempts were performed separated by 4 min to recover from fatigue.  
27 Participants were asked to perform further attempts if the MVIC of their last trial  
28 exceeded the previous trials by at least 10%. However, in no instance the number of  
29 MVIC attempts exceeded four per subject.

1 2) Isokinetic Concentric Contractions. After the MVIC task, the torque–velocity  
2 curve was assessed. Angular velocity values were fixed at 15°, 30°, 60°, 120°, 180°  
3 and 240° s<sup>-1</sup>, and subjects were requested to flex the elbow or to extend the knee “as  
4 hard as possible”. The range of motion (ROM) for elbow flexion and knee extension  
5 was 40°-130° and 80°-170° respectively, which included the angle at which the  
6 maximal torque for each joint was reached. The order of the trials was randomized to  
7 minimize the effect of skill acquisition. Each contraction was followed by a 5-min  
8 rest to prevent cumulative fatigue. In each trial strong verbal encouragement was  
9 given by the test leader.

10 3) Isometric Fatiguing Task. An isometric contraction set at 80% of the maximum  
11 force value obtained during the MVIC was performed at the end of the session. A  
12 horizontal target band was displayed on a PC monitor. Participants were requested to  
13 match the target and to hold the force as long as possible (exhaustion). The end of the  
14 exercise was determined when the torque value dropped more than 10% below the  
15 target for 3 s. Trials were also interrupted if participants reported pain or any  
16 discomfort.

17  
18

19 **Data Analysis.** All data collected during the experiments were analyzed off-line (OT  
20 BioLab, OTBioelettronica, Turin, Italy). For the MVIC task, the trial which showed  
21 the highest value for force was chosen for the analysis. For each contraction speed  
22 tested, subjects performed three repetitions, and the one that produced the highest  
23 torque value was used for analysis. Peak torques assessed during MVIC and  
24 isokinetic contractions were used to assess the torque–velocity relationship. EMG  
25 signals were recorded simultaneously to mechanical data. Trials chosen for MFCV  
26 estimation were selected on the basis of maximal force. Maximal MFCV was  
27 estimated from the two double-differentials over 250ms-windows by means of the  
28 cross-correlation technique ([Sbriccoli et al. 2003](#); [Bazzucchi et al. 2005](#)). The cross  
29 correlation function technique was used to estimate the time delay between the two

1 signals (i.e., the amount of time shift that must be applied to one signal to minimize  
2 the mean square error with the other). This time shift is the same, which maximizes  
3 the cross correlation between the signals ([Naeije and Zorn 1982](#)). Estimates of MFCV  
4 were accepted only when cross-correlation values were  $>0.8$ .

5 For the isometric fatiguing task, parameters of interest were the time to fatigue (TTF)  
6 and MFCV slope. A linear regression was applied to the scattered MFCV data. The  
7 rate of change of MFCV ( $\% s^{-1}$ ) was defined as the percentage ratio between the  
8 slopes of these regression lines and their initial values at time 0. The value at time 0  
9 was calculated as the mean of the first three seconds ([Sbriccoli et al. 2003](#)).

10

11 **Reliability.** Previous test-retest reliability from our laboratory for Torque and MFCV  
12 calculated from isokinetic tests performed 2 to 10 days apart, indicated that the  
13 intraclass correlation coefficients (*ICC*) ranged from 0.89 to 0.96 and 0.74 to 0.93  
14 respectively, with no significant ( $p > 0.05$ ) differences between mean test vs retest  
15 values.

16

17 **Statistical analysis.** All statistical analyses were performed with PASW statistics  
18 20.0 (SPSS Inc, Chicago, Illinois, USA). Standard methods of descriptive analysis  
19 were used for calculation of means and standard deviations (SD) and to test the  
20 normal distribution of variables. When the sphericity assumption was violated, the  
21 Greenhouse-Geisser adjustment was performed. Subsequently, a repeated-measures  
22 analysis of variance (RM-ANOVA) with limb [arm vs leg] and angular velocity [0  
23 (MVIC), 15, 30, 60, 120, 180, 240 $^{\circ}\cdot s^{-1}$ ] as within factors, was used to compare the  
24 dependent variables [torque and MFCV] obtained from the two groups [T2D vs H]  
25 following the 16-week training [PRE vs POST]. In addition, a one-way ANOVA with  
26 limb [arm vs leg] as a within factor was used to determine the effect of training on  
27 torque, MFCV, TTF and MFCV slopes obtained by the two groups of individuals  
28 during the isometric fatiguing task. When significant effects were found, T-tests with  
29 Bonferroni correction for multiple comparisons were performed as follow-up

1 analyses. An alpha of  $p < 0.05$  was considered significant for all comparisons. An a  
2 priori analysis was used to determine a sample size that yielded power values of 0.80  
3 or greater. Data are expressed as mean  $\pm$  SE. Regression lines for individual data sets  
4 of torque vs angular velocity were computed using the least-squares method.

## 5 RESULTS

### 6 **Torque-velocity relationships**

7 Torque-velocity relationships of elbow flexors (EF) and knee extensors (KE) muscles  
8 of the two groups before and after the training period are depicted in fig. 1.

9 After the training program, maximal torque of EF in H and T2D was unchanged, in  
10 both static and dynamic conditions. Differently, a significant interaction was found  
11 among angular velocity, training and group for torque ( $p = 0.02$ ) values in KE in T2D.  
12 More specifically, KE torque-velocity relationship of T2D was shifted towards higher  
13 values after the training program, with a higher effect ( $p < 0.05$ ) observed during  
14 MVIC ( $+9.3 \pm 4\%$ ) and during dynamic contractions conducted at 15 ( $+22.9 \pm 8.5\%$ )  
15 and  $30^\circ \text{ s}^{-1}$  ( $+25.2 \pm 8.2\%$ ).

16

17

*FIG.1 app. here*

18

19

20

### 21 **Muscle fiber conduction velocity**

22 MFCV values recorded during static and dynamic contractions are depicted in fig. 2.

23 A significant interaction was found among angular velocity, training and group for  
24 MFCV ( $p < 0.05$ ) values in KE. After training significantly higher values of MFCV  
25 were found at MVIC and  $30^\circ \text{ s}^{-1}$  ( $p < 0.05$ ) in the EF and at 15, 30, 60,  $90^\circ \text{ s}^{-1}$  ( $p < 0.01$ )  
26 in KE of T2D group. In this group, the MFCV enhancement in KE following the  
27 training was  $+11.2 \pm 1.6\%$  on average.

28

29

1

2

*FIG.2 app. here*

3

4

5

**6 Muscular Endurance**

7 Figure 3 shows time to fatigue (TTF) recorded during the isometric fatiguing  
8 contractions. TTF was  $26.9 \pm 1.3$  s for H and  $20.1 \pm 0.7$  for T2D with no effect of the  
9 muscle group (EF vs KE) and of training (PRE vs POST). A significant interaction  
10 was found between muscle group (EF vs KE) and condition (T2D vs H). In particular,  
11 T2D patient showed significant lower TTF values with respect to H subjects in both  
12 PRE and POST trials in KE muscles ( $p < 0.05$ ).

13 MFCV slopes calculated over the isometric fatiguing contractions are depicted in fig.  
14 4. No significant differences were found between PRE and POST session in EF for  
15 both groups (H and T2D). For T2D group, a higher MFCV slopes was found after  
16 training in the KE muscles ( $p < 0.01$ ).

17

18

19

*FIG. 3 and 4 app. here*

20

21

22

23

24

25

## 1 DISCUSSION

2 The present study aimed at comparing the effect of a 16-week whole-body combined  
3 endurance and resistance training on neuromuscular function of the upper and lower  
4 limb in type 2 diabetic patients. The main findings indicated that the training program  
5 was successful in increasing the maximal torque generated by the KE muscles at slow  
6 contraction velocities along with an associated increase in the MFCV in diabetic  
7 individuals. In addition, T2D patients showed lower TTF than H individuals during  
8 isometric fatiguing contractions of KE before and after training, with an associated  
9 increase in the MFCV slopes seen only after training.

10

11 *Muscle strength*

12 One of the most important functional consequences of diabetes is muscle weakness.  
13 Reduced muscle strength of the knee extensors muscles in T2D patients, has been  
14 reported during maximal isokinetic ([Andersen et al. 1996](#); [Park et al. 2007](#); [Kalyani et al. 2013](#)) and isometric tasks ([Ijzerman et al. 2012](#)) while a comparable isokinetic  
15 strength with respect to healthy controls has been found in muscles acting around the  
16 elbow ([Andersen et al. 2004](#)) and the wrist ([Andreassen et al. 2006](#); [Park et al. 2007](#)).  
17 A greater detrimental effect of type II diabetes on the torque-velocity relationship of  
18 leg muscles with respect to arm muscles has been recently provided ([Sacchetti et al., 2013](#)). Moreover, in that study, using a cross-sectional design, it was found that  
19 trained diabetic patients showed a less pronounced neuromuscular impairment with  
20 respect to their sedentary counterparts, which highlighted the need for investigating  
21 the effect of exercise training programs on neuromuscular function of diabetic  
22 patients.  
23 patients.

24  
25 To the best of our knowledge the present is the first study focusing on the effect of  
26 exercise training on the torque-velocity relationship (and myoelectric activity) of  
27 upper and lower limb muscles of T2D patients.

28 In line with several previous investigations ([Dunstan et al. 2002](#); [Ibáñez et al. 2008](#);  
29 [Kwon et al. 2010](#); [Larose et al. 2010](#)), our findings showed an improvement of

1 muscle strength of T2D patients after training. However, in the present study we  
2 demonstrate that the training was only effective in isometric contractions and  
3 dynamic contractions of lower velocity. While many day-to-day functional tasks  
4 require force production at these low velocities, such as rising from a chair or  
5 walking up stairs, there may be occasions when faster velocity contractions are  
6 required such as recovering from a stumble or grab an object that is falling. This may  
7 be a reflection on the training modality in which the resistance training exercises  
8 were completed at the slower velocities rather than in an explosive movement.  
9 However, this is worthy of consideration for exercise practitioners developing a  
10 suitable training program for muscle function in type 2 diabetic individuals.

11

12 An interesting finding in the present study was the lack of improvement in the muscle  
13 function of the EF, despite an improvement in the KE. The upper and lower  
14 extremities have been previously shown to be affected by type 2 diabetes to different  
15 extents, with lower limbs showing a more pronounced impairment of strength and  
16 greater metabolic derangement ([Andersen et al., 2004](#); [Christer . et al., 2006](#); [Ibáñez  
17 et al., 2008](#); [Ijzerman et al., 2012](#); [Won Park et al., 2006, 2007](#); [Olsen et al., 2005](#);  
18 [Sacchetti et al., 2005](#)). The higher neuromuscular impairment in the KE could be  
19 responsible for a lower baseline muscular capacity compared to the EF and, thus, a  
20 greater “gap to fill” with training. This larger gap would make more evident the effect  
21 of a moderate training regime, as the one adopted in the present study. Indeed, the  
22 exercise modality also deserves consideration. The participants were involved in a  
23 combined endurance and resistance training program, with the aerobic exercise  
24 performed on a treadmill or on a cycle ergometer. Therefore, while resistance  
25 exercises were equally distributed between the upper and lower body districts, the  
26 aerobic training was mainly focusing on the lower body, representing an additional  
27 stimulus for the musculature of the lower limbs. This factor could have contributed to  
28 the greater response of the KE to the 16-week combined exercise training compared

1 to the EF, which represents a further consideration for in the exercise prescription for  
2 these patients.

3

#### 4 *Muscle fiber conduction velocity*

5 In the present study, we explored the modifications in the propagation velocity of the  
6 action potential along the muscle fibers during maximal isometric and isokinetic  
7 contractions before and after the 16-week exercise program. To the best of our  
8 knowledge, this is the first longitudinal study investigating the exercise-induced  
9 modifications of the torque-velocity relationship with a concomitant estimation of the  
10 MFCV in T2D patients.

11 MFCV depends, among other factors, on sarcolemmal excitability (Arendt-Nielsen  
12 and Zwarts 1989; [Linssen et al. 1996](#); Chisari et al. 1998), morphological and  
13 functional characteristics of muscle fibers (Arendt-Nielsen and Zwarts 1989; Kupa et  
14 al. 1995; [Almeida et al. 2008](#)) and can be influenced by changes in neuromuscular  
15 recruitment strategies occurring independently of motor nerve dysfunction (Sacchetti  
16 et al., 2013). In particular, the T2D patients involved in the present study did not  
17 present clinical signs of neuropathy and their median duration of diabetes was very  
18 short, so it may be plausible that their baseline MFCV values were not much different  
19 from the ones of healthy subjects. Studies on the  $\text{Na}^+/\text{K}^+$  pump function, in fact,  
20 demonstrated no significant differences between control subjects and T2D patients  
21 with no evidence of neuropathy at baseline and following normal pump activity  
22 ([Arnold et al. 2013](#)) and activity-dependent changes similar to those observed in  
23 healthy men in the post-contraction period. On the other hand, alterations in  $\text{Na}^+/\text{K}^+$   
24 pump function coupled with reductions in nodal  $\text{Na}^+$  currents have been hypothesized  
25 to be responsible for the slower recovery following maximal voluntary contractions  
26 [in diabetic neuropathy \(Krishnan et al. 2008\)](#).

27 After the training period, an increase in MFCV values of diabetic patients in the VL  
28 was obtained, keeping with the increase of the knee extensors muscles' strength. To  
29 note, even if limited to the lowest contraction speeds, we also found an increased



1 MFCV in the BB muscle of T2D patients, which was not associated to changes in the  
2 torque-velocity curve of the elbow flexors after training. This points toward an  
3 exercise-induced restoration of neuromuscular function which is, again, more evident  
4 in the lower limb likely due to the greater impairment of muscle quality of those body  
5 regions (Park et al., 2006). This, together with the training-induced improvements in  
6 MFCV in the BB despite an unchanged torque, suggests that MFCV may be useful  
7 for unmasking early signs of neuromuscular dysfunction and reconditioning in  
8 diabetic patients.

9

### 10 *Muscle fatigue*

11 Besides the reduction of muscle strength, diabetic patients may suffer from a  
12 reduction in muscle endurance, and this higher muscle fatigability may have a  
13 negative impact on activities of daily living, such as carrying shopping bags or  
14 climbing a long set of stairs. The number of studies investigating muscle endurance in  
15 relation with diabetes is limited ([Andersen 1998](#); [Almeida et al. 2008](#); [Shah et al.](#)  
16 [2011](#); [Ijzerman et al. 2012](#)) and the results are conflicting partially due to the different  
17 testing procedure adopted (isometric vs concentric contractions), the limb tested  
18 (upper vs lower) and the diabetic population (type 1 vs type 2) considered.

19 In the present study, T2D patients showed a lower TTF during the isometric fatiguing  
20 contractions of KE muscles compared to H, both pre and post training. Of note, after  
21 the training period, this was also associated to higher MFCV slopes. In normal  
22 conditions, during sustained voluntary contractions MFCV gradually declines, which  
23 is related to modifications in muscle membrane excitability ([Chisari et al. 1998](#)). The  
24 time course of this MFCV decline has been shown to be higher in muscles with a  
25 higher proportion of type II muscle fibers ([Kupa et al. 1995](#)). Taken together, these  
26 findings of a higher fatigability and MFCV slope in T2D appears to support the  
27 previously reported shift in muscle fiber type composition with diabetes, with a  
28 higher proportion of fast-twitch fibers at the expenses of slow-twitch ones ([Saltin](#)  
29 [et al. 1977](#); [Krotkiewski and Bjorntorp 1986](#); [Lillioja et al. 1987](#); [Gaster et al. 2001](#);

1 Oberbach et al. 2006). The change in contractile properties is further described by  
2 animal studies reporting a more marked diabetes-induced strength loss in fast-twitch  
3 muscles ([Cotter et al. 1989](#); [Sanchez et al. 2005](#)) and in skinned fast-twitch single  
4 fibers ([Paulus and Grossie 1983](#); [Sanchez et al. 2005](#)) than in slow-twitch muscles.  
5 Thus, it is tempting to speculate that the KE muscles of our patients could have been  
6 characterized by a higher proportion of type II fibers with respect to the arm muscles  
7 in reason of a greater diabetes-induced impairment as suggested by some ([Gaster et](#)  
8 [al., 2001](#); [Hickey et al., 1995](#); [Mårin et al., 1994](#); [Mogensen et al., 2007](#); [Oberbach et](#)  
9 [al., 2006](#); [Segerström et al., 2011](#); [Stuart et al., 2013](#)) but not all ([Andreassen et al.,](#)  
10 [2014](#); [Cederholm et al., 2000](#); [He et al., 2001](#); [Leenders et al., 2013](#); [Zierath et al.,](#)  
11 [1996](#)) studies. This would be reflected by lower TTF due to the greater fatigability of  
12 glycolytic fibers, by the higher MFCV slopes during the fatiguing contractions, as  
13 well as by the greater training effect on the torque-velocity relationship.

14

15 The present study has several practical implications. The selective effect of our  
16 conventional training program on the torque-velocity relationship highlight the need  
17 for implementing specific exercise protocols adopting different contraction velocities  
18 to selectively counteract the diabetic-induced neuromuscular deficiencies. The  
19 present data support our previous suggestion of considering different angular  
20 velocities when testing neuromuscular function in diabetes patients, as well as the  
21 effect of specific training regimes. Finally, MFCV appears to be a sensitive parameter  
22 for describing some of the modifications induced on neuromuscular functions by type  
23 2 diabetes, even at its early stage.

24

25 Conclusions:

26 In conclusion, the present training study adopting conventional exercise modalities  
27 for diabetic patients indicates that individuals at the initial stage of the disease show a  
28 higher improvement of neuromuscular function in the lower than in the upper limb  
29 muscles. The training effect appears to be angular velocity-specific, which calls for

1 adopting specifically structured exercise modalities. Finally, the velocity of  
2 propagation of the action potential along the muscle fibers appears to be a more  
3 sensitive marker than torque in detecting the early signs of neuromuscular function  
4 reconditioning.

5

6

7

## 1 ACKNOWLEDGEMENTS

2 The authors are grateful to all the volunteers who spent several hours participating in  
3 this study. We appreciate the assistance of Alessandra Conti with the measurements.

4

5

6

ACCEPTED MANUSCRIPT

## REFERENCES

- ACSM. ACSM Guidelines for Exercise Testing and Prescription, 8th Edition. Med. Sci. Sport. Exerc. 2010. p. 2018.
- Almeida S, Riddell MC, Cafarelli E. Slower conduction velocity and motor unit discharge frequency are associated with muscle fatigue during isometric exercise in type 1 diabetes mellitus. Muscle and Nerve. 2008;37:231–40.
- Andersen H. Muscular endurance in long-term IDDM patients. Diabetes Care. 1998;21:604–9.
- Andersen H. Motor dysfunction in diabetes. Diabetes Metab Res Rev. 2012;28 Suppl 1:89–92.
- Andersen H, Nielsen S, Mogensen C, Jakobsen J. Muscle strength in type 2 diabetes. Diabetes. 2004;53:1543–8.
- Andersen H, Poulsen PL, Mogensen CE, Jakobsen J. Isokinetic muscle strength in long-term IDDM patients in relation to diabetic complications. Diabetes. 1996;45:440–5.
- Andersen H, Schmitz O, Nielsen S. Decreased isometric muscle strength after acute hyperglycaemia in Type 1 diabetic patients. Diabet Med. 2005;22:1401–7.
- Andreassen CS, Jakobsen J, Andersen H. Muscle weakness: A progressive late complication in diabetic distal symmetric polyneuropathy. Diabetes. 2006;55:806–12.
- Andreassen CS, Jensen JM, Jakobsen J, Uthøj BP, Andersen H. Striated muscle fiber size, composition, and capillary density in diabetes in relation to neuropathy and muscle strength. J Diabetes. 2014;
- Arendt-Nielsen L, Zwarts M. Measurement of muscle fiber conduction velocity in humans: techniques and applications. J Clin Neurophysiol. 1989;6:173–90.
- Arnold R, Kwai N, Lin CSY, Poynten AM, Kiernan MC, Krishnan A V. Axonal dysfunction prior to neuropathy onset in type 1 diabetes. Diabetes Metab Res Rev. 2013;29:53–9.

- Bazzucchi I, Felici F, Macaluso A, De Vito G. Differences between young and older women in maximal force, force fluctuations, and surface EMG during isometric knee extension and elbow flexion. Muscle and Nerve. 2004;30(5):626–35.
- Bazzucchi I, Marchetti M, Rosponi A, Fattorini L, Castellano V, Sbriccoli P, et al. Differences in the force/endurance relationship between young and older men. Eur J Appl Physiol. 2005;93(4):390–7.
- Brooks N, Layne JE, Gordon PL, Roubenoff R, Nelson ME, Castaneda-Sceppa C. Strength training improves muscle quality and insulin sensitivity in Hispanic older adults with type 2 diabetes. Int J Med Sci. 2007;4:19–27.
- Cauza E, Hanusch-Enserer U, Strasser B, Ludvik B, Metz-Schimmerl S, Pacini G, et al. The relative benefits of endurance and strength training on the metabolic factors and muscle function of people with type 2 diabetes mellitus. Arch Phys Med Rehabil. 2005;86:1527–33.
- Cederholm T, Sylven C, Esbjornsson-Liljedahl M, Jansson E. Insulin treatment increases skeletal muscle fibre area in patients with diabetes mellitus type 2. Clin Physiol. 2000;20:354–9.
- Chisari C, D’Alessandro C, Manca ML, Rossi B. Sarcolemmal excitability in myotonic dystrophy: Assessment through surface EMG. Muscle and Nerve. 1998;21:543–6.
- Cotter M, Cameron NE, Lean DR, Robertson S. Effects of long-term streptozotocin diabetes on the contractile and histochemical properties of rat muscles. Q J Exp Physiol. 1989;74:65–74.
- Drouin P, Blicke JF, Charbonnel B, Eschwege E, Guillausseau PJ, Plouin PF, et al. Diagnosis and classification of diabetes mellitus. Diabetes Care. 2009;32 Suppl 1:S62–7.
- Dunstan DW, Daly RM, Owen N, Jolley D, De Courten M, Shaw J, et al. High-intensity resistance training improves glycemic control in older patients with type 2 diabetes. Diabetes Care. 2002;25:1729–36.
- Eriksson J, Taimela S, Eriksson K, Parviainen S, Peltonen J, Kujala U. Resistance training in the treatment of non-insulin-dependent diabetes mellitus. Int J Sports Med. 1997;18:242–6.

- Fritschi C, Quinn L. Fatigue in patients with diabetes: A review. J. Psychosom. Res. 2010. p. 33–41.
- Gaster M, Staehr P, Beck-Nielsen H, Schrøder HD, Handberg A. GLUT4 is reduced in slow muscle fibers of type 2 diabetic patients: is insulin resistance in type 2 diabetes a slow, type 1 fiber disease? Diabetes. 2001;50:1324–9.
- Harwood B, Edwards DL, Jakobi JM. Age- and sex-related differences in muscle activation for a discrete functional task. Eur J Appl Physiol. 2008;103:677–86.
- He J, Watkins S, Kelley DE. Skeletal muscle lipid content and oxidative enzyme activity in relation to muscle fiber type in type 2 diabetes and obesity. Diabetes. 2001;50:817–23.
- Herman Ijzerman T, Schaper NC, Melai T, Blijham P, Meijer K, Willems PJB, et al. Motor nerve decline does not underlie muscle weakness in type 2 diabetic neuropathy. Muscle and Nerve. 2011;44:241–5.
- Hickey MS, Carey JO, Azevedo JL, Houmard JA, Pories WJ, Israel RG, et al. Skeletal muscle fiber composition is related to adiposity and in vitro glucose transport rate in humans. Am J Physiol. 1995;268:E453–7.
- Holten MK, Zacho M, Gaster M, Juel C, Wojtaszewski JFP, Dela F. Strength Training Increases Insulin-Mediated Glucose Uptake, GLUT4 Content, and Insulin Signaling in Skeletal Muscle in Patients with Type 2 Diabetes. Diabetes. 2004;53:294–305.
- Ibáñez J, Gorostiaga EM, Alonso AM, Forga L, Argüelles I, Larrión JL, et al. Lower muscle strength gains in older men with type 2 diabetes after resistance training. J Diabetes Complications. 2008;22:112–8.
- Ijzerman TH, Schaper NC, Melai T, Meijer K, Willems PJB, Savelberg HHCM. Lower extremity muscle strength is reduced in people with type 2 diabetes, with and without polyneuropathy, and is associated with impaired mobility and reduced quality of life. Diabetes Res Clin Pract. 2012;95:345–51.
- Ishii T, Yamakita T, Sato T, Tanaka S, Fujii S. Resistance training improves insulin sensitivity in NIDDM subjects without altering maximal oxygen uptake. Diabetes Care. 1998;21:1353–5.
- Kalyani RR, Tra Y, Yeh HC, Egan JM, Ferrucci L, Brancati FL. Quadriceps Strength, Quadriceps Power, and Gait Speed in Older US Adults with Diabetes Mellitus:

Results from the National Health and Nutrition Examination Survey, 1999-2002. *J Am Geriatr Soc.* 2013;61:769-75.

Krishnan A V., Lin CSY, Kiernan MC. Activity-dependent excitability changes suggest Na<sup>+</sup>/K<sup>+</sup> pump dysfunction in diabetic neuropathy. *Brain.* 2008;131:1209-16.

Krotkiewski M, Bjorntorp P. Muscle tissue in obesity with different distribution of adipose tissue. Effects of physical training. *Int J Obes.* 1986;10:331-41.

Kupa EJ, Roy SH, Kandarian SC, De Luca CJ. Effects of muscle fiber type and size on EMG median frequency and conduction velocity. *J Appl Physiol.* 1995;79:23-32.

Kwon HR, Han KA, Ku YH, Ahn HJ, Koo B-K, Kim HC, et al. The effects of resistance training on muscle and body fat mass and muscle strength in type 2 diabetic women. *Korean Diabetes J.* 2010;34:101-10.

Larose J, Sigal RJ, Boulé NG, Wells G a, Prud'homme D, Fortier MS, et al. Effect of exercise training on physical fitness in type II diabetes mellitus. *Med Sci Sport Exerc.* 2010;42:1439-47.

LeBrasseur NK, Walsh K, Arany Z. Metabolic benefits of resistance training and fast glycolytic skeletal muscle. *Am J Physiol Endocrinol Metab.* 2011;300:E3-10.

Leenders M, Verdijk LB, van der Hoeven L, Adam JJ, van Kranenburg J, Nilwik R, et al. Patients with type 2 diabetes show a greater decline in muscle mass, muscle strength, and functional capacity with aging. *J Am Med Dir Assoc.* 2013;14:585-92.

Lillioja S, Young AA, Culter CL, Ivy JL, Abbott WG, Zawadzki JK, et al. Skeletal muscle capillary density and fiber type are possible determinants of in vivo insulin resistance in man. *J Clin Invest.* 1987;80:415-24.

Linssen WHJP, Stegeman DF, Merks MJH, Binkhorst RA, Notermans SLH. Electromyographic evidence of delayed fatigue-induced sarcolemmal excitability impairment in McArdle's disease. *J Electromyogr Kinesiol.* 1996;6:147-57.

Mann S, Beedie C, Balducci S, Zanuso S, Allgrove J, Bertiatto F, et al. Changes in insulin sensitivity in response to different modalities of exercise : a review of the evidence. *Diabetes Metab Res Rev.* 2014;30:257-68.



- Mårin P, Andersson B, Krotkiewski M, Björntorp P. Muscle fiber composition and capillary density in women and men with NIDDM. Diabetes Care. 1994;17:382–6.
- Mogensen M, Sahlin K, Fernström M, Glintborg D, Vind BF, Beck-Nielsen H, et al. Mitochondrial respiration is decreased in skeletal muscle of patients with type 2 diabetes. Diabetes. 2007;56:1592–9.
- Naeije M, Zorn H. Relation between EMG power spectrum shifts and muscle fibre action potential conduction velocity changes during local muscular fatigue in man. Eur. J. Appl. Physiol. Occup. Physiol. 1982. p. 23–33.
- Oberbach A, Bossenz Y, Lehmann S, Niebauer J, Adams V, Paschke R, et al. Altered fiber distribution and fiber-specific glycolytic and oxidative enzyme activity in skeletal muscle of patients with type 2 diabetes. Diabetes Care. 2006;29:895–900.
- Park SW, Goodpaster BH, Lee JS, Kuller LH, Boudreau R, de Rekeneire N, et al. Excessive Loss of Skeletal Muscle Mass in Older Adults With Type 2 Diabetes. Diabetes Care. 2009a. p. 1993–7.
- Park SW, Goodpaster BH, Lee JS, Kuller LH, Boudreau R, de Rekeneire N, et al. Excessive loss of skeletal muscle mass in older adults with type 2 diabetes. Diabetes Care. 2009b;32:1993–7.
- Park SW, Goodpaster BH, Strotmeyer ES, Kuller LH, Broudeau R, Kammerer C, et al. Accelerated loss of skeletal muscle strength in older adults with type 2 diabetes: the health, aging, and body composition study. Diabetes Care. 2007;30:1507–12.
- Park SW, Goodpaster BH, Strotmeyer ES, de Rekeneire N, Harris TB, Schwartz A V, et al. Decreased muscle strength and quality in older adults with type 2 diabetes: the health, aging, and body composition study . Diabetes. 2006a;55:1813–8.
- Park SW, Goodpaster BH, Strotmeyer ES, De Rekeneire N, Harris TB, Schwartz A V., et al. Decreased muscle strength and quality in older adults with type 2 diabetes: The health, aging, and body composition study. Diabetes. 2006b;55:1813–8.
- Paulus SF, Grossie J. Skeletal muscle in alloxan diabetes. A comparison of isometric contractions in fast and slow muscle. Diabetes. 1983;32:1035–9.

- Sacchetti M, Olsen DB, Saltin B, Van Hall G. Heterogeneity in limb fatty acid kinetics in type 2 diabetes. *Diabetologia*. 2005;48:938–45.
- Sacchetti MS, Balducci S, Bazzucchi I, Carlucci F, Di Palumbo AS, Haxhi J, et al. Neuromuscular dysfunction in diabetes: Role of nerve impairment and training status. *Med Sci Sports Exerc*. 2013;45:52–9.
- Saltin B, Henriksson J, Nygaard E, Andersen P, Jansson E. Fiber types and metabolic potentials of skeletal muscles in sedentary man and endurance runners. *Ann N Y Acad Sci*. 1977;301:3–29.
- Sanchez OA, Snow LM, Lowe DA, Serfass RC, Thompson L V. Effects of endurance exercise-training on single-fiber contractile properties of insulin-treated streptozotocin-induced diabetic rats. *J Appl Physiol*. 2005;99:472–8.
- Sbriccoli P, Bazzucchi I, Rosponi A, Bernardi M, De Vito G, Felici F. Amplitude and spectral characteristics of biceps Brachii sEMG depend upon speed of isometric force generation. *J Electromyogr Kinesiol*. 2003;13(2):139–47.
- Segerström AB, Elgzyri T, Eriksson K-F, Groop L, Thorsson O, Wollmer P. Exercise capacity in relation to body fat distribution and muscle fibre distribution in elderly male subjects with impaired glucose tolerance, type 2 diabetes and matched controls. *Diabetes Res Clin Pract*. 2011;1–7.
- Shah S, Sonawane P, Nahar P, Buge K, Vaidya S. Are we ignoring diabetic disability: a cross sectional study of diabetic myopathy. *Indian J Med Sci*. 2011;65:186–92.
- Stuart CA, McCurry MP, Marino A, South MA, Howell MEA, Layne AS, et al. Slow-twitch fiber proportion in skeletal muscle correlates with insulin responsiveness. *J Clin Endocrinol Metab*. 2013;98:2027–36.
- Theou O, Edwards D, Jones GR, Jakobi JM. Age-related increase in electromyography burst activity in males and females. *J Aging Res*. 2013;2013.
- Volpato S, Bianchi L, Lauretani F, Lauretani F, Bandinelli S, Guralnik JM, et al. Role of muscle mass and muscle quality in the association between diabetes and gait speed. *Diabetes Care*. 2012;35:1672–9.
- Zierath JR, He L, Guma A, Odegaard WE, Klip A, Wallberg-Henriksson H. Insulin action on glucose transport and plasma membrane GLUT4 content in skeletal muscle from patients with NIDDM. *Diabetologia*. 1996;39:1180–9.

## CAPTIONS

**Fig.1.** EF (panel A and B) and KE (panel C and D) torque-velocity curves in H and T2D groups during PRE (closed circles) and POST (open circles) training. Data are expressed as percentage of the MVIC values recorded during the PRE trial. Exponential regression lines were fitted. \* $p < 0.05$  PRE vs POST.

**Fig. 2.** MFCV values recorded on biceps brachii (panel A and B) and vastus lateralis (panel C and D) muscles in H and T2D groups during PRE (closed circles) and POST (open circles) training. \* $p < 0.05$  PRE vs POST.

**Fig. 3.** TTF values recorded during the isometric fatiguing task for elbow flexion (panel A) and knee extension (panel B) in H and T2D group before (black bars) and after (white bars) the 16-week training program. # $p < 0.05$  T2D vs H.

**Fig. 4.** MFCV slopes values recorded during the isometric fatiguing task for biceps brachii (panel A) and vastus lateralis (panel B) muscles in H and T2D group before (black bars) and after (white bars) the 16-week training program. \* $p < 0.05$  PRE vs POST.

**Table 1.** Anthropometric and metabolic variables of diabetic and control subjects before (PRE) and after (POST) the 16-week training program. Values are reported as mean  $\pm$  SD \*different from POST  $p < 0.05$ .



Ilenia Bazzucchi received her PhD in Sport Science and Health and she is currently an Assistant Professor in Human Physiology at the Department of Movement, Human and Health Sciences at University of Rome "Foro Italico", Italy. She has recently been part of the Organizing Committee of the XX Congress of ISEK, International Society of Electrophysiology and Kinesiology. Her research work mainly concerns the study of neuromuscular control by means of surface electromyography, non-invasive assessment of muscle damage, neuromuscular effects of exercise and nutritional interventions in healthy individuals, athletes and patients.



Professor Giuseppe De Vito, MD and Specialist in Sport Medicine earned a PhD in Human and Exercise Physiology at the University of Rome "Sapienza", Italy. He is currently the Head of School of Public Health, Physiotherapy and Population Sciences in University College of Dublin (UK) and he previously held academic positions at the University of Strathclyde in Glasgow (UK) and at the University of Rome "Foro Italico", Italy. He is a member of both British and Italian Physiological Societies and Associate Editor of the Journal of Aging and Physical Activity and Member of the Editorial Board of the Journal of Electromyography and Kinesiology. His main research topic concerns: neuro-mechanical adaptations of muscle, tendons and joints

to aging and training; effect of muscle temperature on power production; neuromuscular control studied by surface EMG; assessment of essential tremor; study of sarcopenia in older and diabetic individuals.



Francesco Felici, MD and Specialist in Sport Medicine, is an Associate Professor of Human Physiology and Exercise Physiology at the University of Rome "Foro Italico" Italy. He has chaired the XX Congress of ISEK and is a member of the Italian Physiological Society of the Council of the International Society of Electrophysiology and Kinesiology, of the European College of Sport Sciences. He is an Associate Editor (Section Sport and Exercise) of the Journal of Electromyography. He has authored more than 100 papers and scientific communications and his research activity focuses mostly in the human movement area – from energetic to control - in collaboration with

national and international institutions. The main topics are: neuromuscular control, non linear analysis of surface electromyograms, neuromuscular effects of exercise and sport, exercise physiology in healthy and pathological subjects.



Susan Dewhurst has an undergraduate degree in Sport and Exercise Science and a PhD in Exercise Physiology both from the University of Strathclyde in Glasgow. Prior to starting her lecturing career, she spent time as a researching fellow at Aalborg University in Denmark and the University of Rome "Foro Italico", Italy. Her research interest are: effects of various physical activity interventions on muscle performance in older individuals, postural stability and gait analysis in older individuals, neuromuscular control in different populations, effects of temperature on neuromuscular control.



Antonio Sgadari earned his MD degree and specialization in Geriatrics and Gerontology at the Catholic University of Sacred Heart, Rome. He is currently Aggregate Professor at the Department of Geriatrics, Neurosciences and Orthopedics and Chief of the Physical Activity and Functional Rehabilitation Unit of the same University. His research interests are mainly focused on the effects of physical exercise in the elderly and in special populations (patients with diabetes, obesity, cardiovascular diseases, movement disorders). He is also interested in the validation and adaptation to the elderly of cardiorespiratory tests as well as in the development of specific training programs for special populations.



Massimo Sacchetti earned an MSc and a PhD in Physiology at the University of Copenhagen. He is currently Associate Professor at the Department of Movement, Human and Health Sciences at University of Rome "Foro Italico". At present, his research interests include the neuromuscular and metabolic responses to specific exercise sessions and training protocols for health and fitness in young and older individuals as well as in diabetic patients. He is also interested in studying the physiological aspects related to training and performance in endurance sports, with a special focus on cycling.

ACCEPTED MANUSCRIPT



ACCEPTED MANUSCRIPT







Figure 2


[Click here to download high resolution image](#)



Figure3

[Click here to download high resolution image](#)







Figure4

[Click here to download high resolution image](#)



	Diabetic subjects		Control subjects	
	PRE	POST	PRE	POST
Age (years)	61.5±2.8		63.6±3.6	
BMI (kg·m <sup>-2</sup> )	36.0±2.9	29.8±3.7*	27.4±2.6	26.3±3.1
BB Skinfold Thickness (mm)	9.2±2.7	8.6±3.4	8.2±4.1	7.8±3.2
VL Skinfold Thickness (mm)	28.6±10.5	26.5±9.8	25.7±9.3	25.1±8.7
HbA1C (%)	6.9±0.6	6.3±0.7	5.7±0.1	5.6±0.2
Total Cholesterol (mg·dl)	194.0±19.3	178.4±18.0*	189.6±21.7	181.1±1.0
HDL (mg·dl)	44.8±7.5	49.2±11.2	49.2±6.6	52.8±6.9
LDL (mg·dl)	180.0±23.9	139.0±22.8*	163.0±21.6	149.2±15.2
Triglycerides (mg·dl)	153.8±29.0	118.6±32.9*	113.0±10.1	104.3±11.0