

Regioselective Lithiation of Chiral 3-Acylamino-2-alkylquinazolin-4(3*H*)-ones: Application in Synthesis

Keith Smith,^{a,*} Gamal A. El-Hiti,^{a†} Mohamed F. Abdel-Megeed^b

^a Centre for Clean Chemistry, Department of Chemistry, University of Wales Swansea, Singleton Park, Swansea SA2 8PP, UK

^b Department of Chemistry, Faculty of Science, Tanta University, Tanta 31527, Egypt

Fax: +44(1792)295261

E-mail: k.smith@swansea.ac.uk

Received:

Abstract: Reaction of 3-amino-2-alkylquinazolin-4(3*H*)-ones with several chiral acid chlorides was found to be dependent on the molar proportions. When a 1:1 molar mixture was heated under reflux, the corresponding 3-(diacylamino)- derivatives were obtained in poor yields. However, when a 2:1 molar mixture was reacted in refluxing toluene, the 3-acylamino- derivatives were obtained in good yields based on the acid chloride. Lithiation of the 3-acylamino-2-alkylquinazolin-4(3*H*)-ones was achieved by the use of LDA in anhydrous THF at -78 °C and the reaction was regioselective at the carbon α to position 2 of the quinazolin-4(3*H*)-one moiety. The dilithio reagents thus obtained reacted with electrophiles to give the corresponding 2-substituted derivatives in very good yields. The NMR spectra of the products show the expected diastereotopism for all the CH₂ groups and provide evidence for long-range asymmetric induction.


Key words: lithiation, quinazolin-4(3*H*)-ones, dilithio reagent, asymmetric induction, diastereotopism

Directed metallation^{2,3} has found wide use in regioselective introduction of functional substituents into aromatic and heterocyclic compounds. However, there are relatively few examples concerning directed metallation of quinazolin-4(3*H*)-ones to afford more complex substituted derivatives.⁴⁻¹³ Compounds possessing this ring system are of interest because they show a variety of biological activities.¹⁴ In continuation of our own interest in the use of lithiation reactions for organic synthesis,¹⁵ we have demonstrated the lithiation of various quinazolin-4(3*H*)-ones,⁶⁻⁸ including simple 3-acylamino derivatives.^{9,10} We now report a study of the scope of the lithiation reaction for more complex 3-acylamino-2-alkylquinazolin-4(3*H*)-ones containing chiral acyl units.

We have been able to achieve regioselective lithiation of


a range of such quinazolin-4(3*H*)-ones by the use of LDA, and the synthesis of more complex 2-substituted derivatives in good yields. The products also show evidence of long-range asymmetric induction.

The first stage of this study required the synthesis of representative chiral 3-acylamino-2-alkylquinazolin-4(3*H*)-ones. Initially, reactions of 3-amino-2-ethylquinazolin-4(3*H*)-one (**1**)¹⁶ with chiral but racemic acid chlorides in the presence of triethylamine at 0 °C or room temperature were attempted. However, **1** was recovered unchanged, suggesting that no reaction had occurred under these conditions. A series of experiments was conducted in which the reaction conditions were varied in an attempt to produce the corresponding acylamino derivatives. When a 1:1 molar mixture of **1** and an acid chloride in the presence of triethylamine was heated under reflux in anhydrous toluene for 1 h the corresponding diacylamino derivative **2** or **3** was obtained in 41% or 25% isolated yield, respectively (Scheme 1). The monoacylamino derivative (**5** or **8**) was formed in only low yield and a substantial quantity of unreacted **1** remained.

**Scheme 1**

Product **2** appeared from its NMR spectra as a mixture of two racemic diastereoisomers in unequal proportions, while product **3** appeared from its ^1H NMR spectrum as a pair of racemic diastereoisomers in approximately equal proportions. The ^1H NMR spectrum of **3** showed diastereotopism for the hydrogens of the CH_2 groups at position 2 and in the acyl units.

In a modified procedure for the attempted synthesis of a chiral 3-acylamino-2-alkylquinazolin-4(3*H*)-one a 2:1 molar mixture of 3-amino-2-ethylquinazolin-4(3*H*)-one (**1**) and 2-phenoxypropanoyl chloride was heated under reflux in anhydrous toluene for 1 h in the presence of triethylamine. Work-up gave the desired product **5** in 78% yield. Therefore, these conditions were applied in the synthesis of a range of products (**5-10**, Scheme 2) derived from both **1** and 3-amino-2-propylquinazolin-4(3*H*)-one (**4**)¹⁷ with a variety of acid chlorides. This procedure afforded products **5-10** in very good yields based on the acid chloride (Table 1).

**Scheme 2****Table 1** Synthesis of products **5-10** According to Scheme 2


Compound	R	R^1	R^2	Yield (%) ^a
5	Me	Me	OPh	78
6	Me	Me	1-Naphthyl	74
7	Me	Et	OPh	80
8	Me	Et	Ph	79
9	Et	Me	OPh	90
10	Et	Et	OPh	93

^a Yields of isolated, purified products based on acid chloride added.

Products **5-10** appear from their NMR spectra as mixtures of two diastereoisomers in unequal proportions, due to restricted rotation around the N-N axis.¹⁸ The spectra also showed diastereotopism for the CH_2 protons of the ethyl and propyl groups (see experimental section for details). The ^1H NMR spectra were temperature dependent and showed some evidence of equilibration at 55-60 °C in CDCl_3 . In $\text{DMSO}-d_6$, the spectra showed a single set of signals at 150 °C in all cases.

In order to introduce more complex substituents at position 2, lithiations of **5-10** were carried out. Lithiation was achieved by the use of LDA in anhydrous THF at -78 °C under nitrogen and the reaction was regioselective at the carbon α to position 2 of the quinazolin-4(3*H*)-one moiety. Addition of one equivalent of LDA presumably produced the monolithio reagents **11**, which were converted into the dilithio reagents **12** on addition of a second equivalent of LDA. Reactions of the dilithio reagents **12** with several electrophiles (benzophenone, benzaldehyde, acetophenone, 2-butanone) afforded the corresponding 2-substituted quinazolin-4(3*H*)-ones **13-27** (Scheme 3) in very good yields (Table 2) following column chromatography.

The NMR spectra of all products containing a CH₂ group either in the side chain at position 2 or in the acylamino group (compounds **17-27**) showed the expected diastereotopism for the CH₂ protons. In addition, the spectra (of the total product obtained following chromatography to remove coloured impurities) revealed the presence of diastereoisomers.



Scheme 3

Table 2 Synthesis of Products **13-27** According to Scheme 3

Compound	R	R ¹	R ²	E	Yield (%) ^a
13	Me	Me	OPh	Ph ₂ C(OH)	80
14	Me	Me	OPh	PhCH(OH)	82
15	Me	Me	OPh	PhC(OH)Me	78
16	Me	Me	1-naphthyl	Ph ₂ C(OH)	79
17	Me	Et	OPh	Ph ₂ C(OH)	76
18	Me	Et	OPh	PhCH(OH)	80
19	Me	Et	OPh	PhC(OH)Me	79
20	Me	Et	Ph	Ph ₂ C(OH)	77
21	Me	Et	Ph	EtC(OH)Me	75
22	Et	Me	OPh	Ph ₂ C(OH)	88
23	Et	Me	OPh	PhCH(OH)	76
24	Et	Me	OPh	PhC(OH)Me	80
25	Et	Et	OPh	Ph ₂ C(OH)	90
26	Et	Et	OPh	PhCH(OH)	80
27	Et	Et	OPh	PhC(OH)Me	78

^a Yields of isolated, purified (column chromatography) products.

The number of diastereoisomers of the product expected when benzophenone is used as the electrophile is four. Indeed, it was found that the NMR spectra of products **13**, **22** and **25** showed a mixture of the expected four diastereoisomers in unequal proportions. However, the NMR spectra of products **17** and **20** showed the predominant presence of two substantial diastereoisomers in unequal proportions, while in the case of product **16**, there were two major diastereoisomers in approximately equal proportions.

Product **13** was separated into two fractions by crystallisation. The first fraction (**13a**, 60%) showed the presence of a single diastereoisomer. The x-ray crystallography of this compound (Figure 1) indicated the presence of one THF molecule for each molecule of **13a**. The THF appeared to be hydrogen-bonded to the NH proton of the acylamino group. The crystal structure also showed that this isomer was 2-((1*S**)-2,2-diphenyl-2-hydroxy-1-methyl)-3-((2*R**)-2-phenoxypropionylamino)quinazolin-4(3*H*)-one with (R_a*) configuration about the N-N axis. The second

fraction (40%) showed the presence of all four diastereoisomers, with **13a** being a minor component.




Figure 1 X-ray crystal structure of **13a** with numbered crystallographic atoms

Variable temperature ^1H NMR of **13a** up to 60 °C in CDCl_3 showed only minimal line broadening. However, a sample of **13a** left at room temperature for many months, showed a ^1H NMR spectrum corresponding to a mixture of **13a** and one other diastereoisomer, presumably the one involving the alternative stereochemistry ($S_{\text{a}*}$) about the N-N axis. This enabled identification of the two diastereoisomers having the same configuration at the new asymmetric carbon atom. The ^1H NMR spectrum of the second fraction (mixture) showed that these two components accounted for about 18% of the fraction, i.e. about 7% of the total product, bringing the total amount of the isomers having that stereochemistry at the new asymmetric carbon atom to around 67%. The other two diastereoisomers, having the alternative configuration at the new asymmetric carbon atom, accounted for the remaining 33% or so. Therefore, the new asymmet-

ric centre is formed with a 2:1 stereoselectivity. Although this selectivity is relatively small, it is nevertheless noteworthy, since it represents a long-range asymmetric inductive effect (the two asymmetric carbon atoms have a 1,6-relationship).

For product **17**, the two major diastereoisomers both had the same configuration at the new asymmetric carbon atom. This was deduced by the fact that the newly created CH signals for those two isomers in the ^1H NMR spectrum began to coalesce at 60 °C. Furthermore, the chemical shift positions were virtually identical to those for **13a** and its N-N isomer. The two major diastereoisomers accounted for about 80% of the total, leading to the conclusion that the long-range asymmetric induction produced a ratio of configurations at the new centre of *ca.* 4:1.

For products **16** and **20**, the NMR signals were more difficult to resolve and it was therefore much more difficult to identify the individual diastereoisomers with any confidence, although it was clear, particularly from the ^{13}C NMR spectra, that two diastereoisomers were predominant. At 60 °C the ^1H NMR spectra showed some line broadening, but it was still not clear whether the two major isomers had different configurations at the new asymmetric carbon atoms or whether they differed about the N-N axis.

When an unsymmetrical carbonyl compound (benzaldehyde, acetophenone or 2-butanone) is used as electrophile, the number of diastereoisomers expected to be

formed is eight. The ambient temperature NMR spectra of **21** (2-butanone used as the electrophile) showed the presence of at least seven racemic diastereoisomers in unequal proportions. However, the ambient temperature NMR spectra of products **14**, **15**, **18**, **19** and **24** showed mixtures of no more than four substantial diastereoisomers in unequal proportions. Furthermore, for this group of products, the ¹H NMR spectra recorded at 100 °C showed the presence of only two major diastereoisomers in unequal proportions, which indicates equilibration *via* rotation about the N-N axis. In the case of products **23**, **26** and **27** the ambient temperature NMR spectra showed mixtures of only two significant diastereoisomers in unequal proportions, which indicates a considerable long-range asymmetric induction. Unfortunately, an attempt to bring about equilibration in these cases by recording the ¹H NMR at 150 °C resulted in some decomposition to give bezaldehyde or acetophenone.

In conclusion, lithiation of chiral 3-acylamino-2-alkylquinazolin-4(3H)-ones followed by reactions with carbonyl compounds is useful for the elaboration of more complex 2-substituted 3-acylaminoquinazolin-4(3H)-ones, and in some cases gives considerable long-range asymmetric induction at the newly created asymmetric centre(s). This opens up possibilities for novel synthetic approaches to certain types of chiral compounds, which we intend to investigate.

Melting point determinations were performed by the open capillary method using a Gallenkamp melting point apparatus and are reported uncorrected. The laboratories of the University of Wales Cardiff carried out micro-

analyses. ¹H and ¹³C NMR spectra were recorded on a Bruker AC400 spectrometer operating at 400 MHz for ¹H and 100 MHz for ¹³C measurements. Chemical shifts are reported relative to tetramethylsilane. Assignments of signals are based on coupling patterns and expected chemical shift values and have not been rigorously confirmed. Signals with similar characteristics might be interchanged. Low-resolution mass spectra were recorded on a VG 12-253 spectrometer, electron impact (EI) at 70 eV and chemical ionization (CI) by the use of ammonia as ionization gas. FAB mass spectra were recorded on a VG-Autospec instrument. Accurate mass data were obtained on a VG ZAB-E instrument. Column chromatography was carried out using Merck Kieselgel 60 (230-400 mesh). Lithium diisopropylamide (LDA) and other chemicals were obtained from Aldrich Chemical Company and used without further purification. THF was distilled from sodium benzophenone ketyl.^{19,20} Other solvents were purified by standard procedures.

3-(Diacylamino)-2-ethylquinazolin-4(3H)-ones (2) and (3); General Procedure

To a stirred mixture of **1** (1.9 g, 10 mmol) and Et₃N (5 mL) in anhydrous toluene (20 mL), was added a solution of the appropriate acid chloride (11 mmol) in anhydrous toluene (5 mL). The mixture was heated under reflux for 30 min, allowed to cool, washed with sat. aq. NaHCO₃ (2 x 10 mL) and H₂O (2 x 15 mL), dried (MgSO₄), and evaporated under reduced pressure. The residue obtained was purified by column chromatography on silica gel (Et₂O-hexane, 1:4) to give **2** or **3** as a white powder.

2-Ethyl-3-[di(2-phenoxypropionyl)amino]quinazolin-4(3H)-one (2)

mp 104-107 °C; pair of racemic diastereoisomers; **2a/2b** = 3:5 (¹H NMR).

FAB-MS: *m/z* (%) = 508 (M⁺ + Na, 20), 486 (MH⁺, 67), 392 (10), 338 (31), 216 (33), 121 (100).

HRMS: *m/z* calcd for C₂₈H₂₈N₃O₅ (MH⁺): 486.2029; found: 486.2017.

Compound 2a

¹H NMR (CDCl₃): δ = 8.17 (dd, *J* = 8, 1 Hz, 1 H, H5), 7.69 (app. dt, *J* = 8, 1 Hz, 1 H, H7), 7.48 (d, *J* = 8 Hz, 1 H, H8), 7.41 (app. dt, *J* = 8, 1 Hz, 1 H, H6), 7.26-6.74 (m, 10 H, 2 OPh), 5.43 (q, *J* = 6.5 Hz, 2 H, 2 CH), 2.27 (q, *J* = 7 Hz, 2 H, CH₂), 1.52 (d, *J* = 6.5 Hz, 6 H, 2 CH₃CH), 1.09 (t, *J* = 7 Hz, 3 H, CH₃CH₂).

¹³C NMR (CDCl₃): δ = 172.6 (s, C=O), 160.2 (s, C4), 157.1 (s, C2), 156.8 (s, C1 of 2 OPh), 146.6 (s, C8a), 135.2 (d, C7), 129.7 (d, C3 of OPh), 127.6 (d, C5), 127.0 (d, C6), 126.8 (d, C8), 122.4 (d, C4 of OPh), 120.3 (s, C4a), 115.3 (d, C2 of OPh), 72.9 (d, CH), 25.5 (t, CH₂), 17.8 (q, CH₃CH), 9.7 (q, CH₃CH₂).

Compound 2b

¹H NMR (CDCl₃): δ = 8.17 (dd, *J* = 8, 1 Hz, 1 H, H5), 6.69 (app. dt, *J* = 8, 1 Hz, 1 H, H7), 7.53 (d, *J* = 8 Hz, 1 H, H8), 7.41 (app. dt, *J* = 8, 1 Hz, 1 H, H6), 7.26-6.60 (m, 10 H, 2 OPh), 5.83 (q, *J* = 6.5 Hz, 2 H, 2 CH), 2.31 (q, *J* = 7 Hz, 2 H, CH₂), 1.64 (d, *J* = 6.5 Hz, 6 H, 2 CH₃CH), 1.18 (t, *J* = 7 Hz, 3 H, CH₃CH₂).

¹³C NMR (CDCl₃): δ = 171.8 (s, C=O), 160.2 (s, C4), 157.2 (s, C2), 155.7 (s, C1 of OPh), 146.6 (s, C8a), 135.3 (d, C7), 129.6 (d, C3 of OPh), 127.7 (d, C5), 127.1 (d, C6), 126.8 (d, C8), 122.0 (d, C4 of OPh), 120.3 (s, C4a), 115.0 (d, C2 of OPh), 73.4 (d, CH), 25.6 (t, CH₂), 17.8 (q, CH₃CH), 9.8 (q, CH₃CH₂).

2-Ethyl-3-[di(2-phenylbutyryl)amino]quinazolin-4(3*H*)-one (3)

mp 155-156 °C; pair of racemic diastereoisomers (NMR).

FAB-MS: *m/z* (%) = 504 (M⁺ + Na, 23), 482 (MH⁺, 69), 336 (86), 216 (36), 190 (36), 119 (100).

HRMS: *m/z* calcd for C₃₀H₃₂N₃O₃ (MH⁺): 482.2444; found: 482.2459.

¹H NMR (CDCl₃): δ = 8.30 (dd, *J* = 8, 1 Hz, 1 H, H5), 7.80 (app. dt, *J* = 8, 1 Hz, 1 H, H7), 7.68 (d, *J* = 8 Hz, 1 H, H8), 7.50 (app. dt, *J* = 8, 1 Hz, 1 H, H6), 7.33 (m, 4 H, H3 of 2 Ph), 7.12 (m, 4 H, H2 of 2 Ph), 6.70 (d, *J* = 7.5 Hz, 2 H, H4 of 2 Ph), 4.98, 3.23 (2 t, *J* = 7 Hz, 2 H, 2 CH), 2.24, 2.14 (2 dq, *J* = 14, 7 Hz, 2 H, CH₂CH₃), 1.85-1.68 (m, 2 H, CH₂CH), 1.30-0.93 (m, 2 H, CH₂CH), 0.93, 0.80 (2 t, *J* = 7 Hz, 6 H, 2 CH₃CH₂CH), 0.57 (t, *J* = 7 Hz, 3 H, CH₃CH₂).

¹³C NMR (CDCl₃): δ = 174.2, 173.9 (2 s, C=O), 160.4 (s, C4), 158.2 (s, C2), 147.0 (s, C8a), 137.8, 136.6 (2 s, C1 of Ph), 135.3 (d, C7), 129.0, 128.9 (2 d, C3 of Ph), 128.5, 128.4 (2 d, C4 of Ph), 127.8, 127.7 (2 d, C2 of Ph), 127.5 (d, C5), 127.4 (d, C6), 126.8 (d, C8), 120.5 (s, C4a), 54.0, 53.5 (2 d, CH), 27.7, 27.0 (2 t, CH₂CH), 24.3 (t, CH₃CH₂), 11.9, 11.7 (2 q, CH₃CH₂CH), 9.5 (q, CH₃CH₂).

3-Acylamino-2-alkylquinazolin-4(3*H*)-ones (5-10); General Procedure

To a stirred mixture of **1** or **4** (10 mmol) and Et₃N (3 mL) in anhydrous toluene (20 mL), was added a solution of the appropriate acid chloride (5 mmol) in anhydrous toluene (5 mL). The mixture was heated under reflux for 30 min, allowed to cool, washed with sat. aq. NaHCO₃ (2 x 10 mL) and H₂O (2 x 15 mL), dried (MgSO₄), and evaporated under reduced pressure. The residue obtained was purified by column chromatography on silica gel (Et₂O-hexane, 1:4) to give the corresponding 3-acylamino derivatives **5-10**. The yields obtained are recorded in Table 1.

2-Ethyl-3-(2-phenoxypropionylamino)quinazolin-4(3*H*)-one (5)

mp 90-92 °C; **5a/5b** = 1:2 (¹H NMR).

Anal. Calcd for C₁₉H₁₉N₃O₃: C, 67.65; H, 5.64; N, 12.46. Found: C, 67.6; H, 5.6; N, 12.6.

EI-MS: *m/z* (%) = 337 (M⁺, 22), 244 (13), 216 (100), 173 (32), 130 (34), 121 (81), 77 (99).

HRMS: *m/z* calcd for C₁₉H₁₉N₃O₃ (M⁺): 337.1426; found: 337.1426.

Compound 5a

¹H NMR (CDCl₃): δ = 8.73 (s, exch., 1 H, NH), 8.16 (dd, *J* = 8, 1 Hz, 1 H, H5), 7.72 (m, 1 H, H7), 7.64 (d, *J* = 8 Hz, 1 H, H8), 7.41 (app. dt, *J* = 8, 1 Hz, 1 H, H6), 7.35 (t, *J* = 7.5 Hz, 2 H, H3 of OPh), 7.07 (t, *J* = 7.5 Hz, 2 H, H2 of OPh), 6.99 (d, *J* = 7.5 Hz, 1 H, H4 of OPh), 4.97 (q, *J* = 7 Hz, 1 H, CH), 2.44 (2 dq, *J* = 15, 7.5 Hz, 2 H, CH₂), 1.81 (d, *J* = 7 Hz, 3 H, CH₃CH), 1.06 (t, *J* = 7.5 Hz, 3 H, CH₃CH₂).

¹³C NMR (CDCl₃): δ = 172.8 (s, C=O), 159.9 (s, C4), 158.5 (s, C2), 156.8 (s, C1 of OPh), 147.0 (s, C8a), 134.8 (d, C7), 129.9 (d, C3 of OPh), 127.5 (d, C5), 126.9 (d, C6), 126.7 (d, C8), 122.5 (d, C4 of OPh), 120.6 (s, C4a), 115.2 (d, C2 of OPh), 74.7 (d, CH), 27.0 (t, CH₂), 18.8 (q, CH₃CH), 10.5 (q, CH₃CH₂).

Compound 5b

¹H NMR (CDCl₃): δ = 8.93 (s, exch., 1 H, NH), 8.12 (d, *J* = 8 Hz, 1 H, H5), 7.72 (m, 1 H, H7), 7.64 (d, *J* = 8 Hz, 1 H, H8), 7.41 (app. dt, *J* = 8, 1 Hz, 1 H, H6), 7.35 (t, *J* = 7.5 Hz, 2 H, H3 of OPh), 7.07 (t, *J* = 7.5 Hz, 2 H, H2 of OPh), 6.99 (d, *J* = 7.5 Hz, 1 H, H4 of OPh), 4.95 (q, *J* = 7 Hz, 1 H, CH), 2.81 (2 dq, *J* = 15, 7.5 Hz, 2 H, CH₂), 1.68 (d, *J* = 7 Hz, 3 H, CH₃CH), 1.34 (t, *J* = 7.5 Hz, 3 H, CH₃CH₂).

¹³C NMR (CDCl₃): δ = 172.6 (s, C=O), 159.9 (s, C4), 158.4 (s, C2), 156.8 (s, C1 of OPh), 147.0 (s, C8a), 134.8 (d, C7), 129.9 (d, C3 of OPh), 127.5 (d, C5), 126.9 (d, C6), 126.7 (d, C8), 122.9 (d, C4 of OPh), 120.7 (s, C4a), 116.5 (d, C2 of OPh), 75.8 (d, CH), 26.8 (t, CH₂), 18.7 (q, CH₃CH), 10.6 (q, CH₃CH₂).

2-Ethyl-3-[2-(1-naphthalene)propionylamino]quinazolin-4(3*H*)-one (6)

mp 180-182 °C; **6a/6b** = 5:6 (¹³C NMR).

EI-MS: *m/z* (%) = 371 (M⁺, 5), 216 (40), 182 (38), 155 (100), 153 (34), 130 (15), 115 (9), 90 (8), 76 (10).

CI-MS: *m/z* (%) = 372 (MH⁺, 7), 217 (9), 175 (100), 159 (7), 91 (6).

HRMS: *m/z* calcd for C₂₃H₂₁N₃O₂ (MH⁺): 371.1634; found: 371.1634.

¹H NMR (CDCl₃): δ = 8.20-7.37 (m, 12 H, ArH and NH), 4.71-4.62 (m, 1 H, CH), 2.54-2.28 (m, 2 H, CH₂),

1.89, 1.74 (2 d, $J = 7$ Hz, 3 H, CH_3CH), 0.99 (t, $J = 7.5$ Hz, 3 H, CH_3CH_2).

Compound 6a

^{13}C NMR (CDCl_3): $\delta = 175.0$ (s, C=O), 160.2 (s, C4), 158.8 (s, C2), 146.9 (s, C8a), 136.0 (s, C1'), 134.8 (d, C7), 133.8 (s, C4'a), 131.1 (s, C8'a), 129.1 (d, C5'), 128.7 (d, C5), 127.8 (d, C6), 126.8 (d, C8), 126.7 (d, C3'), 126.5 (d, C2'), 126.1 (d, C4'), 125.8 (d, C7'), 124.9 (d, C6'), 123.2 (d, C8'), 120.6 (s, C4a), 42.0 (d, CH), 26.8 (t, CH₂), 17.7 (q, CH_3CH), 10.5 (q, CH_3CH_2).

Compound 6b

^{13}C NMR (CDCl_3): $\delta = 174.9$ (s, C=O), 160.0 (s, C4), 158.7 (s, C2), 146.9 (s, C8a), 136.0 (s, C1'), 134.7 (d, C7), 134.2 (s, C4'a), 131.1 (s, C8'a), 129.2 (d, C5'), 128.5 (d, C5), 127.3 (d, C6), 126.8 (d, C8), 126.7 (d, C3'), 126.5 (d, C2'), 126.2 (d, C4'), 125.9 (d, C7'), 125.6 (d, C6'), 122.8 (d, C8'), 120.5 (s, C4a), 42.0 (d, CH), 26.9 (t, CH₂), 17.6 (q, CH_3CH), 10.5 (q, CH_3CH_2).

2-Ethyl-3-(2-phenoxybutyrylamino)quinazolin-4(3*H*)-one (7)

mp 93–94 °C; **7a/7b** = 1:2 (^1H NMR).

Anal. Calcd for $\text{C}_{20}\text{H}_{21}\text{N}_3\text{O}_3$: C, 68.36; H, 6.02; N, 11.96. Found: C, 68.4; H, 6.2; N, 11.8.

EI-MS: m/z (%) = 351 (M⁺, 10), 258 (7), 230 (13), 216 (60), 173 (40), 135 (100), 107 (65), 94 (31), 77 (75), 41 (43).

CI-MS: m/z (%) = 352 (MH⁺, 10), 197 (25), 176 (18), 175 (100), 136 (5), 105 (7), 58 (5).

HRMS: m/z calcd for $\text{C}_{20}\text{H}_{21}\text{N}_3\text{O}_3$ (M⁺): 351.1583; found: 351.1583.

Compound 7a

^1H NMR (CDCl_3): $\delta = 8.69$ (s, exch., 1 H, NH), 8.16 (d, $J = 8$ Hz, 1 H, H5), 7.73 (app. dt, $J = 8$, 1 Hz, 1 H, H7), 7.60 (d, $J = 8$ Hz, 1 H, H8), 7.40 (t, $J = 8$ Hz, 1 H, H6), 7.36 (t, $J = 7.5$ Hz, 2 H, H3 of OPh), 7.18–6.99 (m, 3 H, H2 and H4 of OPh), 4.78 (t, $J = 7$ Hz, 1 H, CH), 2.41 (dq, $J = 14$, 7 Hz, 2 H, CH_2CH), 2.18 (dq, $J = 15$, 7.5 Hz, 2 H, CH₂), 1.20 (t, $J = 7$ Hz, 3 H, $\text{CH}_3\text{CH}_2\text{CH}$), 1.02 (t, $J = 7.5$ Hz, 3 H, CH_3CH_2).

^{13}C NMR (CDCl_3): $\delta = 172.1$ (s, C=O), 159.9 (s, C4), 158.5 (s, C2), 157.2 (s, C1 of OPh), 146.9 (s, C8a), 134.8 (d, C7), 129.9 (d, C3 of OPh), 127.0 (d, C5), 126.6 (d, C6), 126.5 (d, C8), 122.4 (d, C4 of OPh), 120.5 (s, C4a), 115.1 (d, C2 of OPh), 79.5 (d, CH), 26.9 (t, CH₂CH), 26.5 (t, CH₂), 10.5 (q, $\text{CH}_3\text{CH}_2\text{CH}$), 9.5 (q, CH_3CH_2).

Compound 7b

^1H NMR (CDCl_3): $\delta = 8.88$ (s, exch., 1 H, NH), 8.16 (d, $J = 8$ Hz, 1 H, H5), 7.43 (app. dt, $J = 8$, 1 Hz, 1 H, H7), 7.60 (d, $J = 8$ Hz, 1 H, H8), 7.40 (t, $J = 8$ Hz, 1 H, H6), 7.36 (t, $J = 7.5$ Hz, 2 H, H3 of OPh), 7.18–6.99 (m, 3 H,

H2 and H4 of OPh), 4.78 (t, $J = 7.4$ Hz, 1 H, CH), 2.87 (dq, $J = 14$, 7 Hz, 2 H, CH_2CH), 2.08 (dq, $J = 15$, 7.5 Hz, 2 H, CH₂), 1.33 (t, 3 H, $J = 7$ Hz, $\text{CH}_3\text{CH}_2\text{CH}$), 1.11 (t, $J = 7.5$ Hz, 3 H, CH_3CH_2).

^{13}C NMR (CDCl_3): $\delta = 172.1$ (s, C=O), 159.8 (s, C4), 158.3 (s, C2), 157.5 (s, C1 of OPh), 146.9 (s, C8a), 134.8 (d, C7), 127.4 (d, C3 of OPh), 127.0 (d, C5), 126.8 (d, C6), 126.7 (d, C8), 122.9 (d, C4 of OPh), 120.6 (s, C4a), 116.7 (d, C2 of OPh), 80.9 (d, CH), 26.9 (t, CH₂CH), 26.1 (t, CH₂), 10.3 (q, $\text{CH}_3\text{CH}_2\text{CH}$), 9.2 (q, CH_3CH_2).

2-Ethyl-3-(2-phenylbutyrylamino)quinazolin-4(3*H*)-one (8)

mp 180–183 °C; **8a/8b** = 1:3 (^1H NMR).

EI-MS: m/z (%) = 335 (M⁺, 5), 216 (40), 189 (17), 173 (12), 119 (65), 91 (100).

HRMS: m/z calcd for $\text{C}_{20}\text{H}_{21}\text{N}_3\text{O}_2$ (M⁺): 335.1634; found: 335.1634.

Compound 8a

^1H NMR (DMSO-*d*₆): $\delta = 11.66$ (s, exch., 1 H, NH), 8.10 (d, $J = 8$ Hz, 1 H, H5), 7.99 (t, $J = 8$ Hz, 1 H, H7), 7.75 (t, $J = 8$ Hz, 1 H, H6), 7.52–7.30 (m, 6 H, H8 and Ph), 3.98 (t, $J = 7$ Hz, 1 H, CH), 3.00, 3.62 (2 dq, $J = 14$, 7 Hz, 2 H, CH_2CH), 2.22, 1.91 (2 dq, $J = 15$, 7.5 Hz, 2 H, CH₂), 1.05 (t, $J = 7$ Hz, 3 H, $\text{CH}_3\text{CH}_2\text{CH}$), 1.00 (t, $J = 7.5$ Hz, 3 H, CH_3CH_2).

^{13}C NMR (DMSO-*d*₆): $\delta = 173.7$ (s, C=O), 167.8 (s, C4), 155.9 (s, C2), 138.7 (s, C8a), 138.0 (s, C1 of Ph), 136.7 (d, C7), 129.5 (d, C5), 128.8 (d, C3 of Ph), 128.3 (d, C6), 128.0 (d, C2 of Ph), 127.6 (d, C8), 120.1 (d, C4 of Ph), 118.5 (s, C4a), 52.3 (d, CH), 26.6 (t, CH₂CH), 25.4 (t, CH₂), 12.1 (q, $\text{CH}_3\text{CH}_2\text{CH}$), 11.3 (q, CH_3CH_2).

Compound 8b

^1H NMR (DMSO-*d*₆): $\delta = 11.38$ (s, exch., 1 H, NH), 8.10 (d, $J = 8$ Hz, 1 H, H5), 7.99 (t, $J = 8$ Hz, 1 H, H7), 7.75 (t, $J = 8$ Hz, 1 H, H6), 7.52–7.30 (m, 6 H, H8 and Ph), 3.90 (t, $J = 7$ Hz, 1 H, CH), 3.34, 3.10 (2 dq, $J = 14$, 7 Hz, 2 H, CH_2CH), 2.22, 1.91 (2 dq, $J = 15$, 7.5 Hz, 2 H, CH₂), 1.44 (t, $J = 7$ Hz, 3 H, $\text{CH}_3\text{CH}_2\text{CH}$), 1.03 (t, $J = 7.5$ Hz, 3 H, CH_3CH_2).

^{13}C NMR (DMSO-*d*₆): $\delta = 173.8$ (s, C=O), 167.6 (s, C4), 155.7 (s, C2), 138.7 (s, C8a), 138.0 (s, C1 of Ph), 136.4 (d, C7), 129.6 (d, C5), 128.6 (d, C3 of Ph), 128.3 (d, C6), 128.2 (d, C2 of Ph), 127.6 (d, C8), 120.2 (d, C4 of Ph), 118.5 (s, C4a), 52.7 (d, CH), 26.60 (t, CH₂CH), 25.1 (t, CH₂), 12.2 (q, $\text{CH}_3\text{CH}_2\text{CH}$), 11.9 (q, CH_3CH_2).

2-Propyl-3-(2-phenoxypropionylamino)quinazolin-4(3*H*)-one (9)

mp 87–89 °C; **9a/9b** = 1:2 (^1H NMR).

EI-MS: m/z (%) = 352 (M⁺ + 1, 12), 337 (M⁺, 20), 323 (100), 285 (35).

CI-MS: m/z (%) = 352 (MH^+ , 16), 189 (100), 183 (32), 91 (17).

HRMS: m/z calcd for $C_{20}H_{22}N_3O_3$ (MH^+): 352.1661; found: 352.1664.

Compound 9a

1H NMR ($CDCl_3$): δ = 8.98 (s, exch., 1 H, NH), 8.17 (dd, J = 8, 1 Hz, 1 H, H5), 7.76-7.60 (m, 2 H, H7 and H8), 7.42 (app. dt, J = 8, 1 Hz, 1 H, H6), 7.37-7.33 (m, 1 H, H3 of OPh), 7.10-7.06 (m, 1 H, H4 of OPh), 7.00 (d, J = 7.5 Hz, 1 H, H2 of OPh), 4.96 (q, J = 7 Hz, 1 H, CH), 2.79-2.73 (m, 2 H, CH₂), 1.79 (d, J = 7 Hz, 3 H, CHCH₃), 1.63-1.49 (m, 2 H, CH₂), 1.05 (t, J = 7.5 Hz, 3 H, CH₃CH₂).

^{13}C NMR ($CDCl_3$): δ = 172.4 (s, C=O), 159.9 (s, C4), 157.5 (s, C1 of OPh), 156.8 (s, C2), 146.9 (s, C8a), 134.9 (d, C7), 129.9 (d, C3 of OPh), 126.9 (d, C5), 126.8 (d, C6), 126.6 (d, C8), 122.8 (d, C4 of OPh), 120.7 (s, C4a), 116.5 (d, C2 of OPh), 75.7 (d, CH), 35.6 (t, CH₂), 20.0 (t, CH₂), 18.7 (q, CH₃CH), 13.9 (q, CH₃CH₂).

Compound 9b

1H NMR ($CDCl_3$): δ = 8.75 (s, exch., 1 H, NH), 8.17 (dd, J = 8, 1 Hz, 1 H, H5), 7.76-7.60 (m, 2 H, H7 and H8), 7.42 (app. dt, J = 8, 1 Hz, 1 H, H6), 7.37-7.33 (m, 1 H, H3 of OPh), 7.10-7.06 (m, 1 H, H4 of OPh), 7.00 (d, J = 7.5 Hz, 1 H, H2 of OPh), 4.94 (q, J = 7 Hz, 1 H, CH), 2.44-2.38 (m, 2 H, CH₂), 1.83 (d, J = 7 Hz, 3 H, CH₃CH), 1.63-1.49 (m, 2 H, CH₂), 0.80 (t, J = 7.4 Hz, 3 H, CH₃CH₂).

^{13}C NMR ($CDCl_3$): δ = 172.9 (s, C=O), 159.9 (s, C4), 157.7 (s, C1 of OPh), 156.9 (s, C2), 146.9 (s, C8a), 134.9 (d, C7), 130.0 (d, C3 of OPh), 127.4 (d, C5), 126.8 (d, C6), 126.6 (d, C8), 122.5 (d, C4 of OPh), 120.5 (s, C4a), 116.5 (d, C2 of OPh), 74.6 (d, CH), 35.5 (t, CH₂), 19.9 (t, CH₂), 18.9 (q, CH₃CH), 13.6 (q, CH₃CH₂).

2-Propyl-3-(2-phenoxybutyrylamino)quinazolin-4(3*H*)-one (10)

mp 115-117 °C; **10a/10b** = 1:2 (1H NMR).

EI-MS: m/z (%) = 365 (M^+ , 4), 337 (6), 244 (15), 230 (30), 187 (20), 173 (20), 160 (60), 135 (45), 117 (34), 107 (40), 94 (43), 77 (100), 65 (35).

CI-MS: m/z (%) = 366 (MH^+ , 30), 197 (32), 189 (100), 173 (13), 136 (5), 105 (33).

HRMS: m/z calcd for $C_{21}H_{24}N_3O_3$ (MH^+): 366.1819; found: 366.1819.

Compound 10a

1H NMR ($CDCl_3$): δ = 8.39 (s, exch., 1 H, NH), 8.16 (m, 1 H, H5), 7.71 (m, 1 H, H7), 7.60 (d, J = 8 Hz, 1 H, H8), 7.44-7.00 (m, 6 H, H6 and OPh), 4.81-4.77 (m, 1 H, CH), 2.74-2.70 (m, 2 H, CH₂), 2.13-2.81 (m, 2 H, CH₂), 1.84-1.79 (m, 2 H, CH₂), 1.12 (t, J = 7 Hz, 3 H, CH₃), 1.04 (t, J = 7.5 Hz, 3 H, CH₃).

^{13}C NMR ($CDCl_3$): δ = 172.1 (s, C=O), 159.8 (s, C4), 157.5 (s, C1 of OPh), 157.5 (s, C2), 146.9 (s, C8a), 134.8 (d, C7), 129.8 (d, C3 of OPh), 127.4 (d, C5), 126.8 (d, C6), 126.6 (d, C8), 122.4 (d, C4 of OPh), 120.7 (s, C4a), 116.7 (d, C2 of OPh), 80.9 (d, CH), 35.7 (t, CH₂), 26.1 (t CH₂), 20.0 (t, CH₂), 13.8 (q, CH₃), 9.1 (q, CH₃).

Compound 10b

1H NMR ($CDCl_3$): δ = 8.73 (s, exch., 1 H, NH), 8.16 (m, 1 H, H5), 7.71 (m, 1 H, H7), 7.60 (d, J = 8 Hz, 1 H, H8), 7.44-7.00 (m, 6 H, H6 and OPh), 4.81-4.77 (m, 1 H, CH), 2.35-2.31 (m, 2 H, CH₂), 2.25-2.20 (m, 2 H, CH₂), 1.57-1.52 (m, 2 H, CH₂), 1.21 (t, J = 7 Hz, 3 H, CH₃), 0.76 (t, J = 7.5 Hz, 3 H, CH₃).

^{13}C NMR ($CDCl_3$): δ = 172.2 (s, C=O), 159.9 (s, C4), 157.7 (s, C1 of OPh), 157.3 (s, C2), 146.9 (s, C8a), 134.8 (d, C7), 129.9 (d, C3 of OPh), 127.0 (d, C5), 126.8 (d, C6), 126.6 (d, C8), 122.9 (d, C4 of OPh), 120.5 (s, C4a), 115.0 (d, C2 of OPh), 79.8 (d, CH), 35.5 (t, CH₂), 26.6 (t CH₂), 19.8 (t, CH₂), 13.6 (q, CH₃), 9.6 (q, CH₃).

2-Substituted 3-Acylaminoquinazolin-4(3*H*)-ones (13-27); General Procedure

A solution of LDA in pentane (1.6 M; 1.3 mL, 2.2 mmol) was added under N_2 in a dropwise manner to a stirred solution of the appropriate 3-acylamino-2-alkylquinazolin-4(3*H*)-ones (**5-10**, 1 mmol) in anhydrous THF (10 mL) at -78 °C. Formation of the dilithio reagent was observed as a very deep red solution. The mixture was stirred at -78 °C for 30 min, after which an electrophile (1.1 mmol) (as a solution in anhydrous THF for benzophenone, otherwise neat) was added. The mixture was stirred for 2 h at -78 °C then removed from the cooling bath and allowed to warm to r.t., diluted with Et₂O (10 mL), and quenched with sat. aq. NH₄Cl solution (10 mL). The organic layer was washed with H₂O (2 x 20 mL), dried (MgSO₄), and evaporated under reduced pressure. The products obtained were purified by column chromatography on silica gel (Et₂O-hexane, 1:4) to give the corresponding 2-substituted derivatives **13-27** as white powders. The yields obtained are recorded in Table 2.

(R_a*)-2-((1S*)-2,2-Diphenyl-2-hydroxy-1-methylethyl)-3-((2R*)-2-phenoxypropionylamino)quinazolin-4(3*H*)-one (13a)

mp 172 °C; a single diastereoisomer (NMR).

Anal. Calcd for $C_{32}H_{29}N_3O_4$.THF: C, 73.08; H, 6.30; N, 7.10. Found: C, 73.2; H, 6.2; N, 7.3.

EI-MS: m/z (%) = 355 (5), 337 (10), 216 (42), 182 (49), 173 (37), 121 (50), 105 (100), 77 (95), 51 (42).

CI-MS: m/z (%) = 520 (MH^+ , 33), 357 (100), 338 (40), 262 (20), 228 (15).

HRMS: m/z calcd for $C_{32}H_{30}N_3O_4$ (MH^+): 520.2236; found: 520.2206.

¹H NMR (CDCl₃): δ = 8.34 (s, exch., 1 H, NH), 8.09 (dd, J = 8, 1 Hz, 1 H, H5), 7.73 (app. dt, J = 8, 1 Hz, 1 H, H7), 7.61 (d, J = 8 Hz, 1 H, H8), 7.51-6.92 (m, 17 H, ArH and OH), 5.02 (q, J = 6.7 Hz, 1 H, CH), 4.00 (q, J = 7 Hz, 1 H, CH), 3.75 (t, J = 6.5 Hz, 4 H, H2 of THF), 1.86 (t, J = 6.5 Hz, 4 H, H3 of THF), 1.82 (d, J = 6.7 Hz, 3 H, CH₃), 1.30 (d, J = 7 Hz, 3 H, CH₃).

¹³C NMR (CDCl₃): δ = 173.7 (s, C=O), 161.6 (s, C4), 159.0 (s, C2), 157.2 (s, C1 of OPh), 147.6 (s, C8a), 145.3, 144.3 (2 s, C1 of 2 Ph), 135.2 (d, C7), 130.4 (d, C3 of OPh), 128.1 (d, C5), 127.2 127.0 (2 d, C3 of 2 Ph), 126.9 (d, C6), 126.6 (d, C8), 126.5, 125.5 (2 d, C2 of 2 Ph), 124.7 (d, C4 of 2 Ph), 122.7 (d, C4 of OPh), 120.2 (s, C4a), 114.4 (d, C2 of OPh), 79.2 (s, C-OH), 74.5 (d, CH), 68.0 (t, C2 of THF), 42.1 (d, CH), 25.6 (t, C3 of THF), 19.2 (q, CH₃), 15.1 (q, CH₃).

2-(2,2-Diphenyl-2-hydroxy-1-methylethyl)-3-(2-phenoxypropionylamino)quinazolin-4(3*H*)-one (13)

mp 171-173 °C; four diastereoisomers (NMR).

Anal. Calcd for C₃₂H₂₉N₃O₄: C, 73.97; H, 5.63; N, 8.09. Found: C, 74.2; H, 5.6; N, 8.0.

EI-MS: *m/z* (%) = 355 (6), 337 (12), 216 (42), 182 (30), 173 (20), 121 (55), 105 (90), 77 (100), 51 (41).

CI-MS: *m/z* (%) = 520 (MH⁺, 45), 357 (90), 338 (100).

HRMS: *m/z* calcd for C₃₂H₃₀N₃O₄ (MH⁺): 520.2236; found: 520.2198.

¹H NMR (CDCl₃): δ = 8.58, 8.55 (2 s, exch., 1 H, NH), 8.09, 7.85 (2 dd, J = 8, 1 Hz, 1 H, H5), 7.80-6.85 (m, 19 H, ArH and OH), 5.12, 5.02 (2 q, J = 7 Hz, 1 H, CH), 4.36, 4.01, 3.96, 3.73 (4 q, J = 7 Hz, 1 H, CH), 1.94, 1.86, 1.76 (3 d, J = 7 Hz, 3 H, CH₃), 1.36, 1.30, 0.75 (3 d, J = 7 Hz, 3 H, CH₃).

¹³C NMR (CDCl₃): δ = 173.6, 173.4 (2 s, C=O), 161.5 (s, C4), 159.1 (s, C2), 156.8 (s, C1 of OPh), 147.9 (s, C8a), 145.4, 145.0 (2 s, C1 of 2 Ph), 135.1 (d, C7), 130.5, 130.2 (2 d, C3 of OPh), 128.4, 128.3, 128.1, 128.0 (4 d, C3 of 2 Ph), 127.2, 127.1 (2 d, C5), 126.9, 126.8 (2 d, C6), 126.4, 126.3 (2 d, C8), 126.0, 125.5 (2 d, C2 of 2 Ph), 125.4, 125.0 (2 d, C4 of 2 Ph), 123.1, 122.5 (2 d, C4 of OPh), 120.2 (s, C4a), 116.6, 114.4 (2 d, C2 of OPh), 80.0, 79.3 (2 s, C-OH), 75.9, 74.6 (2 d, CH), 42.4, 41.9 (2 d, CH) 19.2, 18.9 (2 q, CH₃), 15.8, 14.9, 13.8 (3 q, CH₃).

2-(2-Hydroxy-1-methyl-2-phenylethyl)-3-(2-phenoxypropionylamino)quinazolin-4(3*H*)-one (14)

mp 90-93 °C; four diastereoisomers (NMR).

EI-MS: *m/z* (%) = 337 (3), 216 (11), 173 (8), 130 (15), 121 (33), 94 (29), 77 (100), 39 (41).

CI-MS: *m/z* (%) = 444 (MH⁺, 10), 338 (5), 281 (41), 183 (40), 175 (56), 122 (48), 91 (100), 74 (38).

HRMS: *m/z* calcd for C₂₆H₂₆N₃O₄ (MH⁺): 444.1923; found: 444.1926.

¹H NMR (CDCl₃): δ = 8.85, 8.75, 8.67, 8.61 (4 s, exch., 1 H, NH), 8.11-6.82 (m, 14 H, ArH), 4.90 (m, 1 H, CH), 4.79, 4.57 (2 d, J = 6 Hz, 1 H, CH), 4.03, 3.89 (2 br s, exch., 1 H, OH), 3.46, 3.15, 2.84 (3 pentet, J = 7 Hz, 1 H, CH), 1.71, 1.61 (2 d, J = 7 Hz, 3 H, CH₃), 1.22, 1.17, 0.61 (3 d, J = 7 Hz, 3 H, CH₃).

¹³C NMR (CDCl₃): δ = 172.9, 172.6, 172.1, 171.1 (4 s, C=O), 160.2, 160.1, 159.6 (3 s, C4), 159.6, 159.1 (2 s, C2), 156.8, 156.7 (2 s, C1 of OPh), 146.3, 146.2 (2 s, C8a), 142.7, 142.3, 141.7 (3 s, C1 of Ph), 135.1, 135.0, 134.8 (3 d, C7), 128.6, 128.4 (2 d, C3 of OPh), 128.2, 128.1 (2 d, C3 of Ph), 127.6, 127.2 (2 d, C5), 127.2, 127.1 (2 d, C6), 127.0, 126.8 (2 d, C2 of Ph), 126.2, 125.8 (2 d, C8), 122.7, 122.5, 122.4, 122.3 (4 d, C4 of Ph), 120.9, 120.7 (2 s, C4a), 116.3 (d, C4 of OPh), 114.9, 114.6 (2 d, C2 of OPh), 78.5, 77.8, 77.2 (3 d, CH), 76.1, 75.2, 74.6, 74.3 (4 d, CH), 43.5, 42.8, 42.6, 42.0 (4 d, CH) 19.0, 18.9, 18.4 (3 q, CH₃), 16.9, 16.5, 16.3 (3 q, CH₃).

2-(2-Hydroxy-1-methyl-2-phenylpropyl)-3-(2-phenoxypropionylamino)quinazolin-4(3*H*)-one (15)

mp 148-150 °C; four diastereoisomers (¹H NMR).

EI-MS: *m/z* (%) = 337 (6), 216 (37), 173 (20), 130 (15), 121 (38), 105 (53), 94 (29), 77 (100).

CI-MS: *m/z* (%) = 458 (MH⁺, 88), 338 (25), 295 (100).

HRMS: *m/z* calcd for C₂₇H₂₈N₃O₄ (MH⁺): 458.2080; found: 458.2073.

¹H NMR (CDCl₃): δ = 9.16, 9.11, 8.92, 8.88 (4 s, exch., 1 H, NH), 8.23-6.90 (m, 14 H, ArH), 6.51, 6.15 (2 br s, exch., 1 H, OH), 5.06, 4.96 (2 q, J = 7 Hz, 1 H, CH), 3.50, 3.30, 3.20, 3.00 (4 q, J = 7 Hz, 1 H, CH), 1.91, 1.89, 1.78 (3 d, J = 7 Hz, 3 H, CH₃) 1.64, 1.48, 1.21 (3 s, 3 H, CH₃), 1.11, 1.07, 1.02, 0.60 (4 d, J = 7 Hz, 3 H, CH₃).

¹³C NMR (CDCl₃): δ = 172.3, 172.2, 171.9 (3 s, C=O), 162.1, 161.8 (2 s, C4), 159.5, 159.4 (2 s, C2), 157.0, 156.7, 156.6 (3 s, C1 of OPh), 146.3, 146.1 (2 s, C8a), 145.5, 145.4, 145.3 (3 s, C1 of Ph), 135.3 (d, C7), 130.2, 130.1 (2 d, C3 of OPh), 129.9, 128.2, 128.1 (3 d, C3 of Ph), 127.3, 127.2, 127.1 (3 d, C2 of Ph), 127.0, 126.8 (2 d, C5), 126.5, 126.4 (2 d, C6), 124.9, 124.8, 124.7 (3 d, C8), 122.9, 122.6, 122.5 (3 d, C4 of Ph), 120.5 (s, C4a), 116.5, 116.4 (2 d, C2 of OPh), 114.7, 114.6 (2 d, C2 of OPh), 77.4, 75.8, 75.4 (3 d, CH), 74.9, 74.6 (2 s, C-OH), 44.6, 44.2 (2 d, CH), 30.0, 29.9 (2 q, CH₃), 20.7, 19.2, 19.1, 18.9 (4 q, CH₃), 15.2, 14.9, 14.7, 14.5 (4 q, CH₃).

2-(2,2-Diphenyl-2-hydroxy-1-methylethyl)-3-[2-(1-naphthalene)propionylamino]quinazolin-4(3*H*)-one (16)

mp 131-133 °C; two diastereoisomers in equal proportions (¹H NMR).

EI-MS: *m/z* (%) = 372 (38), 371 (100), 356 (15), 355 (61).

CI-MS: *m/z* (%) = 554 (MH⁺, 100), 372 (20), 200 (75), 183 (30).

HRMS: *m/z* calcd for C₃₆H₃₂N₃O₃ (MH⁺): 554.2444; found: 554.2444.

¹H NMR (CDCl₃): δ = 8.45-6.50 (m, 23 H, ArH, OH and NH), 4.75, 4.66 (2 q, *J* = 7 Hz, 1 H, CH), 3.78, 3.66 (2 q, *J* = 7 Hz, 1 H, CH), 2.03, 1.77 (2 d, *J* = 7 Hz, 3 H, CH₃), 1.26, 1.24 (2 d, *J* = 7 Hz, 3 H, CH₃).

¹³C NMR (CDCl₃): δ = 177.3, 174.9 (2 s, C=O), 161.5, 161.4 (2 s, C4), 159.6, 159.1 (s, C2), 147.2, 147.1 (2 s, C8a), 145.4, 145.3 (2 s, C1 of 2 Ph), 136.5 (s, C1'), 135.2, 135.1 (2 d, C7), 134.2, 134.03 (2 s, C4'a), 132.1, 130.9 (2 s, C8'a), 129.7, 129.5 (2 d, C5'), 129.2, 128.8 (2 d, C5), 128.1, 127.8 (2 d, C3 of 2 Ph), 127.7, 127.2 (2 d, C2 of Ph), 127.1, 126.9 (2 d, C6), 126.9, 126.8 (2 d, C8), 126.5, 126.4 (2 d, C3'), 126.2, 126.0 (2 d, C2'), 126.0, 125.7 (2 d, C4'), 125.4, 125.3 (2 d, C7'), 124.4, 124.2 (2 d, C6'), 124.3 (d, C4 of 2 Ph), 123.1, 122.6 (2 d, C8'), 120.2 (s, C4a), 79.0, 78.9 (2 s, C-OH), 43.0, 42.1 (2 d, CH), 41.6, 41.6 (2 d, CH), 17.2, 17.1 (2 q, CH₃), 15.0 (q, CH₃).

2-(2,2-Diphenyl-2-hydroxy-1-methylethyl)-3-(2-phenoxybutyrylamino)quinazolin-4(3*H*)-one (17)

mp 183-184 °C; **17a/17b** = 1:4 (¹H NMR).

Anal. Calcd for C₃₃H₃₁N₃O₄: C, 74.28; H, 5.86; N, 7.87. Found: C, 74.2; H, 5.7; N, 8.1.

EI-MS: *m/z* (%) = 351 (4), 216 (12), 182 (35), 135 (22), 105 (100), 77 (85), 51 (40).

CI-MS: *m/z* (%) = 534 (MH⁺, 19), 357 (100).

HRMS: *m/z* calcd for C₃₃H₃₂N₃O₄ (MH⁺): 534.2393; found: 534.2393.

Compound 17a

¹H NMR (DMSO-d₆): δ = 11.45 (s, exch., 1 H, NH), 8.07 (dd, *J* = 8, 1 Hz, 1 H, H5), 7.77-6.86 (m, 19 H, ArH and OH), 4.94 (t, *J* = 7 Hz, 1 H, CH), 3.99 (q, *J* = 7 Hz, 1 H, CH), 2.20 (m, 2 H, CH₂), 1.16 (t, *J* = 7 Hz, 3 H, CH₃), 0.90 (d, *J* = 7 Hz, 3 H, CH₃).

¹³C NMR (DMSO-d₆): δ = 171.6 (s, C=O), 161.8 (s, C4), 159.0 (s, C2), 157.5 (s, C1 of OPh), 148.1 (s, C8a), 145.7, 144.6 (2 s, C1 of 2 Ph), 135.0 (d, C7), 129.1 (d, C3 of OPh), 127.9, 127.7 (2 d, C3 of 2 Ph), 127.0 (d, C5), 126.5 (d, C6), 126.4 (d, C8), 125.5 (d, C4 of 2 Ph), 125.0, 124.8 (2 d, C2 of 2 Ph), 121.6 (d, C4 of OPh), 120.2 (s, C4a), 115.5 (d, C2 of OPh), 78.7 (s, C-OH), 77.7 (d, CH), 41.8 (d, CH), 26.1 (t, CH₂), 14.0 (q, CH₃), 9.5 (q, CH₃).

Compound 17b

¹H NMR (DMSO-d₆): δ = 11.74 (s, exch., 1 H, NH), 8.07 (dd, *J* = 8, 1 Hz, 1 H, H5), 7.77-6.86 (m, 19 H, ArH and OH), 4.82 (t, *J* = 7 Hz, 1 H, CH), 3.95 (q, *J* = 7 Hz, 1 H, CH), 2.50 (m, 2 H, CH₂), 1.22 (t, *J* = 7 Hz, 3 H, CH₃), 0.95 (d, *J* = 7 Hz, 3 H, CH₃).

¹³C NMR (DMSO-d₆): δ = 172.6 (s, C=O), 162.3 (s, C4), 158.2 (s, C2), 157.5 (s, C1 of OPh), 148.2 (s, C8a), 145.1, 144.7 (2 s, C1 of 2 Ph), 135.0 (d, C7), 129.5 (d, C3 of OPh), 128.0, 127.8 (2 d, C3 of 2 Ph), 127.0 (d, C5), 126.5 (d, C6), 126.4 (d, C8), 125.9 (d, C4 of 2 Ph), 125.1, 124.8 (2 d, C2 of 2 Ph), 121.7 (d, C4 of OPh), 120.1 (s, C4a), 114.9 (d, C2 of OPh), 78.9 (s, C-OH), 78.8 (d, CH), 41.1 (d, CH), 26.5 (t, CH₂), 14.1 (q, CH₃), 9.6 (q, CH₃).

2-(2-Hydroxy-1-methyl-2-phenylethyl)-3-(2-phenoxybutyrylamino)quinazolin-4(3*H*)-one (18)

mp 100-102 °C; four diastereoisomers (¹H NMR).

EI-MS: *m/z* (%) = 230 (6), 216 (28), 173 (32), 135 (44), 105 (49), 94 (33), 77 (100).

CI-MS: *m/z* (%) = 458 (MH⁺, 33%), 281 (70), 197 (40), 175 (53), 136 (46), 114 (41), 105 (100), 94 (56).

HRMS: *m/z* calcd for C₂₇H₂₈N₃O₄ (MH⁺): 458.2080; found: 458.2074.

¹H NMR (CDCl₃): δ = 9.19, 9.08, 8.98, 8.72 (4 s, exch., 1 H, NH), 8.04-6.51 (m, 14 H, ArH), 5.65, 5.52, 5.30, 5.20 (4 s, exch., 1 H, OH), 4.79-4.19 (m, 2 H, 2 CH), 3.37, 3.02, 2.71, 2.51 (4 pentet, *J* = 7 Hz, 1 H, CH), 2.24-1.98 (m, 2 H, CH₂), 1.32-0.96 (m, 3 H, CH₃), 0.91, 0.87, 0.46, 0.35 (4 d, *J* = 7 Hz, 3 H, CH₃).

¹³C NMR (CDCl₃): δ = 171.2, 171.0 (2 s, C=O), 161.0, 160.1 (2 s, C4), 159.0, 158.7 (2 s, C2), 157.5, 157.4 (2 s, C1 of OPh), 145.2, 145.0 (2 s, C8a), 143.9, 143.6, 143.6 (3 s, C1 of Ph), 135.1 (d, C7), 129.6 (d, C3 of OPh), 127.9, 127.8 (2 d, C3 of Ph), 127.7 (d, C5), 127.2 (d, C6), 127.0, 126.7 (2 d, C2 of Ph), 126.6, 125.4 (2 d, C4 of Ph), 126.3 (d, C8), 121.6, 121.5 (2 d, C4 of OPh), 121.4 (s, C4a), 115.6, 115.4, 115.1 (3 d, C2 of OPh), 78.2, 78.0, 77.4 (3 d, CH), 73.7, 72.9 (2 d, CH), 42.9, 42.6 (2 d, CH), 26.1, 26.0, 25.8 (3 t, CH₂), 13.5, 13.2, 13.1 (3 q, CH₃), 9.4, 9.3 (2 q, CH₃).

2-(2-Hydroxy-1-methyl-2-phenylpropyl)-3-(2-phenoxybutyrylamino)quinazolin-4(3*H*)-one (19)

mp 157-158 °C; four diastereoisomers (NMR).

EI-MS: *m/z* (%) = 351 (54), 258 (29), 220 (52), 216 (100), 202 (98), 189 (64).

CI-MS: *m/z* (%) = 472 (MH⁺, 6), 295 (12), 175 (23), 138 (100), 105 (32).

HRMS: *m/z* calcd for C₂₈H₃₀N₃O₄ (MH⁺): 472.2236; found: 472.2239.

¹H NMR (CDCl₃): δ = 8.91, 8.89, 8.76, 8.71 (4 s, exch., 1 H, NH), 8.15-6.79 (m, 14 H, ArH), 5.60, 6.04 (2 br s, exch., 1 H, OH), 4.79-4.63 (m, 1 H, CH), 3.48, 3.21, 3.11, 2.90 (4 q, *J* = 7 Hz, 1 H, CH), 2.14-1.98 (m, 2 H, CH₂), 1.56, 1.54, 1.37, 1.35 (4 s, 3 H, CH₃), 1.15, 1.13 (2 t, *J* = 7 Hz, 3 H, CH₃), 0.92, 0.51 (2 d, *J* = 7 Hz, 3 H, CH₃).

¹³C NMR (CDCl₃): δ = 172.9, 172.8, 172.6 (3 s, C=O), 162.1, 161.9, 161.8 (3 s, C4), 159.5, 159.4 (2 s, C2), 157.4, 157.1 (2 s, C1 of OPh), 146.2, 146.1 (2 s, C8a), 145.5, 145.3 (2 s, C1 of Ph), 135.3 (d, C7), 130.2, 130.1, 130.0 (3 d, C3 of OPh), 128.2, 128.1 (2 d, C3 of Ph), 127.3, 127.2 (2 d, C2 of Ph), 127.0 (d, C5), 126.5, 126.4 (2 d, C6), 124.9, 124.8 (2 d, C8), 123.1, 122.6, 122.4 (3 d, C4 of Ph), 120.6 (s, C4a), 116.8, 116.7 (2 d, C4 of OPh), 114.7, 114.6 (2 d, C2 of OPh), 81.0, 80.8, 79.9, 79.6 (4 d, CH), 75.5, 75.4 (2 s, C-OH), 44.7, 44.6, 44.2 (3 d, CH), 30.0, 29.8 (2 q, CH₃), 26.9, 26.8, 26.2 (3 t, CH₂), 15.0, 14.9, 14.4 (3 q, CH₃), 9.8, 9.7, 9.0 (3 q, CH₃).

2-(2,2-Diphenyl-2-hydroxy-1-methylethyl)-3-(2-phenylbutyrylamino)quinazolin-4(3*H*)-one (20)

mp 182-183 °C; **20a/20b** = 1:4 (¹H NMR).

Anal. Calcd for C₃₃H₃₁N₃O₃: C, 76.59; H, 6.00; N, 8.12. Found: C, 76.6; H, 6.0; N, 8.1.

CI-MS: *m/z* (%) = 518 (MH⁺, 90), 357 (12), 336 (100), 200 (22), 183 (50), 175 (33), 105 (21).

HRMS: *m/z* calcd for C₃₃H₃₂N₃O₃ (MH⁺): 518.2444; found: 518.2444.

Compound 20a

¹H NMR (DMSO-*d*₆): δ = 11.71 (br s, exch., 1 H, NH), 8.14 (dd, *J* = 8, 1 Hz, 1 H, H5), 7.77-6.86 (m, 18 H, ArH), 6.73 (s, exch., 1 H, OH), 4.12 (q, *J* = 7 Hz, 1 H, CH), 3.66 (t, *J* = 7 Hz, 1 H, CH), 2.34-2.07 (m, 2 H, CH₂), 1.33 (d, *J* = 7 Hz, 3 H, CH₃), 1.07 (t, *J* = 7 Hz, 3 H, CH₃).

¹³C NMR (DMSO-*d*₆): δ = 172.8 (s, C=O), 162.4 (s, C4), 158.6 (s, C2), 148.5 (s, C8a), 146.2, 145.1 (2 s, C1 of Ph's), 135.2 (d, C7), 129.5, 128.3, 128.1 (3 d, C3 of Ph's), 128.0 (d, C5), 127.4 (d, C6), 127.3 (d, C8), 126.6, 126.0 (2 d, C4 of Ph's), 125.8, 125.7, 125.4 (3 d, C2 of Ph's), 120.3 (s, C4a), 79.1 (s, C-OH), 52.3 (d, CH), 41.6 (d, CH), 25.8 (t, CH₂), 14.3 (q, CH₃), 12.7 (q, CH₃).

Compound 20b

¹H NMR (DMSO-*d*₆): δ = 11.71 (br s, exch., 1 H, NH), 8.12 (dd, *J* = 8, 1 Hz, 1 H, H5), 7.77-6.86 (m, 18 H, ArH), 6.73 (s, exch., 1 H, OH), 3.86 (q, *J* = 7 Hz, 1 H, CH), 3.63 (t, *J* = 7 Hz, 1 H, CH), 2.28-1.87 (m, 2 H, CH₂), 1.21 (d, *J* = 7 Hz, 3 H, CH₃), 0.90 (t, *J* = 7 Hz, 3 H, CH₃).

¹³C NMR (DMSO-*d*₆): δ = 173.3 (s, C=O), 162.7 (s, C4), 158.5 (s, C2), 148.3 (s, C8a), 145.9, 145.0 (s, C1 of Ph's), 135.4 (d, C7), 129.2, 128.8, 128.5 (3 d, C3 of

Ph's), 127.8 (d, C5), 127.6 (d, C6), 126.8 (d, C8), 126.3, 126.1 (2 d, C4 of Ph's), 125.0, 124.8, 124.7 (3 d, C2 of Ph's), 120.1 (s, C4a), 79.0 (s, C-OH), 51.6 (d, CH), 41.6 (d, CH), 24.7 (t, CH₂), 14.0 (q, CH₃), 12.4 (q, CH₃).

2-(1,2-Dimethyl-2-hydroxybutyl)-3-(2-phenylbutyrylamino)quinazolin-4(3*H*)-one (21)

mp 90-93 °C; at least seven diastereoisomers (¹H NMR).

CI-MS: *m/z* (%) = 408 (MH⁺, 93), 336 (58), 273 (12), 247 (84), 175 (100).

HRMS: *m/z* calcd for C₂₄H₃₀N₃O₃ (MH⁺): 408.2287; found: 408.2287.

¹H NMR (CDCl₃): δ = 8.94, 8.84, 8.83, 8.76, 8.64, 8.46, 8.39 (7 s, exch., 1 H, NH), 8.19 (m, 1 H, H5), 7.77-6.90 (m, 8 H, ArH), 6.13, 6.06, 5.81, 5.67, 5.23, 5.10, 4.56 (7 s, exch., 1 H, OH), 3.65-3.58 (m, 1 H, CH), 3.11, 2.88, 2.74, 2.70, 2.59, 2.46, 2.37 (7 q, *J* = 7 Hz, 1 H, CH), 2.30-2.18 (m, 2 H, CH₂), 1.92, 1.88, 1.85, 1.80 (4 s, 3 H, CH₃), 1.69-0.52 (m, 11 H, CH₂ and 3 CH₃).

¹³C NMR (CDCl₃): δ = 174.3, 174.1, 173.2 (3 s, C=O), 162.7, 162.5, 161.9, 161.7 (4 s, C4), 160.3, 160.1, 160.0 (3 s, C2), 146.5, 146.3 (2 s, C8a), 138.9, 138.8, 138.7 (3 s, C1 of Ph), 135.1, 135.0 (2 d, C7), 129.0, 128.9 (2 d, C3 of Ph), 128.1, 128.0 (2 d, C2 of Ph), 127.9, 127.8 (2 d, C5), 127.7, 127.6 (2 d, C6), 127.3, 127.2 (2 d, C8), 127.1, 127.0, 126.9 (3 d, C4 of Ph), 120.5, 120.4 (2 s, C4a), 74.6, 73.9, 73.8, 73.6 (4 s, C-OH), 53.3, 52.9, 52.8 (3 d, CH), 41.4, 41.1, 40.6, 40.4 (4 d, CH), 34.4, 33.8, 31.9, 26.8, 25.7, 25.6 (6 t, 2 CH₂), 23.0, 22.6, 20.0 (3 q, CH₃), 14.3, 14.1, 13.7, 13.5 (4 q, CH₃), 12.4, 12.1, 12.0, 10.6 (4 q, CH₃), 8.4, 8.2, 8.1, 7.7 (4 q, CH₃).

2-(1-Diphenylhydroxymethylpropyl)-3-(2-phenoxypropionylamino)quinazolin-4(3*H*)-one (22)

mp 112-114 °C; four diastereoisomers (NMR).

EI-MS: *m/z* (%) = 369 (12), 351 (43), 173 (18), 323 (100), 258 (44).

CI-MS: *m/z* (%) = 534 (MH⁺, 33), 371 (19), 352 (100), 260 (16), 217 (18).

HRMS: *m/z* calcd for C₃₃H₃₂N₃O₄ (MH⁺): 534.2393; found: 534.2397.

¹H NMR (DMSO-*d*₆): δ = 11.47, 11.24 (2 br s, exch., 1 H, NH), 8.07 (m, 1 H, H5), 7.79-6.83 (m, 19 H, ArH and OH), 5.35, 5.24 (2 q, *J* = 7 Hz, 1 H, CH), 4.28, 4.06, 3.96, 3.84 (4 t, *J* = 7.5 Hz, 1 H, CH), 1.98-1.44 (m, 2 H, CH₂), 1.82, 1.78, 1.75, 1.68 (4 d, *J* = 7 Hz, 3 H, CH₃), 0.71, 0.41, 0.35, 0.30 (4 t, *J* = 7.5 Hz, 3 H, CH₃).

¹³C NMR (DMSO-*d*₆): δ = 173.5, 173.3, 172.6, 172.4 (4 s, C=O), 161.8, 161.7, 161.4 (3 s, C4), 158.3, 158.2, 158.1, 158.0 (4 s, C2), 157.3, 157.2, 157.1 (3 s, C1 of OPh), 148.6, 148.5, 148.4, 148.3 (4 s, C8a), 145.7, 145.4, 144.7, 144.5 (4 s, C1 of 2 Ph), 135.5, 135.4 (2 d, C7), 129.9, 129.8, 129.7 (3 d, C3 of OPh), 128.3, 128.0

(2 d, C3 of 2 Ph), 127.9, 127.8, 127.5 (3 d, C2 of 2 Ph), 126.8, 126.5 (2 d, C5), 126.2, 126.1 (2 d, C6), 125.5, 125.3 (2 d, C8), 125.1, 125.0 (2 d, C4 of 2 Ph), 121.9, 121.6 (2 d, C4 of OPh), 119.9, 119.8 (2 s, C4a), 115.4, 115.3, 114.6 (3 d, C2 of OPh), 79.7, 79.5, 79.2 (3 s, C-OH), 73.6, 73.4, 71.8 (3 d, CH), 47.9, 74.5, 47.3, 47.1 (4 d, CH), 22.9, 22.7 (2 t, CH₂), 19.4, 19.1, 18.8, 18.5 (4 q, CH₃), 12.6, 12.3, 12.1, 12.0 (4 q, CH₃).

2-(1-Phenylhydroxymethylpropyl)-3-(2-phenoxypropionylamino)quinazolin-4(3*H*)-one (23)

mp 158–159 °C; **23a/23b** = 3:5 (¹H NMR).

EI-MS: *m/z* (%) = 323 (10), 230 (24), 187 (12), 160 (23), 121 (44), 77 (100).

CI-MS: *m/z* (%) = 458 (MH⁺, 9), 295 (48), 189 (53), 183 (46), 166 (17), 122 (52), 106 (41), 91 (100).

HRMS: *m/z* calcd for C₂₇H₂₈N₃O₄ (MH⁺): 458.2080; found: 458.2075.

Compound 23a

¹H NMR (DMSO-*d*₆): δ = 11.15 (br s, exch., 1 H, NH), 7.94–6.75 (m, 14 H, ArH), 4.91 (q, *J* = 7 Hz, 1 H, CH), 4.79 (d, *J* = 3.5 Hz, exch., 1 H, OH), 4.74 (d, *J* = 5 Hz, 1 H, CHOH), 2.87 (m, 1 H, CH), 1.63 (m, 1 H, 1 H of CH₂), 1.48 (d, *J* = 7 Hz, 3 H, CH₃), 1.32 (m, 1 H, 1 H of CH₂), 0.40 (t, *J* = 7.5 Hz, 3 H, CH₃).

¹³C NMR (DMSO-*d*₆): δ = 172.0 (s, C=O), 159.5 (s, C4), 158.8 (s, C2), 157.1 (s, C1 of OPh), 145.9 (s, C8a), 143.9 (s, C1 of Ph), 135.1 (d, C7), 129.6 (d, C3 of OPh), 127.7 (d, C3 of Ph), 127.3 (d, C5), 127.0 (d, C2 of Ph), 126.7 (d, C6), 126.4 (d, C8), 126.1 (d, C4 of Ph), 121.5 (d, C4 of OPh), 120.4 (s, C4a), 115.1 (d, C2 of OPh), 73.2 (d, CH), 72.1 (d, CH), 50.1 (d, CH), 20.5 (t, CH₂), 19.1 (q, CH₃), 10.6 (q, CH₃).

Compound 23b

¹H NMR (DMSO-*d*₆): δ = 11.15 (br s, exch., 1 H, NH), 7.94–6.75 (m, 14 H, ArH), 5.15 (d, *J* = 3.5 Hz, exch., 1 H, OH), 5.00 (q, *J* = 7 Hz, 1 H, CH), 4.88 (d, *J* = 4.5 Hz, 1 H, CHOH), 2.98 (m, 1 H, CH), 1.86 (m, 1 H, 1 H of CH₂), 1.43 (d, *J* = 7 Hz, 3 H, CH₃), 1.23 (m, 1 H, 1 H of CH₂), 0.34 (t, *J* = 7.5 Hz, 3 H, CH₃).

¹³C NMR (DMSO-*d*₆): δ = 171.4 (s, C=O), 159.2 (s, C4), 158.4 (s, C2), 157.0 (s, C1 of OPh), 146.3 (s, C8a), 144.1 (s, C1 of Ph), 135.0 (d, C7), 129.5 (d, C3 of OPh), 127.9 (d, C3 of Ph), 127.5 (d, C5), 126.9 (d, C2 of Ph), 126.6 (d, C6), 126.3 (d, C8), 126.2 (d, C4 of Ph), 121.6 (d, C4 of OPh), 120.6 (s, C4a), 115.5 (d, C2 of OPh), 73.0 (d, CH), 72.2 (d, CH), 50.2 (d, CH), 18.7 (t, CH₂), 18.5 (q, CH₃), 11.7 (q, CH₃).

2-(1-Ethyl-2-hydroxy-2-phenylpropyl)-3-(2-phenoxypropionylamino)quinazolin-4(3*H*)-one (24)

mp 79–81 °C; four diastereoisomers (NMR).

EI-MS: *m/z* (%) = 323 (7), 230 (22), 173 (18), 160 (40), 121 (32), 105 (49), 94 (22), 77 (100).

CI-MS: *m/z* (%) = 472 (MH⁺, 5), 352 (17), 309 (18), 189 (76), 138 (100), 122 (22), 91 (49).

HRMS: *m/z* calcd for C₂₇H₂₈N₃O₄ (MH⁺): 472.2235; found: 472.2235.

¹H NMR (DMSO-*d*₆): δ = 11.35 (br s, exch., 1 H, NH), 8.23–6.78 (m, 14 H, ArH), 6.19, 6.11, 5.98 (3 s, exch., 1 H, OH), 5.42–5.04 (m, 1 H, CH), 2.67–2.33 (m, 1 H, CH), 1.92–1.87 (m, 1 H, 1 H of CH₂), 1.81, 1.79, 1.75, 1.66 (4 d, *J* = 7 Hz, 3 H, CH₃), 1.72, 1.70, 1.69, 1.66 (4 s, CH₃), 1.17 (m, 1 H, 1 H of CH₂), 0.95, 0.76, 0.51, 0.37 (4 t, *J* = 7.5 Hz, 3 H, CH₃).

¹³C NMR (DMSO-*d*₆): δ = 172.4, 171.8, 171.3, 170.8 (4 s, C=O), 162.1, 162.0, 160.6, 160.2 (4 s, C4), 159.0, 158.9, 158.8, 158.5 (4 s, C2), 158.2, 158.1, 157.2, 157.0 (4 s, C1 of OPh), 149.3, 147.0, 146.7, 146.6 (4 s, C8a), 146.3, 145.7, 145.6, 145.1 (4 s, C1 of Ph), 135.5, 135.3, 135.0, 134.6 (4 d, C7), 129.6, 129.5, 129.3, 128.1 (4 d, C3 of OPh), 128.0, 127.9, 127.8, 127.5 (4 d, C3 of Ph), 127.4, 127.3, 127.2, 127.1 (4 d, C5), 126.9, 126.8, 126.7, 126.6 (4 d, C2 of Ph), 126.4, 126.3, 126.1, 125.9 (4 d, C6), 125.1, 124.8 (2 d, C8), 124.7, 124.6 (2 d, C4 of Ph), 121.7, 121.6, 121.5, 121.1 (4 d, C4 of OPh), 120.8, 120.6, 120.4, 120.3 (4 s, C4a), 115.5, 115.2, 115.0, 114.9 (4 d, C2 of OPh), 76.3, 76.2, 75.5, 75.4 (4 s, C-OH), 73.4, 72.6, 72.0, 71.2 (4 d, CH), 53.7, 52.1, 51.4, 50.3 (4 d, CH), 31.0, 30.8, 30.0, 29.7 (4 q, CH₃), 22.3, 22.0, 21.4, 20.9 (4 t, CH₂), 19.3, 19.2, 19.0, 18.7 (4 q, CH₃), 13.7, 13.5, 12.3, 12.0 (4 q, CH₃).

2-(1-Diphenylhydroxymethylpropyl)-3-(2-phenoxybutyrylamino)quinazolin-4(3*H*)-one (25)

mp 139–140 °C; four diastereoisomers (NMR)

EI-MS: *m/z* (%) = 365 (51), 337 (100), 308 (20), 277 (38).

CI-MS: *m/z* (%) = 548 (MH⁺, 41), 371 (89), 366 (100).

HRMS: *m/z* calcd for C₃₄H₃₄N₃O₄ (MH⁺): 548.2549; found: 548.2549.

¹H NMR (DMSO-*d*₆): δ = 11.55, 11.51, 11.45, 11.28 (4 s, exch., 1 H, NH), 8.10–6.85 (m, 20 H, ArH and OH), 5.11, 5.02 (2 t, *J* = 7 Hz, 1 H, CH), 4.26, 4.05, 3.93, 3.82 (4 t, *J* = 7.5 Hz, 1 H, CH), 2.22–1.43 (m, 4 H, 2 CH₂), 1.20, 1.09 (2 t, *J* = 7 Hz, 3 H, CH₃), 0.68, 0.34, 0.30, 0.28 (4 t, *J* = 7.5 Hz, 3 H, CH₃).

¹³C NMR (DMSO-*d*₆): δ = 172.8, 171.7, 171.6 (3 s, C=O), 161.8, 161.3 (2 s, C4), 158.2, 158.0 (2 s, C2), 157.7, 157.6 (2 s, C1 of OPh), 148.6, 148.5 (2 s, C8a), 145.7, 144.4 (2 s, C1 of 2 Ph), 135.5, 135.4 (2 d, C7), 129.9, 129.7 (2 d, C3 of OPh), 128.3, 128.1 (2 d, C3 of 2 Ph), 127.9, 127.5 (2 d, C5), 126.8, 126.7 (2 d, C6), 126.4, 125.5 (2 d, C8), 126.1, 125.5 (2 d, C4 of 2 Ph), 125.3, 125.1 (2 d, C2 of 2 Ph), 121.7, 121.6 (2 d, C4 of OPh), 119.9 (s, C4a), 115.5, 114.7 (2 d, C2 of OPh), 79.7, 79.2 (2 s, C-OH), 78.7, 76.9 (2 d, CH), 47.9, 47.8, 47.7 (3 d, CH), 26.3, 25.8 (2 t, CH₂), 22.8, 22.7 (2 t, CH₂), 12.7, 12.0 (2 q, CH₃), 9.7, 9.6, 9.5, 9.4 (4 q, CH₃).

2-(1-Phenylhydroxymethylpropyl)-3-(2-phenoxybutyrylamino)quinazolin-4(3*H*)-one (26)

mp 136–137 °C; **26a/26b** = 3.5 (^1H NMR).

EI-MS: m/z (%) = 365 (62), 337 (100), 308 (22), 272 (45).

CI-MS: m/z (%) = 472 (MH^+ , 16), 295 (33), 197 (70), 189 (100), 180 (18), 139 (18), 136 (23), 105 (82), 88 (27).

HRMS: m/z calcd for $\text{C}_{27}\text{H}_{28}\text{N}_3\text{O}_4$ (MH^+): 472.2235; found: 472.2230.

Compound 26a

^1H NMR (DMSO- d_6): δ = 11.26 (br s, exch., 1 H, NH), 8.13–6.97 (m, 15 H, ArH and OH), 5.11 (d, J = 4 Hz, 1 H, CHO), 4.94 (t, J = 7 Hz, 1 H, CH), 3.08 (m, 1 H, CH), 2.14–1.38 (m, 4 H, 2 CH_2), 1.17 (t, J = 7 Hz, 3 H, CH_3), 0.58 (t, J = 7.5 Hz, 3 H, CH_3).

^{13}C NMR (DMSO- d_6): δ = 171.3 (s, C=O), 159.6 (s, C4), 158.8 (s, C2), 157.5 (s, C1 of OPh), 145.9 (s, C8a), 143.9 (s, C1 of Ph), 135.2 (d, C7), 129.6 (d, C3 of OPh), 127.7 (d, C3 of Ph), 127.3 (d, C5), 126.9 (d, C2 of Ph), 126.6 (d, C6), 126.4 (d, C8), 126.3 (d, C4 of Ph), 121.6 (d, C4 of OPh), 120.4 (s, C4a), 115.4 (d, C2 of OPh), 78.2 (d, CH), 72.1 (d, CH), 48.4 (d, CH), 26.1 (t, CH_2), 20.5 (t, CH_2), 15.5 (q, CH_3), 10.5 (q, CH_3).

Compound 26b

^1H NMR (DMSO- d_6): δ = 11.42 (br s, exch., 1 H, NH), 8.13–6.97 (m, 15 H, ArH and OH), 5.34 (d, J = 4 Hz, 1 H, CHO), 4.98 (t, J = 7 Hz, 1 H, CH), 3.12 (m, 1 H, CH), 2.14–1.38 (m, 4 H, 2 CH_2), 1.09 (t, J = 7 Hz, 3 H, CH_3), 0.54 (t, J = 7.5 Hz, 3 H, CH_3).

^{13}C NMR (DMSO- d_6): δ = 170.7 (s, C=O), 159.2 (s, C4), 158.4 (s, C2), 157.5 (s, C1 of OPh), 146.3 (s, C8a), 144.0 (s, C1 of Ph), 135.0 (d, C7), 129.5 (d, C3 of OPh), 127.9 (d, C3 of Ph), 127.5 (d, C5), 127.0 (d, C2 of Ph), 126.7 (d, C6), 126.4 (d, C8), 126.3 (d, C4 of Ph), 121.7 (d, C4 of OPh), 120.6 (s, C4a), 115.7 (d, C2 of OPh), 77.3 (d, CH), 72.9 (d, CH), 50.1 (d, CH), 25.6 (t, CH_2), 18.5 (t, CH_2), 11.7 (q, CH_3), 9.4 (q, CH_3).

2-(1-Ethyl-2-hydroxy-2-phenylpropyl)-3-(2-phenoxybutyrylamino)quinazolin-4(3*H*)-one (27)

mp 105–107 °C; **27a/27b** = 4.5 (^1H NMR).

EI-MS: m/z (%) = 365 (19), 337 (25), 272 (10), 244 (53), 230 (100), 214 (24), 202 (27).

EI-MS: m/z (%) = 486 (MH^+ , 33), 366 (15), 309 (100).

HRMS: m/z calcd for $\text{C}_{29}\text{H}_{32}\text{N}_3\text{O}_4$ (MH^+): 486.2393; found: 486.2394.

Compound 27a

^1H NMR (DMSO- d_6): δ = 11.34 (br s, exch., 1 H, NH), 8.21–6.94 (m, 14 H, ArH), 5.27 (s, exch., 1 H, OH), 5.06–5.00 (m, 1 H, CH), 3.44–3.41 (m, 1 H, CH), 2.15–1.56

(m, 4 H, 2 CH_2), 1.19 (s, 3 H, CH_3), 0.85 (t, J = 7 Hz, 3 H, CH_3), 0.23 (t, J = 7.5 Hz, 3 H, CH_3).

^{13}C NMR (DMSO- d_6): δ = 172.2 (s, C=O), 160.0 (s, C4), 158.4 (s, C2), 157.5 (s, C1 of OPh), 147.0 (s, C8a), 145.7 (s, C1 of Ph), 135.3 (d, C7), 129.7 (d, C3 of OPh), 127.9 (d, C3 of Ph), 127.4 (d, C5), 127.2 (d, C6), 126.6 (d, C8), 124.7 (d, C2 of Ph), 121.5 (d, C4 of Ph), 120.3 (d, C4 of OPh), 119.4 (s, C4a), 115.1 (d, C2 of OPh), 76.9 (d, CH), 75.3 (s, C-OH), 50.2 (d, CH), 29.5 (q, CH_3), 25.9 (t, CH_2), 21.9 (t, CH_2), 11.9 (q, CH_3), 9.7 (q, CH_3).

Compound 27b

^1H NMR (DMSO- d_6): δ = 11.34 (br s, exch., 1 H, NH), 8.21–6.94 (m, 14 H, ArH), 6.01 (s, exch., 1 H, OH), 5.06–5.00 (m, 1 H, CH), 3.10–3.08 (m, 1 H, CH), 2.15–1.56 (m, 4 H, 2 CH_2), 1.28 (s, 3 H, CH_3), 1.15 (t, J = 7 Hz, 3 H, CH_3), 0.33 (t, J = 7.5 Hz, 3 H, CH_3).

^{13}C NMR (DMSO- d_6): δ = 170.6 (s, C=O), 162.1 (s, C4), 158.6 (s, C2), 157.6 (s, C1 of OPh), 146.7 (s, C8a), 145.1 (s, C1 of Ph), 135.5 (d, C7), 129.5 (d, C3 of OPh), 128.0 (d, C3 of Ph), 127.5 (d, C5), 127.1 (d, C6), 126.5 (d, C8), 125.1 (d, C2 of Ph), 121.5 (d, C4 of Ph), 120.6 (d, C4 of OPh), 120.4 (s, C4a), 115.4 (d, C2 of OPh), 78.2 (d, CH), 75.5 (s, C-OH), 52.1 (d, CH), 30.8 (q, CH_3), 26.5 (t, CH_2), 23.6 (t, CH_2), 11.3 (q, CH_3), 9.5 (q, CH_3).


Acknowledgment

We thank the EPSRC Mass Spectroscopy Service, University of Wales Swansea, for recording the mass spectra. We also thank the EPSRC, the Higher Education Funding Council for Wales (ELWA-HEFCW) and the University of Wales Swansea for grants that enabled the purchase and upgrading of NMR equipment used in the course of this work. G. A. El-Hiti thanks the University of Wales Swansea for financial support and the Royal Society of Chemistry for an international author grant.

References

- (1) Permanent address; Department of Chemistry, Faculty of Science, Tanta University, Tanta 31527, Egypt.
- (2) (a) Beak, P.; Zajdel, W. J.; Reitz, D. B. *Chem. Rev.* **1984**, *84*, 471. (b) Snieckus, V. *Chem. Rev.* **1990**, *90*, 879. (c) El-Hiti, G. A. *Heterocycles* **2000**, *53*, 1839. (d) Mongin, F.; Quéguiner, G. *Tetrahedron* **2001**, *57*, 4059. (e) Turck, A.; Plé, N.; Mongin, F.; Quéguiner, G. *Tetrahedron* **2001**, *57*, 4489. (f) Anctil, E. J.-G.; Snieckus, V. *J. Organomet. Chem.* **2002**, *653*, 150. (g) Smith, K.; El-Hiti, G. A. *Curr. Org. Synth.* **2004**, *1*, in press. (h) Chinchilla, R.; Nájera, C.; Yus, M. *Chem. Rev.* **2004**, in press.
- (3) See for example: (a) Choppin, S.; Gros, P.; Fort, Y. *Org. Lett.* **2000**, *2*, 803. (b) Kondo, Y.; Asai, M.; Miura, T.; Uchiyama, M.; Sakamoto, T. *Org. Lett.* **2001**, *3*, 13. (c) Rebstock, A.-S.; Mongin, F.; Trécourt, F.; Quéguiner, G. *Tetrahedron Lett.* **2002**, *43*, 767. (d) Rhonstad, P.; Wensbo, D. *Tetrahedron Lett.* **2002**, *43*, 3137. (e) Mukherjee, C.; De, A. *Synlett* **2002**, 325. (f) Cuperly, D.; Gros, P.; Fort, Y. *J. Org. Chem.* **2002**, *67*, 238. (g) Gros, P.; Louérat, F.; Fort, Y. *Org. Lett.* **2002**, *4*, 175. (h) Toudic, F.; Plé, N.; Turck, A.; Quéguiner, G. *Tetrahedron* **2002**, *58*, 238. (i)

- Bunch, L.; Krogsgaard-Larsen, P.; Madsen, U. *J. Org. Chem.* **2002**, *67*, 2375. (j) Katritzky, A. R.; Manju, K.; Steel, P. J. *J. Org. Chem.* **2003**, *68*, 407. (k) Gros, P.; Choppin, S.; Fort, Y. *J. Org. Chem.* **2003**, *68*, 2243. (l) El-Hiti, G. A. *Synthesis* **2004**, *363*. (m) Cabiddu, M. C.; Cabiddu, S.; Cadoni, E.; De Montis, S.; Fattuoni, C.; Melis, S. *Tetrahedron* **2004**, *60*, 3915.
- (4) Murray, T. P.; Hay, J. V.; Protlock, D. E.; Wolfe, J. F. *J. Org. Chem.* **1974**, *39*, 595.
- (5) Rathman, T. L.; Sleevi, M. C.; Krafft, M. F.; Wolfe, J. F. *J. Org. Chem.* **1980**, *45*, 2169.
- (6) Smith, K.; El-Hiti, G. A.; Abdel-Megeed, M. F.; Abdo, M. A. *Collect. Czech. Chem. Commun.* **1999**, *64*, 515.
- (7) Smith, K.; El-Hiti, G. A.; Abdel-Megeed, M. F.; Abdo, M. A. *J. Org. Chem.* **1996**, *61*, 656.
- (8) El-Hiti, G. A. *Monatsh. Chem.* **2004**, *135*, 323.
- (9) Smith, K.; El-Hiti, G. A.; Abdel-Megeed, M. F.; Abdo, M. A. *J. Org. Chem.* **1996**, *61*, 647.
- (10) Smith, K.; El-Hiti, G. A.; Abdo, M. A.; Abdel-Megeed, M. F. *J. Chem. Soc., Perkin Trans. I* **1995**, *1029*.
- (11) Plé, N.; Turck, A.; Chapoulaud, V.; Quéguiner, G. *Tetrahedron* **1997**, *53*, 2871.
- (12) Dai, X.; Virgil, S. *Tetrahedron Asymmetry* **1999**, *10*, 25.
- (13) Chapoulaud, V. G.; Salliot, I.; Plé, N.; Turck, A.; Quéguiner, G. *Tetrahedron* **1999**, *55*, 5389.
- (14) See for example: (a) Honda, G.; Tabata, M.; Tsuda, M. *Planta Med.* **1979**, *37*, 172. (b) Johne, S. *Prog. Drug Res.* **1982**, *26*, 259. (c) Johne, S. *The Alkaloids* **1986**, *29*, 99. (d) Schlecker, R.; Treiber, H. J.; Behl, B.; Hofmann, H. P. Ger. Offen 4 241 563, **1994**; *Chem. Abstr.* **1994**, *121*, 230787. (e) Barker, A. J. Eur. Patent 635 498, **1995**; *Chem. Abstr.* **1995**, *122*, 214099. (f) Barker, A. J.; Johnstine, C. PCT Int Appl WO 300 44, **1997**; *Chem. Abstr.* **1995**, *127*, 220671. (g) Hour, M.-J.; Huang, L.-J.; Kuo, S.-C.; Xia, Y.; Bastow, K.; Nakanishi, Y.; Hamel, E.; Lee, K.-H. *J. Med. Chem.* **2000**, *43*, 4479. (h) Ho, N.; Harapanhalli, R. S.; Dahman, B. A.; Chen, K.; Wang, K. *Bioconjugate Chem.* **2002**, *13*, 357.
- (15) See for example: (a) Smith, K.; El-Hiti, G. A.; Hamilton, A. *J. Chem. Soc., Perkin Trans. I* **1998**, *4041*. (b) Smith, K.; El-Hiti, G. A.; Pritchard, G. J.; Hamilton, A. *J. Chem. Soc., Perkin Trans. I* **1999**, *2299*. (c) Smith, K.; El-Hiti, G. A.; Shukla, A. P. *J. Chem. Soc., Perkin Trans. I* **1999**, *2305*. (d) Smith, K.; El-Hiti, G. A.; Abdel-Megeed, M. F. *Russ. J. Org. Chem.*, **2003**, *39*, 430. (e) Smith, K.; El-Hiti, G. A.; Hawes, A. C. *Synthesis* **2003**, *2047*. (f) Smith, K.; El-Hiti, G. A.; Mahgoub, S. A. *Synthesis* **2003**, *2345*. (g) El-Hiti, G. A. *Synthesis* **2003**, *2799*.
- (16) Sammour, A.; Selim, M.; El Kasaby, M.; Abdalla, M. *Acta Chim. (Budapest)* **1973**, *78*, 293; *Chem. Abstr.* **1973**, *79*, 137061.
- (17) (a) Essawy, A.; El-Hashash, M. A.; El-Gendy, A. M.; Hamad, M. M. M. *Indian J. Chem.* **1982**, *21B*, 593.
- (18) El-Hiti, G. A. *Spectrosc. Lett.*, **1999**, *32*, 671.
- (19) *Vogel's Textbook of Practical Organic Chemistry*: 5th Edn, Longman, Harlow **1989**.
- (20) Perrin, D. D.; Armarego, W. L. F.; Perrin, D. R. *Purification of Laboratory Chemicals*, Pergamon, 3rd Edn, Butterworth Heinemann, Oxford, **1988**.

Lithiation of Chiral 3-Acylamino-2-alkylquinazolin-4(3*H*)-ones**Manuscript submission checklist**

- Statement of significance of work.
- Full mailing address, telephone and fax numbers, and email address of the corresponding author.
- Paper save as a PDF file.
- Original Word file.
- Original graphic files.
- Graphical abstract.