

# Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <http://orca.cf.ac.uk/13557/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Liu, Xin-Ge, Tang, Mei-Lan and Martin, Ralph Robert 2008. Periodic solutions for a kind of Liénard equation. *Journal of Computational and Applied Mathematics* 219 (1) , pp. 263-275. 10.1016/j.cam.2007.07.024 file

Publishers page: <http://dx.doi.org/10.1016/j.cam.2007.07.024>  
<<http://dx.doi.org/10.1016/j.cam.2007.07.024>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



# Existence and Uniqueness of Periodic Solutions of a Liénard Equation with Delay

Xin-Ge Liu<sup>a\*</sup>, Mei-Lan Tang<sup>a</sup>, Ralph R. Martin<sup>b</sup>

<sup>a</sup> School of Mathematical Science and Computing Technology, Central South University

Changsha, Hunan 410083, China

<sup>b</sup>School of Computer Science, Cardiff University, Cardiff, UK

## Abstract

Using inequality techniques and coincidence degree theory, new results are provided concerning the existence and uniqueness of  $T$ -periodic solutions for a Liénard equations with delay. An illustrative example is provided to demonstrate that the results in this paper hold under weaker conditions than existing results, and are more effective.

**MSC:** 34D 40.

**Keywords:** Liénard equation; Delay; Periodic solution; Existence; Uniqueness; Coincidence degree.

## 1 Introduction

There has been a great deal of work on Liénard equations, which have been used to describe fluid-mechanical and nonlinear elastic mechanical phenomena. For example, in [2, 3, 4, 8] and [13], time maps and phase plane analysis were used to examine the existence of periodic solutions to Liénard equations, and several sufficient conditions for this existence were established. Recently, Liu and Huang [9] discussed the existence and uniqueness of periodic solutions for a Liénard equation with delay, of form

$$x''(t) + f(x(t))x'(t) + g(t, x(t - \tau(t))) = p(t) \quad (1)$$

where  $f, \tau, p : R \rightarrow R$  and  $g : R \times R \rightarrow R$  are continuous functions,  $\tau$  and  $p$  are  $T$ -periodic (i.e. periodic with period  $T$ ),  $g$  is  $T$ -periodic in its first argument, and  $T > 0$ .

In recent years, periodic solutions to Eq.(1) have been extensively studied in the literature (see, for example, [1] and [5, 7, 10, 11, 12, 14]). However, to the best of our knowledge, most authors have only considered the *existence* of periodic solutions, and few results exist concerning both existence and *uniqueness* of periodic solutions to Eq.(1). Liu and Huang [9] provide a sufficient condition for such existence and uniqueness, but their result leaves space for improvement. In their work, they required that the following constraint should be imposed on the Liénard equation:

$$C_1 D \frac{T}{2\pi} + C_2 \frac{T}{2\pi} + \frac{bT^2}{2\pi} < 1.$$

Here,  $C_1, C_2$  and  $b$  are nonnegative constants determined by

$$|f(x_1) - f(x_2)| \leq C_1 |x_1 - x_2|, \quad |f(x)| \leq C_2, \quad C_2 \frac{T}{2\pi} + b \frac{T^2}{4\pi} < 1,$$

$$|g(t, x_1) - g(t, x_2)| \leq b |x_1 - x_2|, \quad \forall t, x_1, x_2, x \in R;$$

and  $d$  is a positive constant such that one of the following conditions holds:

$$x(g(t, x) - p(t)) < 0, \quad \forall t \in R, |x| \geq d \quad \text{or} \quad x(g(t, x) - p(t)) > 0, \quad \forall t \in R, |x| \geq d.$$

---

\*Corresponding Author. Tel.: +86-731-8830812. E-mail address: liuxgliuhua@163.com

Furthermore,

$$D = \frac{[bd + \max\{|g(t, 0)| : 0 \leq t \leq T\} + |p|_\infty]T}{1 - (C_2 \frac{T}{2\pi} + b \frac{T^2}{2\pi})}.$$

Upon examining their proof of Lemma 2.5 and Theorem 3.1 in [9], we have found certain errors. The corrected version of this constraint should read

$$C_1 D \frac{\sqrt{T^3}}{2\pi} + C_2 \frac{T}{2\pi} + \frac{bT^2}{2\pi} < 1.$$

In this paper, we reconsider periodic solutions of a Liénard equation with delay as given in Eq.(1). The main purpose of this paper is to establish a new sufficient condition for the existence and uniqueness of  $T$ -periodic solutions of Eq.(1). Using inequality techniques, we obtain sharp *a priori* estimates for a periodic solution to Eq.(1). Furthermore, by using improved estimates for  $|x|_\infty$ ,  $|x'|_\infty$  and  $|x'|_2$  and coincidence degree theory, both the existence and uniqueness of  $T$ -periodic solutions of Eq.(1) under this sufficient condition are proved. This sufficient condition improves upon the main result obtained in [9], as we demonstrate using an illustrative example.

For simplicity, throughout this paper, we adopt the following notation:

$$|x|_k = \left( \int_0^T |x(t)|^k dt \right)^{1/k}$$

and

$$|x|_\infty = \max_{t \in [0, T]} |x(t)|.$$

In order to use Mawhin's continuation theorem to study the existence of periodic solution of Eq.(1), we introduce the following spaces and operators. Let

$$X = \{x|x \in C^1(R, R), x(t+T) = x(t), \forall t \in R\}$$

and

$$Y = \{x|x \in C(R, R), x(t+T) = x(t), \forall t \in R\}$$

be two Banach spaces with norms

$$\|x\|_X = \max\{|x|_\infty, |x'|_\infty\} \quad \text{and} \quad \|x\|_Y = |x|_\infty.$$

Let  $D(L) = \{x|x \in X, x'' \in C(R, R)\}$ . Define a linear operator  $L : D(L) \subset X \rightarrow Y$  by setting

$$Lx = x'' \tag{2}$$

We also define a nonlinear operator  $N : X \rightarrow Y$  by setting

$$Nx = -f(x(t))x'(t) - g(t, x(t - \tau(t))) + p(t). \tag{3}$$

Obviously,  $\text{Ker}L = R$ , and  $\text{Im}L = \{x|x \in Y, \int_0^T x(s)ds = 0\}$ . Thus the operator  $L$  is a Fredholm operator with index zero. Define the continuous projector  $P : X \rightarrow \text{Ker}L$  and the averaging projector  $Q : Y \rightarrow Y$  by setting  $Px(t) = x(0) = x(T)$  and  $Qx(t) = \frac{1}{T} \int_0^T x(s)ds$ . Hence,  $\text{Im}P = \text{Ker}L$  and  $\text{Ker}Q = \text{Im}L$ . Denoting by  $L_P^{-1} : \text{Im}L \rightarrow D(L) \cap \text{Ker}P$  the inverse of  $L|_{D(L) \cap \text{Ker}P}$ , we have

$$L_P^{-1}y(t) = -\frac{t}{T} \int_0^T (t-s)y(s)ds + \int_0^t (t-s)y(s)ds. \tag{4}$$

It is convenient to introduce the following assumption

(A<sub>0</sub>): there exist nonnegative constants  $C_1$  and  $C_2$  such that

$$|f(x_1) - f(x_2)| \leq C_1|x_1 - x_2|, \quad |f(x)| \leq C_2$$

for all  $x_1, x_2, x \in R$ .

## 2 Preliminaries

In view of Eqs.(2) and (3), the operator equation  $Lx = \lambda Nx$  is equivalent to the following:

$$x''(t) + \lambda[f(x(t))x'(t) + g(t, x(t - \tau(t)))] = \lambda p(t) \quad (5)$$

where  $\lambda \in (0, 1)$ .

For convenience of use, we introduce the Continuation Theorem [5] as follows.

**Lemma 1.** *Let  $X$  and  $Y$  be two Banach spaces. Suppose that  $L : D(L) \subset X \rightarrow Y$  is a Fredholm operator with index zero and  $N : X \rightarrow Y$  is  $L$ -compact on the closure  $\bar{\Omega}$  of  $\Omega$ , where  $\Omega$  is an open bounded subset of  $X$ . Moreover, assume that each of the following conditions is satisfied:*

1.  $Lx \neq \lambda Nx, \forall x \in \partial\Omega \cap D(L), \lambda \in (0, 1)$ ;
2.  $Nx \notin \text{Im}L, \forall x \in \partial\Omega \cap \text{Ker}L$ ;
3. The Brouwer degree  $\deg\{QN, \Omega \cap \text{Ker}L, 0\} \neq 0$ .

Then equation  $Lx = Nx$  has at least one solution on  $\bar{\Omega}$ .

The following lemmas will be useful to prove our main results in Section 3.

**Lemma 2.** *See [6, 14]. If  $x \in C^2(\mathbb{R}, \mathbb{R})$  with  $x(t + T) = x(t)$ , then*

$$|x'(t)|_2^2 \leq \left(\frac{T}{2\pi}\right)^2 |x''(t)|_2^2. \quad (6)$$

**Lemma 3.** *Suppose that there exists a constant  $d > 0$  such that one of the following conditions holds:*

- (A<sub>1</sub>):  $x(g(t, x) - p(t)) < 0, \forall t \in \mathbb{R}, |x| \geq d$ ;
- (A<sub>2</sub>):  $x(g(t, x) - p(t)) > 0, \forall t \in \mathbb{R}, |x| \geq d$ .

If  $x(t)$  is a  $T$ -periodic solution of (5), then

$$|x|_\infty \leq d + \frac{\sqrt{T}}{2} |x'|_2. \quad (7)$$

*Proof.* Let  $x(t)$  be a  $T$ -periodic solution of Eq.(5). Then, integrating Eq.(5) from 0 to  $T$ , we have

$$\int_0^T [g(t, x(t - \tau(t))) - p(t)] dt = 0. \quad (8)$$

This implies that there exists  $\xi \in [0, T]$  such that

$$g(\xi, x(\xi - \tau(\xi))) - p(\xi) = 0. \quad (9)$$

Taking this together with (A<sub>1</sub>) or (A<sub>2</sub>) as appropriate, we have

$$|x(\xi - \tau(\xi))| < d. \quad (10)$$

Let  $\xi - \tau(\xi) = mT + t_0$ , where  $t_0 \in [0, T]$  and  $m$  is an integer. Then,

$$\begin{aligned} |x(t)| &= \left| x(t_0) + \int_{t_0}^t x'(s) ds \right| \\ &= \left| x(mT + t_0) + \int_{t_0}^t x'(s) ds \right| \\ &\leq |x(\xi - \tau(\xi))| + \left| \int_{t_0}^t x'(s) ds \right| \\ &< d + \int_{t_0}^t |x'(s)| ds \end{aligned} \quad (11)$$

where  $t \in [t_0, t_0 + T]$ .

Since  $x(t)$  is the  $T$ -periodic solution, for  $t \in [t_0, t_0 + T]$ ,

$$\begin{aligned}
|x(t)| &= |x(t_0 + T) + \int_{t_0+T}^t x'(s)ds| \\
&= |x(t_0 + T) - \int_t^{t_0+T} x'(s)ds| \\
&\leq |x(t_0 + T)| + |\int_t^{t_0+T} x'(s)ds| \\
&\leq |x(t_0 + T)| + \int_t^{t_0+T} |x'(s)|ds \\
&\leq d + \int_t^{t_0+T} |x'(s)|ds.
\end{aligned} \tag{12}$$

Combining Eqs.(11) and (12) gives

$$\begin{aligned}
|x(t)| &\leq d + \frac{1}{2} \int_{t_0}^{t_0+T} |x'(s)|ds \\
&\leq d + \frac{1}{2} \int_0^T |x'(s)|ds.
\end{aligned} \tag{13}$$

Using the Schwartz inequality yields

$$\begin{aligned}
|x(t)| &\leq d + \frac{1}{2} \sqrt{T} (\int_0^T |x'(s)|^2 ds)^{1/2} \\
&= d + \frac{1}{2} \sqrt{T} |x'|_2.
\end{aligned} \tag{14}$$

Therefore,

$$|x|_\infty = \max_{t \in [0, T]} |x(t)| \leq d + \frac{1}{2} \sqrt{T} |x'|_2. \tag{15}$$

This completes the proof of Lemma 3.  $\square$

**Lemma 4.** *Let  $(A_0)$  and either  $(A_1)$  or  $(A_2)$  hold. Assume that the following condition is satisfied:*

*$(A_3)$ : There exists a nonnegative constant  $b$  such that*

$$C_2 \frac{T}{2\pi} + b \frac{T^2}{4\pi} < 1, \quad \text{and} \quad |g(t, x_1) - g(t, x_2)| \leq b|x_1 - x_2|, \quad \forall t, x_1, x_2 \in R.$$

*If  $x(t)$  is a  $T$ -periodic solution of Eq.(1), then*

$$|x'|_2 \leq D_1 \tag{16}$$

where

$$D_1 = \frac{[bd + \max\{|g(t, 0)| : 0 \leq t \leq T\}] + |p|_\infty T}{2(1 - C_2 \frac{T}{2\pi} - b \frac{T^2}{4\pi})}.$$

*Proof.* Let  $x(t)$  be a  $T$ -periodic solution of Eq.(1). From  $(A_1)$  or  $(A_2)$ , we can easily show that Inequality (7) also holds. Multiplying Eq.(1) by  $x''(t)$  and then integrating from 0 to  $T$ , in view of Inequalities (6) and (7),  $(A_3)$  and the Schwartz inequality, we have

$$|x''|_2^2 = - \int_0^T f(x(t))x'(t)x''(t)dt - \int_0^T g(t, x(t - \tau(t)))x''(t)dt + \int_0^T p(t)x''(t)dt$$

$$\begin{aligned}
&\leq \int_0^T |f(x(t))||x'(t)||x''(t)|dt + \int_0^T |p(t)||x''(t)|dt \\
&\quad + \int_0^T [|g(t, x(t - \tau(t))) - g(t, 0)| + |g(t, 0)||x''(t)|]dt \\
&\leq C_2 \int_0^T |x'(t)||x''(t)|dt + |p|_\infty \int_0^T |x''(t)|dt \\
&\quad + b \int_0^T |x(t - \tau(t))||x''(t)|dt + \int_0^T |g(t, 0)||x''(t)|dt \\
&\leq C_2 |x'|_2 |x''|_2 + |p|_\infty \sqrt{T} |x''|_2 \\
&\quad + b \int_0^T |x(t - \tau(t))||x''(t)|dt + \int_0^T |g(t, 0)||x''(t)|dt \\
&\leq C_2 \frac{T}{2\pi} |x''|_2^2 + |p|_\infty \sqrt{T} |x''|_2 \\
&\quad + b |x|_\infty \sqrt{T} |x''|_2 + \max\{|g(t, 0)| : 0 \leq t \leq T\} \sqrt{T} |x''|_2 \\
&\leq C_2 \frac{T}{2\pi} |x''|_2^2 + |p|_\infty \sqrt{T} |x''|_2 \\
&\quad + b(d + \frac{\sqrt{T}}{2} |x'|_2) \sqrt{T} |x''|_2 + \max\{|g(t, 0)| : 0 \leq t \leq T\} \sqrt{T} |x''|_2 \\
&\leq C_2 \frac{T}{2\pi} |x''|_2^2 + |p|_\infty \sqrt{T} |x''|_2 \\
&\quad + b(d + \frac{\sqrt{T}}{2} \frac{T}{2\pi} |x''|_2) \sqrt{T} |x''|_2 + \max\{|g(t, 0)| : 0 \leq t \leq T\} \sqrt{T} |x''|_2 \\
&\leq [C_2 \frac{T}{2\pi} + b \frac{T^2}{4\pi}] |x''|_2^2 \\
&\quad + [bd + \max\{|g(t, 0)| : 0 \leq t \leq T\} + |p|_\infty] \sqrt{T} |x''|_2. \tag{17}
\end{aligned}$$

Thus,

$$[1 - C_2 \frac{T}{2\pi} - b \frac{T^2}{4\pi}] |x''|_2 \leq [bd + \max\{|g(t, 0)| : 0 \leq t \leq T\} + |p|_\infty] \sqrt{T}, \tag{18}$$

so

$$|x''|_2 \leq \frac{[bd + \max\{|g(t, 0)| : 0 \leq t \leq T\} + |p|_\infty] \sqrt{T}}{1 - C_2 \frac{T}{2\pi} - b \frac{T^2}{4\pi}}. \tag{19}$$

Since  $x(0) = x(T)$ , there exists a constant  $\zeta \in [0, T]$  such that  $x'(\zeta) = 0$ , and

$$\begin{aligned}
|x'(t)| &= |x'(\zeta) + \int_\zeta^t x''(s)ds| \\
&\leq |x'(\zeta)| + \int_\zeta^t |x''(s)|ds \\
&= \int_\zeta^t |x''(s)|ds \tag{20}
\end{aligned}$$

where  $t \in [\zeta, T + \zeta]$ . Again,

$$\begin{aligned}
|x'(t)| &= |x'(\zeta + T) + \int_{\zeta+T}^t x''(s)ds| \\
&= |x'(\zeta + T) - \int_t^{\zeta+T} x''(s)ds| \\
&\leq |x'(\zeta + T)| + \int_t^{\zeta+T} |x''(s)|ds
\end{aligned}$$

$$\begin{aligned}
&\leq |x'(\zeta + T)| + \int_t^{\zeta+T} |x''(s)|ds \\
&= |x'(\zeta)| + \int_t^{\zeta+T} |x''(s)|ds \\
&= \int_t^{\zeta+T} |x''(s)|ds
\end{aligned} \tag{21}$$

where  $t \in [0, T]$ . Inequalities (20) and (21) imply that

$$\begin{aligned}
2|x'(t)| &\leq \int_{\zeta}^t |x''(s)|ds + \int_t^{\zeta+T} |x''(s)|ds \\
&= \int_{\zeta}^{\zeta+T} |x''(s)|ds \\
&= \int_0^T |x''(s)|ds \\
&\leq \sqrt{T}|x''|_2
\end{aligned} \tag{22}$$

where  $t \in [0, T]$ .

Obviously,

$$|x'(t)| \leq \frac{1}{2}\sqrt{T}|x''|_2 \quad \text{for all } t \in [0, T], \tag{23}$$

so

$$|x'|_{\infty} \leq \frac{1}{2}\sqrt{T}|x''|_2. \tag{24}$$

From Inequalities (19) and (24), we have

$$|x'|_{\infty} \leq \frac{[bd + \max\{|g(t, 0)| : 0 \leq t \leq T\} + |p|_{\infty}]T}{2(1 - C_2 \frac{T}{2\pi} - b \frac{T^2}{4\pi})}. \tag{25}$$

This completes the proof of Lemma 4.  $\square$

**Lemma 5.** *Let  $(A_1)$  or  $(A_2)$  hold. Assume that the following condition is satisfied:*

*$(A_4)$  Suppose that  $(A_0)$  holds,  $g(t, x)$  is a strictly monotone function in  $x$ , and there exists a nonnegative constant  $b$  such that*

$$C_1 D_1 \frac{\sqrt{T^3}}{4\pi} + C_2 \frac{T}{2\pi} + \frac{bT^2}{4\pi} < 1, \quad \text{and} \quad |g(t, x_1) - g(t, x_2)| \leq b|x_1 - x_2|$$

where  $C_1, C_2, D_1$  are defined as before.

Then Eq.(1) has at most one  $T$ -periodic solution.

*Proof.* Suppose that  $x_1(t)$  and  $x_2(t)$  are two  $T$ -periodic solutions of Eq.(1). Then

$$(x_1(t) - x_2(t))'' + (f(x_1(t))x_1'(t) - f(x_2(t))x_2'(t)) + (g(t, x_1(t - \tau(t))) - g(t, x_2(t - \tau(t)))) = 0. \tag{26}$$

Set  $Z(t) = x_1(t) - x_2(t)$ . Then, from Eq.(26), we obtain

$$Z''(t) + (f(x_1(t))x_1'(t) - f(x_2(t))x_2'(t)) + (g(t, x_1(t - \tau(t))) - g(t, x_2(t - \tau(t)))) = 0. \tag{27}$$

Since  $x_1(t)$  and  $x_2(t)$  are  $T$ -periodic, integrating Eq.(27) from 0 to  $T$ , we obtain

$$\int_0^T (g(t, x_1(t - \tau(t))) - g(t, x_2(t - \tau(t))))dt = 0. \tag{28}$$

Using the integral mean value theorem, it follows that there exists a constant  $\gamma \in [0, T]$  such that

$$g(\gamma, x_1(\gamma - \tau(\gamma))) - g(\gamma, x_2(\gamma - \tau(\gamma))) = 0. \quad (29)$$

Let  $\gamma - \tau(\gamma) = nT + \bar{\gamma}$ , where  $\bar{\gamma} \in [0, T]$  and  $n$  is an integer. Then, Eq.(29), together with (A4), implies that there exists a constant  $\bar{\gamma} \in [0, T]$  such that

$$Z(\bar{\gamma}) = x_1(\bar{\gamma}) - x_2(\bar{\gamma}) = x_1(\gamma - \tau(\gamma)) - x_2(\gamma - \tau(\gamma)) = 0. \quad (30)$$

Thus,

$$\begin{aligned} |Z(t)| &= |Z(\bar{\gamma}) + \int_{\bar{\gamma}}^t Z'(s)ds| \\ &= \left| \int_{\bar{\gamma}}^t Z'(s)ds \right| \\ &\leq \int_{\bar{\gamma}}^t |Z'(s)|ds. \end{aligned} \quad (31)$$

Again

$$\begin{aligned} |Z(t)| &= |Z(\bar{\gamma} + T) + \int_{\bar{\gamma}+T}^t Z'(s)ds| \\ &= |Z(\bar{\gamma} + T) - \int_t^{\bar{\gamma}+T} Z'(s)ds| \\ &= \left| - \int_t^{\bar{\gamma}+T} Z'(s)ds \right| \\ &\leq \int_t^{\bar{\gamma}+T} |Z'(s)|ds. \end{aligned} \quad (32)$$

Hence

$$\begin{aligned} 2|Z(t)| &\leq \int_{\bar{\gamma}}^t |Z'(s)|ds + \int_t^{\bar{\gamma}+T} |Z'(s)|ds \\ &\leq \int_{\bar{\gamma}}^{\bar{\gamma}+T} |Z'(s)|ds \\ &= \int_0^T |Z'(s)|ds \\ &\leq \sqrt{T} \left( \int_0^T |Z'(s)|^2 ds \right)^{1/2} \\ &= \sqrt{T} |Z'|_2. \end{aligned} \quad (33)$$

Hence

$$|Z|_\infty \leq \frac{1}{2} \sqrt{T} |Z'|_2. \quad (34)$$

Multiplying Eq.(27) by  $Z''(t)$ , then integrating from 0 to  $T$ , from Inequalities (6) and (34), and the Schwartz inequality, we get

$$\begin{aligned} |Z''|_2^2 &= - \int_0^T (f(x_1(t))x_1'(t) - f(x_2(t))x_2'(t))Z''(t)dt \\ &\quad - \int_0^T (g(t, x_1(t - \tau(t))) - g(t, x_2(t - \tau(t))))Z''(t)dt \end{aligned}$$



$$\begin{aligned}
&\leq \int_0^T |f(x_1(t))| |x'_1(t) - x'_2(t)| |Z''(t)| dt \\
&\quad + \int_0^T |f(x_1(t)) - f(x_2(t))| |x'_2(t)| |Z''(t)| dt \\
&\quad + b \int_0^T |x_1(t - \tau(t)) - x_2(t - \tau(t))| |Z''(t)| dt \\
&\leq \int_0^T C_2 |Z'(t)| |Z''(t)| dt \\
&\quad + \int_0^T C_1 |Z(t)| |x'_2(t)| |Z''(t)| dt \\
&\quad + b \int_0^T |Z(t - \tau(t))| |Z''(t)| dt \\
&\leq C_2 \left( \int_0^T |Z'(t)|^2 dt \right)^{1/2} \left( \int_0^T |Z''(t)|^2 dt \right)^{1/2} \\
&\quad + C_1 |Z|_\infty \int_0^T |x'_2(t)| |Z''(t)| dt \\
&\quad + b |Z|_\infty \int_0^T |Z''(t)| dt \\
&\leq C_2 |Z'|_2 |Z''|_2 + C_1 |Z|_\infty |x'_2|_2 |Z''|_2 + b |Z|_\infty \sqrt{T} |Z''|_2 \\
&\leq [C_2 + \frac{1}{2} C_1 \sqrt{T} D_1 + \frac{1}{2} b T] |Z'|_2 |Z''|_2 \\
&\leq [\frac{1}{2} C_1 \sqrt{T} D_1 + C_2 + \frac{1}{2} b T] \frac{T}{2\pi} |Z''|_2^2 \\
&= [C_1 D_1 \frac{\sqrt{T^3}}{4\pi} + C_2 \frac{T}{2\pi} + \frac{b T^2}{4\pi}] |Z''|_2^2. \tag{35}
\end{aligned}$$

Since  $Z(t)$ ,  $Z'(t)$  and  $Z''(t)$  are  $T$ -periodic and continuous functions, in view of (A<sub>4</sub>) and Inequalities (6), (30) and (35), we have

$$Z(t) = Z'(t) = Z''(t) \equiv 0, \quad \forall t \in R.$$

Thus,

$$x_1(t) \equiv x_2(t), \quad \forall t \in R.$$

Therefore, Eq.(1) has at most one  $T$ -periodic solution. □

### 3 Main result

**Theorem 1.** *Let (A<sub>1</sub>) or (A<sub>2</sub>) hold. Assume that condition (A<sub>4</sub>) is satisfied. Then Eq. (1) has a unique  $T$ -periodic solution.*

*Proof.* Lemma 5 states that Eq. (1) has at most one  $T$ -periodic solution. Thus, to prove Theorem 1, it suffices to show that Eq.(1) has at least one  $T$ -periodic solution. To do this, we apply Lemma 1. Firstly, we claim that the set of all possible  $T$ -periodic solutions of Eq.(5) is bounded. Let  $x(t)$  be a  $T$ -periodic solution of Eq.(5). Multiplying Eq.(5) by  $x''(t)$ , then integrating from 0 to  $T$ , and using Lemmas 2 and 3, Assumption (A<sub>4</sub>) and the Schwartz inequality, we have

$$\begin{aligned}
|x''|_2^2 &= -\lambda \int_0^T f(x(t)) x'(t) x''(t) dt - \lambda \int_0^T g(t, x(t - \tau(t))) x''(t) dt + \lambda \int_0^T p(t) x''(t) dt \\
&\leq \int_0^T |f(x(t))| |x'(t)| |x''(t)| dt + \int_0^T |p(t)| |x''(t)| dt
\end{aligned}$$

$$\begin{aligned}
& + \int_0^T [|g(t, x(t - \tau(t))) - g(t, 0)| + |g(t, 0)|] |x''(t)| dt \\
\leq & C_2 \int_0^T |x'(t)| |x''(t)| dt + |p|_\infty \int_0^T |x''(t)| dt \\
& + b \int_0^T |x(t - \tau(t))| |x''(t)| dt + \int_0^T |g(t, 0)| |x''(t)| dt \\
\leq & C_2 |x'|_2 |x''|_2 + |p|_\infty \sqrt{T} |x''|_2 \\
& + b |x|_\infty \sqrt{T} |x''|_2 + \max\{|g(t, 0)| : 0 \leq t \leq T\} \sqrt{T} |x''|_2 \\
\leq & C_2 \frac{T}{2\pi} |x''|_2^2 + |p|_\infty \sqrt{T} |x''|_2 \\
& + b(d + \frac{\sqrt{T}}{2} |x'|_2) \sqrt{T} |x''|_2 + \max\{|g(t, 0)| : 0 \leq t \leq T\} \sqrt{T} |x''|_2 \\
\leq & C_2 \frac{T}{2\pi} |x''|_2^2 + |p|_\infty \sqrt{T} |x''|_2 \\
& + b(d + \frac{\sqrt{T}}{2} \frac{T}{2\pi} |x''|_2) \sqrt{T} |x''|_2 + \max\{|g(t, 0)| : 0 \leq t \leq T\} \sqrt{T} |x''|_2 \\
\leq & [C_2 \frac{T}{2\pi} + b \frac{T^2}{4\pi}] |x''|_2^2 \\
& + [bd + \max\{|g(t, 0)| : 0 \leq t \leq T\} + |p|_\infty] \sqrt{T} |x''|_2
\end{aligned} \tag{36}$$

which, together with (A<sub>4</sub>), implies that there exist positive constants  $D_3$  and  $D_4$  such that

$$|x''|_2 < D_3, \quad |x'|_2 < D_4, \quad |x|_\infty < D_4. \tag{37}$$

Since  $x(0) = x(T)$ , there exists a constant  $\xi \in [0, T]$  such that

$$x'(\xi) = 0$$

and

$$\begin{aligned}
|x'(t)| &= |x'(\xi) + \int_\xi^t x''(s) ds| \\
&= |\int_\xi^t x''(s) ds| \\
&\leq \int_\xi^t |x''(s)| ds
\end{aligned} \tag{38}$$

where  $t \in [\xi, T + \xi]$ . Similarly,

$$\begin{aligned}
|x'(t)| &= |x'(\xi + T) + \int_{\xi+T}^t x''(s) ds| \\
&= |x'(\xi + T) - \int_t^{\xi+T} x''(s) ds| \\
&= |\int_t^{\xi+T} x''(s) ds| \\
&\leq \int_t^{\xi+T} |x''(s)| ds
\end{aligned} \tag{39}$$

where  $t \in [\xi, T + \xi]$ . So

$$|x'(t)| \leq \frac{1}{2} (\int_\xi^t |x''(s)| ds + \int_t^{\xi+T} |x''(s)| ds)$$

$$\begin{aligned}
&= \frac{1}{2} \int_{\xi}^{\xi+T} |x''(s)| ds \\
&= \frac{1}{2} \int_0^T |x''(s)| ds \\
&\leq \frac{1}{2} \sqrt{T} \|x''\|_2 \\
&\leq \frac{1}{2} \sqrt{T} D_3.
\end{aligned} \tag{40}$$

From Inequalities (37) and (40), there exists a positive constant  $M_1 > \frac{1}{2} \sqrt{T} D_3 + D_4$  such that

$$\|x\|_X \leq |x|_{\infty} + |x'|_{\infty} < M_1.$$

If  $x \in \Omega_1 = \{x | x \in \text{Ker}L \cap X, \text{ and } Nx \in \text{Im}L\}$ , then there exists a constant  $M_2$  such that

$$x(t) \equiv M_2,$$

and

$$\int_0^T [g(t, M_2) - p(t)] dt = 0. \tag{41}$$

Thus

$$\int_0^T M_2 [g(t, M_2) - p(t)] dt = 0. \tag{42}$$

Hence,  $\forall x(t) \in \Omega_1$ ,

$$|x(t)| \equiv |M_2| < d. \tag{43}$$

Let  $M = M_1 + d + 1$ . Set

$$\Omega = \{x | x \in X, |x|_{\infty} < M, |x'|_{\infty} < M\}.$$

It is easy to see from Eqs.(3) and (4) that  $N$  is  $L$ -compact on  $\bar{\Omega}$ . It follows from Eqs.(41) and (43), and the fact  $M > \max\{M_1, d\}$ , that conditions (1) and (2) in Lemma 1 hold.

Furthermore, suppose we define continuous functions  $H_1(x, \mu)$  and  $H_2(x, \mu)$  as

$$\begin{aligned}
H_1(x, \mu) &= (1 - \mu)x - \mu \frac{1}{T} \int_0^T [g(t, x) - p(t)] dt; \mu \in [0, 1], \\
H_2(x, \mu) &= -(1 - \mu)x - \mu \frac{1}{T} \int_0^T [g(t, x) - p(t)] dt; \mu \in [0, 1].
\end{aligned} \tag{44}$$

If  $(A_1)$  holds, then

$$xH_1(x, \mu) \neq 0 \tag{45}$$

where  $x \in \partial\Omega \cap \text{Ker}L$ . Hence, using the homotopy invariance theorem, we have

$$\begin{aligned}
\deg\{QN, \Omega \cap \text{Ker}L, 0\} &= \deg\left\{-\frac{1}{T} \int_0^T [g(t, x) - p(t)] dt, \Omega \cap \text{Ker}L, 0\right\} \\
&= \deg\{x, \Omega \cap \text{Ker}L, 0\} \\
&\neq 0.
\end{aligned} \tag{46}$$

If  $(A_2)$  holds, then

$$xH_2(x, \mu) \neq 0 \tag{47}$$

where  $x \in \partial\Omega \cap \text{Ker}L$ . Hence, using the homotopy invariance theorem, we have

$$\begin{aligned}
\deg\{QN, \Omega \cap \text{Ker}L, 0\} &= \deg\left\{-\frac{1}{T} \int_0^T [g(t, x) - p(t)] dt, \Omega \cap \text{Ker}L, 0\right\} \\
&= \deg\{-x, \Omega \cap \text{Ker}L, 0\} \\
&\neq 0.
\end{aligned} \tag{48}$$

From Lemma 1, we conclude that Theorem 1 is proved.  $\square$

## 4 Example

In this section, we provide an example to illustrate that the sufficient condition given in Theorem 1 in this paper is superior to the recently reported result in [9].

**Example 1.** Let  $g(t, x) = x/(6\pi) \forall t, x \in R$ . Consider the existence and uniqueness of a  $2\pi$ -periodic solution to the following Liénard equation:

$$x''(t) + \frac{3}{8}(\sin x(t))x'(t) + g(t, x(t - \sin^2 t)) = \frac{1}{6\pi}e^{\cos(t)-1}. \quad (49)$$

In this example,  $d = 1, b = 1/(6\pi), C_1 = C_2 = 3/8, \tau(t) = \sin^2 t, T = 2\pi$  and  $p(t) = \frac{1}{6\pi}e^{\cos(t)-1}$ . It is obvious that assumptions  $(A_2)$  and  $(A_4)$  both hold. Using the method in Theorem 1 in [9], we find that

$$\begin{aligned} D &= \frac{[bd + \max\{|g(t, 0)| : 0 \leq t \leq T\}] + |p|_\infty T}{1 - (C_2 \frac{T}{2\pi} + b \frac{T^2}{2\pi})} \\ &= \frac{[\frac{1}{6\pi} + \frac{1}{6\pi}] \cdot 2\pi}{1 - \frac{3}{8} - \frac{1}{3}} \\ &= 2.2857, \end{aligned}$$

$$C_1 D \frac{\sqrt{T^3}}{2\pi} + C_2 \frac{T}{2\pi} + b \frac{T^2}{2\pi} = 2.8569 > 1.$$

Since  $2.8569 > 1$ , the condition in Theorem 1 in [9] is not satisfied and hence it cannot provide any result. Therefore, Theorem 1 in [9] fails, while, our criterion in Theorem 1 in this paper remains applicable, as we now show.

In fact,

$$\begin{aligned} D_1 &= \frac{[bd + \max\{|g(t, 0)| : 0 \leq t \leq T\}] + |p|_\infty T}{2(1 - C_2 \frac{T}{2\pi} - b \frac{T^2}{4\pi})} \\ &= \frac{[\frac{1}{6\pi} + \frac{1}{6\pi}]2\pi}{2 - \frac{3}{4} - \frac{1}{3}} \\ &= 0.7273 \end{aligned}$$

and

$$C_1 D_1 \frac{\sqrt{T^3}}{4\pi} + C_2 \frac{T}{2\pi} + b \frac{T^2}{4\pi} = 0.8835 < 1.$$

Thus, Theorem 1 here shows that Eq.(49) has a unique  $2\pi$ -periodic solution.

This example demonstrates that the condition in our Theorem 1 is weaker than that in [9], and is able to demonstrate existence of unique solutions to certain Liénard equations which the latter cannot decide about. Therefore our results substantially improve the works in [9].

## 5 Acknowledgements

The authors are very grateful to the referee for valuable suggestions which improved this article.

## References

- [1] T.A. Burton, Stability and periodic solution of ordinary and functional differential equations, *Academic Press*, Orland, FL., 1985.

- [2] M.A. Del Pino, R.F. Mansevich, Multiple solutions for the p-Laplacian under global non-resonance, *Proc. Amer. Math. Soc.* 112 (1991) 131–138.
- [3] M.A. Del Pino, R.F. Mansevich, A. Murua, Existence and multiple of solutions with prescribed period for a second order quasi-linear ordinary differential equation, *Nonlinear Anal. TMA* 18 (1992) 79–92.
- [4] C. Fabry, D. Fayyad, Periodic solutions of second order differential equations with a p-Laplacian and asymmetric nonlinearities, *Rend. Ist. Univ. Trieste.* 24 (1992) 207–227.
- [5] R.E. Gaines, J. Mawhin, Coincidence degree and nonlinear differential equations, *Lecture Notes in Math.* Vol.568 Spring-Verlag, 1977.
- [6] G.H. Hardy, J.E. Littlewood, G. Polya, Inequalities, *Cambridge Univ. Press* London 1964.
- [7] X. Huang, Z. Xiang, On existence of  $2\pi$ -periodic solutions for delay Duffing equation  $x'' + g(t, x(t - \tau(t))) = p(t)$ , *Chinese Science Bulletin* 39(1994) 201–203.
- [8] B. Liu, Multiplicity results for periodic solutions of a second order quasi-linear ODE with asymmetric nonlinearities, *Nonlinear Anal. TMA* 33 (1998) 139–160.
- [9] B. Liu, L. Huang, Existence and uniqueness of periodic solutions for a kind of Liénard equation with a deviating argument. *Appl. Math. Lett.* in press 2007.
- [10] S. Lu, W. Ge, Periodic solutions for a kind of Liéneard equations with deviating arguments, *J. Math. Anal. Appl.* 249(2004) 231–243 .
- [11] S. Lu, W. Ge, Sufficient conditions for the existence of periodic solutions to some second order differential equations with a deviating argument, *J. Math. Anal. Appl.* 308(2005) 393–419.
- [12] S. Lu, W. Ge, Periodic solutions for a kind of second order differential equation with multiple deviating arguments, *Appl. Math. Comput.* 146(2003) 195–209.
- [13] R.F. Manásevich, J. Mawhin, Periodic solutions for nonlinear systems with p-Laplacian like operators, *J. Differ. Equations* 145 (1998) 367–393.
- [14] J. Mawhin, Periodic solutions of some vector retarded functional differential equations, *J. Math. Anal. Appl.* 45(1974) 588–603.