

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: https://orca.cardiff.ac.uk/id/eprint/96482/

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

White, James ORCID: https://orcid.org/0000-0001-8371-8453, Greene, Giles ORCID: https://orcid.org/0000-0001-9326-8740, Farewell, Daniel ORCID: https://orcid.org/0000-0002-8871-1653, Rodgers, Sarah, Lyons, Ronan A, Humphreys, Ioan, John, Ann, Webster, Chris, Phillips, Ceri J and Fone, David Lawrence ORCID: https://orcid.org/0000-0002-6476-4881 2016. Improving mental health through the regeneration of deprived neighbourhoods: a prospective controlled quasi-experimental study. The Lancet 388 (Supp 2), S110. 10.1016/S0140-6736(16)32346-7 file

Publishers page: http://dx.doi.org/10.1016/S0140-6736(16)32346-7 http://dx.doi.org/10.1016/S0140-6736(16)32346-7

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

http://orca.cl.ac.uk/policies.html for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Improving mental health through the regeneration of deprived neighborhoods: A prospective controlled quasi-experimental study

```
James White, PhD <sup>12</sup>
Giles Greene, PhD <sup>3</sup>
Daniel Farewell, PhD <sup>3</sup>
Sarah Rodgers, PhD <sup>4</sup>
Ronan A Lyons, MD <sup>4</sup> (professor)
Ioan Humphreys, MSc <sup>5</sup> (early career researcher)
Ann John, MD <sup>4</sup>
Chris Webster, PhD <sup>6</sup> (professor)
Ceri J Phillips, PhD <sup>5</sup> (professor)
David Fone, MD <sup>3</sup> (professor)
```

¹ South East Wales Trials Unit (SEWTU), School of Medicine, Cardiff University, Cardiff. CF14 4YS. Wales, United Kingdom.

² Centre for the Development and Evaluation of Complex Interventions for Public Health Improvement (DECIPHer), School of Medicine, Cardiff University, Cardiff. CF14 4YS. Wales, United Kingdom.

³ Institute of Primary Care & Public Health, School of Medicine, Neuadd Meirionnydd, Cardiff University, Heath Park, Cardiff. CF14 4YS. Wales, United Kingdom.

⁴ Farr Institute, College of Medicine, Swansea University, Swansea. SA2 8PP. Wales, United Kingdom.

⁵ Swansea Centre for Health Economics, College of Human and Health Sciences, Swansea University, Singleton Park, Swansea. SA2 8PP. Wales, United Kingdom.

⁶ The University of Hong Kong, Faculty of Architecture, Department of Urban Planning and

Design, Pokfulam, Hong Kong.

Correspondence: James White, PhD, white 11@cf.ac.uk, South East Wales Trials Unit

(SEWTU), School of Medicine, Cardiff University, Cardiff. CF14 4YS. Wales, United

Kingdom. Dr White is not an early career researcher.

Online protocol: http://bmjopen.bmj.com/content/4/10/e006530.short

Word count: 348 words in abstract

Background Policymakers often target deprived neighborhoods for regeneration with the expectation that population health will improve, since housing and neighborhoods of low quality, as well as the social and economic determinants of poor health, are concentrated in the most deprived areas. Our aim was to examine the effects of Communities First, a Welsh Assembly Government community-led program of neighborhood regeneration targeted at the

100 most deprived electoral wards in Wales (United Kingdom) on mental health.

Methods Information on Communities First regeneration activities in 35 intervention lower super output areas (LSOAs) (n=4,197 subjects) and 75 control LSOAs (n=6,695 subjects) were linked to data from the Caerphilly Health & Social Needs Electronic Cohort (eCATALyST) study, a prospective cohort study in 2001 (before regeneration) and 2008

(after regeneration). Communities First was delivered through multi-agency partnership

boards in each of the 22 local authorities. Boards worked with residents to identify and secure

funding for regeneration activities. The primary outcome was the change in Mental Health

Inventory (a population-based measure of anxiety and depressive symptoms) between 2001

and 2008 recorded in eCATALyST. We examined the changes in mental health in

intervention LSOAs in comparison to control LSOAs, using propensity score matching (1:1

ratio) to balance the level of socioeconomic disadvantage across groups. Sensitivity analysis

examined the impact of length of residence in an intervention area and six types of regeneration activity.

Findings 1,500 regeneration projects were funded 2001 to 2008. Before regeneration mental health was worse in the intervention (mean = 66.6, SD = 22.3) than control group (mean = 71.0, SD = 20.8). After propensity score matching, regeneration was associated with an improvement in the mental health of intervention compared to control group residents (β coefficient = 1.54; 95% CI 0.50 to 2.59), suggesting inequalities in mental health narrowed. We found evidence of a dose response relationship between length of residence and improvements in mental health (p = 0.05 for trend). We could not attribute improvements to any one type of regeneration activity.

Interpretation

Targeted regeneration directed by the residents of deprived urban communities may help to reduce inequalities in mental health.

Funding This study was funded by grants from the National Institute for Social Care and Health Research (RFS-12-05). This work is undertaken with the support of The Centre for the Development and Evaluation of Complex Interventions for Public Health Improvement (MR/KO232331/1) and the Farr Institute of Health Informatics Research.

Contributors

GG, JW, and DF critically appraised the research papers. CW, DF designed the methods classifying interventions funded by Communities First. GG, DF, FD, SR and DF designed the statistical analysis. RAL, AJ and SR led the use of SAIL methodology. JW is principal investigator. DF is the principal investigator of the eCATALyST study. JW and GG wrote the manuscript and all authors contributed to critical revision of the final version. All authors have contributed to the scientific content of the abstract and have seen and approved the final version for publication.

Declarations of interest

We declare no competing interests.