

Psychology of Music

**Maximising self-care through familiarity: The role of practice effects in enhancing music-listening and progressive muscle relaxation for pain management**

Journal:	<i>Psychology of Music</i>
Manuscript ID:	POM-13-577.R2
Manuscript Type:	Original Empirical Investigations
Keyword:	Audio-analgesia, Cognitive coping, Distraction, Acute pain, Pain tolerance, Pain threshold
Abstract:	<p>Distraction and relaxation are regularly recommended as part of Pain Management Programmes, with increasing research highlighting the inclusion of music as part of pain management toolkit. However minimal research has assessed the role of practice effects or familiarity with these techniques when used consistently over time. Passive distraction (participant-selected preferred music) and active distraction (Progressive Muscle Relaxation; PMR) were compared against a no distraction control on the cold-pressor test (CPT). Seventy healthy participants completed the CPT with and without distraction at baseline and one week later. Experimental participants practised their distraction strategies daily between trials (7 days), with control participants keeping an activity log. Familiarity with and preference for distractors increased significantly over time, enhancing pain threshold. PMR and music reduced anxiety, enhanced pain tolerance, minimised pain perception and pain ratings. The active distraction of PMR enhanced self-efficacy to a greater extent than music and also regulated heart rate. Repeated exposure to distraction and relaxation approaches enhanced optimal arousal and complexity, maximising pain management. It is suggested that both PMR, and music, are used together as part of a multidimensional toolkit for pain management.</p>

SCHOLARONE™  
Manuscripts

Running head: Familiarity and Pain Management

1

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Maximising self-care through familiarity: The role of practice effects in enhancing music  
listening and progressive muscle relaxation for pain management

For Peer Review

## Familiarity and Pain Management

2

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

A cumulative amount of research has demonstrated the positive effects that psychological treatment can have upon pain (see Eccleston et al., 2009 for review). Psychological therapies are a fundamental part of pain management programmes (PMPs), with multimodal biopsychosocial strategies collectively used to improve self-care in patients. Coping methods such as distraction and relaxation are employed alongside approaches which attempt to normalise pain through Mindfulness or Acceptance and Commitment Therapy (Eccleston et al., 2009). Together these methods are used as part of a multimodal pain management 'kit' (Cepeda et al., 2006) and are privileged by their ease of use, minimal side-effects, cost-effectiveness and universal applicability (Eccleston et al., 2002; Cepeda et al., 2006).

Limited capacity models of attention posit that attention is a finite cognitive resource and that the processing of pain signals can be mediated by attentional redirection away from pain and towards absorptive tasks/stimuli (Shiffrin and Schneider, 1977; Verhoeven et al., 2010). Hence pain perception can be minimised if the distractor's attentional demands are of sufficient magnitude. However, the efficacy of distraction may be dependent upon whether it is passive or active (Dahlquist et al., 2007). Active distractors redirect attention through the active involvement of the individual in a task (Windich-Biermeier et al., 2007). They aim to alter the physiological response system to pain, through the lessening of the pain-tension cycle, using techniques such as Progressive Muscle Relaxation (PMR; Jacobson, 1938), jaw relaxation (Good et al., 1999; Schaffer and Yucha, 2004), electronic games (Forys and Dahlquist, 2007) mental arithmetic (Mitchell et al., 2006), and breath therapy (Mehling et al., 2005). Passive distractors redirect attention without active or deliberate involvement (Chambers et al., 2009), for example

## Familiarity and Pain Management

3

1  
2  
3 through watching television (MacLaren and Cohen, 2005), listening to music (Mitchell et al.,  
4  
5 2008), or conversation (Zelikovsky et al., 2000). Dahlquist et al (2007) argue that active  
6  
7 distraction is more effective than passive distraction due to increased attentional load, modelled  
8  
9 biologically through the activation of descending inhibitory pain pathways (Edwards, Campbell,  
10  
11 Jamison & Wiech, 2009). There is therefore a significant need to assess the impact of active and  
12  
13 passive distractors on pain management. The most commonly recommended approaches are  
14  
15 music listening (passive) and progressive muscle relaxation (active; Pain Toolkit, 2012), and  
16  
17 these therefore should be priority targets for comparative research.  
18  
19  
20  
21  
22  
23  
24

25 Multiple-resource theory (Wickens 1984) has provided a theoretical perspective which  
26  
27 may explain potential differences between active and passive distractors. Wickens (2008) argued  
28  
29 that when multi-sensory attentional resources are used collectively they are more demanding of  
30  
31 attention. Distractors may be visual, auditory, spatial or verbal, each viewed in multiple-resource  
32  
33 theory as separate attentional stores, giving a prospective rationale for why in pain management  
34  
35 multimodal attentional distractors are more absorbing (Wickens, 2008). Johnson et al (1998)  
36  
37 investigated Multiple-resource theory and demonstrated that the more processing resources a  
38  
39 distractor shares with those of pain perception, the more interference occurs and thus less pain is  
40  
41 felt. This theoretical perspective meshes with the established gold-standard recommended in  
42  
43 healthcare, a desire for a biopsychosocial model of care (following Engel, 1977), in which  
44  
45 biological, psychological and sociological factors involved in the pain experience are valued  
46  
47 equally and managed simultaneously.  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Familiarity and Pain Management

4

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

The level of involvement of the person in their pain management has a significant impact on their self-efficacy: an individual's belief that they are capable to achieve certain levels of performance to control events that affect their lives (Bandura, 1997). Individuals' beliefs regarding their self-efficacy enable human functioning through a number of processes: cognitive, motivational, affective and decision-making (Bandura, 2001). With differential levels of self-efficacy, individuals can self-enhance or self-debilitate. Individuals with high self-efficacy are able to better alleviate stress and anxiety, therefore enabling them to implement better coping strategies. Research suggests that high scores of self-efficacy are inversely related to pain intensity (Borsbo et al, 2010), with findings demonstrating that self-efficacy is a highly accurate predictor of pain-related disability (Sharma et al, 2003) and functioning (Woby et al, 2007). It is possible that active distractors may better initiate increases in self-efficacy than passive distractors, and this may further affect the moderating ability of the pain management technique to minimise pain. Therefore PMR may be more effective than music listening as a result of its promotion of active distraction

PMPs teach the use of distraction/relaxation strategies, and it is expected that these are employed at home, yet research has minimally investigated how repeated use of such strategies can alter their efficacy. Finlay (in press) found that increased familiarity with passive distraction through music, improved patient satisfaction with treatment. Similarly, Mitchell et al (2006) found familiar music reduced pain perception to a greater extent than unfamiliar music. Music was also more effective than other distractors such as art, mathematics or humour (Mitchell et al, 2004; Villareal et al, 2012). Music that is chosen by the participant (preferred music) is also more effective than music provided by the experimenter (non-preferred music), or music chosen

## Familiarity and Pain Management

1  
2  
3 by the participant from a pre-prepared experimenter-defined selection (quasi-preferred music;  
4 following Hekmat & Hertel, 1993; Mitchell et al, 2006). Investigating PMR, Persson,  
5 Veenhuizen, Zachrisson & Gard (2008) found regular practice allows the patient to effectively  
6 identify the difference between muscular tension and release, therefore promoting self-initiated  
7 biofeedback. In addition, repeated exposure to distractors enhances emotional engagement,  
8 enhancing adaptive neuroaffective mechanisms useful in pain management (Koelsch, 2010).  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19

20 Berlyne's (1971) inverted-U theory argues that one's ability to tolerate stimuli is a  
21 function of a trade-off between familiarity and complexity, with moderate levels of familiarity  
22 and complexity enhancing the efficacy of the stimulus. Potentially therefore, distractors that  
23 initiate moderate levels of arousal may be most effective for patients and pain sufferers may  
24 choose to continue using these techniques. To date, research comparing active and passive  
25 distractors has been limited, and familiarity effects have not been considered. This study aims to  
26 explore the role of familiarity in the use of active distraction (PMR) and passive distraction  
27 (music), for the management of acute, laboratory-induced pain in a health community-dwelling  
28 sample. As music listening and PMR are often recommended as useful for self-care in PMPs  
29 (Pain Toolkit, 2012), it is essential that their relative efficacy is investigated in the context of  
30 daily, self-initiated use in a home-based environment. The majority of pain self-management  
31 occurs in the familial environment, beyond the immediate reach of monitored clinical care and  
32 medical attention. It is hypothesised that both active and passive distractors will significantly  
33 enhance pain tolerance (the length of time that pain can be tolerated) and pain threshold (the  
34 length of time before the first reported onset of pain sensation), and reduce subjective measures  
35 of pain intensity during experimental trials. It is expected that regular use/practice of  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Familiarity and Pain Management

6

1  
2  
3  
4 active/passive distraction will also prove beneficial, therefore there will be improvements in all  
5  
6 measures after seven days of practice, at week 2. PMR necessitates greater participation during  
7  
8 use than music listening as it involves wilful contraction and relaxation of specific muscle groups  
9  
10 (de Paula, de Carvalho & dos Santos, 2002). Therefore, it is hypothesised that self-efficacy will  
11  
12 be enhanced to a greater extent by PMR than by music listening.  
13  
14  
15  
16  
17  
18  
19

**Method***Participants*

20  
21  
22  
23  
24  
25 70 healthy participants completed the full study, both male (N = 31) and female (N = 39).  
26  
27 Participants' ages ranged from between 18 to 73 years, (M = 37.50 years; SD = 14.63). All were  
28  
29 recruited from a community-based sample from Hertfordshire and Buckinghamshire. Exclusion  
30  
31 criteria consisted of diabetes, circulation disorders, smoking, pre-existing pain conditions and  
32  
33 claustrophobia due to blindfolding (following Jackson et al, 2005). All participants completed  
34  
35 both assessment phases of the research study.  
36  
37

*Design*

38  
39  
40  
41 A mixed design was employed, using the between-subjects factor of Group (three  
42  
43 conditions); active distraction through Progressive Muscle Relaxation, passive distraction  
44  
45 through preferred music listening and a no distraction (silent) control. Participants were  
46  
47 randomly assigned to one of the three conditions through a computerised randomisation  
48  
49 schedule. Within-subjects factors of Time of Testing (2 levels; Week 1 and Week 2) and Type of  
50  
51 Testing (2 levels; baseline and experimental) were used to assess all outcome measures.  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Familiarity and Pain Management

*Materials**Apparatus*

*Circulatory Water Bath.* A JeioTech circulatory water bath (RW-3025G) was used, which cooled circulating water to 0°C. Consistency in water temperature is important for comparable and reliable results (Mitchell et al, 2004). There is a lack of consensus regarding the maximum length of time the hand should be kept in the water. For safety, maximum length of time participants were allowed to keep their hand in the water was 240 seconds (4 minutes; Jackson et al, 2005).

*Thermometer.* Hand temperature was measured using a digital thermometer (Omron Smart Digital), temperatures being recorded in degrees Celsius.

*Music Player.* Participant-provided preferred music was played through the music software application iTunes, on a MacBook Pro for all participants. Volume level was self-selected by participants as appropriate to personal taste.

*Heart Rate (HR).* HR was recorded using an Omron M6 sphygmomanometer. Heart rate is an objective measure of physiological stress induced by clinical context (Thayer, Åhs, Fredrikson, Sollers & Wager, 2009).

*Stimuli*

*Music.* Individuals were required to bring with them a chosen piece of music in CD format. Previous findings demonstrate the efficacy in pain reduction of chosen (preferred) music over pre-selected (non-preferred) music by the experimenter (see Mitchell et al, 2006). Chosen tracks

## Familiarity and Pain Management

8

were played on the music player (detailed above) during the experimental trials (as appropriate to Group Allocation), without repetition of, adjustment or alteration to the chosen track. All track lengths exceeded the length of time tolerated by participants undertaking the cold pressor test.

*PMR.* 'Progressive Muscle Relaxation' from the album Guided Relaxation by Edna Reinhardt was used. The track described progressive tightening and relaxing of each set of muscles throughout the body, and the physical feelings associated with the reduction of tension in the body. This PMR track was chosen as there was no background music behind the spoken instructions, ensuring the effect of PMR was being measured and not the effect of music. The track lasted for 8 mins and 37 secs.

*Performance Assessments*

*Pain Threshold.* This was calculated as the timespan between participants' immersion of their hand into the water and when they first report experiencing sensations of pain, measured in seconds, using a stopwatch (Duschek et al, 2008). Pontinen (1998) reports that pain threshold is a valid and reliable measure for quantitative evaluation of pain.

*Pain Tolerance.* This was calculated as the timespan between initial immersion of hand into the water, and when participants withdrew their hand, measured in seconds (following Duschek et al, 2008; Mitchell et al, 2008). Edwards et al (2001) reports that pain tolerance provides a quantitative benchmark for objective evaluation of pain.

*Questionnaire Measures*

## Familiarity and Pain Management

*Demographic Questionnaire.* A short 4-item demographic questionnaire was used to assess age, gender and educational status. Additionally, one item was used to confirm whether or not the participant was a smoker or not, in accordance with exclusion criteria. Education status was measured by a fixed choice selection of highest qualification achieved (GCSE, AS-level, A-level, Undergraduate, Masters, Doctoral or other equivalent qualification).

*Short Form State Trait Anxiety Inventory (STAI): State scale.* A six-item scale assessing state anxiety on four-point likert scales, addressing statements such as “I felt calm”, “I felt relaxed” was measured using the endpoints strongly agree and strongly disagree. Originally developed by Spielberger (1983), the STAI is a measure of individual differences in anxiety (Taylor & Deane, 2002). It continues to be widely used in research and applied settings, with literature finding it useful in pain research (Mitchell et al., 2008). The short-form scale has been well-validated and found to be reliable with clinical populations (Marteau & Bekker, 1992).

*Numerical Rating Scales (NRS).* Five 11-point Likert-type numerical ratings scales were used:

1. *NRS-Pain Perception (NRS-PP).* The NRS-PP utilised the end points ‘no pain’ and ‘extreme pain’. The NRS-PP, as developed by Jensen and Karoly (2001) is endorsed by The British Pain Society, and has been used in numerous studies (e.g. Gustavsson & von Koch, 2006).
2. *NRS-Current Pain Intensity (NRS-CPI).* The NRS-CPI is a variation of the NRS-PP, but with the wording altered to reflect a demographic assessment of baseline pain at the present time at the outset of the research study. This was used as a between-groups screening tool, and is validated and used as the NRS-PP.

## Familiarity and Pain Management

10

3. *NRS-Efficacy of Distraction (NRS-EOD)*. The NRS-EOD contained end points ‘not at all’ and ‘completely’, following the statement “I felt able to take my mind off the pain”. The EOD has been effectively used in previous pain research (Mitchell et al, 2008).
4. *NRS-Familiarity (NRS-F)*. The NRS-F was used to assess level of familiarity with the musical selection used by the participants in weeks 1 and 2. The selected end-points were ‘not at all familiar’ and ‘extremely familiar’.
5. *NRS-Preference (NRS-P)*. The NRS-P was used to assess level of preference for the musical selection chosen by the participants. The end-points ‘not at all liked and ‘extremely liked’ were included to anchor the scale.

*Short Form McGill Pain Questionnaire (SF-MPQ; Melzack, 1983)*. Fifteen pain descriptors (11 sensory; 4 affective) were measured using a 4-point Likert scale, where 0 = none, 1 = mild, 2 = moderate and 3 = severe. Total Pain Score (TPS) was used as a principal measure, with subscales for sensory and affective pain. The SF-MPQ has been used in previous literature, showing reliability and validity (Katz & Melzack, 2011).

*The General Self-Efficacy Scale (GSE)*. Ten items were measured using the representative end points ‘not at all true’ and ‘exactly true’. Item scores are totalled to yield an overall score ranging from 10-40: the higher the score, the greater the perceived self-efficacy. Developed by Schwarzer and Jerusalem (1995), the GSE has been used internationally and is accepted to be highly reliable.

*Procedure*

## Familiarity and Pain Management

11

1  
2  
3  
4 Participants willing to take part in the study were recruited through advertisements in the  
5  
6 local community and given an instruction booklet outlining what their participation would  
7  
8 require. Participants were also asked if they were happy to be blindfolded, to ensure visual  
9  
10 distraction did not occur, and were given the opportunity to ask questions about the research.  
11  
12 Those willing to continue were screened against the exclusion criteria and asked to provide  
13  
14 written, informed consent. Once participants had agreed to take part, they were allocated to one  
15  
16 of the three conditions. Participants allocated to the music group were asked to bring with them a  
17  
18 chosen piece of music, and were reassured that their musical taste was not under examination  
19  
20 and they could chose any music that they felt they would enjoy listening to.  
21  
22  
23  
24

25  
26  
27 Following recruitment and consent, participants were asked to complete all  
28  
29 questionnaire/NRS assessment measures to provide baseline scores. From time of consent to the  
30  
31 conclusion of the experiment, all further interactions with participants were scripted to minimise  
32  
33 any influence of the presence of the experimenter and to ensure that the repeated cold pressor  
34  
35 tests (at weeks 1 and week 2) were comparable (following Saab et al., 1993). Firstly, a week 1  
36  
37 baseline trial was completed. Participants were seated next to the cold pressor bath, asked to  
38  
39 remove any jewellery from their right arm, they were then blindfolded and the  
40  
41 sphygmomanometer attached to their left arm. Participants were asked to submerge their right  
42  
43 hand in the water and to give a verbal indication of when they first felt pain, and pain tolerance  
44  
45 was monitored by stopwatch. Immediately on withdrawal of the hand from the water,  
46  
47 participants' completed the NRS measures (excepting the NRS-CPI baseline assessment), STAI,  
48  
49 SF-MPQ and their HR was recorded. The participant's hand was then allowed to return to  
50  
51 baseline temperature and recovery time was provided.  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Familiarity and Pain Management

12

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

The cold pressor test was next repeated as a week 1 experimental trial. According to group allocation, participants listened to music, used progressive muscle relaxation or had no distraction during the course of their experimental trial. Pre- and post-test measures were taken as before. After completion of the experimental cold pressor test, participants in the music listening and PMR groups were asked to listen/practice their distraction every day (1-6) and were provided with a CD of the PMR where appropriate. Participants were asked to focus on their music/PMR and to refrain from doing other tasks whilst listening to their music or PMR CD during the practice days. From days 1-6 participants tracked their levels of familiarity with their distraction, daily pain ratings, and their preference for the music/PMR through a personal information booklet. Practice was considered to be completed if participants had listened to their chosen musical selection or the full PMR CD (approx. 8mins). Control group participants completed a short reflection activity, diarising the content of their day in the information booklet.

Participants returned at day 7 (week 2), again completing two cold pressor tests - at baseline and with distraction as appropriate to their grouping. The same measures were taken as at Week 1. At the completion of the study, all participants were debriefed and asked to give feedback regarding the experiment.

*Ethics*

This research met all standards of the British Psychological Society and the University of Buckingham granted ethics approval. Participants were informed that data would remain anonymous, and they could withdraw from the study at any stage, with data being discarded

## Familiarity and Pain Management

13

1  
2  
3 immediately. The cold pressor test has been found to cause short-term laboratory-induced pain  
4  
5 (Saab et al., 1993), with no lasting effects, and no long-term damage, where exclusion criteria  
6  
7 (listed above) are upheld. Participants were told prior to starting the experiment that they could  
8  
9 remove their arm at any stage during the test. Participants were asked to remove their arm if they  
10  
11 remained in the cold pressor for the full four minutes.  
12  
13  
14  
15  
16

*Statistics and Analysis*

17  
18  
19 The primary measure outcomes were Tolerance and Threshold (measured in seconds).  
20  
21 Secondary measure outcomes were Anxiety, Pain Perception Ratings (NRS-PP), Total Pain  
22  
23 Score (TPS; obtained from the SF-MPQ), Self-efficacy (obtained through the GSE and NRS-  
24  
25 EOD) and Heart Rate, used as physiological measure of pain impact. Statistical analysis was  
26  
27 carried out using SPSS v. 20.  
28  
29  
30  
31  
32  
33

34 The following analyses were computed to investigate between group differences at  
35  
36 baseline: a one-way ANOVA (3 levels; music, PMR and control) on age; and chi-square  
37  
38 analyses on the outcome measures of Level Of Education and Gender. Paired samples *t*-tests  
39  
40 were undertaken for music and PMR groups, comparing preference and familiarity at week 1 and  
41  
42 week 2.  
43  
44  
45  
46  
47

48 To investigate the impact of familiarity and preference on between group performance  
49  
50 the following were computed: repeated-measures ANOVAs were computed to determine the  
51  
52 impact of Group, Week Of Testing and Type Of Testing on all outcome variables. To assess  
53  
54 Heart Rate, an ANCOVA was computed, with preliminary baseline HR used as a covariate.  
55  
56  
57  
58  
59  
60

## Results

### *Internal Consistency, Validity, Normality*

Cronbach's alpha was used to assess internal consistency, with results demonstrating high internal consistency in all standardised questionnaire outcome measures (GSE  $\alpha = .94$ ; SF-MPQ  $\alpha = .87$ ; STAI  $\alpha = .88$ ). These results were within the recommended ranges for the standardised questionnaires: GSE,  $\alpha = .76 - .9$  (Schwarzer & Jerusalem, 1995); SF-MPQ  $\alpha = .73 - .89$  (Burckhardt & Bielle, 1994); STAI  $\alpha = .85 - .95$  (Tanaka-Matsumi & Kameoka, 1986). To assess normality of results, outcome measures were assessed using Kolmogorov-Smirnov test. As results attained significance, Greenhouse-Geisser adjustments were used in all further analyses.

### *Between-Group Comparability*

To assess comparability between Groups (Control, Music or PMR) at baseline, one-way ANOVAs were used. All groups were comparable in age ( $F(2, 69) = .48, p = .62$ ) and current pain intensity (NRS-CPI;  $F(2, 69) = 1.71, p = .19$ ) at baseline. Results from chi-squared analyses also showed that Groups were comparable in gender distribution ( $\chi^2 = 3.21, df = 2, p = .2$ ) and educational levels ( $\chi^2 = 3.21, df = 2, p = .2$ ).

### *Familiarity and Preference*

Mean familiarity significantly increased from week 1 to week 2 for both Music ( $t(23) = -5.3, p < .001$ ) and PMR ( $t(22) = -41.1, p < .001$ ), as shown in Table 1. Mean preference showed the same pattern, increasing by week 2 for both Music ( $t(23) = -4.1, p < .001$ ) and PMR ( $t(22) = -42.4, p < .001$ ).

## Familiarity and Pain Management

Insert Table 1: *Mean Outcomes of Familiarity and Preference for Music and PMR Distractors.*

*Primary Outcome Measures*

*Pain Tolerance.* A significant interaction between Type of Trial and Group was found, with experimental trials inducing significantly greater pain tolerance than baseline trials ( $F(2,67)=9.1, p < .0001, \eta^2=.23$ ), and with the music ( $p < .0001$ ) and PMR ( $p < .001$ ) groups showing significantly greater improvements over the control group (see Figure 1). A significant main effect of Type of Trial was found ( $F(1, 67) = 38.77, p < .001, \eta^2=.37$ ), with experimental trials improving pain tolerance for all participants. Similarly, a significant main effect of Week Of Testing was found ( $F(1,67)= 12.36, p < .001, \eta^2= .16$ ) with mean tolerance at week 2 significantly higher than mean tolerance at week 1. Descriptive Statistics for all outcome measures are contained in Table 2.

*Insert Figure 1:* The influence of week of testing and type of trial on pain tolerance

*Pain Threshold.* There was a significant interaction between Type of Trial and Group ( $F(2, 67) = 4.13, p < .05, \eta^2= .11$ ). Post-hoc tests revealed that the PMR significantly enhanced pain threshold over control ( $p < .05$ ), but there was no benefit of music over control or difference between PMR and Music listening (see Figure 2). A significant main effect of Week of Testing was found ( $F(1,67)= 11.14, p < .001, \eta^2=.14$ ), with mean threshold enhanced at week 2 over week 1. Experimental trials also enhanced pain threshold over baseline trials ( $F(1,67) = 27.71, p < .0001, \eta^2= .13$ ), with a significant main effect of Group allocation ( $F(2,67) = 3.63, p < .05, \eta^2= .10$ ).

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Insert *Figure 2*: The influence of week of testing and type of trial on pain threshold

Insert Table 2: *Mean and Standard Deviations for Week of Testing (week 1 vs. Week 2) and Type of Trial (Baseline and Experimental), for Control, Music and PMR Groups, on Outcome Variables.*

### *Secondary Outcome Measures*

*Anxiety.* A significant interaction was found between Week of Testing and Type of Trial ( $F(1, 67) = 8.98, p < .005, \eta^2 = .12$ ), with mean anxiety for all participants showing the greatest reduction after the experimental trial in week 2. Results also showed that there was a main effect of Group ( $F(1, 67) = 4.16, p < .05, \eta^2 = .11$ ), with the PMR group showing significantly lower anxiety ( $p < .05$ ) than the control group, and a trend towards an advantage for the music group over control ( $p = .06$ ), but with no difference between the music and PMR groups. No further main effects were shown.

*Pain Perception Rating (NRS-PP).* Though no overall main effect of Group was found ( $F(1, 67) = .455, p = .64, \eta^2 = .01$ ), an interaction was found between Week of Testing and Group ( $F(2, 67) = 11.40, p < .001, \eta^2 = .25$ ) and between Type of Trial and Group ( $F(1, 67) = 4.92, p < .01, \eta^2 = .13$ ) with experimental groups broadly showing greater reductions than controls in Week 2 in experimental trials (3-way interaction: Group, Week of Testing, Type of Trial: ( $F(2, 67) = 3.34, p < .05, \eta^2 = .09$ )). Week of Testing significantly influenced Pain Perception Ratings, with scores at Week 1 significantly higher than those at week 2 ( $F(1, 67) = 10.76, p < .01, \eta^2 = .01$ ).

## Familiarity and Pain Management

A main effect of Type of Trial was also shown ( $F(1, 67)=57.91, p < .001, \eta^2=.24$ ), with experimental trials reducing pain perception in comparison to baseline trials.

*Total Pain Score (TPS)*. No significant main effect of Week of Testing was found, however there was a significant interaction between Week of Testing and Group ( $F(2,67) = 10.53, p < .0001, \eta^2=.24$ ), with both the music and PMR groups showing reduced pain scores at week 2 in comparison to the control group. A significant interaction was also found between Week of Testing and Type of Trial ( $F(1, 67) = 17.02, p < .0001, \eta^2=.20$ ), with experimental trials in Week 2 showing the greatest Total Pain Score reductions. No further main effects were found.

*Self-efficacy*. No significant main effect of Week Of Testing was found ( $p=.52$ ), however a significant main effect of Group was found ( $F(2,67) = 7.83, p < .001, \eta^2=.19$ ), with post-hoc tests revealing that self-efficacy scores when listening to music were significantly greater than those of the control group ( $p < .001$ ), as were those of the PMR group ( $p < .05$ ). Music and PMR showed different patterns, however, with a trend towards the greatest self-efficacy for the music group shown in week 1 and for the PMR group shown after practice and familiarity with the technique, at week 2 ( $F(2, 67) = 2.65, p = .078, \eta^2=.07$ ). A significant main effect of Type Of Trial was found ( $F(1,67)= 49.5, p < .001, \eta^2=.06$ ), with mean self-efficacy scores increasing during experimental trials compared to baseline trials, however no main effect of Group was found ( $p=.25$ ; See Figure 3). No further interaction effects were shown.

Insert *Figure 3*: The effect of Week of Testing on Self-efficacy

## Familiarity and Pain Management

18

1  
2  
3  
4 *Heart Rate.* There was no significant impact of baseline HR on subsequent HR during any  
5  
6 further trials, since the covariate was found to be non-significant,  $p = .11$ . No significant main  
7  
8 effect of Week of Testing was found ( $p = .51$ ), however a significant main effect of Group was  
9  
10 found ( $F(2,66) = 3.3, p < .05, \eta^2 = .09$ ). Post-hoc analysis revealed a significant difference  
11  
12 between PMR group HR and control HR ( $p < .05$ ), with PMR HR decreasing over time and  
13  
14 control HR increasing over time. No further main effects or interactions were found.  
15  
16  
17  
18  
19

**Discussion**

20  
21  
22 This study aimed to investigate whether active and passive distractors had differential  
23  
24 effects in reducing individuals' perceptions of pain and associated symptoms, in the context of  
25  
26 increased familiarity with and preference for the distractors. Results showed that asking  
27  
28 experimental participants to practice PMR or to listen to their chosen music daily did  
29  
30 significantly enhance participants' preference for and familiarity with the distraction techniques.  
31  
32 Both groups showed comparable increases in liking and familiarity. Asking participants to  
33  
34 familiarise themselves with their pain management technique was beneficial in enhancing pain  
35  
36 control, with PMR delaying the onset of pain by increasing the time before participants first  
37  
38 reported pain (pain threshold). Similarly, both music and PMR enabled participants to tolerate  
39  
40 pain for longer than when exposed to pain with no distraction.  
41  
42  
43  
44  
45  
46  
47  
48

49 PMR was privileged over music in its ability to reduce anxiety, with anxiety at its lowest  
50  
51 at week two after participants had increased familiarity with the pain management techniques.  
52  
53 Music also reduced anxiety, but to a lesser extent than PMR, suggesting that both active and  
54  
55 passive distractors have a positive anxiolytic effect in the context of laboratory-induced pain.  
56  
57  
58  
59  
60

## Familiarity and Pain Management

19

1  
2  
3 Both Music and PMR enhanced self-efficacy over no pain management intervention, however  
4  
5 familiarity with music listening decreased self-efficacy by week two, whereas familiarity with  
6  
7 PMR increased self-efficacy, showing that practice was beneficial for PMR but not for preferred  
8  
9 music. Pain perception was also modestly influenced by distraction, with both PMR and Music  
10  
11 groups showing decreased pain perception ratings in the second week of testing, when  
12  
13 employing their active or passive intervention. Total Pain score, similarly showed a similar  
14  
15 pattern, with interventional group participants reporting reduced pain at week two in comparison  
16  
17 with the control group. The role of distraction in pain management was also modelled  
18  
19 physiologically, specifically for the PMR group, with Heart Rate reduced when the active  
20  
21 distraction was employed, whereas the control group showed an increase in Heart Rate. Overall,  
22  
23 these results demonstrated that active and passive distractors did influence experimentally-  
24  
25 induced pain in differing ways. PMR reduced anxiety, pain perception, total pain score and HR,  
26  
27 and increased pain tolerance and pain threshold. Importantly, repeated exposure to PMR  
28  
29 enhanced locus of control by increasing participants' self-efficacy. Music also reduced anxiety,  
30  
31 but to a lesser extent, and did not impact physiologically on Heart Rate. However, music  
32  
33 listening did enhance pain tolerance, pain threshold, reduce pain perception and total pain score.  
34  
35 Music, as an already familiar preferred distractor, did not enhance self-efficacy after repeated  
36  
37 exposure despite preference for the music and familiarity with the music increasing after regular  
38  
39 listening.  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50

51 Results demonstrated that listening to music and practicing PMR each day for a week,  
52  
53 did increase individuals' preference for and familiarity with both distraction approaches. These  
54  
55 results are concordant with previous research; increased exposure to distractors results in  
56  
57  
58  
59  
60

## Familiarity and Pain Management

20

1  
2  
3 improved ratings of familiarity (Schellenberg, 2008). It is possible therefore, that the short-term  
4  
5 longitudinal profile of this research served to enhance the arousal potential of music and PMR in  
6  
7 line with Berlyne's (1971) inverted-U theory, with the arousal potential being of significant  
8  
9 magnitude to affect psychological and (for PMR) physiological markers of pain. However, in  
10  
11 some aspects, music was less effective than PMR, and this requires further explanation. Heyduk  
12  
13 (1975) proposed the optimal complexity model, which suggested that individuals will prefer  
14  
15 stimuli fitting with their perception of optimal complexity, rather than stimuli that they consider  
16  
17 to be too complex, or not complex enough. Repeated exposure to distractors increases  
18  
19 familiarity, which in turn potentially decreases the distractor's perceived complexity. This could  
20  
21 be extended to explain the findings in relation to anxiety modulation, self-efficacy, and Heart  
22  
23 Rate, in which music listening was less effective: results suggested that familiarity and  
24  
25 preference were reaching optimum for preferred music listening in week one and for the  
26  
27 unfamiliar PMR in week two. As the music was self-selected by participants, this may already  
28  
29 have been at or near the peak of optimum arousal, therefore with repeated hearings its arousal  
30  
31 potential and efficacy as a distractor waned. Whereas PMR's arousal potential improved with  
32  
33 regular practice by week two. Further research could broaden the length of longitudinal follow-  
34  
35 up to firmly establish at which point, if any, perceived complexity of active and passive  
36  
37 distraction peaks and then wanes, therefore reducing its efficacy as an analgesic. Whilst these  
38  
39 results would suggest that efficacy of music and PMR distractors increase over a week-long trial,  
40  
41 long term effects need to be tested, in order to further apply to a long-term clinical population of  
42  
43 chronic pain sufferers.  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Familiarity and Pain Management

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Increased familiarity and preference for music and PMR, through listening and practice were found to delay pain threshold and increase pain tolerance (though to a lesser extent). With regard to PMR, Persson et al (2008) state that for the intervention to work effectively, regular practice needs to be undertaken. Repetitive practice of PMR allows the patient to quickly and effectively learn to identify the difference between muscle tension and muscle release, therefore enabling them to acquire perceptual calmness and maintain muscle relaxation (Persson et al, 2008). Concerning music, Mitchell et al (2006) found greater efficacy for the use of familiar music for reducing perceptions of pain, than for unfamiliar music. Therefore, as music familiarity was rated greater during week two testing than week one, it is appropriate that pain threshold and tolerance increased over time. With increased familiarity with music, emotional engagement is also found to rise (Silva-Pereira et al, 2011), this shown to be influential in the enhanced efficacy of emotionally-salient music as a reliever of pain (Villarreal et al, 2012). The different findings in self-efficacy between music and PMR could potentially be explained by the emotional valence of the distractor. Music, as a preferred selection, is likely to already hold emotional relevance, whereas PMR may develop emotional salience with increasing practice and recognition of psychophysiological benefits.

Such emotional salience is potentially both positive and negative. Though some research has shown positive effects of preferred (emotionally-salient) music (e.g. Mitchell & MacDonald, 2006), pain theory suggests emotional processing of distractors could be detrimental. Leventhal et al (1983) proposed the parallel process theory of pain, which consists of two competing components; the informational and the emotional mechanisms. The informational module encompasses sensory facets of the physical stimulus, whereas the emotional module comprises

## Familiarity and Pain Management

22

1  
2  
3 feelings such as fear and anxiety. According to Leventhal and colleagues, an individual will feel  
4 pain proportional to the amount of attention placed on the emotional component of the stimulus,  
5 rather than the informational component; if something is processed emotionally, it will lead to  
6 greater perceptions of pain (Broucek & Bartholomew, 1993). Broucek and Bartholomew (1993)  
7 state that the role of distraction is to help the individual to ignore the affective response to the  
8 painful stimulus. Due to the highly emotional aspect of music, using this technique may fail to  
9 allow the individual to ignore the affective component of the painful stimulus, due to this system  
10 being highly activated whilst listening to music. It could be that PMR has a greater effect on the  
11 informational component of the painful stimulus, and therefore, for individuals in the PMR  
12 group, pain was not processed emotionally, leading to reduced perceptions of pain. Further  
13 research could consider the emotional aspect of distractors on individual's perceptions of pain,  
14 and consider whether there is an optimum level of emotional engagement needed in order for the  
15 greatest efficacy of distractors to be obtained. Research needs to consider the impact that  
16 emotional engagement has over both music and PMR, and the influence this has on perception of  
17 pain.  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40

41 The anxiolytic properties of both music listening and PMR found in this study have been  
42 demonstrated in previous research. Relaxation techniques have commonly been found to reduce  
43 individuals' levels of anxiety, with distraction/relaxation procedures easily grasped, undertaken  
44 and managed by patients (Seers & Carroll 2001; Kristine et al, 2006). According to Wong et al  
45 (2010), during relaxation, the individual feels that they can exercise control over the painful  
46 stimulus, resulting in threat reduction, and therefore, reducing anxiety levels. Leubbert et al.  
47 (2001) suggest that increased anxiety results in an increase in muscle tension, which in turn leads  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Familiarity and Pain Management

1  
2  
3  
4 to an increased perception of pain. Therefore, through reducing muscle tension and anxiety,  
5  
6 perceptions of pain should also be reduced. The fact that PMR had a stronger anxiolytic effect  
7  
8 may be due to the active element of the technique allowing the individual to feel that they are  
9  
10 instrumental in controlling their perceptions of pain. This concurs with the findings regarding  
11  
12 self-efficacy, where regular practice of PMR enhanced locus of control to a greater extent than  
13  
14 music listening. According to Roykulcharoen and Good (2004), distraction/relaxation can be  
15  
16 implemented as a coping strategy that can enable the individual to feel they can govern the pain;  
17  
18 thus increasing self-efficacy. Zusman (2005) states that the valuable effects of enhancing self-  
19  
20 efficacy of chronic pain patients in a clinical setting are commonly noted in current literature,  
21  
22 with findings consistently demonstrating that increasing levels of self-efficacy reduce  
23  
24 individual's levels of pain (Keefe et al, 2004). Mastery experiences, which are gained through  
25  
26 performance accomplishments, have been found to have the greatest influence on creating and  
27  
28 reinforcing individuals' self-efficacy (Turk & Okifuji, 2002). In fitting with this theory of  
29  
30 mastery experiences, as individuals practiced their PMR each day, they may have felt as though  
31  
32 they were accomplishing set tasks, thus increasing their levels of self-efficacy. Through daily  
33  
34 practice, individuals were likely to experience high levels of performance accomplishments.  
35  
36 Increase in preference and familiarity for music was found to reduce levels of self-efficacy, and  
37  
38 this may be due to the passive task leaving individuals lacking in mastery experiences. Music  
39  
40 listening facilitates reduced anxiety as participants have control in selecting their own distractor,  
41  
42 but as it is already familiar and requires no additional learning and is therefore passive, the  
43  
44 additional activity required by It is possible that future research could allow participants to self-  
45  
46 select music for individually perceived biopsychosocial benefits as appropriate to their daily  
47  
48 needs, therefore enhancing the active component of music and enabling familiar music to  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Familiarity and Pain Management

24

1  
2  
3  
4 become more active in its distractive properties. In the same way, it may be argued that the  
5  
6 multi-dimensional nature of music and music listening processes may be conferring additional  
7  
8 benefits upon the listener, in a more active way than may originally appear. Future research  
9  
10 could aim to investigate the extent to which participants are immersing themselves in potentially  
11  
12 active components of music listening, such as analysis of the instrumentation, recognition of  
13  
14 compositional patterns, and communication through lyrics. Think-aloud protocols could be used  
15  
16 to assess the extent to which preferred music may be initiating enhanced self-efficacy, though to  
17  
18 a lesser extent than PMR. in its psychological impact. Self-efficacy is a variable that should not  
19  
20 be underestimated in pain management as heightened beliefs of self-efficacy are directly related  
21  
22 to patients' willingness to continue with pain management, regardless of potential difficulties  
23  
24 that they may face (Turk, 2004).  
25  
26  
27  
28  
29  
30  
31

32 The current results demonstrate that PMR has a positive effect on reducing individuals'  
33  
34 heart rate, whereas music listening does not. This negative finding for music does not fit with  
35  
36 previous research, such as that by Mimi et al (2005), who found individuals who listened to  
37  
38 relaxing music had lower heart rates compared to controls. Similarly, research by Tse, Chan &  
39  
40 Benzie (2005) and Cadigan et al (2001) found that music listening regulated or reduced heart rate  
41  
42 intra-operatively and post-operatively. The ability to lower heart rate is important in a pain  
43  
44 management context: Reeves and Shapiro (1983) found that individuals who were able to lower  
45  
46 their heart rate demonstrated lower levels of pain perceptions during a cold pressor task. Yet  
47  
48 more recent suggested that an increase in heart rate can enhance pain tolerance through  
49  
50 activating fight or flight responding, for example when people swear in response to pain (see  
51  
52 Stephens & Umland, 2011). As music enhanced pain tolerance, it may be that the increase in  
53  
54  
55  
56  
57  
58  
59  
60

## Familiarity and Pain Management

25

1  
2  
3 heart rate was positive for these participants. However, interestingly, the opposite effect was  
4  
5 modelled with PMR and PMR showed enhancements in pain threshold in addition to pain  
6  
7 tolerance. This may suggest that reducing heart rate (e.g. through active distraction) could have  
8  
9 greater concomitant analgesic effects than increasing heart rate (e.g. through passive distraction).  
10  
11 Future research could target Heart Rate as a primary outcome measure to assess this. has  
12  
13 demonstrated that However additional research could explain why Heart Rate was elevated in  
14  
15 response to music listening in this study: Kenntner-Mabiala et al (2007) found that music of fast,  
16  
17 moderate or slow tempi affected pain perception in different ways. Music with a fast tempo  
18  
19 increased Heart Rate, respiration rate and arousal, therefore its regulatory and analgesic effect  
20  
21 was poor. It is possible that in the current study, music chosen by participants was variable in  
22  
23 tempo and chosen for a variety of different reasons, for example for its uplifting properties or  
24  
25 high emotional significance, resulting in a greater affective component and therefore, physical  
26  
27 components of the painful experience, such as heart rate, were not impacted. Further research  
28  
29 could investigate requesting that participants provide music of different tempi to investigate the  
30  
31 interruptive function of musical constructs on physiological markers of arousal and pain.  
32  
33  
34  
35  
36  
37  
38  
39  
40

41 The current research is limited by its inclusion of healthy community-dwelling  
42  
43 volunteers, and it would be useful to extend the trial to chronic pain sufferers who have been  
44  
45 recommended relaxation/distraction techniques as part of a pain management programme. An  
46  
47 increased sample size and clinical population would further establish the efficacy of active and  
48  
49 passive approaches recommended by pain management services. The duration of this study was  
50  
51 limited to one week, and whilst promising results were found, its applicability to chronic pain is  
52  
53 limited by its short-term longitudinal profile. Chronic pain is, by definition considered to be  
54  
55  
56  
57  
58  
59  
60

## Familiarity and Pain Management

26

1  
2  
3  
4 benign or progressive pain extending beyond 12 weeks, therefore it is essential to assess whether  
5  
6 distractors have a lasting impact over longer time periods (The British Pain Society, 2012). In  
7  
8 addition, this experimental study showed that all participants improved in psychological markers  
9  
10 of pain between week one and week two of testing. It is likely that because participants have  
11  
12 already undertaken the cold pressor test during week one, their previous encounter with the  
13  
14 painful stimulus may have had a natural effect on reducing anxiety and pain perception, therefore  
15  
16 memory for pain is contributing to these results. It has been noted that pain-inhibitory systems  
17  
18 are influenced by memory processes (Flor, 2003), therefore research could further consider the  
19  
20 relationship between memory and inhibition of pain, and question whether prior exposure to a  
21  
22 painful stimuli causing the participant to effectively cope with pain may in some instances have a  
23  
24 positive effect.  
25  
26  
27  
28  
29  
30  
31

32 This research intended to consider the efficacy of active and passive distractors on  
33  
34 individuals' perceptions of pain. Overall findings would suggest that active distractors, such as  
35  
36 PMR, are more effective at reducing individuals' perceptions of pain than passive distractors  
37  
38 (following Villarreal et al, 2012). These results reflect the concepts of multiple resource theory  
39  
40 of attention (Wickens, 2002). Wickens (1984) argued that active and passive distractors are more  
41  
42 or less effective as a result of the number of multi-sensory attentional resources that they occupy.  
43  
44 Distractors that work best are those that saturate a number of attentional stores (e.g. auditory and  
45  
46 spatial) as in the case of active distraction PMR, and to a lesser extent for passive distraction  
47  
48 through music. Current literature suggests that a multidimensional approach towards pain  
49  
50 management appears to have the greatest and most positive efficacy in reducing perceptions of  
51  
52 pain (Turk & Okifuji, 2002). The positive effect of both music and PMR suggest that it would be  
53  
54  
55  
56  
57  
58  
59  
60

Familiarity and Pain Management

appropriate to combine both active and passive distractors, as a holistic tool kit for pain management.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review

## References

- Bandura, A. (1997). *Self-efficacy: the exercise of control*. San Francisco: WH Freeman and Co.
- Bandura, A. (2001). Social cognitive theory: An agentic perspective. *Annual review of psychology*, 52(1), 1-26.
- Berlyne, D. E. (1971). *Aesthetics and psychobiology*. New York: Appleton-Century-Crofts.
- Börsbo, B., Gerdle, B., & Peolsson, M. (2010). Impact of the interaction between self-efficacy, symptoms and catastrophising on disability, quality of life and health in with chronic pain patients. *Disability & Rehabilitation*, 32(17), 1387-1396.  
doi:10.3109/09638280903419269
- Broucek, M. W., & Bartholomew, J. B. (1993). The effects of relaxation with a warning cue on pain tolerance. *Journal Of Sport Behavior*, 16(4), 239.
- Burckhardt, C. S., & Bjelle, A. (1994). A Swedish version of the short-form McGill Pain Questionnaire. *Scandinavian journal of rheumatology*, 23(2), 77-81.
- Cadigan, M. E., Haldeman, S. M., McNamara, M. E., Noyes, D. A., Spadafora, M. A., & Carroll, D. L. (2001). The effects of music on cardiac patients on bed rest. *Progress in Cardiovascular Nursing*, 16(1), 5-13.
- Cepeda, M. S., Carr, D.B., Lau, J. & Alvarex, H. (2010). Music for pain relief (Review). *The Cochrane Collaboration*. Wiley
- Dahlquist, L. M., McKenna, K. D., Jones, K. K., Dillinger, L., Weiss, K. E., & Ackerman, C. (2007). Active and passive distraction using a head-mounted display helmet: Effects on cold pressor pain in children. *Health Psychology*, 26(6), 794-801. doi:10.1037/0278-6133.26.6.794

## Familiarity and Pain Management

29

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Duschek, S. S., Schwarzkopf, W. W., & Schandry, R. R. (2008). Increased pain sensitivity in low blood pressure. *Journal Of Psychophysiology*, 22(1), 20-27. doi:10.1027/0269-

8803.22.1.20

Eccleston, C., Morley, S., Williams, A., Yorke, L., & Mastroiannopoulou, K. (2002). Systematic review of randomised controlled trials of psychological therapy for chronic pain in children and adolescents, with a subset meta-analysis of pain relief. *Pain*, 99, 157–165.

Eccleston, C., Williams, A. C. D. C., & Morley, S. (2009). Psychological therapies for the management of chronic pain (excluding headache) in adults. *Cochrane Database of Systematic Reviews*, 2. DOI: 10.1002/14651858.CD007407.pub2.

Edwards, R.R., Campbell, C., Jamison, R.N. & Wiech, K. (2009). The neurobiological underpinnings of coping with pain. *Current Directions In Psychological Science*, 18, 237-241.

Edwards, R. R., Doleys, D. M., Fillingim, R. B., & Lowery, D. (2001). Ethnic differences in pain tolerance: clinical implications in a chronic pain population. *Psychosomatic medicine*, 63(2), 316-323.

Engel, G. L. (1977). The need for a new medical model: A challenge for biomedicine. *Science*, 196, 129-135.

Finlay, K. A. (in press). Familiarity with music in post-operative clinical care. Chapter 8 in Daynes, H. & King, E. (Eds.) *Music and Familiarity*.

Flor, H. (2003). Cortical reorganisation and chronic pain: Implication for rehabilitation. *Journal of Rehabilitation Medicine*, 41, 66-72.

## Familiarity and Pain Management

30

- 1  
2  
3  
4 Forys, K. L., & Dahlquist, L. M. (2007). The Influence of Preferred Coping Style and Cognitive  
5 Strategy on Laboratory-Induced Pain. *Health Psychology, 26*(1), 22–29. doi:10.1037/0278-  
6 6133.26.1.22  
7  
8  
9  
10 Gustavsson, C., & von Koch, L. (2006). Applied Relaxation in the treatment of long-lasting neck  
11 pain: A randomized controlled pilot study. *Journal Of Rehabilitation Medicine, 38*(2), 100-  
12 107.  
13  
14  
15  
16  
17 Good, M., Stanton-Hicks, M., Grass, J. A., Anderson, G. C., Choi, C., Schoolmeesters, L. J., &  
18 Salman, A. (1999). Relief of postoperative pain with jaw relaxation, music and their  
19 combination. *Pain, 81*(1-2), 163–172.  
20  
21  
22  
23  
24 Hekmat, H. & Hertel, J. (1993) Pain attenuating effects of preferred versus non-preferred music  
25 interventions. *Psychology of Music, 21*(2), 163-173.  
26  
27  
28  
29 Henschke, N., Ostelo, R. W., van Tulder, M. W., Vlaeyen, J. W., Morley, S., Assendelft, W. J.,  
30 & Main, C. J. (2010). Behavioural treatment for chronic low-back pain. *Cochrane*  
31 *Database Syst Rev, 7*(7).  
32  
33  
34  
35  
36 Heyduk, R. G. (1975). Rated preference for music composition as it relates to complexity and  
37 exposure frequency. *Perception and Psychophysics, 17*, 84-91  
38  
39  
40  
41 Jacobson, E. (1938). *Progressive relaxation*. Chicago: University of Chicago Press.  
42  
43  
44 Jackson, T. D., Pope, L. E., Nagasaka, T. O., Fritch, A. L., Iezzi, T. Y., & Chen, H. G. (2005).  
45 The impact of threatening information about pain on coping and pain tolerance. *British*  
46 *Journal Of Health Psychology, 10*(3), 441-451.  
47  
48  
49  
50 Jensen, M.P., & Karoly, P. (2001) Self-report scales and procedures for assessing pain in adults,  
51 in Turk DC and Melzack R (eds.), *Handbook of Pain Assessment, 2nd edition*, New York:  
52 Guilford Press,  
53  
54  
55  
56  
57  
58  
59  
60

## Familiarity and Pain Management

31

- 1  
2  
3 Johnson, M. H., Breakwell, G., Douglas, W., & Humphries, S. (1998). The effects of imagery  
4 and sensory detection distractors on different measures of pain: How does distraction  
5 work? *British Journal of Clinical Psychology*, 37(2), 141–154. doi:10.1111/j.2044-  
6 8260.1998.tb01290.x  
7  
8  
9  
10  
11  
12 Katz, J., & Melzack, R. (2011). The McGill Pain Questionnaire: Development, psychometric  
13 properties, and usefulness of the long form, short form, and short form-2. In D. C. Turk &  
14 R. Melzack (Eds.), *Handbook of pain assessment* (3rd ed.). (pp. 45–66). New York, NY  
15 US: Guilford Press.  
16  
17  
18  
19  
20  
21  
22 Kenntner-Mabiala, R., Gorges, S., Alpers, G. W., Lehmann, A. C., & Pauli, P. (2007). Musically  
23 induced arousal affects pain perception in females but not in males: A psychophysiological  
24 examination. *Biological Psychology*, 75(1), 19–23.  
25  
26  
27  
28  
29 Koelsch, S. (2010). Towards a neural basis of music-evoked emotions. *Trends in cognitive*  
30 *sciences*, 14(3), 131–137. doi:10.1016/j.tics.2010.01.002  
31  
32  
33  
34 Kristine L., Kwekkeboom K. & Gretarsdottir E. (2006) Systematic review of relaxation  
35 intervention for pain. *Clinical Scholarship 3rd quarter*, 269–277.  
36  
37  
38  
39 Leubbert K., Dahme B. & Hasenbring M. (2001) The effectiveness of relaxation training in  
40 reducing treatment related symptoms and improving emotional adjustment in acute non-  
41 surgical cancer treatment: a meta-analytical review. *Psycho-Oncology* 10(6), 490–502.  
42  
43  
44  
45 Leventhal, H., Safer, M. A., & Panagis, D. M. (1983). The impact of communications on the  
46 self-regulation of health beliefs, decisions, and behavior. *Health Education &*  
47 *Behavior*, 10(1), 3-29.  
48  
49  
50  
51  
52  
53 MacLaren, J. E., & Cohen, L. L. (2005). A comparison of distraction strategies for venipuncture  
54 distress in children. *Journal of Pediatric Psychology*, 30, 387–396.  
55  
56  
57  
58  
59  
60

## Familiarity and Pain Management

32

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
- Marteau, T. M., & Bekker, H. (1992). The development of a six-item short-form of the state scale of the Spielberger State-Trait Anxiety Inventory (STAI). *The British journal of clinical psychology / the British Psychological Society*, 31 ( Pt 3), 301–306.
- Mehling, W.E., Hamel, K.A., Acree, M., Byl, N., Hecht, F.M. (2005) Randomized, controlled trial of breath therapy for patients with chronic low-back pain. *Alternative Therapies in Health and Medicine*, 11, 44-52
- Melzack, R. (1983) *Pain Measurement and Assessment*. Raven Publishing, pp 15-34.
- Mimi, M. Y, Chan, M. F. and Benzie, I. F. F (2005) the effect of music therapy in postoperative pain, heart rate, systolic blood pressure and analgesic use following nasal surgery, *Journal of Pain and Palliative Care Pharmacotherapy*, 19 (3), 21-29
- Mitchell, L. A., MacDonald, R. A. R., & Brodie, E. E. (2004). Temperature and the Cold Pressor Test. *Journal of Pain*, 5(4), 233–237.
- Mitchell, L. A., & MacDonald, R. A. R. (2006). An experimental investigation of the effects of preferred and relaxing music listening on pain perception. *Journal of music therapy*, 43(4), 295–316.
- Mitchell, L. A., MacDonald, R. A., & Brodie, E. E. (2006). A comparison of the effects of preferred music, arithmetic and humour on cold pressor pain. *European journal of Pain*, 10(4), 343-343.
- Mitchell, L. A., MacDonald, R. R., & Knussen, C. (2008). An investigation of the effects of music and art on pain perception. *Psychology Of Aesthetics, Creativity, And The Arts*, 2(3), 162-170.

1 Familiarity and Pain Management

2  
3 De Paula, A. A. D., de Carvalho, E. C., & dos Santos, C. B. (2002). The use of the “progressive  
4 muscle relaxation” technique for pain relief in gynecology and obstetrics. *Revista latino-*  
5 *americana de enfermagem*, 10(5), 654–659.  
6  
7

8  
9  
10  
11  
12 Pain Toolkit (2012). Retrieved from [http://www.paintoolkit.org/assets/downloads/Pain-Toolkit-](http://www.paintoolkit.org/assets/downloads/Pain-Toolkit-Booklet-Nov-2012.pdf)  
13 [Booklet-Nov-2012.pdf](http://www.paintoolkit.org/assets/downloads/Pain-Toolkit-Booklet-Nov-2012.pdf) on 1 May, 2013.  
14

15  
16  
17 Persson, A. L., Veenhuizen, H., Zachrison, L., & Gard, G. (2008). Relaxation as treatment for  
18 chronic musculoskeletal pain – a systematic review of randomised controlled studies.  
19  
20  
21  
22 *Physical Therapy Reviews*, 13(5), 355-365. doi:10.1179/174328808X356366  
23

24  
25 Pontinen, P.J. (1998) Reliability, validity, reproducibility of algometry in diagnosis of active and  
26 latent tender sports and trigger points, *Journal of Musculoskeletal Pain*, 6(1), 61-71  
27

28  
29 Reeves, J. L., & Shapiro, D. (1983). Heart-rate reactivity to cold pressor stress following  
30 biofeedback training. *Applied Psychophysiology and Biofeedback*, 8(1), 87-99.  
31

32  
33  
34 Roykulcharoen V. & Good M. (2004) Systematic relaxation to relieve postoperative pain.  
35  
36  
37 *Journal of Advanced Nursing* 48(2), 140–148.  
38

39 Saab, P. G., Llabre, M. M., Hurwitz, B. E., Schneiderman, N., Wohlgenuth, W., Durel, L. A.,  
40 Nagel, J. (1993). The cold pressor test: vascular and myocardial response patterns and their  
41 stability. *Psychophysiology*, 30(4), 366–373.  
42  
43  
44

45  
46 Schellenberg, E. (2008). Liking for happy- and sad-sounding music: Effects of  
47 exposure. *Cognition & Emotion*, 22(2), 218-237.  
48  
49

50  
51 Schaffer S.D. & Yucha C.B. (2004) Relaxation and pain management: the relaxation response  
52 can play a role in managing chronic and acute pain. *American Journal of Nursing*, 104(8),  
53 75–76, 78–79, 81–82.  
54  
55  
56  
57  
58  
59  
60

## Familiarity and Pain Management

34

- 1  
2  
3  
4 Schwarzer, R., & Jerusalem, M. (1995). Generalized Self-Efficacy scale. In J. Weinman,  
5  
6 S. Wright, & M. Johnston, *Measures in health psychology: A user's portfolio. Causal and*  
7  
8 *control beliefs* (pp. 35-37). Windsor, UK: NFER-NELSON.
- 9  
10 Seers K. & Carroll D. (2001) Relaxation techniques for acute pain management: a systematic  
11  
12 review. *Journal of Advanced Nursing* 27(3), 466–475.
- 13  
14  
15 Sharma, L., Cahue, S., Song, J., Hayes, K., Pai, Y. C., Dunlop, D. (2003). Physical functioning  
16  
17 over three years in knee osteoarthritis: role of psychosocial, local mechanical, and  
18  
19 neuromuscular factors. *Arthritis Rheum*, 48, 3359–3370.
- 20  
21  
22 Shiffrin, R. M., & Schneider, W. (1977). Controlled and automatic human information  
23  
24 processing II: perceptual learning, automatic attending and a general theory. *Psychology*  
25  
26 *Review*, 84, 127–90.
- 27  
28  
29 Silva-Pereira C, Teixeira J, Figueiredo P, Xavier J, Castro S, et al. (2011) The role of familiarity  
30  
31 in the emotional engagement with music: an fMRI study. Submitted.
- 32  
33  
34 Spielberger, C. D. (1983). *State-Trait Anxiety Inventory*. Palo Alto, CA: Consulting  
35  
36 Psychologists Press.
- 37  
38  
39 Stephens, R., & Umland, C. (2011). Swearing as a response to pain-effect of daily swearing  
40  
41 frequency. *The journal of pain: official journal of the American Pain Society*, 12(12),  
42  
43 1274–1281. doi:10.1016/j.jpain.2011.09.004
- 44  
45  
46 Tanaka-Matsumi, J. & Kameoka, V. A. (1986). Reliabilities and concurrent validities of popular  
47  
48 self-report measures of depression, anxiety and social desirability. *Journal of Consulting*  
49  
50 *and Clinical Psychology*, 54, 328-333
- 51  
52  
53 Taylor, J., & Deane, F. (2002). Development of a Short Form of the Test Anxiety Inventory  
54  
55 (TAI). *The Journal Of General Psychology*, 129(2), 127-136.
- 56  
57  
58  
59  
60

## Familiarity and Pain Management

- 1  
2  
3  
4 Thayer, J. F., Åhs, F., Fredrikson, M., Sollers, J. J. & Wager, T. D. (2011). Review: A meta-  
5  
6 analysis of heart rate variability and neuroimaging studies: Implications for heart rate  
7  
8 variability as a marker of stress and health. *Neuroscience and Biobehavioral Reviews*, 36,  
9  
10 747–756. doi:10.1016/j.neubiorev.2011.11.009  
11  
12  
13 The British Pain Society. (2012) *Pain Management Programmes*, accessed on 1<sup>st</sup> November  
14  
15 2012: [http://www.britishpainsociety.org/patient\\_pmp.htm](http://www.britishpainsociety.org/patient_pmp.htm)  
16  
17  
18 Tse, M. M., Chan, M. F., & Benzie, I. F. (2005). The effect of music therapy on postoperative  
19  
20 pain, heart rate, systolic blood pressures and analgesic use following nasal surgery. *J Pain*  
21  
22 *Palliat Care Pharmacother*, 19(3), 21–29.  
23  
24  
25 Turk, D. C. (2004). Understanding pain sufferers: the role of cognitive processes. *Spine*, 4, 1–7  
26  
27  
28 Turk, D.C. & Okifuji, A. (2002) Psychological factors in chronic pain: Evolution and revolution.  
29  
30 *Journal of Consulting and Clinical Psychology*, 70 (3), 678-690  
31  
32  
33 Verhoeven, K., Crombez, G., Eccleston, C., Van Ryckeghem, D. M. L., Morley, S., & Van  
34  
35 Damme, S. (2010). The role of motivation in distracting attention away from pain: An  
36  
37 experimental study. *Pain*, 149(2), 229–234. doi:10.1016/j.pain.2010.01.019  
38  
39  
40 Villarreal, E., Brattico, E., LenêVase, Østergaard, L., & Vuust, P. (2012). Superior Analgesic  
41  
42 Effect of an Active Distraction versus Pleasant Unfamiliar Sounds and Music: The  
43  
44 Influence of Emotion and Cognitive Style. *Plos ONE*, 7(1), 1-8.  
45  
46 doi:10.1371/journal.pone.0029397  
47  
48  
49 Wickens, C. D. "Processing Resources in Attention." *Varieties of attention* (1984): 63.  
50  
51  
52 Wickens, C. D. (2002). Multiple resources and performance prediction. *Theoretical Issues in*  
53  
54 *Ergonomic Science*, 3, 150 –177.  
55  
56  
57  
58  
59  
60

1 Familiarity and Pain Management 36

2  
3 Wickens, C. (2008). Multiple Resources and Mental Workload. *Human Factors: The Journal of*  
4 *the Human Factors and Ergonomics Society*, 50(3), 449–455

5  
6  
7  
8 Windich-Biermeier, A., Sjoberg, I., Dale, J. C., Eshelman, D., & Guzzetta, C. E. (2007). Effects  
9  
10 of Distraction on Pain, Fear, and Distress During Venous Port Access and Venipuncture in  
11  
12 Children and Adolescents With Cancer. *Journal of Pediatric Oncology Nursing*, 24(1), 8–  
13  
14 19. doi:10.1177/104345420629601


15  
16  
17 Woby, S., Roach, N., Urmston, M., Watson, P. (2007) The relation between cognitive factors  
18  
19 and levels of pain and disability in chronic low back pain patients presenting for  
20  
21 physiotherapy. *European Journal of Pain*, 11, 869–77.

22  
23  
24  
25 Wong, E., Chan, S., & Chair, S. (2010). Effectiveness of an educational intervention on levels of  
26  
27 pain, anxiety and self-efficacy for patients with musculoskeletal trauma. *Journal Of*  
28  
29 *Advanced Nursing*, 66(5), 1120-1131. doi:10.1111/j.1365-2648.2010.05273.x

30  
31  
32 Zelikovsky, N., Rodrigue, J. R., Gidycz, C. A., & Davis, M. A. (2000). Cognitive behavioral and  
33  
34 behavioral interventions help young children cope during a voiding cystourethrogram.  
35  
36 *Journal of Pediatric Psychology*, 25, 535–543.

37  
38  
39 Zusman, M. (2005). Cognitive-behavioural components of musculoskeletal physiotherapy: the  
40  
41 role of control. *Physical Therapy Reviews*, 10(2), 89-98.


Figure 1: The influence of week of testing and type of trial on pain tolerance



Peer Review


1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Figure 2: The influence of week of testing and type of trial on pain threshold



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Figure 3: The effect of Week of Testing on Self-efficacy



Peer Review

Table 1

	Familiarity		Preference	
	Week 1	Week 2	Week 1	Week 2
Music	6.8 ± 1.4	8.8 ± 1.0	7.3 ± 1.3	8.8 ± 1.1
PMR	0.6 ± 0.8	9.4 ± 0.7	1.1 ± 0.6	9.1 ± 0.7

For Peer Review

Table 2

		Control				Music				PMR				<i>p</i> -values	
		Week 1		Week 2		Week 1		Week 2		Week 1		Week 2		Week of	Type of
		B	E	B	E	B	E	B	E	B	E	B	E	Testing	Trial
Threshold	M	17.74	19.61	17.57	32.39	32.71	64.90	31.88	84.88	30.26	58.70	35.43	112.00	<.0001	<.0001†
	SD	11.75	17.10	16.91	65.75	55.66	88.60	53.75	96.76	48.35	73.83	47.13	97.06		
Tolerance	M	122.43	107.39	98.70	114.39	104.13	167.96	114.50	188.79	82.65	113.22	109.65	188.74		†
	SD	99.82	93.24	87.81	93.92	91.68	97.25	95.67	88.67	86.71	95.54	90.39	85.34		
Anxiety	M	20.22	21.04	19.22	18.96	17.71	18.25	18.63	17.17	18.17	17.43	18.61	16.61	<.05	<.05
	SD	3.47	4.27	3.55	3.21	3.14	2.49	4.16	4.29	3.75	4.07	3.06	2.71		
NRS-PP	M	5.13	5.43	5.07	4.30	5.75	4.54	6.71	4.46	5.89	6.52	5.91	3.74	†	
	SD	2.28	2.50	2.36	2.49	2.01	2.48	2.26	3.08	1.90	2.02	2.02	2.16		
Self-efficacy	M	5.39	5.96	5.70	6.13	4.33	7.92	4.33	8.29	3.91	5.57	4.52	7.70	<.01	
	SD	3.23	3.01	2.98	3.20	3.36	2.26	2.75	1.73	2.64	2.45	2.84	2.67		
HR	M	67.52	68.22	65.70	68.83	72.67	71.63	74.82	70.88	71.78	68.57	70.00	66.22	<.05	<.05
	SD	9.97	11.27	8.14	10.47	12.76	14.67	13.37	9.82	16.39	10.00	12.34	9.56		
TPS	M	7.65	8.30	6.22	5.09	5.71	5.42	9.00	6.13	8.09	9.52	8.26	6.39	†	

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

SD	4.89	5.46	3.25	3.07	4.54	4.18	8.82	6.07	4.03	5.18	5.60	4.39
----	------	------	------	------	------	------	------	------	------	------	------	------

---

For Peer Review

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Table 1: *Mean Outcomes of Familiarity and Preference for Music and PMR*

*Distractors. Results shown as mean ± standard deviation.*

Table 2: *Mean and Standard Deviations for Week of Testing (week 1 vs. Week 2) and*

*Type of Trial (Baseline and Experimental), for Control, Music and PMR Groups, on*

*Outcome Variables. Between-Groups differences are shown as \*  $p < .05$ ; \*\*  $p < .01$ ;*

*\*\*\*  $p < .001$ ; Interactions with Group are shown as †.*

For Peer Review