



Alterations in Circulating Adhesion Molecules in Acute Myocardial Infarction before and after Thrombolysis with Streptokinase

Hossein Saidi¹, Maryam Vakilian², Gholam Hosein Noori¹, Hamed-Basir Ghafouri^{3*}, Niloofar Abazarian³

¹Department of Emergency, Rasul-Akram Hospital, Iran University of Medical Sciences, Tehran, Iran

²Department of Emergency, Sina Hospital, Tehran University of Medical Sciences, Tehran, Iran

³Department of Emergency, 7th Tir Martyrs Hospital, Iran University of Medical Sciences, Tehran, Iran

ARTICLE INFO

Article Type:

Original Article

Article History:

Received: 9 June 2013

Accepted: 8 September 2013

Keywords:

Adhesion Molecules
Myocardial Infarction
Streptokinase

ABSTRACT

Introduction: The role of adhesion molecules in the development and progression of coronary atherosclerosis is inevitable. It is not clear yet whether these molecules increase or decrease in level after thrombolytic therapy. This study was designed to compare concentrations of soluble forms of adhesion molecules in patients with acute myocardial infarction before and after reperfusion by thrombolysis with streptokinase (SK).

Methods: In this study, in 40 patients with acute myocardial infarction who were admitted in our Emergency Department undergoing thrombolysis with SK, plasma concentrations of six adhesion molecules [soluble L-selectin, P-selectin, E-selectin, intercellular adhesion molecule-1 (ICAM-1), vascular cell adhesion molecule-1 (VCAM-1) and platelet endothelial cell adhesion molecule-1 (PECAM-1)] were measured by enzyme linked immunosorbent assay (ELISA), before and 3 hours after intervention.

Results: While soluble E-selectin and PECAM-1 concentrations did not differ within the 3 hours after interventions ($P > 0.05$), the level of P-selectin, L-selectin, ICAM-1, VCAM-1 were significantly reduced after thrombolysis with SK ($P < 0.05$).

Conclusion: Adhesion molecules which mediate the interactions in leukocyte endothelium vary in levels after reperfusion with SK. It was shown that 4 out of 6 adhesion molecules significantly reduced after thrombolysis with SK.

Introduction

Cardiovascular disease is the most prevalent and major cause of death world wide and burdens huge costs on the health care system.¹⁻⁴ Mortality due to Ischemic Heart Diseases (IHD) decreases by early treatment and long term survival in patients increases subsequently.⁵ Coronary artery reperfusion is accepted as a method to achieve reduction in size of myocardial infarction.⁶ Thrombolysis technique with streptokinase (SK) is a confirmed method for reopening of an occluded vessel in patients with acute myocardial infarction.⁶ Cellular adhesion molecules (such as: P-selectin, E-selectin, L-selectin, VCAM-1, PECAM-1, ICAM-1) have an important role to play in progression of atherosclerotic plaque.^{7,8}

Besides, those mononuclear cells which are responsible for release of matrix metalloproteinases are activated by adhesion molecules leading to plaque rupture and initiation of acute coronary syndrome. Consequently, it can be concluded that an intervention leads to decline in adhesion molecules level could postpone the process of atherosclerotic plaque formation or rupture. On the other hand, it has been shown that patients with high

levels of these adhesion molecules have poor prognosis in future.⁹ Some studies imply that adhesion molecules would decrease¹⁰ or increase¹¹ in level, after reperfusion with different types of thrombolytic drugs.

Here in our hospitals in Iran, the only cost-benefit and available thrombolytic drug for reperfusion of coronary arteries in patients with acute myocardial infarction is intravenous thrombolysis with streptokinase (SK), 1.5 million IU over 60 minutes. The aim of this study was to compare concentrations of six soluble forms of adhesion molecules in patients with acute myocardial infarction before and 3 hour after attempted reperfusion by thrombolysis with streptokinase (SK).

Materials and Methods

From January to June 2011, 40 patients admitted to the emergency department of Rasul-Akram Hospital with the diagnosis of acute myocardial infarction (based on a history of typical chest pain and electrocardiographic changes) and with no contraindication of streptokinase administration were included in this study. The patients were selected using a convenient sampling method.

*Corresponding author: Hamed-Basir Ghafouri, E-mail: hbgssl@yahoo.com

Blood samples from all 40 patients were taken prior to thrombolysis with SK, and then 3 hours after treatment. Plasma samples were extracted by centrifugation at 4°C, at 3000 rpm for 10 min. Samples were later stored at -80°C before measurement of plasma levels of the soluble forms sE-selectin, sL-selectin, sP-selectin, sICAM-1, sVCAM-1, and sPECAM-1 using commercially available enzyme-linked immunosorbent assays (ELISAs). Results were presented as mean±standard deviation (SD) for serum levels of adhesion molecules before and after thrombolysis with SK. The Statistical test used in this study was Paired-Sample *t*-Test. A *P* ≤ 0.05 was considered statistically significant. The SPSS 17 (SPSS Inc., Chicago, Illinois, USA) statistical software package was used for all calculations. Our study was approved by the ethics committee of Iran University of Medical Sciences (IUMS). Informed written consent was obtained from each patient or his or her guardian.

Results

All 40 patients who were included in the study were in the age range of 46 to 83 years old (65.90 ± 10.6 years). There were 30 male patients (75%) and 10 female patients (25%). The levels of 6 adhesion molecules were measured before and 3 hours after thrombolysis with streptokinase (1.5 million IU over 60 min). The results have been presented in Table 1.

Discussion

Adhesion molecules are considered to be indicators of activation of leukocytes, platelets and endothelial cells. It has been shown that in patients with acute coronary syndrome, these molecules would increase in atherosclerotic plaque rupturing, ischemic and reperfused areas.⁸ We found significantly decreased serum levels for P-Selectin after SK reperfusion. The concentration of L-Selectin was also reduced by 3 hours after SK lysis. VCAM-1 plays an important role in atherosclerosis plaque progression and even rupture of the plaque leading to thrombosis. VCAM-1 is related to the aggregation of leukocytes, on the other

hand, adherence of leukocytes to vascular endothelial cells causes endothelium dysfunction.¹² It has also been shown that plasma levels of VCAM-1 were significantly decreased. After thrombolysis in patient with acute myocardial ischemia, it has been shown that both soluble ICAM-1 and PECAM-1 increase in levels within the initial hours after reperfusion.^{13,14}

In the present study, we found PECAM-1 to be stable after reperfusion, but significant decline in ICAM-1 level was observed. Endothelial cells are the only source of E-selectin expression and an increase in E-selectin could reveal endothelial activity.¹⁵ Some studies have shown that this molecule rises in patients with myocardial ischemia and hypertension.¹⁶⁻²⁰ No significant differences were observed in the concentration of E-selectin within the 3 hours after intervention. In a 2001 study by Kerner *et al.*, the concentration of 6 adhesion molecules in 30 patients with acute myocardial infarction was measured and it was seen that E-selectin, ICAM-1, VCAM-1 and PECAM-1 had no significant difference in levels after streptokinase administration; however, L-selectin and P-selectin significantly decreased after thrombolysis with SK. These findings are not consistent with results of our study.⁶ In other separated studies by Postadzhiyan⁹, Squadrito,¹⁰ Parise,²¹ and Zeitler,²² it has been shown that adhesion molecules would decrease in levels after thrombolytic therapy. These findings have a relative conformity to our findings.

Conclusion

It seems that adhesion molecules vary in levels following reperfusion therapy with streptokinase. Overall, 4 out of 6 adhesion molecules showed significant decline in levels after reperfusion by SK thrombolytic therapy in acute myocardial infarction. The clinical importance of these results should be investigated in further studies. Besides, these results indicate that thrombolysis with streptokinase would decrease adhesion molecule levels and may lead to decline in atherosclerosis progress and recurrence of myocardial infarction.

Table 1. Serum levels of soluble adhesion molecules

	P-Selectin (ng/ml)	L-Selectin (ng/ml)	VCAM-1 (ng/ml)	ICAM-1 (ng/ml)	E-Selectin (ng/ml)	PECAM-1 (ng/ml)
Before reperfusion	127.15 ± 22.84	479.7 ± 148.91	1012.75 ± 402.9	446.35 ± 106.02	87.71 ± 6.13	67.95 ± 15.29
Three hours after reperfusion with SK	108 ± 16.43	389.6 ± 106.24	858 ± 322.9	348.45 ± 75.6	101.32 ± 7.95	67.1 ± 17.81
<i>P</i>	0.002	0.006	0.002	0.0001	0.117	0.681

Limitations

Blood samples of the patients were taken in only two time intervals; once before thrombolysis with SK, and then 3 hours after SK administration due to funding shortage.

Ethical issues: This study was reviewed and confirmed by the ethics committee of Iran University of Medical Sciences.

Conflict of interests: The authors declare no conflicts of interest.

References

- Xie J, Wu EQ, Zheng ZJ, Sullivan PW, Zhan L, Labarthe DR. Patient-Reported Health Status in Coronary Heart Disease in the United States Age, Sex, Racial, and Ethnic Differences. *Circulation* 2008;118:491-7.
- Lloyd-Jones D, Adams RJ, Brown TM, Carnethon M, Dai S, De Simone G, *et al.* Heart disease and stroke statistics-2010 update: A report from the American Heart Association. *Circulation* 121:e46-e215.
- Yaghoubi A, Safaie N, Azarfarin R, Alizadehasl A, Golzari SE. Evaluation of Cardiovascular Diseases and Their Risk Factors in Hospitalized Patients in East Azerbaijan Province, Northwest Iran: A Review of 18323 Cases. *J Teh Univ Heart Ctr* 2013;8:101-5.
- Farb A, Kolodgie FD, Jenkins M, Virmani R. Myocardial infarct extension during reperfusion after coronary artery occlusion: pathologic evidence. *J Am Coll Cardiol* 1993;21:1245-1253.
- Ma XL, Lefer DJ, Lefer AM, Rothlein R. Coronary endothelial and cardiac protective effects of a monoclonal antibody to intercellular adhesion molecule-1 in myocardial ischemia and reperfusion. *Circulation* 1992;86:937-946.
- Schleiffenbaum B, Spertini O, Tedder TF. Soluble L-selectin is present in human plasma at high levels and retains functional activity. *J Cell Biol* 1992;119:229-238.
- Haverslag R, Pasterkamp G, Hoefer IE. Targeting adhesion molecules in cardiovascular disorders. *Cardiovasc Hematol Disord Drug Targets* 2008;8:252-60.
- Blankenberg S, Barbaux S, Tiret L. Adhesion molecules and atherosclerosis. *Atherosclerosis* 2003;170:191.
- Postadzhiyan AS, Tzontcheva AV, Kehayov I, Finkov B. Circulating soluble adhesion molecules ICAM-1 and VCAM-1 and their association with clinical outcome, troponin T and C-reactive protein in patients with acute coronary syndromes. *Clin Biochem* 2008;41:126-33.
- Squadrito F, Saitta A, Altavilla D, Ioculano M, Canale P, Campo GM, *et al.* Thrombolytic therapy with urokinase reduces increased circulating endothelial adhesion molecules in acute myocardial infarction. *J Eur His Res* 1996; 45:14-9.
- Ramachandran A, Jha S, Lefer DJ. REVIEW paper: pathophysiology of myocardial reperfusion injury: the role of genetically engineered mouse models. *Vet Pathol* 2008;45:698-706.
- Lawson C, Wolf S. ICAM-1 signaling in endothelial cells. *Pharmacol Rep* 2009;61:22-32.
- Miwa K, Igawa A, Inoue H. Soluble E-selectin, ICAM-1 and VCAM-1 levels in systemic and coronary circulation in patients with variant angina. *Cardiovasc Res* 1997;36:37-44.
- Gurbel PA, Serebruany VL. Soluble vascular cell adhesion molecule-1 and E-selectin in patients with acute myocardial infarction treated with thrombolytic agents. *Am J Cardiol* 1998;81:772-5.
- Hwang SJ, Ballantyne CM, Sharrett AR, Smith LC, Davis CE, Gotto AM Jr, *et al.* Circulating adhesion molecules VCAM-1, ICAM-1, and E-selectin in carotid atherosclerosis and incident coronary heart disease cases. *Circulation* 1997;96:4219-25.
- Ghaisas NK, Shahi CN, Foley B, Goggins M, Crean P, Kelly A, *et al.* Elevated levels of circulating soluble adhesion molecules in peripheral blood of patients with unstable angina. *Am J Cardiol* 1997;80:617-9.
- Miwa K, Igawa A, Inoue H. Soluble E-selectin, ICAM-1 and VCAM-1 levels in systemic and coronary circulation in patients with variant angina. *Cardiovasc Res* 1997;36:37-44.
- Buemi M, Allegra A, Aloisi C, Corica F, Alonci A, Ruello A, *et al.* Cold pressor test raises serum concentrations of ICAM-1, VCAM-1, and E-selectin in normotensive and hypertensive patients. *Hypertension* 1997;30:845-7.
- Mulvihill N, Foley JB, Ghaisas N, Murphy R, Crean P, Walsh M. Early temporal expression of soluble cellular adhesion molecules in patients with unstable angina and subendocardial myocardial infarction. *Am J Cardiol* 1999;83:1265-7.
- Semaan HB, Gurbel PA, Anderson JL, Muhlestein JB, Carlquist JF, Horne BD, *et al.* Soluble VCAM-1 and E-selectin, but not ICAM-1 discriminate endothelial injury in patients with documented coronary artery disease. *Cardiology* 2000;93:7-10.
- Parise P, Hauert J, Iorio A, Callegari P, Agnelli G. Streptokinase and rt-PA activate platelets by a different way: implications on the rethrombosis rate after their administration in myocardial infarction. *J Lab Clin Med* 1995;125:212-21.
- Zeitler H, Ko Y, Zimmermann C, Nickenig G, Glänzer K, Walger P, *et al.* Elevated serum concentrations of soluble adhesion molecules in coronary artery disease and acute myocardial infarction. *Eur J Med Res* 1997;29;2:389-94.