



Please download and read the [Referee Guidelines](#)

## Alleviation of pain in painful diabetic neuropathy

Journal:	<i>Expert Opinion On Drug Metabolism and Toxicology</i>
Manuscript ID	EOMT-2016-0021.R1
Manuscript Type:	Review
Keywords:	antidepressants, antiepileptics, diabetes, metabolism, painful neuropathy, pharmacodynamics, pharmacokinetics, treatment

SCHOLARONE™  
Manuscripts

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Alleviation of pain in painful diabetic neuropathy**

For Peer Review Only

**Abstract**

**Introduction:** Painful diabetic neuropathy (PDN) is a disabling pain condition. Its pathomechanism remains unknown, but a sensitization and neuronal hyperexcitability have been suggested. Only symptomatic pharmacological pain-alleviation treatment is currently available.

**Areas covered:** The origin of PDN is enigmatic, and the evidence-based therapeutic guidelines therefore consist only of antidepressants and antiepileptics as first-line recommended drugs. This article relates to a MEDLINE/PubMed systematic search (2005-2015).

**Expert opinion:** The results of the meta-analysis from the aspect of the efficacy of amitriptyline, duloxetine, venlafaxine, gabapentin and pregabalin are favorable, but the placebo response rate is relatively high in patients with neuropathic pain. For personalization of the medication of PDN patients, the optimum dosing, the genotyping of the metabolizing enzymes and optimum biomarkers are needed. As concerns the future perspectives, specific sodium channel subtype inhibitors acting on peripheral nociceptive neurons or modified T-type voltage-gated calcium channel blockers may be promising targets for pharmaceutical innovations. Another attractive strategy for the treatment is based on the effects of monoclonal antibodies against nerve growth factor, sodium channel, specific receptor and cytokines. Botulinum toxin A, capsaicin patch and spinal cord stimulation therapies are the nearest future therapeutic options for the treatment of PDN patients.

**Keywords:** antidepressants, antiepileptics, diabetes, metabolism, painful neuropathy, pharmacodynamics, pharmacokinetics, treatment

**Article highlights**

- The pathomechanism of painful diabetic neuropathy is unknown, and therefore only symptomatic non-specific treatment is available.
- As a tricyclic antidepressant, amitriptyline has multimodal action, and is effective with a favorable number needed to treat, but it exerts many unfavorable adverse events related to its pharmacological profile.
- The selective serotonin-norepinephrine reuptake inhibitor duloxetine has been subjected to pharmacogenomic studies.
- Venlafaxine has a strong dose-dependent effect and due to its extended release formulation, the treatment-related adverse events are low.
- Gabapentin as an antiepileptic has no metabolism and no drug-drug interactions with a high adherence level.
- Pregabalin as a gabapentinoid with linear kinetics has been strongly recommended for the treatment of painful diabetic neuropathy.

## 1. Introduction

Neuropathic pain (NP), one of the most devastating pain conditions by definition caused by a lesion or disease of the somatosensory system [1]. It has distinct clinical features with a heterogeneity of different causes [2, 3]. Neuropathic sensory symptoms characterized by spontaneous or evoked pain is described as electric shock-like or burning, and other sensory disturbances such as paraesthesia, dysaesthesia, allodynia, hyperpathia or hyperalgesia. The symptoms of painful diabetic neuropathy (PDN) occur symmetrically in the distal parts of the extremities in an ascending manner [4, 5]. The aetiology of NP involves a wide range of peripheral or central nervous system (CNS) disorders, including diabetic and other metabolic neuropathies, dysimmune neuropathies, postherpetic neuralgia, trigeminal neuralgia, human immunodeficiency virus (HIV)-associated neuropathy or pain after spinal cord injury, post-stroke pain, multiple sclerosis-associated pain or cancer-related pain [5].

Diabetes mellitus is a modern global medical challenge with an estimated prevalence of 366 million patients in 2030 [6]. Recent epidemiological data showed that the estimated presence of NP was around 7% of the general population in Europe [7]. Diabetic neuropathy occurs in 59-66% of insulin-dependent (type 1) and non-insulin-dependent (type 2) diabetic patients [8]. The adjusted incidence of diabetic polyneuropathy is 54 per 100,000 persons per year [9]. Several studies have reported a prevalence of PDN of 10.9-20.0% [10]. Diabetic neuropathy and pain occur more frequently in type 2 diabetes mellitus, in women and in non-Caucasians [6].

The personal and socio-economic impact of PDN is high, influencing the quality of life, the work productivity and the functional disability of the patients [11]. The total annual direct medical costs for PDN were recently estimated as 27,931 USD and for severe PDN as 30,755 USD [11]. PDN has a significant impact on the health-related quality of life [12].

The pathomechanism of NP has not yet been completely clarified, but the sensitization process is widely accepted as playing an important role. Hyperexcitability and sensitization in NP similar to migraine as a primary headache disorder involve changes in the peripheral nervous system and CNS [5, 13-16]. One of the key elements of the peripheral sensitization is the increased expression of voltage-gated sodium channels  $Na_v1.8$  and  $Na_v1.9$  on lesioned peripheral nerves, which are the sources of ectopic pain impulse generation. The upregulation of sodium ion channels is a consequence of nerve growth factor (NGF) release from the damaged peripheral sensory nerve fibres [17]. **NGF is the first discovered member of the neurotrophin family. Other members of this family include brain-derived neurotrophic factor (BDNF), neurotrophin-3 (NT-3), and neurotrophin-4/5 (NT-4/5) [18]. The receptors**

1  
2  
3 of these molecules belong to the tropomyosin receptor kinase (Trk) family. NGF binds to  
4 TrkA with high affinity, whereas BDNF and NT-4 bind to TrkB, and NT-3 binds to  
5 TrkC [19]. These molecules have a common Trk downstream pathway involved in the  
6 induction of neuropathic pain [19]. Recent clinical findings have shown that anti-NGF  
7 antibody and small molecule TrkA inhibitors alleviate neuropathic pain [18, 19].  
8

9  
10  
11 Ectopic primary afferent firing clinically explains the devastating spontaneous burning pain  
12 and unpleasant electric shock-like sensations [20]. Another important mechanism of the  
13 peripheral sensitization is the presence of the proinflammatory cytokines, **bradykinin,**  
14 **prostaglandins, 5-hydroxytryptamine and histamine** ('inflammatory soup') along the  
15 damaged peripheral nerve elements from the dorsal root ganglion (DRG) to the nociceptors in  
16 the target tissue [17, 20-23]. **Human genetic studies revealed the important role of Nav1.7 in**  
17 **pain sensation [24]. Nav1.7 is expressed in small-sized nociceptive DRG neurons [25] and**  
18 **takes part in spinal cord nociceptive synaptic transmission involved in neuropathic pain**  
19 **processing [26, 27]. The importance of Nav1.7 is also highlighted by the findings that its**  
20 **expression level was found increased in DRG neurons in diabetic rats [28, 29].**  
21  
22  
23

24  
25  
26  
27  
28 Recent findings demonstrate the importance of the microglia function in the central  
29 sensitization process. The central terminals of the damaged primary sensory neurons in the  
30 spinal cord release adenosine triphosphate (ATP), which is able to activate the microglia via  
31 purinergic (P2X4) receptors. The activated microglia releases the BDNF, which activates the  
32 TrkB receptors, resulting in the down-regulation of the K<sup>+</sup> - Cl<sup>-</sup> cotransporter 2 of the  
33 nociceptive second-order neurons in spinal lamina I. These nerve cells convey the information  
34 to the thalamus (the third-order neurons) and project to the somatosensory cortex [21, 30].  
35 Excitatory amino acids, mainly glutamate and its receptors (N-methyl-D-aspartate (NMDA)  
36 receptors), have also been implicated in the pathomechanism of the NP [31]. The membrane  
37 depolarization caused by nociceptive stimuli permits NMDA receptor-mediated synaptic  
38 transmission. Calcium influx into the cells activates the non-receptor tyrosine kinase, which  
39 leads to phosphorylation of the NMDA receptors [31]. This process leads to the glutamate-  
40 related hypersensitivity of the neuronal cells [31] (**Figure 1**).  
41  
42  
43  
44  
45  
46  
47  
48

49  
50 PDN is an underdiagnosed and an undertreated condition in the daily clinical practice [4]. The  
51 aim of this article is to give an overview of the current evidence-based symptomatic treatment  
52 of PDN, focusing on the metabolism and toxicological properties of the strongly  
53 recommended first-line drugs (antidepressants and antiepileptics) in order to promote good  
54 clinical practice [3, 6, 32]. **We also provide data on future therapeutic options (i.e.**  
55 **neutrophins, NO-donors, sodium channel blockers, monoclonal antibodies (mAb)**  
56  
57  
58  
59  
60

1  
2  
3 against Nav1.7, integrins and NGF, botulinum toxin type A, topical capsaicin patch,  
4 TRPV1 and TRPA1 antagonists, and neuromodulation approaches), and give an outlook  
5 toward personalized medicine. The realization of drug-drug and drug-gene interactions  
6 (i.e. observed or inferred interactions between gene products and drug compounds) are  
7 fundamental elements of novel pharmaceutical innovation and modern therapeutic  
8 approaches for PDN patients [33].  
9  
10  
11  
12

## 13 14 15 **2. Symptomatic pain treatment of PDN**

16 The exact pathomechanism of the origin of the pain, associated with diabetic neuropathy is  
17 still lacking, and specific pharmacons for NP do not exist. The currently used drugs for NP  
18 were originally developed for depression, epilepsy and cancer-related pain, but they are  
19 effective at different levels in the alleviation of NP. This type of therapy is considered to be  
20 successful in achieving at least a 50% pain intensity reduction [34]. In this field, it is an  
21 important fact that the placebo effect for NP is higher than those for other disorders [4].  
22  
23  
24  
25

26 Over recent decades, several therapeutic recommendations and guidelines for NP and PDN  
27 have been put forward [6, 35, 36]. The recent evidence-based recommendations for the oral and  
28 topical pharmacotherapy of NP were published by the Neuropathic Pain Special Interest  
29 Group (NeuPSIG) of the International Association for the Study of Pain [3, 7]. The guideline of  
30 NeuPSIG used the Grading of Recommendation Assessment, Development and Evaluation  
31 (GRADE) system, which is based on the consensus on the rating of the quality of evidence  
32 and strength of recommendations ranging from weak to strong for or against the treatment [3, 7,  
33 37].  
34  
35  
36  
37  
38

39 This current survey analyses the pharmacological (pharmacokinetic and pharmacodynamic)  
40 and toxicological properties (absorption, distribution, metabolism and excretion) of the first-  
41 line (strong recommendation) drugs for PDN suggested by the NeuPSIG recommendations.  
42  
43  
44  
45

### 46 **2.1. Antidepressants**

47 From the aspect of the efficacious treatment of PDN, among the antidepressive drugs tricyclic  
48 antidepressants (TCAs) and serotonin-norepinephrine reuptake inhibitors (SNRIs) are  
49 beneficial. Their analgesic effect is independent of their antidepressant properties [7].  
50  
51  
52  
53

#### 54 **2.1.1. Tricyclic antidepressants (TCAs)**

1  
2  
3 The group of TCAs involves amitriptyline (AMI), clomipramine, desipramine, imipramine  
4 and nortriptyline. TCAs generally have similar efficacy as concerns PDN [3]. Particularly AMI  
5 has been investigated in most clinical trials for PDN [6].  
6  
7  
8

#### 9 10 **2.1.1.1. Amitriptyline (AMI)**

11 AMI is a more than 50-year-old drug, which was approved for the indication of depression by  
12 the US Food and Drug Administration (FDA) in 1961, and is still in use for the treatment of  
13 PDN world-wide [38].  
14

15 The chemical structure of AMI is 3-(10,11-dihydro-5*H*-dibenzo[*a,d*]cycloheptene-5-ylidene)-  
16 *N,N*-dimethylpropan-1-amine (**Figure 2**).  
17

18 The mode of action of AMI is the inhibition of the re-uptake of norepinephrine and serotonin  
19 at the adrenergic nerve endings in the synaptic cleft [39]. It has multimodal actions such as an  
20 anticholinergic effect, histaminergic down-regulation, gamma-aminobutyric acid (GABA)-B  
21 receptor up-regulation,  $\alpha$ -adrenergic receptor antagonism and sodium, L-type calcium and  
22 Kv1.1, Kv7.2, and Kv7.3 voltage-gated potassium ion channel blockade [34, 40, 41]. Importantly,  
23 AMI is an agonist of the TrkA and TRkB receptors, so it influences the activation of the NGF  
24 and BDNF [42]. It enhances the neuronal sensitivity to substance P, but does not inhibit the re-  
25 uptake of dopamine or monoamine oxidase [39]. AMI increases the activation of the adenosine  
26 A1 receptor, which leads to antinociception [40].  
27

28 The pharmacokinetic parameters of 75 mg of an orally administered osmotic controlled  
29 release tablet of AMI revealed that the maximum plasma concentration ( $C_{max}$ ) was 15.3 ( $\pm 7.0$   
30 standard deviation, S.D.) ng/L, the area under the concentration vs. time curve (AUC) was  
31 593 ( $\pm 229$  S.D.) ng/mL x h, the time to  $C_{max}$  ( $T_{max}$ ) was 25.7 ( $\pm 5.8$  S.D.) h and the plasma  
32 elimination half-life ( $T_{1/2}$ ) was 20.4 ( $\pm 4.3$  S.D.) h [39]. The bioavailability of AMI is 30-60%,  
33 the protein binding is 96%. Its active metabolite is nortriptyline [43]. It is metabolized on the  
34 first pass through the liver by cytochrome P450 (CYP450) isoenzymes (CYP2C19 and  
35 CYP2D6) and is excreted renally (around 30-50%) (**Table 1**) [39].  
36

37 As regards the pharmacogenetic aspects of AMI, the risk of the lack of efficacy or the  
38 occurrence of side-effects depends on the types of metabolizer (poor, intermediate, extensive  
39 or ultrarapid), and the genetic variations of CYP2C19 or CYP2D6 enzymes. The Clinical  
40 Pharmacogenetics Implementation Consortium recommends therapeutic drug monitoring for  
41 the guided dose adjustment [44].  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 The results of clinical trials for managing PDN revealed a superior effect of AMI to placebo  
4 [45]. The number needed to treat (NNT) of AMI is 1.3, which means an extremely good rank  
5 (Table 2) [34]. The recommended daily dose of AMI for PDN is 25-100 mg [4].  
6  
7

8 Due to its multimodal actions, AMI resulted in several unfavorable adverse events (AEs)  
9 regarding anticholinergic, histaminergic, adrenergic, dopaminergic and serotonergic effects  
10 [43]. The common (>1% frequency) AEs of AMI are dry mouth, fatigue, somnolence, weight  
11 gain and dizziness [7, 46]. Contraindications for AMI use are a history of myocardial infarction,  
12 arrhythmias, heart block, congestive heart failure, coronary artery insufficiency and severe  
13 liver disease [47]. An increased risk of sudden cardiac death mainly occurs at a dosage of more  
14 than 100 mg per day [3]. Caution is needed for AMI users with comorbidity, e.g. narrow angle  
15 glaucoma, urinary retention, prostatic hypertrophy, a history of epilepsy or hyperthyroidism [7,  
16 47].  
17  
18  
19  
20  
21  
22

23 The Cochrane Database revealed that AMI was effective for the treatment of PDN, but its  
24 wide-range use was limited by its unfavorable safety and tolerability profile [48]. The NeuSPIG  
25 consensus declares that TCAs reached the strong recommendations for use in the treatment of  
26 PDN [3].  
27  
28  
29  
30

### 31 **2.1.2. Serotonin-norepinephrine reuptake inhibitors (SNRIs)**

32 SNRIs have dual effects: they are strong blockers of the reuptake of serotonin and  
33 norepinephrine. The pharmacological group of SNRIs includes duloxetine (DUL), venlafaxine  
34 (VEN), desvenlafaxine, milnacipran, levomilnacipran and sibutramine. For the treatment of  
35 PDN, DUL and VEN have shown good efficacy [34].  
36  
37  
38  
39  
40

#### 41 **2.1.2.1. Duloxetine (DUL)**

42 DUL was developed in 1988, and FDA approval for depression and PDN was dated in 2004  
43 [34, 49]. The chemical structure of DUL is (+)-(S)-N-methyl-3-(naphthalen-1-yloxy)-3-  
44 (thiophen-2-yl) propan-1-amine. It is a potent and selective dual neuronal SNRI  
45 (serotonin:norepinephrine effects 10:1) [50]. It is a weak inhibitor of dopamine transporters and  
46 does not have a strong effect on cholinergic, histaminergic, opioid, glutamate and GABA  
47 reuptake transporters [51-53]. It does not block the monoamine oxidase types A or B [53]. DUL  
48 has the capability to block the production and/or release of pro-inflammatory cytokines and/or  
49 can also provoke the production of anti-inflammatory cytokines [54]. DUL has properties for  
50 the modulation of nitroxidative stress by influencing the balance between pro-oxidant and  
51 anti-oxidant processes [55]. One of the basic modes of action of DUL in the alleviation of pain  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 in PDN is based on the blockade of the neuronal sodium ion channel <sup>[56]</sup>. DUL blocks  
4 persistent late Nav1.7 Na<sup>+</sup> currents preferentially, which may account in part for its analgesic  
5 action <sup>[56]</sup>. In a preclinical study it was shown that the impaired norepinephrine homeostasis is  
6 one of the major factors of the PDN model in rats, and the analgesic effect of DUL might be  
7 due to the enhancement of noradrenergic signals <sup>[57]</sup>.

8  
9  
10 The pharmacokinetic parameters of **single oral dose of 60 mg of DUL tablet revealed that**  
11 **C<sub>max</sub> was 39-40.1 ng/L, AUC was 584-591 ng x h/mL, T<sub>max</sub> was 6 h and T<sub>1/2</sub> was 10.2 h** <sup>[53,</sup>  
12 <sup>58]</sup>. The bioavailability of DUL is 50%, and the protein binding (mainly to albumin) is high  
13 (>90%), and widely distributed throughout the tissues (the volume of distribution is  
14 approximately 1640 L) <sup>[43, 53, 58]</sup>. The steady state of DUL is usually achieved by day 3 of  
15 administration <sup>[53]</sup>. Its circulating metabolites are pharmacologically inactive. It is metabolized  
16 by the liver (CYP1A2 and CYP2D6) and is excreted 70% in the urine and 20% in the faeces  
17 <sup>[43]</sup>.

18  
19 The pharmacokinetics of DUL is influenced by the demographic features of the patients, such  
20 as sex, smoking, age, ethnicity, hepatic and renal function <sup>[58]</sup>. The genotype of the CYP2D6  
21 enzyme also influences the pharmacokinetics of DUL <sup>[58]</sup>. A recent pharmacogenomic study  
22 concluded that interleukin-6 variants (rs2066992 and rs10242595) may take part in DUL  
23 response in patients with major depression <sup>[59]</sup>. In DUL-treated patients with major depressive  
24 disorder, tests of the single-nucleotide polymorphisms demonstrated that variants in  
25 corticotropin-releasing hormone receptor 1 were associated with the DUL response <sup>[60]</sup>.

26  
27 As concerns the drug-drug interactions of DUL, it should be underlined that activated  
28 charcoal diminishes DUL exposure and CYP1A2 blockade increases DUL exposure. In the  
29 presence of fluvoxamine, the AUC of DUL is increased more than 400% and C<sub>max</sub> is  
30 increased more than 100% <sup>[58]</sup>. Smoking decreases the DUL plasma concentration (by 30%) <sup>[58]</sup>.  
31 In the presence of DUL with CYP2D6 inhibitors or in CYP2D6 poor metabolizers, the  
32 exposure of DUL is less increased as compared with CYP1A2 inhibition <sup>[58]</sup>. DUL has the  
33 capability to enhance the exposure of pharmaceuticals that are metabolized by CYP2D6, but not  
34 CYP1A2 <sup>[58]</sup>. Pharmacodynamic trials suggested that DUL may enhance the effects of  
35 benzodiazepines, but not alcohol or warfarin <sup>[58]</sup>.

36  
37 The investigation of the AmpliChip CYP450 Genotyping Test for genes CYP2D6 and  
38 CYP2C19 concluded that poor metabolizers accounted for around 7% of Caucasians for  
39 CYP2D6 and around 25% of East Asians for CYP2C19, and ultra-rapid metabolizers for  
40 CYP2D6 of around 29% in North Africa and the Middle-East <sup>[61, 62]</sup>.

1  
2  
3 The recent Cochrane Collaboration meta-analysis pointed out that 60 mg and 120 daily doses  
4 of DUL were efficacious in the treatment of PDN [63]. The related NNT of benefit of DUL at  
5 60 mg is 5 (95% confidential interval, CI 4 to 7) in the alleviation of pain in diabetic  
6 neuropathy [63].  
7  
8

9 The recommended daily dose of DUL for PDN is 60-120 mg once a day [3, 4].

10  
11 Mild AEs of DUL were frequent at a high dose, such as nausea, dry mouth, headache,  
12 dizziness and fatigue [7, 53, 63]. The common AEs of DUL (60 mg per day) are somnolence  
13 (20%) and constipation (14%) [34, 64]. Severe AEs of DUL were rare [63]. Precautions for the  
14 use of DUL are needed in cases of hepatic disorder, hypertension and concomitant medication  
15 with tramadol [7].  
16  
17

18 The NeuSPIG consensus declared that DUL reached the strong recommendations for use in  
19 the treatment of PDN [3].  
20  
21  
22  
23

#### 24 25 **2.1.2.2. Venlafaxine (VEN)**

26 VEN was launched on the pharmaceutical market after FDA approval in 1993, and VEN  
27 extended release (XR) in 1997 [50]. Its chemical structure is (*RS*)-1-[2-dimethylamino-1-(4-  
28 methoxyphenyl)ethyl]cyclohexanol.  
29  
30

31 VEN, as parent drug, and its active and structurally-similar major metabolite desvenlafaxine  
32 (*O*-desmethylvenlafaxine) are potent and selective dual neuronal SNRIs (VEN:  
33 serotonin:norepinephrine effects 30:1; desvenlafaxine: 10:1) [50, 65]. VEN has a strong dose-  
34 dependent action: in low doses it inhibits serotonin reuptake, while in higher doses it blocks  
35 both serotonin and norepinephrine reuptake [66]. VEN and desvenlafaxine modulate the  
36 dopaminergic neurotransmission at low affinity level [50]. They may not influence the  
37 adrenergic, muscarinic, cholinergic, histaminergic, benzodiazepine or opioids receptors, or  
38 monoamine oxidase [65]. There are later data regarding the antinociceptive effect of VEN,  
39 which revealed that this analgetic effect may be influenced by opioid receptors and the  
40 alpha2-adrenergic receptor [67]. There are findings which proved that VEN XR can modify  
41 pituitary adenylate cyclase-activating peptide (PACAP) and its PAC1 receptor genes in  
42 generalized anxiety disorder [68].  
43  
44  
45  
46  
47  
48  
49  
50

51 The pharmacokinetic parameters of orally administered VEN XR tablets of 150 mg revealed  
52 that  $C_{max}$  was 150 ug/L, the AUC was 1877 ug/L x h,  $T_{max}$  was 5.5 h and  $T_{1/2}$  was 5 h [65]. The  
53 plasma protein binding of VEN XR at 150 mg is 27% and its major metabolic pathway  
54 proceeds primarily through the liver by enzymes CYP2D6 and CYP3A3/4 [50]. It is mainly  
55 excreted in the urine (87%) [65].  
56  
57  
58  
59  
60

1  
2  
3 A recent Cochrane meta-analysis based on six randomized clinical trials showed a positive  
4 benefit for a daily dose of VEN above 150 mg for the treatment of neuropathic pain (mostly  
5 PDN) [69]. In the largest multicentre study for PDN, the VEN treatment group (150-225 mg  
6 daily dose) reached a significantly greater improvement than that for the placebo group at  
7 week 6 (50% versus 27%) [70]. The NNT of benefit of VEN in the treatment of PDN is 3.1 [34].

8  
9 The recommended daily dose of VEN XR for PDN is 150-225 mg once a day [3].

10  
11 The most common treatment-emergent AEs of VEN XR were nausea, somnolence and  
12 hypertension at high dosage [7, 70]. Nausea showed a dose-proportional manner (at 75 mg 22%  
13 and at 150-225 mg 10%) [71]. Precautions for the use of VEN include cardiac disease,  
14 hypertension and the administration of tramadol [7].

15  
16 The NeuSPIG consensus declared that VEN reached the strong recommendations for use in  
17 the treatment of PDN [3].

18  
19 Overall, the above-mentioned antidepressants are all effective in the treatment of PDN. As  
20 concerns the safety profile of SNRIs, it is a very important fact that the SNRIs have low  
21 affinity for different neuronal receptors, which results in low AEs. In spite of this, TCAs are  
22 held to be „dirty drugs” as regards their multireceptorial effects, resulting in high amounts of  
23 side-effects. In the daily clinical practice, besides the effectiveness and the safety profile of  
24 the first-line drugs for PDN, the economic aspects are also very important factors. In most  
25 countries, TCAs are readily available with low charge. DUL is a cost-effective pharmacoon for  
26 the treatment of PDN in the UK [34]. The results of randomized clinical trials and real-world  
27 studies showed DUL to be more cost-effective than antiepileptics like pregabalin (PREG) in  
28 PDN therapy [72].

## 2.2. Antiepileptics

29  
30 Among antiepileptics, only the GABA analogues, gabapentin (GBP), GBP XR and its  
31 prodrug GBP enacarbil and pregabalin (PREG) have reached the strong recommendation level  
32 in the treatment of PDN [3, 73]. Each has a similar mechanism of action to inhibit the release of  
33 presynaptic excitatory neurotransmitters [74]. A wide variety of other antiepileptic drugs were  
34 inconclusive for recommendation, e.g. carbamazepine, lacosamide, lamotrigine, levetiracetam,  
35 oxcarbazepine, topiramate, valproate and zonisamide [3, 73].

### 2.2.1. Gabapentin (GBP)

1  
2  
3 GBP was developed in 1993 and was first marketed in the US in 1994. The FDA later  
4 approved it for PDN and postherpetic neuralgia [75]. Its chemical structure is 1-  
5 (aminomethyl)cyclohexaneacetic acid.  
6

7  
8 The mechanism of analgesic action of GBP in humans has not been clarified, but in animal  
9 models has been proved that GBP binds to the  $\alpha_2\delta$  subunit of L-type voltage-dependent  
10 calcium channels [34, 76]. The consequences of this effect are the reduction of the calcium  
11 currents and inhibition of the presynaptic release of neurotransmitters, such as glutamate and  
12 substance P [76]. Interestingly, it is structurally related to GABA, but does not bind to GABA-  
13 A or GABA-B receptors, and does not block the uptake or degradation of GABA [76]. The  
14 main signs of the central sensitization are allodynia and hyperalgesia, which could be  
15 diminished by GBP [76].  
16

17  
18 The pharmacokinetic parameters of orally administered GBP tablets of 600 mg three times  
19 daily revealed that  $C_{max}$  was 8.46 mg/L, the AUC was 53.1 mg/L x h,  $T_{max}$  was 2 h and  $T_{1/2}$   
20 was 6 h [76]. The plasma protein binding of GBP is less than 3% , it is not significantly  
21 metabolized, and it is renally excreted in unchanged form [76]. The bioavailability of GBP is  
22 inversely proportional to the dose: 60% at 900 mg/day and 27% at 4800 mg/day (administered  
23 in three divided doses) [76]. **In contrast, its prodrug, enacarbil, showed essentially linear**  
24 **dose-proportional pharmacokinetic properties** [77, 78]. GBP has a volume of distribution of  
25 58 L after 150 mg intravenous administration [76].  
26

27  
28 In the available literature, there are no *in vitro* data on GBP and CYP450 interactions in  
29 humans, so GBP is neither an inducer nor an inhibitor of the hepatic metabolism of different  
30 drugs [79]. However, there are data that hydrocodone and morphine can increase GBP  
31 absorption and the plasma concentration of GBP [80]. Naproxen as a frequently used non-  
32 steroidal antiinflammatory drug (NSAID) can increase  $C_{max}$  and the AUC of GBP [80].  
33

34  
35 The recent Cochrane Database meta-analysis revealed that the efficacy of GBP for the therapy  
36 of PDN was superior to placebo (38% versus 21%, NNT 5.9, 95% CI 4.6 to 8.3) [81]. A similar  
37 effect was seen for standard GBP and a GBP XR formulation [81].  
38

39  
40 The recommended daily dose of GBP for PDN is 1200-3600 mg in three divided doses, while  
41 that of GBP XR or enacarbil is 1200-3600 mg in two divided doses [3].  
42

43  
44 A recent large retrospective clinical study in the United States revealed that among newly  
45 diagnosed PDN patients the most commonly prescribed drug was GBP (45%) [82]. From the  
46 aspect of the adherence to treatment during the first year, GBP showed a high adherence level  
47 versus PREG and DUL [82].  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 The most common AEs of GBP were dizziness (19%), somnolence (14%), peripheral oedema  
4 (7%), gait disturbance (9%), and also sedation and weight gain [7, 81]. Serious AEs were no  
5 more frequent as compared with placebo [81]. A precaution concerning the application of GBP  
6 in a case of renal insufficiency is that a reduced dosage is recommended [7].  
7  
8

9  
10 In a safety issue based on patient withdrawals due to AEs, a meta-analysis revealed that there  
11 was less withdrawal among patients who take GBP, and the most patient withdrawals due to  
12 AEs were in the case of AMI [83]. That study concluded in the focus of the benefit-risk  
13 analysis between six antidepressants and antiepileptics that the best benefit-risk ratio was  
14 shown by GBP, and the worst one by AMI [83].  
15  
16

17 The NeuSPIG consensus declares that GBP, GBP XR and enacarbil reached the strong  
18 recommendations for use in the treatment of PDN [3].  
19  
20  
21

### 22 23 **2.2.2. Pregabalin (PREG)**

24 The GABA analogue PREG received the US FDA approval for the treatment of PDN and  
25 postherpetic neuralgia in 2004 [84]. Its chemical structure is (S)-3-(aminomethyl)-5-  
26 methylhexanoic acid.  
27  
28

29 The mode of anti-nociceptive action of PREG is based on the molecular binding to the  $\alpha_2\delta$   
30 subunit of the P/Q type of the voltage-gated calcium channel in the CNS, resulting in a  
31 decreased influx of calcium ion into the neuronal cell [85]. The lack of the required quantity of  
32 calcium ion in the nerve cells results in the blockade of the propagation of neurotransmission  
33 due to inadequate synaptic vesicle fusion to the presynaptic membrane of the synapses [85].  
34 PREG has a high affinity for  $\alpha_2\delta$  type 1 and 2 proteins versus TCAs (e.g. AMI), SNRIs (e.g.  
35 VEN) or other antiepileptics (e.g. carbamazepine) [85]. PREG shows strong binding to the pain  
36 matrix-like cortex or dorsal horn of the spinal cord [85]. The PREG-induced blockade of  
37 neurotransmitters such as glutamate, glycine, norepinephrine, substance P, calcitonin gene-  
38 related peptide or acetylcholine resulted in inhibition of neuronal excitability and diminished  
39 the process of the central sensitization in NP [85].  
40  
41  
42  
43  
44  
45  
46  
47

48 As a gabapentinoid, PREG does not bind directly to GABA or benzodiazepine or opioid  
49 receptors. PREG as a glutamic acid decarboxylase (GAD) activator can increase neural  
50 GABA levels [86].  
51  
52

53 The pharmacokinetic parameters of an orally administered single dose of a PREG capsule of  
54 150 mg revealed that  $C_{max}$  was 3.99 ng/L, AUC was 28.31 ng/mL x h,  $T_{max}$  was 1.00 h and  
55  $T_{1/2}$  was 5.66 h [87]. The kinetics of PREG showed linear features (proportional to the dosage  
56 up to 900 mg per day) [88]. The plasma protein binding of PREG is absent and its metabolism  
57  
58  
59  
60



1  
2  
3 is negligible, with no effect on the CYP450 system [79, 83, 88]. The bioavailability of PREG is  
4 more than 90% and the volume of distribution 0.5 L/kg [83]. 90% is excreted in the urine as the  
5 unchanged drug with minor metabolites (e.g. N-methyl-PREG) [83]. It has no interactions  
6 metabolically with other antiepileptic drugs [79, 80].  
7

8  
9  
10 The results of randomized clinical trials for the use of PREG (300-600 mg per day) in PDN  
11 revealed significantly better efficacy as compared with placebo [84]. The evidence-based  
12 guidelines of the American Academy of Neurology announced that PREG is the only drug  
13 which reached the Level A (top-tier) of evidence in the treatment of PDN [36]. The NNT of  
14 PREG in the treatment of PDN is 5.0 [34].  
15

16  
17 The recommended daily dose of PREG for PDN is 300-60 mg in two divided doses [3].

18  
19 The common major AEs of PREG are sedation, somnolence (less than or around 50%),  
20 dizziness (less than or around 49%), headache (less than or around 29%), peripheral oedema  
21 and weight gain [7, 84].  
22

23  
24 There are data for predictive factors for AEs associated with the use of PREG in PDN patients,  
25 e.g. long duration of therapy – somnolence; advanced age – unsteadiness; elevated serum  
26 creatinine level – weight gain and oedema [89].  
27

28  
29 Precautions for PREG therapy include a reduced dosage in the case of renal failure patients [7].

30  
31 The NeuSPIG consensus declares, that PREG reached the strong recommendations for use in  
32 the treatment of PDN [3].  
33

34  
35  
36 Taken together, the effectiveness and safety profile of antiepileptics involved in the treatment  
37 of PDN are based on their pharmacokinetic and pharmacodynamic properties. From their  
38 licence permission for clinical use their prescription has been increasing, but many PDN  
39 patients do not receive them or take antiepileptics below the recommended dosage [82, 90].  
40  
41  
42  
43

### 44 45 **3. Conclusions**

46 PDN is a debilitating consequence of diabetes, which greatly affects the quality of life of the  
47 patient. The exact pathomechanism of PDN is unknown, and causative treatment is therefore  
48 unavailable. For the symptomatic treatment of PDN, there are first-line strongly  
49 recommended drugs, such as antidepressants (TCAs: AMI; and SNRIs: DUL and VEN) and  
50 antiepileptics (only gabapentinoids: GBP and PREG). Among antidepressants, TCAs and  
51 SNRIs show similar efficacy as concerns PDN, but from the aspect of their AEs, based on  
52 their pharmacokinetic and pharmacodynamic properties, huge discrepancies may be seen.  
53 Gabapentinoids have a similar mode of action and effectiveness in the treatment of PDN, but  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 some of their pharmacological properties, such as their bioavailability, absorption and kinetics  
4 vary, which leads to different features of their therapeutic and safety profiles.  
5  
6  
7

#### 8 **4. Expert opinion**

9  
10 The treatment of PDN patients is a great challenge for primary care physicians, diabetologists,  
11 neurologists and even pain specialists. Epidemiological surveys have established that a huge  
12 number of patients with NP do not receive appropriate treatment [3]. The difficulties in the  
13 treatment of PDN include the unclear pathomechanism and pathogenesis, and there are no  
14 specific efficacious pain relief drugs for PDN. It is interesting that the placebo response rate is  
15 relatively high in NP patients, and especially PDN patients. Focusing on the efficacy of a drug  
16 used for the alleviation of pain in PDN, we underline the fact, that the placebo effects are  
17 usually notably strong. This shows the narrow gap between the achieved effectiveness and the  
18 result of the placebo treatment, which gives a challenge for a novel pharmacological  
19 innovation for the preparation of new and effective drugs for the successful treatment PDN.  
20

21 Unfortunately, the recent evidence-based guidelines consist of only a few first-line strongly  
22 recommended drugs for the treatment of PDN, such as amitriptyline, duloxetine, venlafaxine,  
23 gabapentin and pregabalin. All of these have unfavourable AEs and contraindications. The  
24 correct choice of drugs is a real art of medicine. It is a perpetually recurring question for the  
25 medical staff to decide which PDN patient should receive which effective, well-tolerated and  
26 safe drug. Unfortunately, the demographic data or disease characteristics are not predictors for  
27 the pain alleviation treatment, and they do not help in the choice of the proper medication for  
28 PDN patients. Another problem is that there are no optimum biomarkers of this field at the  
29 present. There are some promising data which showed a positive correlation between plasma  
30 tumor necrosis factor (TNF)alpha levels and macrophages, inducible nitric oxide synthase  
31 (iNOS) and TNFalpha expression and the severity of the pain in PDN patients [91]. **Nitric  
32 oxide (NO) is an indigenous gas, and its elevated level has been implicated in the  
33 pathomechanism of PDN [92, 93]. Other potential biomarkers may include certain  
34 angiogenic and anti-angiogenic factors such as vascular endothelial growth factor,  
35 soluble endoglin, endothelin-1, and NO, molecules with plasma levels found elevated in  
36 diabetic patients compared to controls [94]. The furoxan NO donor, PRG150, was  
37 investigated in the streptozotocin-induced diabetic rat model of PDN and demonstrated  
38 a dose-dependent analgesic effect, which may represent a novel promising approach in  
39 PDN [93]. Furthermore, a NO-donating pregabalin, NCX1404, exerted strong**  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 **antiallodynic response in the streptozocin-induced murine model of PDN. Repeated**  
4 **dosing with NCX1404 re-established normal nociceptive response in this model** [95].  
5 **Oxidative stress has a role in the development of microvascular complications in**  
6 **patients with diabetic neuropathy. A clinical study revealed a decreased serum prolidase**  
7 **enzyme activity, which may be associated with increased NO levels and oxidative stress**  
8 **in diabetic patients with neuropathy** [96]. **In a randomized placebo-controlled study, a**  
9 **NO-releasing patch (NitroSense Derma Protect) showed a significant reduction in pain**  
10 **of PDN patients** [97].

11 We highlight the importance of finding the optimum dose of drugs for PDN, especially in the  
12 elderly due to the genotyping of the CYP450 enzymes, which play a crucial role in the  
13 metabolism of these drugs. It is similarly important for better personalizing treatment of PDN  
14 patients to integrate the CYP genotyping with therapeutic drug monitoring for the safe use of  
15 medications.

16 In the near future it is recommended to guide pharmacogenetic studies based on the genetic  
17 polymorphisms in different genes (e.g. novel techniques such as AmpliChip CYP450  
18 Genotyping Test), which can reflect the treatment response to drugs used in PDN.

19 At the bedside, one of the major challenges is the interindividual variability to the response to  
20 the drug used for PDN. As concerns the comorbidities of PDN, the polymedication and the  
21 genetic polymorphism are other important issues from the aspect of the drug metabolization  
22 and the drug-drug interactions. For the patients who are ultrarapid metabolizers, the  
23 pharmacological medications are usually ineffective, while those who are poor metabolizers  
24 are at high risk of life-threatening toxicity. There is therefore a need for a new aspect of the  
25 treatment of PDN patients such as determining the genotyping of CYP variant alleles.

26 In the coming years, it will be strongly recommended to use the genotyping analysis of  
27 CYP450 drug-drug and drug-gene interactions in the daily clinical practice in order to  
28 estimate the effectiveness and safety of the prescribed drugs for PDN. This advanced  
29 diagnostic method might serve as the basis of the request for personalized medicine.

30 The analysis of drug-drug interactions is a crucial part for pharmacokinetic and  
31 pharmacodynamic studies of the drugs used for PDN. There is a need for methods to evaluate  
32 pharmacons of this type, e.g. micellar electrokinetic chromatography electrospray ionization-  
33 tandem mass spectrometry, which is a cost-effective technique for measuring VEN and  
34 desvenlafaxine enantiomers [98].

35 One of the potential fields for the causal therapy of PDN is the sodium channel isoforms  
36 (Nav1.3, Nav1.7, Nav1.8 and Nav1.9) on the peripheral nociceptive neurons. Preclinical and  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 clinical trials have revealed that the blockers of these subtypes of the sodium channels give us  
4 new fashionable effective drugs against PDN [27].

5  
6 The sodium channel voltage sensor monoclonal antibody which targets the peripheral and  
7 central Nav1.7 channel diminishes itch and neuropathic pain in an animal model [26]. This  
8 channel-specific antibody may therefore be a potential therapeutic option in human PDN [26,  
9 99].

10  
11  
12 Another potential research area may be the modification of the glycosylation of the subclass  
13 of T-type voltage-gated calcium channels (Cav3.2), which are related to the hyperexcitable  
14 state of nociceptors in diabetic animal models [100]. In the preclinical state of this research field,  
15 the classical streptozotocin-induced diabetic rodent model as a reliable way to induce strong  
16 mechanical allodynia and thermal hyperalgesia is not sufficient to mimic all the aspects of the  
17 clinical picture of PDN. For future perspectives there is a need for new animal models which  
18 serve the innovation of the targeted treatment of PDN.

19  
20 One of the most promising therapeutic options arises from the technique of fully humanized  
21 monoclonal antibodies (mAb). **There are preclinical data pinpointing the role of integrins,  
22 receptors for extracellular matrix proteins, in the processes of inflammatory and  
23 neuropathic hyperalgesia and the chronification of pain [101-103]. The mAbs against the  
24 beta1 integrin subunit demonstrated an inhibitory effect on prostaglandin E2- and  
25 epinephrine-induced hyperalgesia in rats [101, 102]. These findings may provide new  
26 therapeutic options for the treatment of PDN. Antibodies against NGF (e.g. tanezumab,  
27 fulranumab and fasinumab) in the treatment of osteoarthritis of the hip or knee showed  
28 superiority in efficacy in pain relief (Western Ontario and McMaster – WOMAC pain)  
29 and Patient's Global Assessment compared to placebo [104]. Unfortunately, unexpected  
30 AEs with tanezumab were reported, such as rapid osteonecrosis with an unknown cause  
31 at present, a complication necessitating joint replacement surgery [105]. Based on this  
32 finding, the US Food and Drug Administration placed the studies of all anti-NGF  
33 monoclonal antibodies on clinical hold [105]. In an early truncated Phase II double-blind  
34 placebo-controlled trial with fulranumab for the treatment of PDN resulted in a dose-  
35 dependent efficacy, and there was no case necessitating joint replacement therapy [106].  
36 Another randomized controlled study with tanezumab for PDN demonstrated pain  
37 reduction versus placebo; however, there was no significant difference in Patient's  
38 Global Assessment scale [107].**

39  
40 Another class of mAb against the receptor for advanced glycation and products, which is  
41 expressed through the sensory nervous system, resulted in a dose-related attenuation of NP in  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 a mouse NP model <sup>[108]</sup>. Neutralized mAb against interleukin-17A diminished the  
4 hyperalgesia in a spinal nerve ligation model in mice <sup>[109]</sup>. Another type of mAb against high-  
5 mobility group box-1 strongly alleviated the mechanical hypersensitivity and the expression  
6 of matrix metalloproteinase-9 in the partial sciatic nerve ligation model in mice <sup>[110]</sup>. On the  
7 basis of the preclinical and early clinical results, we emphasize that mAbs targeting the  
8 different sites of the pain matrix may give a strong possibility for the causal therapy of PDN.  
9  
10 Further perspectives emerge from the intradermal application of the botulinum toxin type A  
11 for PDN. Early clinical studies suggested some effectiveness for benefit for PDN patients <sup>[111]</sup>.  
12  
13 Due to these early clinical data, further larger randomized controlled studies are required for  
14 the evaluation of the efficacy of botulinum toxin type A in this disease.  
15  
16

17  
18  
19 **The Transient Receptor Potential (TRP) cation channels, including Ankyrin repeat**  
20 **domain 1 (TRPA1) and Vanilloid type 1 (TRPV1), have crucial roles in the nociceptive**  
21 **process** <sup>[112, 113]</sup>. Recent reports demonstrated that a topical 8% patch of capsaicin, **the**  
22 **classical TRPV1 agonist**, may have a beneficial outcome in PDN patients <sup>[114]</sup>. The release of  
23 the final results is waited urgently. **In healthy human volunteers, the pharmacokinetic**  
24 **profile of TRPV1 antagonists (ABT-102 and JNJ-38893777) were evaluated, and they**  
25 **also showed a good tolerability** <sup>[115, 116]</sup>. **In a phase 2a (proof-of-concept) study, one of the**  
26 **potent and selective TRPA1 antagonist (GRC17536) demonstrated efficacy in patients**  
27 **with PDN** <sup>[117]</sup>. **Phase 3 studies are warranted in the near future to provide more**  
28 **accurate data. At the experimental level, a novel oxime compound, as a potent TRPA1**  
29 **and TRPV1 antagonist, has been proven to have promising efficacy on primary sensory**  
30 **neurons** <sup>[113]</sup>. **This compound might be a new candidate for drug development for the**  
31 **treatment of neuropathic pain** <sup>[113]</sup>.  
32  
33  
34  
35  
36  
37  
38  
39  
40

41 In the case of pharmacologically refractory PDN, one of the ultimate choices is spinal cord  
42 stimulation. A multicenter randomized clinical trial concluded that spinal cord stimulation  
43 significantly alleviated the pain in PDN patients <sup>[118]</sup>. Neuromodulation methods, such as  
44 spinal cord stimulation are invasive and expensive techniques, and careful patient selection is  
45 therefore needed. From the aspect of drug innovation, the targeting of good clinical efficacy,  
46 tolerability, safety and comparable cost-effectiveness are very important research fields.  
47 Innovative drug technology is additionally needed to achieve these goals. One example in  
48 which immediate release versus an XR formulation of VEN were compared, the XR  
49 formulation showed less AEs (e.g. nausea and dizziness) due to the novel pharmacological  
50 technology <sup>[50]</sup>.  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 Currently prescribed first-line strongly recommended symptomatic treatment for PDN  
4 (amitriptyline, duloxetine, venlafaxine, gabapentin, pregabalin) are efficacious, but all have  
5 limitations due to their precautions and AEs. To attain the better adherence of PDN patients,  
6 the ultimate goal is a large pharmacogenetic study, with the aim of the drug-drug and drug-  
7 gene interactions.  
8  
9  
10

### 11 12 13 **Conflict of interest**

14  
15 The authors declare that they have no conflict of interest and have received no payment in the  
16 preparation of their manuscript.  
17  
18  
19

### 20 21 22 **Acknowledgements**

23  
24 This work was supported by the project TÁMOP-4.2.2.A-11/1/KONV-2012-0052, by the  
25 Hungarian Brain Research Programme (NAP, Grant No. KTIA\_13\_NAP-A-III/9. and by the  
26 MTA-SZTE Neuroscience Research Group of the Hungarian Academy of Sciences and the  
27 University of Szeged.  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Figures

### Figure 1. Scheme of sensitization and hyperexcitability of mechanism of neuropathic pain

Modified (Ref.: [5])

Abbreviations:

ATP: adenosine triphosphate, BDNF: brain-derived neurotrophic factor, DRG: dorsal root ganglion,  $Na_v1.8$  and  $Na_v1.9$ : voltage-gated sodium channels, NMDA-R: N-methyl-D-aspartate receptor, NGF: nerve growth factor, P2X4: purinergic receptor, TrkB: tropomyosin receptor kinase B

### Figure 2. Chemical structures of the first-line drugs for painful diabetic neuropathy

(Ref.: [88, 119-122])

**Table legends****Table 1. Pharmacokinetic and pharmacodynamic properties of the first-line drugs for the treatment of PDN**

Abbreviations: AUC: the area under the concentration vs. time curve,  $C_{\max}$ : the maximum plasma concentration, h: hours, iv: intravenously, L: litre, NA: not available, PDN: painful diabetic neuropathy, Ref.: references,  $T_{\max}$ : the time to  $C_{\max}$ ,  $T_{1/2}$ : the plasma elimination half-life, XR: extended release

**Table 2. Recommended daily doses and adverse events of the first-line drugs for the treatment of PDN**

Abbreviations: NNT: number needed to treat; PDN: painful diabetic neuropathy; XR: extended release

(Ref.: [3, 4, 7, 34, 46, 53, 63, 64, 81, 84])

## Bibliography

Papers of special note have been highlighted as either of interest (●) or of considerable interest (●●) to readers.

1. Jensen TS, Baron R, Haanpaa M, et al. A new definition of neuropathic pain. *Pain* 2011;152:2204-5.
2. Baron R, Binder A, Wasner G. Neuropathic pain: diagnosis, pathophysiological mechanisms, and treatment. *Lancet Neurol* 2010;9:807-19.
3. Finnerup NB, Attal N, Haroutounian S, et al. Pharmacotherapy for neuropathic pain in adults: a systematic review and meta-analysis. *Lancet Neurol* 2015;14:162-73.

●● **An updated review of evidence-based recommendation for the pharmacological treatment of neuropathic pain.**

4. Alabdali M, Qrimli M, Barnett C, et al. Choosing drugs for the treatment of diabetic neuropathy. *Expert Opin Pharmacother* 2015;16:1805-14.

●● **This paper reported the recent therapeutic guidelines of diabetic neuropathy.**

5. Vecsei L, Majlath Z, Balog A, Tajti J. Drug targets of migraine and neuropathy: treatment of hyperexcitability. *CNS Neurol Disord Drug Targets* 2015;14:664-76.
6. Marmioli P, Cavaletti G. Drugs for the treatment of peripheral neuropathies. *Expert Opin Pharmacother* 2015;1-14.

**The authors presented the advanced therapy of specific neuropathies based on the most important published trials and from clinical experience.**

7. Attal N, Bouhassira D. Pharmacotherapy of neuropathic pain: which drugs, which treatment algorithms? *Pain* 2015;156 Suppl 1:S104-14.

●● **This article revealed the treatment algorithms of recommended pharmacotherapy of neuropathic pain.**

8. Dyck PJ, Kratz KM, Karnes JL, et al. The prevalence by staged severity of various types of diabetic neuropathy, retinopathy, and nephropathy in a population-based cohort: the Rochester Diabetic Neuropathy Study. *Neurology* 1993;43:817-24.
9. MacDonald BK, Cockerell OC, Sander JW, Shorvon SD. The incidence and lifetime prevalence of neurological disorders in a prospective community-based study in the UK. *Brain* 2000;123 ( Pt 4):665-76.
10. Sadosky A, McDermott AM, Brandenburg NA, Strauss M. A review of the epidemiology of painful diabetic peripheral neuropathy, postherpetic neuralgia, and less commonly studied neuropathic pain conditions. *Pain Pract* 2008;8:45-56.
11. Sadosky A, Mardekian J, Parsons B, et al. Healthcare utilization and costs in diabetes relative to the clinical spectrum of painful diabetic peripheral neuropathy. *J Diabetes Complications* 2015;29:212-7.
12. Smith SC, Lamping DL, Maclaine GD. Measuring health-related quality of life in diabetic peripheral neuropathy: a systematic review. *Diabetes Res Clin Pract* 2012;96:261-70.
13. Tajti J, Szok D, Csati A, Vecsei L. Prophylactic Drug Treatment of Migraine in Children and Adolescents: An Update. *Curr Pain Headache Rep* 2016;20:1.
14. Tajti J, Szok D, Majlath Z, et al. Migraine and neuropeptides. *Neuropeptides* 2015;52:19-30.
15. Szok D, Csati A, Vecsei L, Tajti J. Treatment of Chronic Migraine with OnabotulinumtoxinA: Mode of Action, Efficacy and Safety. *Toxins (Basel)* 2015;7:2659-73.
16. Tajti J, Tuka B, Botz B, et al. Role of pituitary adenylate cyclase-activating polypeptide in nociception and migraine. *CNS Neurol Disord Drug Targets* 2015;14:540-53.
17. Finnerup NB, Jensen TS. Mechanisms of disease: mechanism-based classification of neuropathic pain—a critical analysis. *Nat Clin Pract Neurol* 2006;2:107-15.



18. Hirose M, Kuroda Y, Murata E. NGF/TrkA Signaling as a Therapeutic Target for Pain. *Pain Pract* 2016;16:175-82.
19. Khan N, Smith MT. Neurotrophins and Neuropathic Pain: Role in Pathobiology. *Molecules* 2015;20:10657-88.
20. Baron R. Mechanisms of disease: neuropathic pain--a clinical perspective. *Nat Clin Pract Neurol* 2006;2:95-106.
21. Biggs JE, Lu VB, Stebbing MJ, et al. Is BDNF sufficient for information transfer between microglia and dorsal horn neurons during the onset of central sensitization? *Mol Pain* 2010;6:44.
22. Calvo M, Dawes JM, Bennett DL. The role of the immune system in the generation of neuropathic pain. *Lancet Neurol* 2012;11:629-42.
23. Sommer C. Painful neuropathies. *Curr Opin Neurol* 2003;16:623-8.
24. Cox JJ, Reimann F, Nicholas AK, et al. An SCN9A channelopathy causes congenital inability to experience pain. *Nature* 2006;444:894-8.
25. Minett MS, Nassar MA, Clark AK, et al. Distinct Nav1.7-dependent pain sensations require different sets of sensory and sympathetic neurons. *Nat Commun* 2012;3:791.
26. Lee JH, Park CK, Chen G, et al. A monoclonal antibody that targets a NaV1.7 channel voltage sensor for pain and itch relief. *Cell* 2014;157:1393-404.
27. Kharatmal SB, Singh JN, Sharma SS. Voltage-Gated Sodium Channels as Therapeutic Targets for Treatment of Painful Diabetic Neuropathy. *Mini Rev Med Chem* 2015;15:1134-47.
28. Zhang JL, Yang JP, Zhang JR, et al. Gabapentin reduces allodynia and hyperalgesia in painful diabetic neuropathy rats by decreasing expression level of Nav1.7 and p-ERK1/2 in DRG neurons. *Brain Res* 2013;1493:13-8.
29. Galloway C, Chattopadhyay M. Increases in inflammatory mediators in DRG implicate in the pathogenesis of painful neuropathy in Type 2 diabetes. *Cytokine* 2013;63:1-5.
30. Trang T, Beggs S, Salter MW. Brain-derived neurotrophic factor from microglia: a molecular substrate for neuropathic pain. *Neuron Glia Biol* 2011;7:99-108.
31. Petrenko AB, Yamakura T, Baba H, Shimoji K. The role of N-methyl-D-aspartate (NMDA) receptors in pain: a review. *Anesth Analg* 2003;97:1108-16.
32. Beniczky S, Tajti J, Timea Varga E, Vecsei L. Evidence-based pharmacological treatment of neuropathic pain syndromes. *J Neural Transm (Vienna)* 2005;112:735-49.
33. Wagner AH, Coffman AC, Ainscough BJ, et al. DGIdb 2.0: mining clinically relevant drug-gene interactions. *Nucleic Acids Res* 2016;44:D1036-44.
34. Javed S, Petropoulos IN, Alam U, Malik RA. Treatment of painful diabetic neuropathy. *Ther Adv Chronic Dis* 2015;6:15-28.
35. **●● The authors gave us a detailed pain management concerning the painful diabetic neuropathy.**
35. Attal N, Cruccu G, Baron R, et al. EFNS guidelines on the pharmacological treatment of neuropathic pain: 2010 revision. *Eur J Neurol* 2010;17:1113-e88.
36. Bril V, England J, Franklin GM, et al. Evidence-based guideline: Treatment of painful diabetic neuropathy: report of the American Academy of Neurology, the American Association of Neuromuscular and Electrodiagnostic Medicine, and the American Academy of Physical Medicine and Rehabilitation. *Neurology* 2011;76:1758-65.
37. Guyatt GH, Oxman AD, Vist GE, et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ* 2008;336:924-6.
38. Fangmann P, Assion HJ, Juckel G, et al. Half a century of antidepressant drugs: on the clinical introduction of monoamine oxidase inhibitors, tricyclics, and tetracyclics. Part II: tricyclics and tetracyclics. *J Clin Psychopharmacol* 2008;28:1-4.
39. Gupta SK, Shah JC, Hwang SS. Pharmacokinetic and pharmacodynamic characterization of OROS and immediate-release amitriptyline. *Br J Clin Pharmacol* 1999;48:71-78.




- 1  
2  
3 40. Dharmshaktu P, Tayal V, Kalra BS. Efficacy of antidepressants as analgesics: a review. *J Clin Pharmacol* 2012;52:6-17.
- 4  
5 41. Punke MA, Friederich P. Amitriptyline is a potent blocker of human Kv1.1 and Kv7.2/7.3  
6 channels. *Anesth Analg* 2007;104:1256-64, tables of contents.
- 7  
8 42. Jang SW, Liu X, Chan CB, et al. Amitriptyline is a TrkA and TrkB receptor agonist that  
9 promotes TrkA/TrkB heterodimerization and has potent neurotrophic activity. *Chem Biol*  
10 2009;16:644-56.
- 11  
12 43. Italiano D, Spina E, de Leon J. Pharmacokinetic and pharmacodynamic interactions  
13 between antiepileptics and antidepressants. *Expert Opin Drug Metab Toxicol* 2014;10:1457-  
14 89.
- 15  
16 44. Hicks JK, Swen JJ, Thorn CF, et al. Clinical Pharmacogenetics Implementation  
17 Consortium guideline for CYP2D6 and CYP2C19 genotypes and dosing of tricyclic  
18 antidepressants. *Clin Pharmacol Ther* 2013;93:402-8.
- 19  
20 45. Max MB, Lynch SA, Muir J, et al. Effects of desipramine, amitriptyline, and fluoxetine on  
21 pain in diabetic neuropathy. *N Engl J Med* 1992;326:1250-6.
- 22  
23 46. Dodick DW, Freitag F, Banks J, et al. Topiramate versus amitriptyline in migraine  
24 prevention: a 26-week, multicenter, randomized, double-blind, double-dummy, parallel-group  
25 noninferiority trial in adult migraineurs. *Clin Ther* 2009;31:542-59.
- 26  
27 47. Gore M, Dukes E, Rowbotham D, et al. Prevalence of contraindicated medical conditions  
28 and use of precluded medications in patients with painful neuropathic disorders prescribed  
29 amitriptyline. *Pain Pract* 2006;6:265-72.
- 30  
31 48. Moore RA, Derry S, Aldington D, et al. Amitriptyline for neuropathic pain in adults.  
32 *Cochrane Database Syst Rev* 2015;7:CD008242.
- 33  
34 **• The importance of the role of amitriptyline for neuropathic pain treatment is  
35 highlighted.**
- 36  
37 49. Bymaster FP, Beedle EE, Findlay J, et al. Duloxetine (Cymbalta), a dual inhibitor of  
38 serotonin and norepinephrine reuptake. *Bioorg Med Chem Lett* 2003;13:4477-80.
- 39  
40 50. Sansone RA, Sansone LA. Serotonin norepinephrine reuptake inhibitors: a  
41 pharmacological comparison. *Innov Clin Neurosci* 2014;11:37-42.
- 42  
43 51. Bymaster FP, Lee TC, Knadler MP, et al. The dual transporter inhibitor duloxetine: a  
44 review of its preclinical pharmacology, pharmacokinetic profile, and clinical results in  
45 depression. *Curr Pharm Des* 2005;11:1475-93.
- 46  
47 52. Pergolizzi JV, Jr., Raffa RB, Taylor R, Jr., et al. A review of duloxetine 60 mg once-daily  
48 dosing for the management of diabetic peripheral neuropathic pain, fibromyalgia, and chronic  
49 musculoskeletal pain due to chronic osteoarthritis pain and low back pain. *Pain Pract*  
50 2013;13:239-52.
- 51  
52 53. Carter NJ, McCormack PL. Duloxetine: a review of its use in the treatment of generalized  
53 anxiety disorder. *CNS Drugs* 2009;23:523-41.
- 54  
55 54. De Berardis D, Conti CM, Serroni N, et al. The effect of newer serotonin-noradrenalin  
56 antidepressants on cytokine production: a review of the current literature. *Int J Immunopathol*  
57 *Pharmacol* 2010;23:417-22.
- 58  
59 55. Salat K, Moniczewski A, Librowski T. Nitrogen, oxygen or sulfur containing heterocyclic  
60 compounds as analgesic drugs used as modulators of the nitroxidative stress. *Mini Rev Med Chem* 2013;13:335-52.
- 56  
57 56. Wang SY, Calderon J, Kuo Wang G. Block of neuronal Na<sup>+</sup> channels by antidepressant  
58 duloxetine in a state-dependent manner. *Anesthesiology* 2010;113:655-65.
- 59  
60 57. Kinoshita J, Takahashi Y, Watabe AM, et al. Impaired noradrenaline homeostasis in rats  
with painful diabetic neuropathy as a target of duloxetine analgesia. *Mol Pain* 2013;9:59.
- 58  
59 58. Knadler MP, Lobo E, Chappell J, Bergstrom R. Duloxetine: clinical pharmacokinetics and  
drug interactions. *Clin Pharmacokinet* 2011;50:281-94.

- 1  
2  
3 59. Maciukiewicz M, Marshe VS, Tiwari AK, et al. Genetic variation in IL-1beta, IL-2, IL-6,  
4 TSPO and BDNF and response to duloxetine or placebo treatment in major depressive  
5 disorder. *Pharmacogenomics* 2015;16:1919-29.
- 6 60. Perlis RH, Fijal B, Dharia S, Houston JP. Pharmacogenetic investigation of response to  
7 duloxetine treatment in generalized anxiety disorder. *Pharmacogenomics J* 2013;13:280-5.
- 8 61. de Leon J, Susce MT, Murray-Carmichael E. The AmpliChip CYP450 genotyping test:  
9 Integrating a new clinical tool. *Mol Diagn Ther* 2006;10:135-51.
- 10 62. de Leon J, Armstrong SC, Cozza KL. Clinical guidelines for psychiatrists for the use of  
11 pharmacogenetic testing for CYP450 2D6 and CYP450 2C19. *Psychosomatics* 2006;47:75-85.
- 12 63. Lunn MP, Hughes RA, Wiffen PJ. Duloxetine for treating painful neuropathy, chronic  
13 pain or fibromyalgia. *Cochrane Database Syst Rev* 2014;1:CD007115.
- 14 **• A metaanalysis of randomized clinical trials demonstrated the effect of duloxetine for**  
15 **painful neuropathies.**
- 16 64. Goldstein DJ, Lu Y, Detke MJ, et al. Duloxetine vs. placebo in patients with painful  
17 diabetic neuropathy. *Pain* 2005;116:109-18.
- 18 65. Barman Balfour JA, Jarvis B. Venlafaxine Extended-Release: A Review of its Clinical  
19 Potential in the Management of Generalised Anxiety Disorder. *CNS Drugs* 2000;14:483-503.
- 20 66. Redrobe JP, Bourin M, Colombel MC, Baker GB. Dose-dependent noradrenergic and  
21 serotonergic properties of venlafaxine in animal models indicative of antidepressant activity.  
22 *Psychopharmacology (Berl)* 1998;138:1-8.
- 23 67. Schreiber S, Bleich A, Pick CG. Venlafaxine and mirtazapine: different mechanisms of  
24 antidepressant action, common opioid-mediated antinociceptive effects--a possible opioid  
25 involvement in severe depression? *J Mol Neurosci* 2002;18:143-9.
- 26 68. Cooper AJ, Narasimhan S, Rickels K, Lohoff FW. Genetic polymorphisms in the PACAP  
27 and PAC1 receptor genes and treatment response to venlafaxine XR in generalized anxiety  
28 disorder. *Psychiatry Res* 2013;210:1299-300.
- 29 69. Gallagher HC, Gallagher RM, Butler M, et al. Venlafaxine for neuropathic pain in adults.  
30 *Cochrane Database Syst Rev* 2015;8:CD011091.
- 31 70. Rowbotham MC, Goli V, Kunz NR, Lei D. Venlafaxine extended release in the treatment  
32 of painful diabetic neuropathy: a double-blind, placebo-controlled study. *Pain* 2004;110:697-  
33 706.
- 34 71. Grothe DR, Scheckner B, Albano D. Treatment of pain syndromes with venlafaxine.  
35 *Pharmacotherapy* 2004;24:621-9.
- 36 72. Bellows BK, Dahal A, Jiao T, Biskupiak J. A cost-utility analysis of pregabalin versus  
37 duloxetine for the treatment of painful diabetic neuropathy. *J Pain Palliat Care Pharmacother*  
38 2012;26:153-64.
- 39 73. Wiffen PJ, Derry S, Moore RA, et al. Antiepileptic drugs for neuropathic pain and  
40 fibromyalgia - an overview of Cochrane reviews. *Cochrane Database Syst Rev*  
41 2013;11:CD010567.
- 42 **• An overview of effectiveness and safety of antiepileptic drugs for the treatment of**  
43 **neuropathic pain.**
- 44 74. Schreiber AK, Nones CF, Reis RC, et al. Diabetic neuropathic pain: Physiopathology and  
45 treatment. *World J Diabetes* 2015;6:432-44.
- 46 75. Stacey BR. Gabapentin: a viewpoint by Brett R. Stacey. *CNS Drugs* 2003;17:983.
- 47 76. Curran MP, Wagstaff AJ. Gabapentin: in postherpetic neuralgia. *CNS Drugs* 2003;17:975-  
48 82.
- 49 77. Chen C. Meta-analyses of dose-exposure relationships for gabapentin following oral  
50 administration of gabapentin and gabapentin enacarbil. *Eur J Clin Pharmacol* 2013;69:1809-  
51 17.
- 52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 78. Zhang L, Rainka M, Freeman R, et al. A randomized, double-blind, placebo-controlled  
4 trial to assess the efficacy and safety of gabapentin enacarbil in subjects with neuropathic pain  
5 associated with postherpetic neuralgia (PXN110748). *J Pain* 2013;14:590-603.
- 6 79. Patsalos PN. Drug interactions with the newer antiepileptic drugs (AEDs)--part 1:  
7 pharmacokinetic and pharmacodynamic interactions between AEDs. *Clin Pharmacokinet*  
8 2013;52:927-66.
- 9 80. Patsalos PN. Drug interactions with the newer antiepileptic drugs (AEDs)--Part 2:  
10 pharmacokinetic and pharmacodynamic interactions between AEDs and drugs used to treat  
11 non-epilepsy disorders. *Clin Pharmacokinet* 2013;52:1045-61.
- 12 81. Moore RA, Wiffen PJ, Derry S, et al. Gabapentin for chronic neuropathic pain and  
13 fibromyalgia in adults. *Cochrane Database Syst Rev* 2014;4:CD007938.
- 14 82. Yang M, Qian C, Liu Y. Suboptimal Treatment of Diabetic Peripheral Neuropathic Pain  
15 in the United States. *Pain Med* 2015;16:2075-83.
- 16 83. Rudroju N, Bansal D, Talakokkula ST, et al. Comparative efficacy and safety of six  
17 antidepressants and anticonvulsants in painful diabetic neuropathy: a network meta-analysis.  
18 *Pain Physician* 2013;16:E705-14.
- 19  
20 **• A network meta-analysis comparing efficacy and safety of different antiepileptics and**  
21 **antidepressants in the treatment of painful diabetic neuropathy.**
- 22 84. Tassone DM, Boyce E, Guyer J, Nuzum D. Pregabalin: a novel gamma-aminobutyric acid  
23 analogue in the treatment of neuropathic pain, partial-onset seizures, and anxiety disorders.  
24 *Clin Ther* 2007;29:26-48.
- 25 85. Mico JA, Prieto R. Elucidating the mechanism of action of pregabalin: alpha(2)delta as a  
26 therapeutic target in anxiety. *CNS Drugs* 2012;26:637-48.
- 27 86. Froestl W. An historical perspective on GABAergic drugs. *Future Med Chem* 2011;3:163-  
28 75.
- 29 87. Tjandrawinata RR, Setiawati E, Putri RS, et al. Pharmacokinetic equivalence study of two  
30 formulations of the anticonvulsant pregabalin. *Clin Pharmacol* 2015;7:69-75.
- 31 88. Schulze-Bonhage A. Pharmacokinetic and pharmacodynamic profile of pregabalin and its  
32 role in the treatment of epilepsy. *Expert Opin Drug Metab Toxicol* 2013;9:105-15.
- 33 89. Kanbayashi Y, Onishi K, Hosokawa T. Factors predicting adverse events associated with  
34 pregabalin administered for neuropathic pain relief. *Pain Res Manag* 2014;19:e164-7.
- 35 90. Hall GC, Morant SV, Carroll D, et al. An observational descriptive study of the  
36 epidemiology and treatment of neuropathic pain in a UK general population. *BMC Fam Pract*  
37 2013;14:28.
- 38 91. Purwata TE. High TNF-alpha plasma levels and macrophages iNOS and TNF-alpha  
39 expression as risk factors for painful diabetic neuropathy. *J Pain Res* 2011;4:169-75.
- 40 92. Mikaili P, Moloudizargari M, Aghajanshakeri S. Treatment with topical nitroglycerine  
41 may promote the healing process of diabetic foot ulcers. *Med Hypotheses* 2014;83:172-4.
- 42 93. Huang LY, Tsui DY, Williams CM, et al. The Furoxan Nitric Oxide Donor, PRG150,  
43 Evokes Dose-Dependent Analgesia in a Rat Model of Painful Diabetic Neuropathy (PDN).  
44 *Clin Exp Pharmacol Physiol* 2015.
- 45 94. Motawi TK, Rizk SM, Ibrahim IA, El-Emady YF. Alterations in circulating angiogenic  
46 and anti-angiogenic factors in type 2 diabetic patients with neuropathy. *Cell Biochem Funct*  
47 2014;32:155-63.
- 48 95. Varani K, Vincenzi F, Targa M, et al. Repeated Dosing with NCX1404, a Nitric Oxide-  
49 Donating Pregabalin, Re-establishes Normal Nociceptive Responses in Mice with  
50 Streptozotocin-Induced Painful Diabetic Neuropathy. *J Pharmacol Exp Ther* 2016;357:240-7.
- 51 96. Sayin R, Aslan M, Kucukoglu ME, et al. Serum prolidase enzyme activity and oxidative  
52 stress levels in patients with diabetic neuropathy. *Endocrine* 2014;47:146-51.
- 53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 97. Agrawal RP, Jain S, Goyal S, et al. A Clinical Trial of Nitrosense patch for the treatment  
4 of patients with painful diabetic neuropathy. *J Assoc Physicians India* 2014;62:385-90.
- 5 98. Liu Y, Jann M, Vandenberg C, et al. Development of an enantioselective assay for  
6 simultaneous separation of venlafaxine and O-desmethylvenlafaxine by micellar  
7 electrokinetic chromatography-tandem mass spectrometry: Application to the analysis of  
8 drug-drug interaction. *J Chromatogr A* 2015;1420:119-28.
- 9 99. Bray N. Analgesia: anti-itch and anti-ouch antibody. *Nat Rev Drug Discov* 2014;13:574.
- 10 100. Orestes P, Osuru HP, McIntire WE, et al. Reversal of neuropathic pain in diabetes by  
11 targeting glycosylation of Ca(V)3.2 T-type calcium channels. *Diabetes* 2013;62:3828-38.
- 12 101. Dina OA, Hucho T, Yeh J, et al. Primary afferent second messenger cascades interact  
13 with specific integrin subunits in producing inflammatory hyperalgesia. *Pain* 2005;115:191-  
14 203.
- 15 102. Dina OA, Parada CA, Yeh J, et al. Integrin signaling in inflammatory and neuropathic  
16 pain in the rat. *Eur J Neurosci* 2004;19:634-42.
- 17 103. Ferrari LF, Levine JD. Plasma membrane mechanisms in a preclinical rat model of  
18 chronic pain. *J Pain* 2015;16:60-6.
- 19 104. Schnitzer TJ, Marks JA. A systematic review of the efficacy and general safety of  
20 antibodies to NGF in the treatment of OA of the hip or knee. *Osteoarthritis Cartilage* 2015;23  
21 Suppl 1:S8-17.
- 22 105. Hochberg MC. Serious joint-related adverse events in randomized controlled trials of  
23 anti-nerve growth factor monoclonal antibodies. *Osteoarthritis Cartilage* 2015;23 Suppl  
24 1:S18-21.
- 25 106. Wang H, Romano G, Frustaci ME, et al. Fulranumab for treatment of diabetic peripheral  
26 neuropathic pain: A randomized controlled trial. *Neurology* 2014;83:628-37.
- 27 107. Bramson C, Herrmann DN, Carey W, et al. Exploring the role of tanezumab as a novel  
28 treatment for the relief of neuropathic pain. *Pain Med* 2015;16:1163-76.
- 29 108. Brederson JD, Strakhova M, Mills C, et al. A monoclonal antibody against the receptor  
30 for advanced glycation end products attenuates inflammatory and neuropathic pain in the  
31 mouse. *Eur J Pain* 2015.
- 32 109. Yao CY, Weng ZL, Zhang JC, et al. Interleukin-17A Acts to Maintain Neuropathic Pain  
33 Through Activation of CaMKII/CREB Signaling in Spinal Neurons. *Mol Neurobiol* 2015.
- 34 110. Zhang FF, Morioka N, Harano S, et al. Perineural expression of high-mobility group  
35 box-1 contributes to long-lasting mechanical hypersensitivity via matrix metalloproteinase-9  
36 upregulation in mice with painful peripheral neuropathy. *J Neurochem* 2015.
- 37 111. Lakhani SE, Velasco DN, Tepper D. Botulinum Toxin-A for Painful Diabetic  
38 Neuropathy: A Meta-Analysis. *Pain Med* 2015;16:1773-80.
- 39 112. Jordt SE, Ehrlich BE. TRP channels in disease. *Subcell Biochem* 2007;45:253-71.
- 40 113. Payrits M, Saghy E, Matyus P, et al. A novel 3-(4,5-diphenyl-1,3-oxazol-2-yl)propanal  
41 oxime compound is a potent Transient Receptor Potential Ankyrin 1 and Vanilloid 1 (TRPA1  
42 and VR1) receptor antagonist. *Neuroscience* 2016;324:151-62.
- 43 114. Burness CB, McCormack PL. Capsaicin 8 % Patch: A Review in Peripheral Neuropathic  
44 Pain. *Drugs* 2016;76:123-34.
- 45 115. Othman AA, Nothhaft W, Awni WM, Dutta S. Pharmacokinetics of the TRPV1  
46 antagonist ABT-102 in healthy human volunteers: population analysis of data from 3 phase 1  
47 trials. *J Clin Pharmacol* 2012;52:1028-41.
- 48 116. Manitpisitkul P, Mayorga A, Shalayda K, et al. Safety, Tolerability and Pharmacokinetic  
49 and Pharmacodynamic Learnings from a Double-Blind, Randomized, Placebo-Controlled,  
50 Sequential Group First-in-Human Study of the TRPV1 Antagonist, JNJ-38893777, in Healthy  
51 Men. *Clin Drug Investig* 2015;35:353-63.
- 52  
53  
54  
55  
56  
57  
58  
59  
60


- 1  
2  
3 117. Preti D, Saponaro G, Szallasi A. Transient receptor potential ankyrin 1 (TRPA1)  
4 antagonists. *Pharm Pat Anal* 2015;4:75-94.  
5 118. de Vos CC, Meier K, Zaalberg PB, et al. Spinal cord stimulation in patients with painful  
6 diabetic neuropathy: a multicentre randomized clinical trial. *Pain* 2014;155:2426-31.  
7 119. Smith TR. Duloxetine in diabetic neuropathy. *Expert Opin Pharmacother* 2006;7:215-23.  
8 120. Ryan NM. A review on the efficacy and safety of gabapentin in the treatment of chronic  
9 cough. *Expert Opin Pharmacother* 2015;16:135-45.  
10 121. Reinen J, Postma G, Tump C, et al. Application of a cocktail approach to screen  
11 cytochrome P450 BM3 libraries for metabolic activity and diversity. *Anal Bioanal Chem*  
12 2016;408:1425-43.  
13 122. Wellington K, Perry CM. Venlafaxine extended-release: a review of its use in the  
14 management of major depression. *CNS Drugs* 2001;15:643-69.  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43




Figure 2. Chemical structures of the first-line drugs for PDN




**Amitriptyline**  
(3-(10,11-Dihydro-5H-dibenzo[a,d]cycloheptene-5-ylidene)-N,N-dimethylpropan-1-amine)




**Duloxetine**  
(+)-(S)-N-Methyl-3-(naphthalen-1-yloxy)-3-(thiophen-2-yl)propan-1-amine



**Venlafaxine**  
(RS)-1-[2-dimethylamino-1-(4-methoxyphenyl)-ethyl]cyclohexanol



**Gabapentin**  
1-(Aminomethyl)cyclohexaneacetic acid



**Pregabalin**  
(S)-3-(Aminomethyl)-5-methylhexanoic acid

Table 1. Pharmacokinetic and pharmacodynamic properties of the first-line drugs for the treatment of PDN

Drugs	C <sub>max</sub>	T <sub>max</sub>	AUC	T <sub>1/2</sub>	Bioavailability	Protein binding	Volume of distribution	Metabolism	Excretion	References
<b>Antidepressants</b>										
Amitriptyline (75 mg tablet)	15.3 ng/L	25.7 h	593 ng/mL x h	20.4 h	30-60%	96%	NA	CYP2C1 9 CYP2D6	urine (30-50%) faeces (NA)	[39, 43]
Duloxetine (60 mg tablet)	39- 40.1 ng/L	6 h	584-591 ng x h/mL	10.2 h	50%	>90%	1640 L	CYP1A2 CYP2D6	urine (70%) faeces (20%)	[50, 53, 58]
Venlafaxine XR (150 mg tablet)	150.0 ug/L	5.5 h	1877 ug/L x h	5.0 h	NA	27%	NA	CYP2D6 CYP3A3 /4	urine (87%) faeces (NA)	[65]
<b>Antiepileptics</b>										
Gabapentin (600 mg tablet)	8.5 mg/L	2.0 h	53 mg/L x h	6.0 h	60% (900 mg/day) 27% (4800	<3%	58 L (150 mg iv.)	nil	urine (NA) faeces (NA)	[76, 79, 83]



					mg/day)					
Pregabalin (150 mg capsule)	4.0 ng/L	1 h	28 ug/mL x h	5.7 h	90%	absent	0.5 L/kg	nil	urine (90%) faeces (NA)	[87, 88]

Abbreviations: AUC: the area under the concentration vs. time curve,  $C_{max}$ : the maximum plasma concentration, h: hours, iv: intravenously, L: litre, NA: not available, PDN: painful diabetic neuropathy, Ref.: references,  $T_{max}$ : the time to  $C_{max}$ ,  $T_{1/2}$ : the plasma elimination half-life, XR: extended release

**Table 2. Recommended daily doses and adverse events of the first-line drugs for the treatment of PDN**

<b>Drugs</b>	<b>Recommended daily dose (mg)</b>	<b>NNT</b>	<b>Adverse events</b>
<b>Antidepressants</b>			
Amitriptyline	25-100 mg	1.3	dry mouth, fatigue, somnolence, weight gain, dizziness
Duloxetine	60-120 mg	6.0	nausea, dry mouth, headache, dizziness, fatigue, somnolence, constipation
Venlafaxine Venlafaxine XR	150-225 mg	3.1	nausea, somnolence, hypertension
<b>Antiepileptics</b>			
Gabapentin Gabapentin XR Gabapentin enacarbil	1200-3600 mg	5.9	dizziness, somnolence, peripheral oedema, gait disturbance, sedation, weight gain
Pregabalin	300-600 mg	5.0	sedation, somnolence, dizziness, headache, peripheral oedema, weight gain

Abbreviations: NNT: number needed to treat; PDN: painful diabetic neuropathy; XR: extended release  
(Ref.: [3, 4, 7, 34, 46, 53, 63, 64, 81, 84])

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



**PUBLISHING AGREEMENT**

This is an agreement under which you, the author, assign copyright in your article to Informa UK Limited registered in England under no. 1072954 trading as Taylor & Francis Group, Registered Office: 5 Howick Place, London SW1P 1WG (hereinafter 'Taylor & Francis') to allow us to publish your article, including abstract, tables, figures, data, and supplemental material hosted by us, as the Version of Record (VoR) in the Journal for the full period of copyright throughout the world, in all forms and all media, subject to the Terms & Conditions below.

Please read this agreement carefully, complete it, and return a copy to us by email, fax, or hard copy immediately, to avoid any delay in the publication of your article.

Postal address: Taylor & Francis Journals Production, 4 Park Square, Milton Park, Abingdon OX14 4RN, UK

Fax: +44 (0) 207 017 6336 Email: [T&Fproduction@tandf.co.uk](mailto:T&Fproduction@tandf.co.uk)

ARTICLE TITLE: (Article) ALLEVIATION OF PAIN IN PAINFUL DIABETIC NEUROPATHY

AUTHOR(S): JAVOS TASTI, BELLA NOK, USFIA HASLATHY AVERT CSFTI, ANKA PETROVICI-BALOG, LAJLOS VESZELY

JOURNAL TITLE: Expert Opinion on Drug Metabolism & Toxicology ('Journal')

Please complete and sign below.

Please tick either box A or box B, BUT NOT BOTH

- A** I own copyright, and I am assigning copyright in my article to Taylor & Francis. In the case of a multi-authored article, I confirm that I am authorized by my co-authors to make this assignment as their agent on their behalf. The co-authors have agreed the priority of the assertion of copyright and the order of names in the publication of the article.
- B** I am a civil servant or an employee of a Government, Government Agency, International Organization, or Commercial Corporation which is granting a non-exclusive licence to publish the article and which hereby recognizes Taylor & Francis as the sole licensee for the publication of the final, definitive, and citable Version of Record (VoR). In the case of a multi-authored article, I confirm that I am authorized by my co-authors to enter into this licence as their agent on their behalf. The co-authors have agreed the priority of the assertion of copyright and the order of names in the publication of the article.


If you have ticked B, please indicate which of the statements below apply to you (and your co-authors):

- I am an employee of the UK, Canadian, Australian, or another Commonwealth Realm Government, and the Crown retains and asserts copyright.
- I am a US Government (including NIH) employee and there is no copyright to transfer.
- I am a contractor of the US Government (includes NIH contractors) under contract number: \_\_\_\_\_
- I am an employee of the European Commission and copyright is asserted and retained by the European Union.
- I am an employee of the World Bank and copyright is asserted and retained by that entity.
- I am an employee of the Food & Agricultural Organization and copyright is asserted and retained by that entity.
- I am an employee of a Government, Agency, or International Organization and copyright is retained by that entity. Name of entity: \_\_\_\_\_
- I am employed and the copyright belongs to my employer (or is a 'work made for hire' under US law). Name of corporation: \_\_\_\_\_

**ASSIGNMENT OF COPYRIGHT**

I hereby assign Taylor & Francis with full title guarantee all rights of copyright and related publishing rights in my article, in all forms and all media (whether known at this time or developed at any time in the future) throughout the world, in all languages, where our rights include but are not limited to the right to translate, create adaptations, extracts, or derivative works and to sublicense such rights, for the full term of copyright (including all renewals and extensions of that term), to take effect if and when the article is accepted for publication.

I confirm that I have read and accept the full Terms & Conditions below including my author warranties, and have read and agree to comply with the Journal's policies on peer review and publishing ethics.

Signed:  Name (Print): JAVOS TASTI

Position: ASSOCIATE PROFESSOR Date: 11/04/2016

THIS FORM IS A LEGALLY BINDING DOCUMENT. WE RECOMMEND THAT YOU RETAIN A COPY OF IT AND CONSULT A LEGAL ADVISOR IF YOU HAVE ANY QUESTIONS.

209x298mm (300 x 300 DPI)

## ASSIGNMENT OF COPYRIGHT, TERMS &amp; CONDITIONS

## DEFINITION

1. Your article is defined as comprising (a) your Accepted Manuscript (AM) in its final form; (b) the final, definitive, and citable Version of Record (VoR) including the abstract, text, bibliography, and all accompanying tables, illustrations, data, and media; and (c) any supplemental material hosted by Taylor & Francis. This and these Terms & Conditions constitute the entire agreement and assignment and these Terms & Conditions shall be taken into account when interpreting your and our rights and obligations under this agreement, unless amended by a written document signed by both of us.

## TAYLOR &amp; FRANCIS' RESPONSIBILITIES

2. If deemed acceptable by the Editors of the Journal, we shall prepare and publish your article in the Journal. We may post your accepted manuscript in advance of the formal publication of the VoR. We reserve the right to make such editorial changes as may be necessary to make the article suitable for publication, or as we reasonably consider necessary to avoid infringing third-party rights or breaching any laws; and we reserve the right not to proceed with publication for whatever reason.
3. Taylor & Francis will deposit your Accepted Manuscript (AM) to any designated institutional repository including PubMedCentral (PMC), with which Taylor & Francis has an article deposit agreement, see 4. iv (j) below.

## RIGHTS RETAINED BY YOU AS AUTHOR

4. These rights are personal to you, and your co-authors, and cannot be transferred by you to anyone else. Without prejudice to your rights as author set out below, you undertake that the fully referenced-linked VoR will not be published elsewhere without our prior written consent. You assert and retain the following rights as author(s):
- The right to be identified as the author of your article, whenever and wherever the article is published, such rights including moral rights arising under § 77, Copyright, Designs & Patents Act 1988, and so far as is legally possible, any corresponding rights we may have in any territory of the world.
  - The right to retain patent rights, trademark rights, or rights to any process, product or procedure described in your article.
  - The right to post and maintain at any time the Author's Original Manuscript (AOM; your manuscript in its original and unrefined form: a 'preprint').
  - The right to post at any time after publication of the VoR your AM (your manuscript in its revised after peer review and accepted for publication form: a 'postprint') as a digital file on your own personal or departmental website, provided that you do not use the VoR published by us, and that you include any amendments or deletions or warnings relating to the article issued or published by us; and with the acknowledgement: 'The Version of Record of this manuscript has been published and is available in <JOURNAL TITLE> <date of publication> <http://www.tandfonline.com/<Article DOI>>'
- a) Please note that embargoes apply with respect to posting the AM to an institutional or subject repository. For further information, please see our list of journals with applicable embargo periods: EDC | Excel. For the avoidance of doubt, you are not permitted to post the final published paper, the VoR published by us, to any site, unless it has been published as Open Access on our website.
- If, following publication, you or your funder pay an Article Publishing Charge for retrospective Open Access publication, you may then opt for one of three licenses: CC BY, CC BY-NC, or CC BY-NC-AD; if you do not respond, we shall assign a CC BY license. All rights in the article will revert to you as author.
  - The right to share with colleagues copies of the article in its published form as supplied to you by Taylor & Francis as a digital print or printed reprint on a non-commercial basis.
  - The right to make printed copies of all or part of the article on a non-commercial basis for use by you for lecture or classroom purposes provided that such copies are not offered for sale or distributed in any systematic way, and provided that acknowledgement to prior publication in the Journal is given.
  - The right, if the article has been produced within the scope of your employment, for your employer to use all or part of the article internally within the institution or company on a non-commercial basis provided that acknowledgement to prior publication in the Journal is given.
  - The right to include the article in a thesis or dissertation that is not to be published commercially, provided that acknowledgement to prior publication in the Journal is given.
  - The right to present the article at a meeting or conference and to distribute printed copies of the article to the delegates attending the meeting provided that this is not for commercial purposes and provided that acknowledgement to prior publication in the Journal is given.
  - The right to use the article in its published form in whole or in part without revision or modification in personal compilations, or other publications of your own work, provided that acknowledgement to prior publication in the Journal is given.
  - The right to expand your article into book-length form for publication provided that acknowledgement to prior publication in the Journal is made explicit (see below). Where permission is sought to revise an article in a book chapter or edited collection on a commercial basis a fee will be payable by the publisher of the new work. Where you are the author of the article have had the lead role in the new work (i.e. you are the author of the new work or the editor of the edited collection), fees will be waived. Acknowledgement to prior publication in the Journal should be made explicit (see below).

**Acknowledgement:** This <chapter or book> is derived in part from an article published in <JOURNAL TITLE> <date of publication> copyright Taylor & Francis, available online: <http://www.tandfonline.com/<Article DOI>>

If you wish to use your article in a way that is not permitted by this agreement, please contact [permissions@tandf.co.uk](mailto:permissions@tandf.co.uk)

## WARRANTIES MADE BY YOU AS AUTHOR

5. You warrant that:
- All persons who have a reasonable claim to authorship are named in the article as co-authors including yourself, and you have not fabricated or misappropriated anyone's identity, including your own.
  - You have been authorized by all such co-authors to sign this agreement on their behalf, and to agree on their behalf the priority of the assertion of copyright and the order of names in the publication of the article.
  - The article is your original work, apart from any permitted third-party copyright material you include, and does not infringe any intellectual property rights of any other person or entity and cannot be construed as plagiarizing any other published work, including your own published work.
  - The article is not currently under submission to, nor is under consideration by, nor has been accepted by any other journal or publication, nor has been previously published by any other journal or publication, nor has been assigned or licensed by you to any third party.
  - The article contains no content that is abusive, defamatory, libellous, obscene, fraudulent, nor in any way infringes the rights of others, nor is in any other way unlawful or in violation of applicable laws.
  - Any patient, service user, or participant (or that person's parent or legal guardian) in any research or clinical experiment or study who is described in the article has given written consent to the inclusion of material, text or image, pertaining to themselves, and that they acknowledge that they cannot be identified via the article and that you have anonymized them and that you do not identify them in any way. Where such a person is deceased, you warrant you have obtained the written consent of the deceased person's family or estate.
  - You have complied with all mandatory laboratory health and safety procedures in the course of conducting any experimental work reported in your article; your article contains all appropriate warnings concerning any specific and particular hazards that may be involved in carrying out experiments or procedures described in the article or involved in instructions, materials, or formulae in the article; your article includes explicitly relevant safety precautions; and cites, if an accepted Standard or Code of Practice is relevant, a reference to the relevant Standard or Code.
  - You have acknowledged all sources of research funding, as required by your research funder, and disclosed any financial interest or benefit you have arising from the direct applications of your research.
  - You have obtained the necessary written permission to include material in your article that is owned and held in copyright by a third party, which shall include but is not limited to any proprietary text, illustration, table, or other material, including data, audio, video, film stills, screenshots, musical notation and any supplemental material.
  - You have read and complied with our policy on publishing ethics.
  - You have read and complied with the Journal's Instructions for Authors.
  - You will keep us and our affiliates indemnified in full against all loss, damages, injury costs and expenses (including legal and other professional fees and expenses) awarded against or incurred or paid by us as a result of your breach of the warranties given in this agreement.
  - You consent to allowing us to use your article for marketing and promotional purposes.

## GOVERNING LAW

6. This agreement (and any dispute, proceeding, claim or controversy in relation to it) is subject to English law and the parties hereby submit to the exclusive jurisdiction of the Courts of England and Wales.

209x298mm (300 x 300 DPI)