

University of Wollongong
Research Online

Faculty of Science, Medicine and Health -
Papers: part A

Faculty of Science, Medicine and Health

1-1-2016

Self-determination in the context of mental health recovery

Ellie K. Taylor

University of Wollongong, elliejo@uow.edu.au

Lorna Moxham

University of Wollongong, lmoxham@uow.edu.au

Dana J. Perlman

University of Wollongong, dperlman@uow.edu.au

Christopher F. Patterson

University of Wollongong, cpatters@uow.edu.au

Renee M. Brighton

University of Wollongong, reeneb@uow.edu.au

See next page for additional authors

Follow this and additional works at: <https://ro.uow.edu.au/smhpapers>



Part of the [Medicine and Health Sciences Commons](#), and the [Social and Behavioral Sciences Commons](#)

Recommended Citation

Taylor, Ellie K.; Moxham, Lorna; Perlman, Dana J.; Patterson, Christopher F.; Brighton, Renee M.; and Liersch, Susan, "Self-determination in the context of mental health recovery" (2016). *Faculty of Science, Medicine and Health - Papers: part A*. 3864.
<https://ro.uow.edu.au/smhpapers/3864>

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library: research-pubs@uow.edu.au

Self-determination in the context of mental health recovery

Abstract

Approximately one in every five Australians will experience a mental illness each year (ABS, 2007). Mental illnesses are not homogenous. There are no clearly established clinical pathways and, as such, care and treatment is necessarily highly individualised.

Disciplines

Medicine and Health Sciences | Social and Behavioral Sciences

Publication Details

Taylor, E., Moxham, L., Perlman, D., Patterson, C., Brighton, R. & Liersch, S. (2016). Self-determination in the context of mental health recovery. *Australian Nursing and Midwifery Journal*, 23 (10), 41-41.

Authors

Ellie K. Taylor, Lorna Moxham, Dana J. Perlman, Christopher F. Patterson, Renee M. Brighton, and Susan Liersch

Title: Self-determination in the context of mental health recovery

Authors: Ellie Taylor¹, Lorna Moxham², Dana Perlman³, Christopher Patterson⁴,
Renee Brighton⁴, Susan Sumskis⁴

¹ PhD Candidate (Health Science), University of Wollongong, Northfields Ave,
Wollongong NSW 2522; PH: (02) 4239 2137; elliejo@uow.edu.au; Qualifications:
BPsych(Hons); MSc(Research).

² Professor of Mental Health Nursing, School of Nursing, University of Wollongong

³ Senior Lecturer, School of Education, University of Wollongong

⁴ Lecturer, School of Nursing, University of Wollongong

Article:

Approximately one in every five Australians will experience a mental illness each year (ABS 2007). Mental illnesses are not homogenous. There are no clearly established clinical pathways and, as such, care and treatment is necessarily highly individualised. It is within this nebulous treatment approach that personal recovery is the goal for many who live with debilitating mental health issues. Personal recovery is not synonymous with cure, but can be defined as “gaining a social identity through engagement in an active life” (Moxham et al 2015, p. 62).

In recent years, self-determination has arisen as an area of importance within mental health care (Craike and Coleman 2005). Carpenter (2002) asserts that there are strong links between increased self-determination and personal recovery. Self-determination can be considered as the propensity of an individual to act in a “self-directed, self-regulated, autonomous” way (Field et al 1998, p. 2). Such an approach is respectful of values and appreciates lived experience – a fundamental tenant of personal recovery. People with a lived experience of mental illness, however, report significantly low levels of self-determination (Okon and Webb 2014). According to Hagger and Chatzisarantis (2009), self-determined motivation is strongly associated with engagement in positive health behaviours such as increased medication and service adherence, and other activities that promote wellbeing (Chang 2011).

The Australian Department of Health (2013), in their *National Framework for Recovery-Oriented Mental Health Services: Guide for Practitioners and Providers*, posits that mental health care should maximise opportunities for personal autonomy and self-determination. This involves maximising choice on the part of the consumer, with less reliance on restrictive methods of treatment, such as restraint and seclusion. However, very little empirical research has been conducted regarding the relationship between practices and outcomes (including self-determination) associated with personal recovery from mental illness (Le Boutillier et al 2011).

This research project explores the influence of a recovery-oriented, strengths-based therapeutic recreation initiative and how it influences self-determination among people with a mental illness. In doing so, a detailed examination will occur to investigate the mechanisms (e.g. shared decision making, addressing social context) by which self-determination influences personal recovery from mental illness. Informed by the lived experience of people with a mental illness, a mixed methods approach has been employed. Surveys will be administered at three time points: before and after the intervention, and again at three-month follow-up. At follow-up, a qualitative approach will be used to collect data through individual semi-structured interviews and focus groups. These interviews will further explore the quantitative findings by incorporating a comprehensive person-centred approach. To date, quantitative findings from this project have supported the efficacy of therapeutic recreation in increasing self-determination among people with a mental illness. The mechanisms by which this contributes to personal recovery are, however, not visible in research literature. It is this concept that will be explored in this project.

Health care providers have a responsibility to respect and emphasise the autonomy and self-determination of all people for whom they provide care. This is no less an expectation for people who live with a mental illness. Their journey towards personal recovery is one that should be heavily reliant on their own expertise and based on their own choices. Results from this research are expected to inform future mental health services and programs.

Word count: 539 words

References:

- Australia. Bureau of Statistics (ABS). 2007. National Survey of Mental Health and Well-being: Summary of results. Canberra, ACT: Australian Bureau of Statistics. Cat.no.4326.0.
- Australia. Department of Health. 2013. A national framework for recovery-oriented mental health services: Guide for practitioners and providers. Canberra, ACT: Australian Department of Health.
- Carpenter, J. 2002. Mental health recovery paradigm: Implications for social work. *Health and Social Work*. 27(2): 86-94.
- Chang, L. 2011. An interaction effect of leisure self-determination and leisure competence of older adults' self-rated health. *Journal of Health Psychology*. 17(3): 324-332.
- Craike, M. and Coleman, D. 2005. Buffering effect of leisure self-determination on the mental health of older adults. *Leisure/Loisir*. 29(2): 310-328.
- Field, S., Martin, J., Miller, R., Ward, M. and Wehmeyer, M. 1998. *A practical guide for teaching self-determination*. Reston, VA: Council for Exceptional Children.
- Hagger, M. and Chatzisarantis, N. 2009. Integrating the theory of planned behavior and self-determination theory in health behavior: A meta-analysis. *British Journal of Health Psychology*. 14: 275-302.
- Le Boutillier, C., Leamy, M., Bird, V. J., Davidson, L., Williams, J. and Slade, M. 2011. What does recovery mean in practice? A qualitative analysis of international recovery-oriented practice guidance. *Psychiatric Services*. 62(12): 1470-1476.
- Moxham, L., Liersch-Sumskis, S., Taylor, E., Patterson, C. and Brighton, R. 2015. Preliminary outcomes of a pilot therapeutic recreation camp for people with a mental illness: Links to recovery. *Therapeutic Recreation Journal*. 49(1): 61-75.
- Okon, S. and Webb, D. 2014. Self-determination: A curriculum of empowerment for health and wellness in a psychosocial rehabilitation clubhouse. *Occupational Therapy in Mental Health*. 30(2): 196-212.