

Evaluation of noni (*Morinda citrifolia*) volatile profile by dynamic headspace and gas chromatography-mass spectrometry

Avaliação do perfil de voláteis em noni (Morinda citrifolia) por headspace dinâmico e cromatografia gasosa-espectrometria de massas

Adriana SOUSA¹, Manoel Alves SOUZA NETO¹, Deborah dos Santos GARRUTI¹, João SOUSA¹, Edy Sousa de BRITO^{1*}

Abstract

Noni is a fruit that has interested the scientific community due to its medicinal and functional activities. Different products that contain noni are already in the market, but their consumption could be impaired by their distinctive unpleasant aroma and flavor. The aim of this work was to evaluate the noni pulp volatile profile by dynamic headspace and gas chromatography-mass spectrometry. Thirty seven volatile compounds were detected, mainly alcohols (63.3%), esters (26.9%), cetones (7.4%), and acids (1.2%).

Keywords: noni; volatile compounds; GC-MS; aroma; flavor.

Resumo

O noni é um fruto que tem interessado à comunidade científica por sua atividade funcional e medicinal. Já se encontram no mercado diferentes produtos que contêm noni em sua composição, mas seu consumo tem sido prejudicado por seu aroma e sabor desagradáveis. O objetivo deste trabalho foi o de avaliar o perfil de voláteis da polpa de noni pela técnica de *headspace* dinâmico e cromatografia gasosa-espectrometria de massas. Foram detectados 37 compostos voláteis, sendo os principais: alcoóis (63,3%), ésteres (26,9%), cetonas (7,4%) e ácidos (1,2%).

Palavras-chave: noni; compostos voláteis; CG-EM; aroma; sabor.

1 Introduction

Noni comes from the Rubiaceae family which contains approximately eighty species, twenty of which have economic or other benefits. Noni stands out among its other family members because of its diverse uses, and it is able to grow in distinctly different environments including sandy areas, fertile soils, and rocky terrains. Noni is found in most South Pacific islands, Malaysia, the West Indies, Indonesia, the Philippines, Taiwan, Vietnam, India, Africa, and Guam (ELKINS, 2002).

The noni plant has traditionally been used by Polynesians for medicinal purposes for more than 2000 years (McCLATCHEY, 2002). The presence of biological compounds has been reported, such as glycosides, polysaccharides, iridoids, alkaloids, lignans, trisaccharide fatty acid esters, anthraquinones, scopoletin, morindin, vitamins, and minerals (WANG et al., 2000; LIU et al., 2001; FURUSAWA et al., 2003; SU et al., 2005). Some in vitro and in vivo analysis demonstrate that noni can confer health benefits in the form of scavenging of free radicals, antimutagenicity, anticarcinoma activity, anticlastogenic activity, inhibition of low-density-lipoprotein oxidation, anti-inflammatory activity, blood purification, stimulation of the immune system, regulation of cell function, and regulation of cholesterol (SALUDES et al., 2002; FURUSAWA et al., 2003; HORNICK et al., 2003; KAMIYA et al.,

2004). It is noteworthy that according to the *Informe Técnico* nº 25 (technical safety report #25) noni products commercialization as food should be not allowed in Brazil. This recommendation was based on the lack of scientific information regarding its safety as food (AGÊNCIA..., 2008).

Volatile compounds are responsible for the characteristic flavor of foods. These compounds are from different chemical classes present in a wide concentration range and are thermal instable. Commonly, the volatile profile is composed by dozens of compounds, but only a few have an effective contribution to the characteristic food aroma and flavor. Despite its richness in functional compounds, the consumption of noni products could be impaired by their sensory properties. The flesh of the fruit is characteristically bitter, and when completely ripe it produces a very distinctive, somewhat offensive, odor and flavor (ELKINS, 2002). Only one study was found in literature on noni fruit volatile compounds. Farine et al. (1996) analyzed a solvent extract and reported 51 volatile compounds on noni collected in Moorea (French Polynesia). These authors found 25 acids, which corresponded to 83% of the total volatile compounds, mostly low molecular weight fatty acids which are widely known as possessing strong sweat, cheese, fat, and rancid odors.

Recebido para publicação em 6/5/2008

Aceito para publicação em 7/7/2009 (003481)

¹ Tecnologia de Alimentos, Embrapa Agroindústria Tropical, CEP 60511-110, Fortaleza - CE, Brasil, E-mail: edy@cnpat.embrapa.br

*A quem a correspondência deve ser enviada

The aim of this study was to characterize the volatile profile of noni pulp, cultivated on the Northeast of Brazil, by headspace analysis since it is generally recognized that the composition of volatile components in the vapor phase above food is more closely related to the aroma than to the composition of these compounds in the food medium itself.

2 Materials and methods

Noni plants were cultivated on a commercial farm in Trairí – State of Ceará, Brazil. The plants were originally from Suriname and were introduced in Brazil by the Federal University of Sergipe. After harvesting, the fruits were transported to the laboratory and stored at ambient temperature for three days before pulp extraction. The pulp was collected on a pulp extractor with screen diameter of 0.2 cm, packed in plastic bags, and stored at $-16\text{ }^{\circ}\text{C}$ until use.

2.1 Isolation of volatile compounds

The volatile compounds from the headspace of the noni pulp were swept by vacuum (70 mmHg) and concentrated into a 0.1 g Porapak Q trap according to the general dynamic headspace methodology described by Franco and Rodriguez-Amaya (1983) for soursop and adapted for other fruits by Franco, Garruti and Da Silva (1998) and Jales et al. (2006).

The Noni pulp (200 g) were diluted with 100 mL water, added 30% w/w NaCl, and put into the collecting apparatus (Figure 1). Suction was kept for 2 hours. Trap desorption was performed by elution with 300 μL of three different solvents (acetone, hexane, and methylene chloride). All solvents employed were pure and of chromatographic grade (Em Science, Merck - Darmstadt, Germany). The Porapak Q polymer (80–100 mesh) was obtained from Waters Associates Inc (Milford, MA, USA). Prior to use in the experiment, the polymer was heated at $170\text{ }^{\circ}\text{C}$ for 48 hours in a flow of pure nitrogen at 30 mL/minute.

2.2 Gas chromatographic analysis

Volatile compounds were separated on a CP-Sil-8CB (Varian, Walnut Creek, CA, USA), fused silica capillary column (30 m length, 0.25 mm id, and 0.25 μm film thickness) in a

Varian model 3800 gas chromatograph. The splitless mode injector was maintained at $220\text{ }^{\circ}\text{C}$ and the flame ionization detector (FID) at $250\text{ }^{\circ}\text{C}$. Hydrogen was the carrier gas at a flow rate of 52 cm/second. The oven temperature was set at $30\text{ }^{\circ}\text{C}$, held for 10 minutes, programmed to $90\text{ }^{\circ}\text{C}$ ($5\text{ }^{\circ}\text{C}/\text{minute}$) and then to $200\text{ }^{\circ}\text{C}$ ($20\text{ }^{\circ}\text{C}/\text{minute}$), held at $200\text{ }^{\circ}\text{C}$ for 10 minutes. The injected volume was 1 μL .

2.3 Gas chromatography/mass spectrometry

The volatile compounds were identified in a Shimadzu gas chromatograph (Kyoto, Japan) coupled to a quadrupole mass spectrometer (GC/MS), model QP-2010, at 70 eV ionization voltage and 1 scan/second MS scan range. The column and oven conditions were the same as those used for the chromatographic analysis. Helium was the carrier gas at a flow rate of 35 cm/second.

2.4 Retention indices

A standard mixture of paraffin homologues C8–C21 (Polyscience 211C kit, Chicago, IL, USA) was prepared using methylene chloride as solvent. 1 μL standard mixture was injected under the same gas chromatographic/mass spectrometry conditions for calculating linear retention indices.

2.5 Identification of compounds

The identification was made by matching the mass spectra of the unknown compound with those present in the computer database (National Institute for Standard Technology – NIST 02 - 147,198 compounds) and also by the comparison of their experimental retention indices with known compounds' indices (LINSTROM; MALLARD, 2008).

3 Results and discussion

Figure 2 shows the chromatograms of noni volatiles collected by dynamic headspace and eluted from the trap with three different solvents (acetone, hexane, and methylene chloride). The methylene chloride extract presented the richest profile with higher number of peaks and larger peaks. Therefore, it was used in further analysis for volatile identification.

Table 1 lists the 37 compounds detected in the methylene chloride extract. The peak area percentage was used to indicate the relative concentration of each compound. They were identified as seven alcohols, corresponding to 63.3% of the volatile compounds, 20 esters (26.9%), three cetones (7.4%), six acids (1.2%), and one aldehyde. The major compounds, based on the relative amount, were 3-methyl-3-buten-1-ol (54.83%), methyl hexanoate (13.04%), methyl butanoate (8.1%), 2-heptanone (6.86%), and benzyl alcohol (5.2%).

Farine et al. (1996), studying the volatile components of ripe noni fruits and their effects on drosophila, also detected significant amount of 3-methyl-3-buten-1-ol (4%). The 3-methyl-3-buten-1-ol was reported to be the aglycone of nonioside A (3-methylbut-3-enyl-6-O- α -D-glucopyranosyl- α -D-glucopyranoside), a glycoside reported to be present in noni pulp (WANG et al., 2000; DALSGAARD et al, 2006).




Figure 1. Entrapment system by suction.




Figure 2. Effect of different desorption solvents on the profile of noni headspace volatiles.

However, Farine et al. (1996) observed a quite diverse volatile profile from that reported in the present work, with the predominance of fatty acids (83%) and small percentage of esters (3%). Only two compounds were found to be responsible for 77% of the total amount, octanoic acid (58%), and hexanoic acid (19%). The differences between these two noni profiles could be due to the volatiles isolation method employed since the cited authors used a direct solvent extraction with methylene chloride while the present work studied the headspace volatile profile. Besides, in Farine's work, the solvent extraction was carried out in a strong basic medium, which certainly liberated the fatty acids from the noni glycosides. These glycosides are esterified with fatty acids, specially hexanoic and octanoic acids, and have a trivial name of noniosides (DALSGAARD et al., 2006).

Table 1. Headspace volatile compounds from noni pulp.

Peak nº	Compound	tR (minute)	IK	Area (%)
Ester				
1	methyl 2-methylpropanoate	3.917	<800	1.20
4	methyl butanoate	5.038	<800	8.10
8	ethyl butanoate	9.462	807	0.45
8b	ethyl lactate	10.373	824	0.17
9	4-pentenyl acetate	15.073	891	0.53
12	methyl hexanoate	17.132	931	13.04
13	ethyl hexanoate	20.374	1000	0.51
17	4-pentenyl butanoate	22.698	1076	0.45
19	methyl benzoate	23.350	1096	0.06
20b	3-methylbutyl pentanoate	23.623	1104	-
21	3-methyl-3-butenyl 3-methylbutanoate	23.709	1114	0.01
23	methyl octanoate	23.927	1128	2.19
24	ethyl benzoate	24.651	1174	0.12
27	3-methyl-2-butenyl hexanoate	26.016	1290	0.04
28	methyl decanoate	26.291	1322	0.02
28a	ethyl 3-phenylpropanoate	26.545	1354	-
32	ethyl phthalate	28.362	1589	0.03
33	di-isobutyl phthalate	31.353	1858	-
34	methyl hexadecanoate	32.406	1920	-
35	dibutyl phthalate	33.089	1952	-
Alcohols				
5	3-methyl-3-buten-1-ol	5.397	<800	54.83
6	dimethyl disulfide + 3-methyl-1-butanol	5.713	<800	-
7	3-methyl-2-buten-1-ol	7.595	<800	2.69
8e	1-hexanol	14.399	883	0.48
15a	Benzyl alcohol	21.648	1043	5.20
11	3-methyl-2-buten-1-ol acetate	17.042	929	-
20a	nonanol	23.587	1103	0.09
Acids				
6a	2-methylpropanoic acid	7.379	<800	-
8a	butanoic acid	9.729	812	0.35
8c	3-methylbutanoic acid	13.201	867	0.06
8d	2-methylbutanoic acid	14.170	880	0.09
14	hexanoic acid	20.629	1009	-
25	octanoic acid	24.729	1179	0.72
Ketones				
2	2-pentanone	3.965	<800	0.44
10	2-heptanone	15.310	894	6.86
18	2-nonanone	23.303	1094	0.08
Aldehyde				
3	pentanal	4.262	<800	-

In the present work, no modification was made in the noniosides, thus the free hexanoic and octanoic acids concentration in the noni pulp were much lower than that found by Farine. Besides, these two fatty acids show very low vapor pressure values (0.02 to 0.2 mmHg), and their concentrations in the headspace become much lower than those found in the food matrix. On the other hand, esters show vapor pressure values a hundred times higher than those of the fatty acids (above 2 mmHg). Such fact can explain why the predominance of this

class of compounds was found in the headspace analysis. In addition, Farine et al. (1996) concentrated the 450 mL methylene chloride extracts to 500 µL by Kuderna-Danish method, which, raising the temperature for a long period of time can cause large loss of esters in their extracts.

Although sensory analysis was not performed in this study, it could seem quite curious that a product with a small percentage of hexanoic and octanoic and high content of esters in its headspace can present such a sweat and cheese-like aroma. Nonetheless, it is important to point out that the impact of a volatile compound on food flavor depends on its threshold value and solubility in water or fat, besides its concentration (SHAHIDI; RUBIM; SOUZA, 1986). Hexanoic and octanoic acids present very low threshold values, 0.6 and 8 ppb, respectively (NATIONAL..., 2004), and so their odor can be perceived even in small concentrations. However, the two approaches are not exclusive, but complementary. Prior to consumption, the volatiles are sampled orthonasally, and this first sniff often has a major influence on the overall acceptability of the product. Volatiles are also released during eating, liberated from the matrix in the masticatory process, and travel to the olfactory receptors by the retronasal route. Together with the basic tastes (sour, salty, sweet, bitter, and umami) and other oral sensations (freshness, astringency, etc) they form the food flavor. Other source of variation could be the species variability. Bicalho et al. (2000), studying cashew apple volatiles concluded that the differences found in the literature depend more on genetic variability than on the method of extraction.

4 Conclusions

The headspace volatiles profile of noni harvested in the Northeastern region of Brazil showed 37 compounds. The major chemical class was alcohols (63.3%), followed by esters (26.9%), cetones (7.4%), and acids (1.2%). The major compounds were 3-methyl-3-buten-1-ol, methyl hexanoate, methyl butanoate, 2-heptanone, and benzyl alcohol. Further studies are underway to identify the compounds responsible by the noni unpleasant flavor and to find a feasible process to remove them.

References

- AGÊNCIA NACIONAL DE VIGILÂNCIA SANITÁRIA – ANVISA. **Informe Técnico no 25:** Esclarecimentos sobre as avaliações de segurança realizadas de produtos contendo *Morinda Citrifolia*, também conhecida como Noni. Disponível em: <http://www.anvisa.gov.br/alimentos/informes/25_290507.htm>. Acesso em: 22 dez. 2008.
- BICALHO, B. et al. Application of high-temperature gas chromatography–mass spectrometry to the investigation of glycosidically bound components related to cashew apple (*Anacardium occidentale* L. var. *nanum*) volatiles. **Journal of Agriculture and Food Chemistry**, v. 48, p. 1167–1174, 2000.
- DALSGAARD, P. W. et al. Noniosides E-H, new trisaccharides fatty acid esters from the fruit of *Morinda citrifolia* (Noni). **Planta Medica**, v. 72, p. 1322-1327, 2006.
- ELKINS, R. M. H. **The noni revolution:** Today's tropical wonder that can battle disease, boost energy and revitalize your health. Woodland, 2002. p. 34-35.
- FARINE, J. P. et al. Volatile components of ripe fruits of *Morinda citrifolia* and their effects on *Drosophila*. **Phytochemistry**, v. 41, p. 433-438, 1996.
- FRANCO, M. R. B.; GARRUTI, D. S.; DA SILVA, M. A. A. P. Time and solvent optimization for the trapping of the volatile compounds of cashew juice and wine by suction on Porapak Q. **Revista Cubana de Química**, v. 10, p. 273–274, 1998.
- FRANCO, M. R. B.; RODRIGUEZ-AMAYA, D. B. Trapping of soursop (*Annona muticata*) juice volatile on Porapak Q by suction. **Journal of the Science of Food and Agriculture**, v. 34, p. 293-299, 1983.
- FURUSAWA, E. et al. Antitumor potential of a polysaccharide-rich substance from the fruit juice of *Morinda citrifolia* (noni) on sarcoma180 ascites tumour in mice. **Phytotherapy Research**, v. 17, p. 1158–1164, 2003.
- HORNICK, C. A. et al. Inhibition of angiogenic initiation and disruption of newly established human vascular networks by juice from *Morinda citrifolia* (noni). **Angiogenesis**, v. 6, p. 143–149, 2003.
- JALES, K. A. et al. Otimização das condições de isolamento dos voláteis do headspace do maracujá amarelo por concentração a vácuo em Porapak Q. **Alimentos e Nutrição**, v. 17, p. 195-201, 2006.
- KAMIYA, K. et al. Chemical constituents of *Morinda citrifolia* fruits inhibit copper-induced low-density lipoprotein oxidation. **Journal of Agricultural and Food Chemistry**, v. 52, p. 5843–5848, 2004.
- LINSTROM, P. J.; MALLARD, W. G. (Eds.). **NIST Chemistry WebBook**. NIST Standard Reference Database Number 69. Gaithersburg MD: National Institute of Standards and Technology, s.d. Disponível em: <<http://webbook.nist.gov>>. Acesso em: 8 set. 2008.
- LIU, G. M. et al. Two novel glycosides from the fruits of *Morinda citrifolia* (noni) inhibit AP-1 transactivation and cell transformation in the mouse epidermal JB6 cell line. **Cancer Research**, v. 61, p. 5749–5756, 2001.
- McCLATCHEY, W. From Polynesian healers to health food stores: Changing perspectives of *Morinda citrifolia* (Rubiaceae). **Integrative Cancer Therapies**, v. 1, p. 110–120, 2002.
- NATIONAL LIBRARY OF MEDICINE. **Hazardous Substances Data Bank**. Bethesda, MD: 2004. Disponível em: <<http://toxnet.nlm.nih.gov/2004>>. Acesso em: fev. 2009.
- SALUDES, J. P. et al. Antitubercular constituents from the hexane fraction of *Morinda citrifolia* Linn. (Rubiaceae). **Phytotherapy Research**, v. 16, p. 683–685, 2002.
- SHAHIDI, F.; RUBIN, L. J.; D'SOUZA, L. A. Meat flavor volatiles: A review of the composition, techniques of analysis, and sensory evaluation **CRC Critical Review in Food Science and Nutrition**, v. 24, n. 2, p.141, 1986.
- SU, B. N. et al. Chemical constituents of the fruits of *Morinda citrifolia* (noni) and their antioxidant activity. **Journal of Natural Products**, v. 68, p. 592–595, 2005.
- WANG, M. F. et al. Novel glycosides from noni (*Morinda citrifolia*). **Journal of Natural Products**, v. 63, p. 1182–1183, 2000.