



Plant & Animal Genomes XIX Conference

January 15-19, 2011
Town & Country Convention Center
San Diego, CA

W153: Coffee Genomics

Analysis Of Nucleotide Diversity In *Coffea* spp.

[Luiz Filipe P Pereira](#)^{1,2} , [Karina Yanagui](#)² , [Lucia P Ferreira](#)² , [Douglas S Domingues](#)² , [Luiz G E Vieira](#)² , [David Pot](#)^{1,3}

¹ Embrapa Café, PqEB s/n, Brasília, DF, 70770-901, Brasil

² Lab. Biotecnologia, IAPAR, Rod. Celso Garcia CID Km 375, Londrina, PR, 86001-970, Brasil

³ CIRAD, Avenue Agropolis, TA-A-96/03, 34398 Montpellier Cedex, France

Marker-assisted selection becomes a reality for many crops; in perennial crops, the utilization of molecular markers in breeding programs can speed up genotype selection. However, the most important commercial coffee species - allotetraploid *C. arabica* - has a restrict number of available polymorphic markers, which is a consequence of the narrow genetic basis and low molecular variability among cultivars. In order to study the nucleotide diversity in *C. arabica*, as well in other diploid *Coffea* relatives, we sequenced PCR amplified fragments of nine genes in 20 *Coffea* genotypes: twelve *C. arabica*, including eight wild genotypes and four commercial cultivars; and eight *C. canephora* genotypes. Genotypes of *C. eugenioides*, *C. racemosa* and *Psyllanthus bengalensis* were also included in this analysis. From a total of 9 Kb analyzed, we found 573 polymorphisms: 500 SNPs; 39 INDELs and 34 SSRs. In *C. canephora* genotypes, we detected 188 polymorphisms (frequency of 2.09/100bp). For *C. arabica* we obtained similar results: 144 polymorphism (frequency of 2.13/100bp). Most of the polymorphism found in *C. arabica* only reflected the differences between ancestral homeologs, and they were monomorphic among different genotypes. However, 19 % of these polymorphisms (27 SNPs) were interespecific for *C. arabica*, and 13 of them were fixed among genotypes. The strategy of this work reflects the importance in using a more diverse panel of genotypes in order to identify SNPs in *C. arabica*, pointing out that the exploitation of wild germplasm will be an important source of genetic variability.