



**EMBRAPA**

EMPRESA BRASILEIRA DE PESQUISA AGROPECUÁRIA

CENTRO NACIONAL DE PESQUISA DE MILHO E SORGO

MAR, I.  
1978

TRENDS IN SORGHUM PRODUCTION IN BRAZIL AND RECENT  
RESEARCH ADVANCES AT THE "CENTRO NACIONAL DE  
PESQUISA DE MILHO E SORGO" (CNPMS/EMBRAPA)

Robert E. Schaffert<sup>1</sup>  
Fernando T. Fernandes<sup>2</sup>  
Laudelino C. Leite<sup>2</sup>  
Ivanildo E. Marriel<sup>2</sup>  
Gilson V. E. Pitta<sup>2</sup>  
José M. Waquil<sup>2</sup>

Presented at the "Reunion Internacional de Sorgo", March 6-11, 1978  
BUENOS AIRES - ARGENTINA

00502

<sup>1</sup> Sorghum Project Leader and Sorghum Breeder - CNPMS, IICA/EMBRAPA  
/World Bank, Caixa Postal 151 - 35.700 - Sete Lagoas, MG, Brazil.

<sup>2</sup> Multidisciplinary Research Staff - CNPMS/EMBRAPA, Caixa Postal 151,  
35.700 - Sete Lagoas, MG, Brazil.

TRENDS IN SORGHUM PRODUCTION IN BRAZIL AND RECENT  
RESEARCH ADVANCES AT THE "CENTRO NACIONAL DE  
PESQUISA DE MILHO E SORGO" (CNPMS/EMBRAPA)

I. Sorghum production trends

Sorghum is a relative new crop in Brazil that has developed considerably in recent years. Grain sorghum accounts for about seventy five percent of the acreage planted and forage sorghum accounts for the remainder. Preliminary economic analyses demonstrate that sweet sorghum can be used economically to produce alcohol for mixing with gasoline at current prices, and may soon be planted in large areas near existing sugar mills and distilleries. As sweet sorghum would be harvested when the mills are normally idle little capital investment would be necessary to produce alcohol from sorghum on a large scale.

The data in Table 1 show that grain sorghum acreage increased rapidly during the first half of this decade and since 1975 has decreased slightly. This decrease has been caused principally by lack of adequate handling and storage infrastructure, marketing problems, and inadequate governmental policy for feed grain production and exportation. The current estimated 25 to 30 percent reduction of corn production and feed grain deficiency again demonstrates the need for a healthy sorghum industry in Brazil. A campaign has been initiated in the state of São Paulo to plant grain sorghum during the second semester of this agricultural year to reduce the current grain deficit; however, the problem lies in an adequate seed supply of improved hybrids to meet the demand. Since 1977 the importation of sorghum seed has been prohibited by the Ministry of Agriculture for phytosanitary reasons. Sorghum seed currently being produced is estimated to be only five hundred tons and with little "carry over" will be inadequate to plant the acreage to produce the two to three million tons of feed grain that will be necessary to eliminate the current estimated deficiency that will occur this year.

During the First Brazilian Sorghum Symposium held in 1977 it was shown that by using grain sorghum in the ration industry as only eight percent of the ration, the demand for sorghum by 1980 would be nearly one million tons. This is more than twice the amount of sorghum and about five percent of the corn produced in 1977. The demand for sorghum exists and

with adequate policy by the government and/or the ration industry sorghum acreage should increase significantly by 1980.

TABLE 1. Estimates of area harvested for grain sorghum and grain production in Brazil from 1971 to 1977.

Year	Area harvested (1.000 hectares)	Production (1.000 tons)
1971(1)	80	170
1972(1)	120	220
1973(1)	210	400
1974(1)	250	500
1975(1)	230	483
1976(1)	210	553
1977(2)	178	435

Source: (1) CFP  
(2) SUPLAN - MA

TABLE 2. Grain sorghum production in Brazil by State in 1977

States	Area Harvested (ha)	Production (t)	Average yield (Kg/ha)
Ceará	2,000	1,600	800
Rio Grande do Norte	4,615	3,733	809
Pernambuco	106	152	1,434
Minas Gerais	2,290	2,740	1,200
Espírito Santo	205	615	3,000
São Paulo	56,540	169,620	3,000
Roraima	855	3,470	4,058
Santa Catarina	450	1,320	2,933
Rio Grande do Sul	91,000	214,000	2,352
Mato Grosso	4,583	8,258	1,802
Goiás	15,000	29,625	1,975
Brasil	177,644	435,446	2,444

Source: SUPLAN - MA

Table 2 shows that grain sorghum is concentrated in the states of Rio Grande do Sul and São Paulo, however production in the semi-arid Northeast is growing. In this region sorghum is one of the best options

for the farmers as sorghum is more drought tolerant than the traditionally grown corn and the great deficit of feed grains in this region.

In 1977 the national yield averaged nearly two and one-half tons per hectare. In the state of São Paulo which has experienced a recent growth in area planted the average yield was three tons per hectare while in the semi-arid Northeast the average were much less. National yield trials of both commercial and experimental hybrids and varieties frequently produce more than double the state average.

Nearly all the grain sorghum acreage is planted with hybrids except in the Northeast where some varieties are used. Until this year this seed was imported or produced by a few commercial companies. Next year, however, the majority of the seed available will be from two companies with two or three other companies supplying a small percentage. Because of the uncertainty, at planting time, of regulatory action by the federal government related to the disease "downy mildew", the supply of grain sorghum seed will probably be inadequate next year. Some companies may produce extra seed in the Northeast this winter considering the current feed grain shortage but they will not be able to meet the total demand.

The only forage sorghum seed being commercially produced in Brazil this year is the variety Sart. Seed of some hybrids that have been imported and sold in the past will be unavailable next year.

The National Corn and Sorghum Center released both grain and forage sorghum hybrids in 1977 and will probably release additional hybrids in 1978. The foundation seed will be produced and sold by the Foundation Seed Production Service of EMBRAPA.

## II. Research Activities at the National Research Center

Sorghum research at the CNPMS has developed around specific problem areas and the research program contains several multidisciplinary projects to resolve specific problems, or develop production systems for specific conditions. The research projects currently include the following:

1. Ecological zoning.
2. Development and testing of new production systems.
3. Survey of insects and diseases.
4. Disease control.
5. Insect control.
6. Development of technology to produce sorghum in acid soils.
7. Development of technology to produce sorghum in droughty areas.
8. Improvement of productivity and quality.
9. Improvement in the efficiency of the use of plant nutrients.

10. Mechanization for sorghum.
11. Development of technology for handling and storage of Grain sorghum.
12. Improvement in the efficiency of the use of plant nutrients.
13. Analysis of systems of marketing and use of sorghum.
14. Germoplasm bank activities.

These research projects are summarized in the bulletin "Atividades e Programa do Centro Nacional de Pesquisa de Milho e Sorgo - 1978" and will not be discussed here in detail. Instead I would like to concentrate on some recent advances in priority areas that have been made at the National Corn and Sorghum Research Center.

In tropical and subtropical regions, the soils frequently are characterized by low levels of calcium, magnesium and phosphorus, high acidity and toxic levels of aluminum and/or manganese. In Central Brazil where sorghum and corn acreage is expanding, this is a problem that is not resolved completely or economically by the application of lime and fertilizer. Our objective at the National Corn and Sorghum Research Center is to adapt the plant to this environment instead of only modifying the environment with the addition of lime and fertilizer. After screening several hundred lines in soils with aluminum saturation above sixty percent five lines have been identified as tolerant. These are SC 112-14 (IS 12612), SC 418 (IS 13356 C), SC 048 (IS 12564 C), SC 283 (IS 7173 C) and SC 175-14 (IS 12666 C). In addition two breeding-lines (CMSXS 101 and CMSXS 903) have been identified as being highly susceptible. Preliminary inheritance studies indicate that a single dominant gene is responsible for tolerance to this soil acidity complex. Many progenies of crosses of these lines with other elite materials are currently being evaluated.

Sorghum midge continues to be one of the most serious pests of sorghum and is being studied in great depth at CNPMS. The most animating aspect of this program is the confirmation that the tropical line, AF-28 the most resistant genotype to this pest, tested to date, and a short early selection (In cooperation with researchers at Texas A and M University) of Tx 2536 x AF-28 appears to have maintained this resistance. Panicles of this selection infested with ten adults per day maintained a high degree of seed production. These data are currently being tabulated and analyzed. Several crosses have been made between this selection and elite materials and the F<sub>2</sub> generation is currently being produced in our winter nursery. Techniques are being studied to rear the midge for controlled infestations. Providing that this resistance holds up and is reasonably simply inherited, CNPMS should have resistant varieties and hybrids in the near future.

Symbiotic-biological nitrogen fixation (Azospirillum lipoferum) has been observed in sorghum, millet and corn at CNPMS. Preliminary experiments have shown genetic variability between the sorghum genotypes tested and probably between these three species. Measurements have indicated that sorghum was

fixing nearly 500 g of nitrogen per hectare per day at anthesis. Indirect evidence indicates that pearl millet has much greater nitrogen fixing capacity than sorghum. Sorghum lines selected in 1977 showing greater fixation capacity are currently being reevaluated as well as several pearl millet cultivares.

The most important diseases for sorghum in Central Brazil are anthracnose and rust. Downy mildew is a potential disease as it has been observed in many areas of the states of Rio Grande do Sul and São Paulo. Several elite lines have been identified as tolerant to anthracnose and rust and other foliar diseases and have been used in our breeding program. One line showing much potential is SC 326-6 (IS 3758 C). Several hundred progenies between this line and several other elite lines are currently being evaluated as varieties and in hybrid combination. Several selections with yellow endosperm, open panicle, large seed and good disease resistance are currently being evaluated in hybrid combinations. CNPMS in cooperation with the Secretary of Agriculture of the state of Rio Grande do Sul and the University of São Paulo in Jaboticabal has set up a downy mildew testing service for both commercial and public institutions. In Jaboticabal the infestations this year were much higher in corn than in sorghum. The incidence of the disease in corn varied from zero to one hundred percent while in sorghum the incidence only varied from zero to about twenty percent.

The sorghum breeding program has as its objective to incorporate the various traits mentioned earlier into high yielding productive varieties and hybrids of good quality. In our program special attention is being given to forage quality and nearly half our program will be devoted to the improvement of forage and sweet sorghum types.

Governmental policy to mix alcohol with gasoline has generated a search for new renewable sources of raw materials for fermentation. Sweet sorghum appears to have much potential in this aspect as it can be milled and fermentated when the sugar mills are normally idle and is a crop that can easily be mechanized from planting to harvest with existing equipment. As a result of this, CNPMS has initiated a sweet sorghum breeding and testing program to identify sweet sorghum cultivares adapted to the various regions having potential. The factor most limiting in varieties from the US is the sensibility to short days. In terms of liters of alcohol per hectare per days, sweet sorghum is even competitive with sugar cane.

The National Corn and Sorghum Research Center in cooperation with the National Center for Genetic Resources maintains a Germplasm Bank for corn, sorghum and millet. Introduced genetic materials as well as improve material will be maintained in the germplasm bank. Currently there are over three thousand sorghum lines and several random mating populations from our breeding program that are being catalogued into the germplasm bank.

One other function of CNPMS is to coordinate sorghum research in all of Brazil. CNPMS evaluates all sorghum projects of other institutions (financed by EMBRAPA) and offers technical assistance when necessary.

A network of national grain sorghum, forage sorghum and sweet sorghum trials as well as experimental trials are organized in all of Brazil. In 1977 more than 165 trials were prepared and distributed to cooperators. The breeding program also prepares genetic materials to support the other sorghum breeding programs in Brazil principally those in Pelotas, Rio Grande do Sul (UEPAE-Pelotas/EMBRAPA) and Recife, Pernambuco (IPA-Secretary of Agriculture of Pernambuco). Several lines and varieties as well as 16 bulks were released in 1977. Additional material will be released in 1978.

In summary these are a few of the highlights of sorghum production in Brazil and some research advancements that we at the National Center consider important. Thank you for the opportunity to present this report.

- 2