

Phenotypic plasticity of composite beef cattle performance using reaction norms model with unknown covariate

M. L. Santana Jr^{1,2†}, J. P. Eler², F. F. Cardoso³, L. G. Albuquerque⁴ and J. B. S. Ferraz²

¹Grupo de Melhoramento Animal de Mato Grosso, Instituto de Ciências Agrárias e Tecnológicas, Universidade Federal de Mato Grosso, Campus Universitário de Rondonópolis, MT-270, Km 06, CEP 78735-901, Rondonópolis, MT, Brazil; ²Grupo de Melhoramento Animal e Biotecnologia, Faculdade de Zootecnia e Engenharia de Alimentos, Universidade de São Paulo, C. Postal 23, CEP 13635-970, Pirassununga, SP, Brazil; ³Embrapa Pecuária Sul, BR 153 – Km 603, C. Postal 242, CEP 96401-970, Bagé, RS, Brazil; ⁴Faculdade de Ciências Agrárias e Veterinárias, Universidade Estadual Paulista, CEP 14884-900, Jaboticabal, SP, Brazil

(Received 10 February 2012; Accepted 5 June 2012; First published online 27 September 2012)

The objective of the present study was to determine the presence of genotype by environment interaction ($G \times E$) and to characterize the phenotypic plasticity of birth weight (BW), weaning weight (WW), postweaning weight gain (PWG) and yearling scrotal circumference (SC) in composite beef cattle using the reaction norms model with unknown covariate. The animals were born between 1995 and 2008 on 33 farms located throughout all Brazilian biomes between latitude -7° and -31° , longitude -40° and -63° . The contemporary group was chosen as the environmental descriptor, that is, the environmental covariate of the reaction norms. In general, higher estimates of direct heritability were observed in extreme favorable environments. The mean of direct heritability across the environmental gradient ranged from 0.05 to 0.51, 0.09 to 0.43, 0.01 to 0.43 and from 0.12 to 0.26 for BW, WW, PWG and SC, respectively. The variation in direct heritability observed indicates a different response to selection according to the environment in which the animals of the population are evaluated. The correlation between the level and slope of the reaction norm for BW and PWG was high, indicating that animals with higher average breeding values responded better to improvement in environmental conditions, a fact characterizing a scale of $G \times E$. Low correlation between the intercept and slope was obtained for WW and SC, implying re-ranking of animals in different environments. Genetic variation exists in the sensitivity of animals to the environment, a fact that permits the selection of more plastic or robust genotypes in the population studied. Thus, the $G \times E$ is an important factor that should be considered in the genetic evaluation of the present population of composite beef cattle.

Keywords: composite beef cattle, genetic parameters, genotype by environment interaction, growth, reaction norms

Implications

The response to selection is expected to be different according to the environment in which the animals of the population are evaluated. Genetic variation exists in the sensitivity of animals to the environment, a fact that permits the selection of more plastic or robust genotypes for growth traits in the composite beef cattle studied. The genotype by environment interaction is an important factor that should be considered in the genetic evaluation of animals of the present population.

Introduction

Phenotypic plasticity or environmental sensitivity is the ability of a genotype to alter its phenotypic expression in response to environmental influences (Bradshaw, 1965).

Phenotypic plasticity often characterizes the so-called genotype by environment interaction ($G \times E$). This interaction provokes alterations in genetic, phenotypic and environmental variations that consequently lead to changes in genetic and phenotypic parameter estimates, thus permitting to change selection criteria depending on the environment where the animals are raised and evaluated (Alencar *et al.*, 2005).

There are various approaches to study and detect $G \times E$. One of the most widely used methods is reaction norm, which permits the visualization of the trajectory of animal performance as a function of the environment, thus describing environmental sensitivity (Via and Lande, 1985; Kirkpatrick and Heckman, 1989). Reaction norms are able to indicate where $G \times E$ occurs across the environmental gradient and their magnitude.

A reaction norm can be obtained by random regression on environmental descriptors (Kolmodin *et al.*, 2002; Fikse *et al.*, 2003). The environmental variable is unknown and is defined

† E-mail: 10mario@gmail.com

as the average phenotypic performance of the animals in each environment. Traditionally, one approximation proposed is to compute the mean phenotypic of the trait studied in each environment and then use these estimates as known covariates in the model (Kolmodin *et al.*, 2002 and 2004). According to Su *et al.* (2006), this approach can lead to bias in the estimation of genetic parameters and breeding values. To overcome this problem, these authors proposed the simultaneous estimation of environmental values and other parameters of the model (reaction norm model with unknown covariate), which provided more accurate estimates.

In the case of composite beef cattle, the different breed compositions provide numerous alternative genetic responses to different environments. Few studies have investigated the effect of $G \times E$ on the performance of these animals. Therefore, the objective of the present study was to determine the presence of $G \times E$ and to characterize the phenotypic plasticity of birth weight (BW), weaning weight (WW), postweaning weight gain (PWG) and yearling scrotal circumference (SC) in composite beef cattle using the reaction norms model with unknown covariate.

Material and methods

Data

Data were taken from animals born between 1995 and 2008 on 33 farms located between latitude -7° and -31° , longitude -40° and -63° , in the Brazilian states of Goiás, Minas Gerais, Mato Grosso do Sul, Mato Grosso, Pará, Rio Grande do Sul e São Paulo (Figure S1). The farms participated in the Montana Tropical Composite Breeding Program, CFM-Leachman Pecuária Ltda. These farms were practically located throughout all Brazilian biomes (Amazon, Cerrado, Pantanal, Atlantic Rain Forest and Pampa), which are characterized by wide variations in the conditions of climate, soil and vegetation.

The objective of Montana Tropical Composite Breeding Program is to produce animals with high level of production, fertility and adaptation to the tropical conditions of climate and management using a combination of multiple breeds (*Bos taurus* \times *Bos indicus*). In the present population, the bulls are selected based on an index including standardized Estimated Progeny Differences for BW, WW, PWG, SC and muscle score, weighted -1 , $+3$, $+4$, $+1$ and 1 , respectively. On the basis of the original program of formation of composite beef cattle, that is, the population used in the present study, the breeds were pre-grouped according to their genetic similarity and performance into four large biological types, which are identified by the abbreviation NABC (Ferraz *et al.*, 1999): group N (*B. indicus*) represented by Zebu breeds, which mainly include Guzarat, Indubrasil, Nellore, Tabapuã and other Zebu breeds of African origin such as Boran; group A (*B. taurus*) represented by breeds adapted to tropical climates such as Afrikaner, Belmont Red, Bonsmara, Caracu, Romosinuano and Senepol; group B (*B. taurus*) represented by European breeds of British origin such as Aberdeen Angus, Red Angus, Devon and Hereford; group C (*B. taurus*) represented by European breeds of continental origin such as Charolais, Gelbvieh, Brown Swiss, Simmental, Limousin and others. In this population, an animal (composite beef cattle) should be composed of at least three different breeds, 12.5% adapted breed to the tropical environment and 25% of Zebu breed (N) plus tropically adapted breeds (A). The maximum acceptable values are 37.5% group N, 87.5% group A, 100% group N plus A, 75% for groups B, C and B plus C. Further details about the population studied can be found in Santana *et al.* (2012a) The schematic diagram of the composite beef cattle program is shown in the Figure 1.

The animals were kept on pasture with or without supplements in the dry season. In general, the pastures were composed of *Brachiaria brizantha*. Especially in the Pantanal




Figure 1 Schematic diagram of the composite beef cattle program showing breed compositions (% N = Zebu; % A = tropically adapted taurine; % B = Britain; % C = Continental).

Table 1 Structure of the data set for BW, WW, PWG and SC in composite beef cattle

Item	BW (kg)	WW (kg)	PWG (kg)	SC (cm)
Number of records	118 007	104 899	42 176	19 249
Animals in the pedigree	183 637	174 279	105 408	59 124
Sires	1586	2139	1666	1378
Dams	75 423	55 308	64 847	38 965
Sires with progeny record	348	888	271	216
Dams with progeny record	70 158	48 755	29 397	15 683
Number of CGs	1139	2321	811	399
Mean of the trait	30.90	196.44	87.24	27.29
Standard deviation of the trait	4.05	36.70	45.93	3.95

BW = birth weight; WW = weaning weight; PWG = postweaning weight gain; SC = scrotal circumference; CG = contemporary groups.

(Midwest) and Pampa (southern Brazil), the native pasture plays an important role in animal feeding. All farms provided mineral supplements and some protein. About 60% of cows were inseminated and 40% were placed in lots with a group of bulls. The cow-to-bull ratio was 30 : 1 or 25 : 1. Calves born between September and December (spring in Brazil, the beginning of the rainy period) remained with their dams up to 7 months of age. Weight recordings were obtained at weaning and at yearling (~205 and 420 days of age, respectively). The SC was measured by placing a tape around the scrotum at the point of maximum diameter.

Model and parameter estimation

The statistical models for BW, WW, PWG and SC included the fixed effects of contemporary group (CG; farm, year of birth, management group and sex) and age of animal at recording (linear effect), age of dam at calving (linear and quadratic effects). For PWG, the model included the age of animal at weaning and age of animal at yearling as linear covariates.

All observations were previously adjusted using adjustment factors developed by the Animal Breeding and Biotechnology Group of University of Sao Paulo (GMAB), for maternal breed composition and individual and maternal heterozygosis. Further details on this procedure can be found in Bocchi *et al.* (2008). Records exceeding 3.5 standard deviations below or above the overall mean, CG containing fewer than 20 animals were excluded from the data set. A summary of the data file is shown in Table 1.

The hierarchical reaction norms model with unknown covariate (Su *et al.*, 2006) was used to estimate genetic parameters and study the $G \times E$. In the present study, the CG was chosen as the environmental descriptor, that is, the environmental covariate of the reaction norms. This covariate was jointly estimated with the reaction norms of the animals.

The reaction norm model used can be described as

$$y_{ij} = x'_i \beta + X_j + a_i + b_i X_{(cg)j} + m_i + mp_i + e_{ij},$$

where y_{ij} is the record of animal i in the environment j ; β the vector of fixed effects; x'_i the corresponding incidence vector; $X_{(cg)j}$ the environment random effect [$X_{(cg)j} \sim N(0, \sigma_{cg}^2)$]; a_i the additive genetic value of the intercept

or random level of the animal reaction norm i ; b_i the random regression coefficient or slope of reaction norm of animal i in environment represented by $X_{(cg)j}$; m_i the maternal additive genetic effect (only BW and WW); mp_i the maternal permanent environmental effect (only WW) and e_{ij} the residual effect, $e_{ij} \sim N(0, \sigma_e^2)$. The covariance between direct and maternal effect was set to zero.

The additive genetic variance in the environment X ($\sigma_a^2|X$) was estimated by $\sigma_a^2|X = \text{var}(a_i + b_i X) = \sigma_a^2 + X^2 \sigma_b^2 + 2X \sigma_{ab}$. The direct heritability for a trait with maternal additive genetic and maternal permanent environment effects was obtained by

$$h_a^2 = \frac{\sigma_a^2|X}{\sigma_a^2|X + \sigma_m^2 + \sigma_{mp}^2 + \sigma_{cg}^2 + \sigma_e^2},$$

where $\sigma_a^2|X$, σ_m^2 , σ_{mp}^2 , σ_{cg}^2 and σ_e^2 are variances due to direct additive genetic effect in the environment X , maternal additive genetic effect, maternal permanent environmental effect and CG effect. The maternal heritability (h_m^2) and the proportion of the total phenotypic variance due to maternal permanent environment effect (c^2) were obtained in a similar manner. More details about the hierarchical reaction norms model with unknown covariate are reported by Su *et al.* (2006).

The (co)variance components were obtained using a Bayesian approach through the InterGen program (Cardoso, 2010). Analysis consisted of a single chain of 550 000 cycles, with a conservative burn-in period of 50 000 cycles and a thinning interval of 50 cycles. Thus, 10 000 samples were effectively used to estimate the parameters and highest posterior density intervals. A random sample of animals was used to illustrate the phenotypic plasticity by the reaction norms approach.

Results and discussion

The environmental gradients obtained based on the solutions for the effect of CG ranged from -6.45 to $+4.75$ kg, -65 to $+65$ kg, -72 to $+112$ kg and -6.5 to $+5.5$ cm for BW, WW, PWG and SC, respectively. According to Mattar *et al.* (2011), the CG combines important environmental information that affects animal performance. Therefore, the CG can be naturally defined as environmental descriptor.

Table 2 Components of (co)variance for BW, WW, PWG and SC in composite beef cattle

Item	σ_a^2	σ_{ab}	σ_b^2	σ_e^2	σ_m^2	σ_{mp}^2	σ_{cg}^2	σ_p^2
BW (kg)								
Mean	5.03	0.67	0.10	8.80	0.42	–	2.84	17.08
s.d.	0.20	0.03	0.00	0.11	0.04	–	0.15	3.81
HPD95%	4.65 to 5.45	0.61 to 0.74	0.09 to 0.12	8.56 to 9.02	0.34 to 0.51	–	2.55 to 3.16	12.63 to 24.92
WW (kg)								
Mean	100.86	1.12	0.07	345.60	51.94	68.28	284.14	969.90
s.d.	7.92	0.00	0.00	4.62	4.03	3.75	12.48	138.99
HPD95%	85.83 to 116.62	0.89 to 1.37	0.06 to 0.09	336.35 to 354.68	43.86 to 59.84	60.82 to 75.47	260.28 to 309.26	823.07 to 1317.72
PWG (kg)								
Mean	229.40	2.80	0.03	307.46	–	–	1431.23	2212.89
s.d.	11.78	0.10	0.00	6.21	–	–	76.98	416.27
HPD95%	205.98 to 252.66	2.59 to 3.02	0.03 to 0.04	295.08 to 319.72	–	–	1287.45 to 1587.04	1684.27 to 3075.22
SC (cm)								
Mean	2.26	0.04	0.03	7.17	–	–	5.05	14.93
s.d.	0.30	0.02	0.01	0.23	–	–	0.41	3.80
HPD95%	1.73 to 2.91	–0.01 to 0.09	0.01 to 0.06	6.67 to 7.60	–	–	4.28 to 5.93	13.82 to 16.45

BW = birth weight; WW = weaning weight; PWG = postweaning weight gain; SC = scrotal circumference; σ_a^2 = direct additive genetic variance of intercept; σ_{ab} = additive genetic covariance between intercept and slope; σ_b^2 = direct additive genetic variance of slope; σ_e^2 = residual variance; σ_m^2 = maternal additive genetic variance; σ_{mp}^2 = maternal permanent environmental variance; σ_{cg}^2 = contemporary group variance; σ_p^2 = phenotypic variance; s.d. = standard deviation; HPD95% = high posterior density interval 95%.

Table 3 Direct heritability (h_a^2), maternal heritability (h_m^2), proportion of the total phenotypic variance due to maternal permanent environment effect (C^2) and correlation between intercept and slope of reaction norm (r_{ab}) for traits studied of composite beef cattle

	h_a^2	h_m^2	C^2	r_{ab}
BW				
Mean	0.26	0.02	–	0.93
s.d.	0.15	0.00	–	0.01
HPD95%	0.05 to 0.51	0.01 to 0.03	–	0.90 to 0.95
WW				
Mean	0.21	0.05	0.07	0.40
s.d.	0.09	0.08	0.00	0.03
HPD95%	0.09 to 0.43	0.03 to 0.06	0.05 to 0.08	0.33 to 0.46
PWG				
Mean	0.18	–	–	0.94
s.d.	0.13	–	–	0.00
HPD95%	0.01 to 0.43	–	–	0.92 to 0.96
SC				
Mean	0.18	–	–	0.14
s.d.	0.03	–	–	0.09
HPD95%	0.12 to 0.26	–	–	–0.04 to 0.31

s.d. = standard deviation; HPD95% = high posterior density interval 95%; BW = birth weight; WW = weaning weight; PWG = postweaning weight gain; SC = scrotal circumference.

Heritability

The direct additive genetic component obtained for all traits indicates that these traits should respond to selection (Table 2). The proportion of phenotypic variance due to maternal permanent environmental effects and the maternal additive genetic variance were relatively low (Table 3). These results agree with those reported by Eler *et al.* (2000) for BW, WW and PWG of Nellore cattle and by Santana *et al.* (2010) for PWG and SC using part of the same population of composite cattle.

In general, higher direct heritabilities were observed in extreme favorable environments (Figure 2). Similar results have been reported by Corrêa *et al.* (2009) and Cardoso *et al.* (2011) who observed an increase of heritability for PWG in Devon and Hereford cattle, respectively, as the environmental gradient became more favorable. As seen in the present study for WW and SC, Pégolo *et al.* (2009) obtained higher heritabilities for weight of Nellore cattle at 450 days of age in extreme environments. In a study involving Swedish Red and White dairy cattle, Kolmodin *et al.* 2004 observed an increase of heritability for protein yield and days open with increasing environmental gradient.

Several authors studying the postweaning performance of beef cattle reported changes in genetic parameters across the environmental gradient, with the observation of a higher proportion of phenotypic variation due to genetic factors in more favorable environments (Corrêa *et al.*, 2009; Cardoso *et al.*, 2011; Mattar *et al.*, 2011). Heritability is a property of the population and of the environment to which the animal is exposed, whereas environmental variance depends on production and management conditions. In this respect, greater

environmental variation reduces heritability and a more uniform environment increases heritability (Falconer and Mackay, 1996). The variation in direct heritability observed indicates a different response to selection according to the environment in which the animals of the population are evaluated.

In general, the mean heritability estimates obtained were similar to those reported for Nellore cattle raised under similar conditions of climate and management. In the present study, the mean of direct and maternal heritability for BW was 0.26 and 0.02, respectively (Table 3). Eler *et al.* (2000) found direct and maternal heritabilities for BW of 0.25 and 0.03, respectively, in Nellore cattle using a model that included the effect of sire \times herd interaction. Direct heritabilities for BW of 0.32 and 0.26 have been reported by Albuquerque and Meyer (2001) and Dias *et al.* (2005), respectively, for Nellore cattle. In an extreme favorable environment, the mean direct heritability for BW was 0.51 (Figure 2), a value similar to that reported for cattle raised under temperate climate conditions. Eriksson *et al.* (2004) obtained direct heritabilities of 0.44 to 0.51 and maternal heritabilities of 0.06 to 0.15 for BW in Swedish Hereford and Charolais cattle. The heritability estimates for BW suggest that this trait should respond to mass selection, especially in environments with better management and feed conditions. The mean of direct and maternal heritability for WW was 0.21 and 0.05, respectively. In Nellore cattle, Albuquerque and Meyer (2001) found a direct heritability for WW of 0.14 to 0.16 and a maternal heritability of 0.07 to 0.08. In contrast, Burrow (2001) reported direct and maternal heritabilities of 0.17 and 0.34, respectively, for WW of a composite cattle population. As observed for BW, in a more




Figure 2 Mean of direct heritability (dots) and high posterior density interval 95% (vertical bars) for the traits studied according to the environmental gradient (CG, contemporary group).

favorable environment the phenotype for WW better expressed the genetic potential of the individuals of this population, with a mean maximum direct heritability of 0.41 (Figure 2).

PWG presented a mean direct heritability of 0.01 in the worst environment and of 0.43 in the most favorable environment. According to DeNise *et al.* (1988) and Toral *et al.* (2004), in distinct environments a trait is influenced by sets of different genes and these genes are expressed at variable intensities according to the degree of similarity or difference within and between environments.

The heritability estimates for PWG obtained in the present study were similar to those reported in the literature (0.18 to 0.59; Gregory *et al.*, 1995; Bennett and Gregory, 1996; Marcondes *et al.*, 2000). For SC, mean heritability ranged from 0.15 to 0.23 across the environmental gradient. These estimates are lower than those reported by Vargas *et al.* (1998), Eler *et al.* (2004) and Santana *et al.* (2012b), who found heritabilities ranging from 0.28 to 0.57 for SC of purebred animals. However, similar estimates were obtained by Santana *et al.* (2010) in a study, with part of the present composite cattle raised in the Midwest and Southeast of Brazil. As can be seen in Figure 2, this trait should respond satisfactorily to selection both in unfavorable and in favorable environments. This finding is an interesting opportunity for selection programs designed to achieve genetic progress in this trait in extreme environments.

G × E parameters and phenotypic plasticity

The correlation between the level and slope of the reaction norm for BW and PWG was high, indicating that animals with higher average breeding values responded better to improvement in environmental conditions, a fact characterizing a scale effect of $G \times E$ (Table 3). A high correlation for PWG has also been observed by Corrêa *et al.* (2009) and Cardoso *et al.* (2011) in Devon and Hereford cattle, respectively, and by Calus *et al.* (2002) in Friesian cattle. According to Namkoong (1985), if scale effects differ for traits that are combined in an economic index, the relative importance of each trait may change and result in the re-ranking of animals based on this index.

The correlation between the intercept and slope was low for WW and SC. According to Su *et al.* (2006), this finding implies the re-ranking of animals in different environments, that is, the best animal in one environment is not necessarily the best in another environment. The reaction norms of animals are illustrated in the Figure 3.

According to Falconer (1990), the variation in the reaction norm slope is directly related to the importance of $G \times E$ and reflects environmental sensitivity. Genetic variation in environmental sensitivity of the animals was observed for all traits studied (Table 2). Most animals presented an intermediate sensitivity, that is, the reaction norm slope was close to zero. As a consequence, moderate responses to variations in environmental conditions are expected for the traits studied.




Figure 3 Individual reaction norms according to the environmental gradient (CG, contemporary group) for a random sample of 10 animals of the composite beef cattle population studied.

According to the scale of phenotypic plasticity proposed by Mattar *et al.* (2011), the individuals can be classified as robust genotypes $|b| < \sigma_b$, plastic genotypes $\sigma_b \leq |b| < 2\sigma_b$ and extremely plastic genotypes $|b| \geq 2\sigma_b$. The distribution of animals in the three classes proposed is




Figure 4 Percentage of robust, plastic and extremely plastic animals for birth weight (BW), weaning weight (WW), postweaning weight gain (PWG) and scrotal circumference (SC) in the composite beef cattle population studied.

shown in the Figure 4. A greater number of extreme plastic genotypes were observed for WW, that is, animals with reaction norm slope far from zero. Thus, selection for greater plasticity should be more effective for this trait.

de Jong and Bijma (2002) reported that the phenotypic plasticity must be taken into consideration in the selection of livestock because it offers the possibility to include the environment in the breeding goal. According to Strandberg *et al.* (2000), for most production systems it would be preferable to select animals that present a high level and flat slope of the reaction norm for production traits, that is, more robust animals that show good performance under different environmental conditions. These authors acknowledge that the identification of such animals might be a very difficult task. In reality, no matter how robust a genotype, it will hardly be able to produce a superior phenotype under any condition. According to Stearns (1989), plastic animals might be able to make the genetic alterations necessary according to the environment to which they are exposed. In terms of adaptation, more plastic genotypes may better adapt to drastic environmental changes, whereas robust genotypes may not even be able to survive under certain conditions.

The Spearman correlations between expected breeding values for low (unfavorable), medium (intermediate) or high (favorable) environments and the other environments demonstrated the occurrence of re-ranking of animals, to a greater or lesser extent, according to the environment in which they are evaluated (Figure 5). In general, the correlation between environments was higher for BW and SC than for WW and PWG. The correlation between opposite environments was >0.60 for BW and >0.70 for SC. For WW, the correlation between breeding values of opposite environments reached negative values, whereas a correlation >0.40 was observed for PWG. Strongly negative to positive correlations have been reported by Strandberg *et al.* (2000 and 2009) for productive and reproductive traits of Dairy Nordic and Holstein cattle. The same was observed by Pollott and Greeff (2004) for fecal egg count and six productive traits in Merino sheep. According to Corrêa *et al.* (2009), these correlation estimates indicate that the genetics necessary for production in adverse environments is not the same as that necessary for




Figure 5 Spearman correlations between expected breeding values for birth weight (BW), weaning weight (WW), postweaning weight gain (PWG) and scrotal circumference (SC) of composite beef cattle. The lines indicate the correlation between one environment (low, medium or high) and all other environments.

production in intermediate and favorable environments. These results have major implications for genetic evaluations and selection decisions in the population studied. For example, the correlation between medium and high environments was >0.8 for all traits. In general terms, the best animals in medium environments also show the best performance in high environments. However, this situation changes when the correlation between medium and low environments is considered, which was <0.40 depending on the trait studied. Thus, it is important to indicate the best environment for the animals and what animal should be selected in each environment (de Jong and Bijma, 2002). In this sense, the reaction norms approach should be suitable for this proposal.

Conclusion

A $G \times E$ was observed for all traits studied. The model used here permitted to identify the type and extent of $G \times E$, which differed depending on the trait evaluated. Superior

genotypes in favorable environments may not be the best genotypes in unfavorable environments, especially when WW is considered. Therefore, $G \times E$ is an important factor that should be included in the genetic evaluation of animals of the present population. Genetic variation exists in the sensitivity of animals to the environment, a fact that permits the selection of more plastic or robust genotypes in the population studied. Because of the present composite beef cattle breeding program providing genetic material for a range of production systems, it can be important to include the phenotypic plasticity as breeding goal. The hierarchical reaction norms model with unknown covariate is a valid and interesting option for the genetic evaluation of animals of the population of composite beef cattle studied here.

Acknowledgments

The authors are grateful to CFM-Leachman Pecuária Ltda. for providing the data set. They thank the Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) for the fellowship granted to the first author (2009/05072-4).

Supplementary materials

For supplementary material referred to in this article, please visit <http://dx.doi.org/doi:10.1017/S1751731112001711>

References

- Albuquerque LG and Meyer K 2001. Estimates of covariance functions for growth from birth to 630 days of age in Nelore Cattle. *Journal of Animal Science* 79, 2776–2789.
- Alencar MM, Mascioli AS and Freitas AR 2005. Evidences of genotype \times environment interaction for growth traits in beef cattle. *Brazilian Journal of Animal Science* 34, 489–495.
- Bennett GL and Gregory KE 1996. Genetic (co)variances among birth weight, 200-day weight, and postweaning gain in composites and parental breeds of beef cattle. *Journal of Animal Science* 74, 2598–2611.
- Bocchi AL, Oliveira HN, Ferraz JBS and Eler JP 2008. Multibreed genetic evaluation for pre-weaning daily gain in a composite cattle population. *Brazilian Journal of Animal Science* 37, 1207–1215.
- Bradshaw AD 1965. Evolutionary significance of phenotypic plasticity in plants. *Advances in Genetics* 13, 115–155.
- Burrow HM 2001. Variances and covariances between productive and adaptive traits and temperament in a composite breed of tropical beef cattle. *Livestock Production Science* 70, 213–233.
- Calus MPL, Groen AF and de Jong G 2002. Genotype \times environment interaction for protein yield in Dutch dairy cattle as quantified by different models. *Journal of Dairy Science* 85, 3115–3123.
- Cardoso FF 2010. Application of Bayesian inference in animal breeding using the Intergen program, Manual of Version 1.2., Embrapa Pecuária Sul, Bagé, RS, 30 pp. Retrieved November 29, 2011, from <http://www.cppsul.embrapa.br/unidade/servicos/intergen>
- Cardoso LL, Braccini Neto J, Cardoso FF, Cobuci JA, Biassus IO and Barcellos JOJ 2011. Hierarchical Bayesian models for genotype \times environment estimates in post-weaning gain of Hereford bovine via reaction norms. *Brazilian Journal of Animal Science* 40, 294–300.
- Corrêa MBB, Dionello NJL and Cardoso FF 2009. Genotype by environment interaction characterization and model comparison for post weaning gain adjustment of Devon cattle via reaction norms. *Brazilian Journal of Animal Science* 38, 1468–1477.
- de Jong G and Bijma P 2002. Selection and phenotypic plasticity in evolutionary biology and animal breeding. *Livestock Production Science* 78, 195–214.

- DeNise SK, Torabi M, Ray DE and Rice R 1988. Genetic parameter estimates for preweaning traits of beef cattle in a stressful environment. *Journal of Animal Science* 66, 1899–1906.
- Dias LT, Albuquerque LG, Tonhati H and Teixeira RA 2005. Estimation of genetic parameters for weight in different ages in Tabapuã cattle. *Brazilian Journal of Animal Science* 34, 1914–1919.
- Eler JP, Ferraz JBS, Golden BL and Pereira E 2000. Influence of sire \times herd interaction on the estimation of correlation between direct and maternal genetic effects in Nellore cattle. *Brazilian Journal of Animal Science* 29, 1642–1648.
- Eler JP, Silva JAIV, Evans JL, Ferraz JBS, Dias F and Golden BL 2004. Additive genetic relationships between heifer pregnancy and scrotal circumference in Nellore cattle. *Journal of Animal Science* 82, 2519–2527.
- Eriksson S, Näsholm A, Johansson K and Philipsson J 2004. Genetic parameters for calving difficulty, stillbirth, and birth weight for Hereford and Charolais at first and later parities. *Journal of Animal Science* 82, 375–383.
- Falconer DS 1990. Selection in different environments: effects on environmental sensitivity (reaction norm) and on mean performance. *Genetical Research* 56, 57–70.
- Falconer DS and Mackay TFC 1996. *Introduction to Quantitative Genetics*, 4th edition. Longman Group Ltd, Essex, UK.
- Ferraz JBS, Eler JP and Golden BL 1999. A formação do composto Montana Tropical. *Revista Brasileira de Reprodução Animal* 23, 115–117.
- Fikse WF, Rekaya R and Weigel KA 2003. Assessment of environmental descriptors for studying genotype by environment interaction. *Livestock Production Science* 82, 223–231.
- Gregory KE, Cundiff LV and Koch RM 1995. Genetic and phenotypic (co)variances for growth and carcass traits of purebred and composite populations of beef cattle. *Journal of Animal Science* 73, 1920–1926.
- Kirkpatrick M and Heckman N 1989. A quantitative genetic model for growth, shape, reaction norms, and other infinite-dimensional characters. *Journal of Mathematical Biology* 27, 429–450.
- Kolmodin R, Strandberg E, Danella B and Jorjani H 2004. Reaction norms for protein yield and days open in Swedish red and white dairy cattle in relation to various environmental variables. *Acta Agriculturae Scandinavica* 54, 139–151.
- Kolmodin R, Strandberg E, Madsen P, Jensen J and Jorjani H 2002. Genotype by environment interaction in Nordic dairy studied by use of reaction norms. *Acta Agriculturae Scandinavica* 52, 11–24.
- Marcondes CR, Bergmann JAG, Eler JP, Ferraz JBS, Pereira JCC and Penna VM 2000. Analysis of some selection criteria for growth traits in Nellore cattle. *Arquivo Brasileiro de Medicina Veterinária e Zootecnia* 52, 83–89.
- Mattar M, Silva LOC, Alencar MM and Cardoso FF 2011. Genotype \times environment interaction for long-yearling weight in Canchim cattle quantified by reaction norm analysis. *Journal of Animal Science* 89, 2349–2355.
- Namkoong G 1985. The influence of composite traits on genotype by environment relations. *Theoretical and Applied Genetics* 70, 315–317.
- Pégolo NT, Oliveira HN, Albuquerque LG, Bezerra LAF and Lôbo RB 2009. Genotype by environment interaction for 450-day weight of Nelore cattle analyzed by reaction norm models. *Genetics and Molecular Biology* 32, 281–287.
- Pollott GE and Greeff JC 2004. Genotype \times environment interactions and genetic parameters for fecal egg count and production traits of Merino sheep. *Journal of Animal Science* 82, 2840–2851.
- Santana ML Jr, Eler JP, Cardoso FF, Albuquerque LG, Bignardi AB and Ferraz JBS 2012a. Genotype by environment interaction for birth and weaning weights of composite beef cattle in different regions of Brazil. *Livestock Science* (in press).
- Santana ML Jr, Eler JP, Ferraz JBS and Mattos EC 2012b. Genetic relationship between growth and reproductive traits in Nellore cattle. *Animal* 6, 565–570.
- Stearns SC 1989. The evolutionary significance of phenotypic plasticity. *BioScience* 39, 436–445.
- Strandberg E, Brotherstone S, Wall E and Coffey MP 2009. Genotype by environment interaction for first-lactation female fertility traits in UK dairy cattle. *Journal of Dairy Science* 92, 3437–3446.
- Strandberg E, Kolmodin R, Madsen P, Jensen J and Jorjani H 2000. Genotype by environment interaction in Nordic dairy cattle studied by use of reaction norms. *Interbull Bulletin* 25, 41–45.
- Su G, Madsen P, Lund MS, Sorensen D, Korsgaard IR and Jensen J 2006. Bayesian analysis of the linear reaction norm model with unknown covariates. *Journal of Animal Science* 84, 1651–1657.
- Toral FLB, Silva LOC, Martins EN, Gondo A and Simonelli SM 2004. Genotype \times environment interaction in growth traits of Nellore cattle of Mato Grosso do Sul. *Brazilian Journal of Animal Science* 33, 1445–1455.
- Vargas CA, Elzo MA, Chase CC Jr, Chenoweth PJ and Olson TA 1998. Estimation of genetic parameters for scrotal circumference, age at puberty in heifers, and hip height in Brahman cattle. *Journal of Animal Science* 76, 2536–2541.
- Via S and Lande R 1985. Genotype–environment interaction and the evolution of phenotypic plasticity. *Evolution* 39, 505–522.