June 23-27, 2013 50TH ANNIVERSARY MEETING

ASSOCIATION FOR TROPICAL BIOLOGY AND CONSERVATION & ORGANIZATION FOR TROPICAL STUDIES

ATBC Online Web Program

S17-4

Comparing tree communities of white-sand and terra-firme forests across Amazonian regions

Tuesday, 25 June 2013: 14:50 Americas B-C (Herradura San Jose)

Hans ter Steege, Ecology and Biodiversity Group, Utrecht University, Netherlands Juliana Stropp, Ecology and Biodiversity Group, Utrecht University, Netherlands

leda Amaral , Coordenação de Biodiversidade/Botânica, Instituto Nacional de Pesquisas da Amazônia, Brazil

Gerardo Aymard , Herbario Universitario PORT, UNELLEZ, Venezuela

Carolina Castilho , EMBRAPA CPAFRR, BRAZIL

Cid Ferreira , Coordenação de Biodiversidade/Botânica, Instituto Nacional de Pesquisas da Amazônia, Brazil

Terry Henkel, Department of Biological Sciences, Humboldt State University

Dairon Cardenas Lopez, Sinchi, Colombia

William Ernest Magnusson, Coordenação de Pesquisas em Ecologia, Instituto Nacional de Pesquisas da Amazônia, Brazil Francisca Almeida Matos, Coordenação de Pesquisas em Ecologia, Instituto Nacional de Pesquisas da Amazônia, Brazil

William Milliken , Royal Botanic Gardens, Kew, United Kingdom

Atila Oliveira , Coordenação de Pesquisas em Ecologia, Instituto Nacional de Pesquisas da Amazônia, Brazil

Daniela Pauletto , Serviço Florestal Brasileiro, Santarem, Brazil

Oliver L Phillips, School of Geography, University of Leeds, United Kingdom

Raquel Thomas, Iwokrama International Centre for Rain Forest Conservation and Development, Guyana

Local processes, such as species interaction, environmental sorting, and random dynamics have remained the focus of studies comparing tree alpha- and beta-diversity of white-sand and terra-firme forests in the Amazon. The focus on local processes can be explained by the design of most floristic studies that analyze data from single sites or individual regions. Yet, patterns observed at a local scale in a given region may not apply to other regions as a distinct evolutionary history shaped the current landscape and tree diversity across the Amazon. To date, the large-scale and long-term processes influencing tree diversity of white-sand and terra-firme forests still remain largely unexplored. Here, we address this knowledge gap by analyzing the extent to which tree alpha- and beta-diversity of white-sand and terra-firme forests vary across Amazonian regions. We base our analysis on a dataset containing 93,081 trees (≥ 10 cm DBH) sampled in 192 tree inventory plots established in white-sand and terra-firme forests of three regions: the surroundings of Manaus, the Guianas, and the upper Rio Negro. We found that white-sand forests in the surroundings of Manaus have a higher average tree alpha-diversity (average Fisher's alpha = 56.78) than those in the upper Rio Negro (30.27) and the Guianas (10.89). Additionally, we found that beta-diversity within each region, of both white-sand and terra-firme forests, is lower than between regions. We discuss our results emphasizing the contribution of landscape evolution and habitat size for the current tree alpha- and beta-diversity. The upper

CORE Westers of tree alpha- and beta-diversity across the three regions analyzed here.

See more of: Floristics, Ecology and Evolution of Vegetation in Oligotrophic White-Sand and Sandstone Habitats in the Neotropics See more of: Symposium Proposal

<< Previous Abstract | Next Abstract >>

Start		
Browse		
Browse by Day		
Author Index		

Meeting Information

When:

June 23 - 27, 2013

Where:

San Jose, Costa Rica