



Analytical standards production for the analysis of pomegranate anthocyanins by HPLC

Produção de padrões analíticos para análise de antocianinas da romã por CLAE

Autores | Authors

✉ **Manuela Cristina Pessanha de Araújo SANTIAGO**

Embrapa Agroindústria de Alimentos
Laboratório de Cromatografia Líquida
Avenida das Américas, 29501, Guaratiba
CEP: 23020-470
Rio de Janeiro/RJ - Brasil
e-mail: manuela.santiago@embrapa.br

Ana Cristina Miranda Senna GOUVÊA

Universidade Federal Rural do Rio de Janeiro (UFRRJ)
Programa em Pós-Graduação em Ciência e Tecnologia de Alimentos
Seropédica/RJ - Brasil
e-mail: acristinagouvea@hotmail.com

Ronoel Luiz de Oliveira GODOY Renata Galhardo BORGUINI Sidney PACHECO

Embrapa Agroindústria de Alimentos
Laboratório de Cromatografia Líquida
Rio de Janeiro/RJ - Brasil
e-mail: ronoel.godoy@embrapa.br
renata.borguini@embrapa.br
sidney.pacheco@embrapa.br

Regina Isabel NOGUEIRA

Embrapa Agroindústria de Alimentos
Planta Piloto de Operações Unitárias I
Rio de Janeiro/RJ - Brasil
e-mail: regina.nogueira@embrapa.br

Luzimar da Silva de Mattos do NASCIMENTO

Embrapa Agroindústria de Alimentos
Laboratório de Cromatografia Líquida
Rio de Janeiro/RJ - Brasil
e-mail: luzimar.mattos@embrapa.br

Suely Pereira FREITAS

Universidade Federal do Rio de Janeiro (UFRJ), Departamento de Engenharia Química, Rio de Janeiro/RJ - Brasil
e-mail: freitasp@eq.ufrj.br

✉ Autor Correspondente | Corresponding Author

Recebido | Received: 22/08/2013
Aprovado | Approved: 11/03/2014
Publicado | Published: mar./2014

■ Summary

Pomegranate (*Punica granatum* L.) is a fruit with a long medicinal history, especially due to its phenolic compounds content, such as the anthocyanins, which are reported as one of the most important natural antioxidants. The analysis of the anthocyanins by high performance liquid chromatography (HPLC) can be considered as an important tool to evaluate the quality of pomegranate juice. For research laboratories the major challenge in using HPLC for quantitative analyses is the acquisition of high purity analytical standards, since these are expensive and in some cases not even commercially available. The aim of this study was to obtain analytical standards for the qualitative and quantitative analysis of the anthocyanins from pomegranate. Five vegetable matrices (pomegranate flower, jambolan, jaboticaba, blackberry and strawberry fruits) were used to isolate each of the six anthocyanins present in pomegranate fruit, using an analytical HPLC scale with non-destructive detection, it being possible to subsequently use them as analytical standards. Furthermore, their identities were confirmed by high resolution mass spectrometry. The proposed procedure showed that it is possible to obtain analytical standards of anthocyanins with a high purity grade (98.0 to 99.9%) from natural sources, which was proved to be an economic strategy for the production of standards by laboratories according to their research requirements.

Key words: *Liquid chromatography; Cyanidin; Delphinidin; Pelargonidin; Punica granatum L.*

■ Resumo

Romã (*Punica granatum* L.) é um fruto com um longo histórico medicinal, especialmente devido aos compostos fenólicos presentes em sua composição, como as antocianinas, as quais são relatadas como um dos mais importantes antioxidantes naturais. A análise de antocianinas por Cromatografia Líquida de Alta Eficiência (CLAE) pode ser considerada uma ferramenta importante para avaliar a qualidade do suco de romã. Para os laboratórios de pesquisa, o maior desafio para a análise quantitativa pela técnica de CLAE é a aquisição de padrões analíticos de alta pureza, uma vez que eles são caros e, em alguns casos, não se encontram disponíveis comercialmente. O objetivo deste estudo foi obter padrões analíticos para a análise qualitativa e quantitativa de antocianinas da romã. Cinco matrizes vegetais (flor de romã e frutos de jambolão, jaboticaba, amora e morango) foram usadas para isolar cada uma das seis antocianinas presentes no fruto da romã, usando a escala analítica com detecção não destrutiva, sendo possível usá-las posteriormente como padrões analíticos. Além disso, as suas identidades foram confirmadas pela técnica de espectrometria de massa de alta resolução. O procedimento proposto mostrou que é possível obter padrões analíticos de antocianinas com elevado grau de pureza (98,0%-99,9%) a partir de fontes naturais e provou ser uma estratégia econômica para os laboratórios que necessitam adquirir padrões, de acordo com as necessidades de suas pesquisas.

Palavras-chave: *Cromatografia líquida; Cianidina; Delfinidina; Pelargonidina; Punica granatum L.*

Analytical standards for the analysis of pomegranate anthocyanins by HPLC

SANTIAGO, M. C. P. A. *et al.*

1 Introduction

The pomegranate (*Punica granatum* L.) fruit, native from Iran and endemic in the Middle East, grows in semiarid climates. In recent years there has been renewed interest in the global nutraceutical and functional benefits of this fruit, both fresh and processed (SUMNER *et al.*, 2005).

The pomegranate fruit has some phenolic compounds such as anthocyanins (delphinidin, cyanidin and pelargonidin) in its composition, and also quercetin, phenolic acids and tannins (punicalagin). The fruit is consumed fresh or processed as juice and can be used in the food industry for the manufacture of juice beverages, soft drinks, confectionary products and also colorants (QU *et al.*, 2011).

Due to the increased demand for healthy products, anthocyanin-rich fruits have great potential as raw materials in food formulations, acting as one of the most important natural antioxidants and being responsible for the intense red colour of pomegranate juice based products. The colour is one of the quality parameters that most promotes sensory acceptability by consumers (GIL *et al.*, 2000; ALIGHOURCHI and BARZEGAR, 2009; BOROCHOV-NEORI *et al.*, 2009; PATRAS *et al.*, 2010). Thus, the analysis of anthocyanins by a reliable technique, such as HPLC, can be considered as an important tool to evaluate pomegranate juice quality. Besides the sensory-organoleptic characteristics, knowledge of the anthocyanin profile of pomegranate materials becomes important, since it allows for the identification of adulteration in these products (ZHANG *et al.*, 2009). Thus the analysis of the anthocyanins by a reliable technique can also be considered as an important tool to evaluate pomegranate juice authenticity.

Studies on the characterization and quantification of the phytochemical and antioxidant properties of fruits have become essential and increase awareness about the different cultivars, whether natural or enhanced. Knowledge of the quality and chemical characteristics of some species provides subsidies to distinguish them from each other, and can also provide information to enable an improvement in their genetics, since the concentration and variety of types of the anthocyanins is what will determine the intensity of colouring of the various fruit cultivars (POMAR *et al.*, 2005; ÖZGEN *et al.*, 2009).

The major challenge in the quantitative analysis of compounds by HPLC, especially of anthocyanins, is the obtaining of analytical standards. Standardization is certainly the largest source of analytical errors, because it directly impacts the final result (KIMURA and RODRIGUEZ-AMAYA, 2002). In some countries the acquisition of high purity analytical standards usually depends on highly expensive importations, and in addition

there are no commercial standards available for many of the anthocyanins found in nature.

The production of analytical standards of anthocyanins by isolation is considered to be a challenge, mainly due to the difficulties of obtaining crystalline anthocyanins, free from impurities, in sufficient amounts to allow for reliable weighing (GIUSTI *et al.*, 1999). Thus the aim of this study was to isolate analytical anthocyanin standards from different matrices, using a practical and reliable method known as the analytical HPLC scale with non-destructive detection, for the qualitative and quantitative analysis of pomegranate fruit anthocyanins.

2 Materials and methods

2.1 Chemicals

HPLC grade acetonitrile, 96% formic acid and methanol were purchased from Tedia (USA). Ultrapure water was obtained from the Milli-Q™ Gradient 10A System (Merck Millipore, USA). Delphinidin-3,5-diglucoside chloride was purchased from Chromadex (USA).

2.2 Samples

Samples with the potential for use as sources of the anthocyanins of interest, based on their high content of these compounds, were selected for this study as follows: pomegranate flower, jambolan (peel), jabuticaba (peel), blackberry (whole fruit) and strawberry (whole fruit) (Table 1). Pomegranate flowers were collected in the western region of Rio de Janeiro city. Pomegranate fruits (from the Brazilian semiarid region), strawberry (*Fragaria* spp.), jabuticaba (*Plinia* spp.) and jambolan (*Syzygium cumini*) fruits were purchased in the Rio de Janeiro market. Pomegranate juice was obtained by extraction from the arils. For the blackberry and strawberry juices, the whole fruits were processed in a blender. The jambolan and jabuticaba peels, blackberry and strawberry juices and the pomegranate flowers were freeze-dried in a Liotop™ L101 (Liobras, Brazil) at -40°C for 24 hours, and stored at -18 °C until extraction.

Table 1. Vegetable matrices selected for the isolation of the anthocyanins.

Anthocyanin	Matrix selected
Delphinidin-3,5-diglucoside	Jambolan fruit
Cyanidin-3,5-diglucoside	Jambolan fruit
Delphinidin-3-glucoside	Jabuticaba fruit
Pelargonidin-3,5-diglucoside	Pomegranate flower
Cyanidin-3-glucoside	Blackberry fruit
Pelargonidin-3-glucoside	Strawberry fruit

Analytical standards for the analysis of pomegranate anthocyanins by HPLC

SANTIAGO, M. C. P. A. *et al.*

2.3 Sample extraction

Two grams of each freeze-dried sample were weighed into four centrifuge tubes with lids for extraction with methanol: formic acid (10:90, v/v) under sonication, following by centrifugation (BRITO *et al.*, 2007). All the materials obtained in the supernatants after extraction were concentrated using a Büchi RE rotatory evaporator (Switzerland) at 38 °C for 4 hours. The dried extract was diluted with 4 mL of a 5% formic acid solution in water: methanol (90:10, v/v) and filtered through a hydrophilic type Millex™ membrane (0.45 µm; Merck Millipore; USA) directly into an automatic chromatograph injector vial.

2.4 HPLC-PDA evaluation of the anthocyanins

Chromatographic analysis was carried out following the methodology described by Brito *et al.* (2007), using a Waters (USA) Alliance™ 2695 system equipped with a Waters 2996 photodiode array detector (at 520 nm). A Thermo Scientific C₁₈ BDS (100 mm × 4.6 mm; 2.4 µm; USA) column was used with an injection volume of 20 µL, mobile phase consisting of 5% aqueous formic acid (solvent A) and acetonitrile (solvent B) in the gradient elution mode (Table 2) with a flow rate of 1.0 mL min⁻¹. The column temperature was 40 °C.

2.5 Conditions for the isolation of the anthocyanins by HPLC-PDA

The anthocyanins were isolated following the methodology described by Brito *et al.* (2007) using a Waters (USA) Alliance™ 2695 system equipped with a Waters 2996 photodiode array detector (at 520 nm). A

Table 2. Gradient elution mode for the evaluation of the anthocyanins.

Time (minutes)	Solvent A (%)	Solvent B (%)
0	95	5
2	93	7
10	90	10
12	95	5
14	90	10

Solvent A: 5% aqueous formic acid. Solvent B: acetonitrile

Table 3. Gradient elution mode for isolation of the anthocyanins.

Time (minutes)	Solvent A (%)	Solvent B (%)
0	95	5
2	95	5
10	90	10
15	87	13
16.5	80	20
18	95	5
20	95	5

Solvent A: 5% aqueous formic acid. Solvent B: acetonitrile

Symmetry™ C₁₈ column (150 mm × 4.6; 3.5 µm; Waters; USA) was used with an injection volume of 50 µL, mobile phase consisting of 5% aqueous formic acid (solvent A) and acetonitrile (solvent B), in the gradient elution mode (Table 3) with a flow rate of 1.0 mL min⁻¹. The column temperature was 40 °C.

2.6 Anthocyanin isolation method

The anthocyanins were isolated by liquid chromatography coupled to a Rheodyne six-channel selection valve (GOUVÊA *et al.*, 2012). The valve was adapted to select output channels rather than possible columns, replacing the traditional fraction collector. Substances of interest were collected on elution, according to the retention time of each anthocyanin, using a selector valve commanded by Empower™ software (Waters, USA).

2.7 Concentration of the standards

For the concentration step, a Waters (USA) Sep-Pak™ C₁₈ cartridge previously packed with methanol was saturated with the aqueous extract of the isolated anthocyanins. The cartridge was then washed with an aqueous solution of 0.01% HCl to remove the more polar compounds present, such as sugars and phenolic acids, and the anthocyanin pigments retained were eluted with methanol. The eluted anthocyanin pigments were dried under a filtered compressed air flow, and diluted with a 5% formic acid solution in water: methanol (90:10, v/v) in a 5 mL amber volumetric flask. Aliquots of each isolated anthocyanin were injected under the same chromatographic conditions described above, and the peak areas used to check the purity.

2.8 MS/MS condition

A high resolution Waters mass spectrometer (USA) Synapt™ ESI-QTOF, with direct injection, was used to confirm the identity of the anthocyanins isolated. The MS source used was positive electrospray ionization (ESI⁺) with the following conditions: source temperature at 120 °C, desolvation gas (N₂) delivered at 12.5 L min⁻¹ at 500 °C, capillary exit set at 3.0 kV, sampling cone energy set at 25.0 V and extraction cone energy set at 4.0V.

2.9 Calculation of the concentrations of the standards

200 µL aliquots of the solutions obtained after concentration were collected, dried under a filtered compressed air flow and diluted with 2.0 mL of the appropriate solution for each anthocyanin, according to the molar absorptivity used (Table 4). The corresponding anthocyanin concentration of these solutions was calculated using the Beer-Lambert law from the

Analytical standards for the analysis of pomegranate anthocyanins by HPLC

SANTIAGO, M. C. P. A. *et al.*

absorbance reading obtained using a Shimadzu UV-1800 spectrophotometer (Japan).

2.10 Product evaluation

The anthocyanin profile of the different pomegranate products such as the juice and microcapsules, obtained in the laboratory using a spray drying process, were evaluated to verify the application of the method.

3 Results and discussion

The chromatogram obtained from the pomegranate juice presented six anthocyanins (delphinidin-3,5-diglucoside; cyanidin-3,5-diglucoside; delphinidin-3-glucoside; pelargonidin-3,5-diglucoside; cyanidin-3-glucoside; pelargonidin-3-glucoside) (Figure 1). The same anthocyanins profile was reported by Gil *et al.* (2000) for pomegranate juice.

The six fruits chosen as anthocyanin sources made it possible to isolate these compounds by the HPLC technique with a great grade of purity at 520nm (Figure 2), which is the same wavelength that the anthocyanins are

detected and quantified using the proposed analytical method.

The purity of the isolated anthocyanins was calculated by evaluating the peak area of each one in relation to the total area of the chromatogram. Verification of the purity of the compounds at different wavelengths ensured an accurate calculation of their concentration. The purity of the isolated anthocyanins was also checked at 280 and 360nm, since at those wavelengths occurs the absorption by other compounds that could cause interference, such as other phenolic compounds (GIUSTI *et al.*, 1999). Values above 90% of purity were obtained for all the six compounds evaluated at these two wavelengths.

The high values for purity obtained ensured there were no interfering substances that could cause bathochromic or hypsochromic effects of the ultraviolet/visible absorption spectra. The compounds isolated in the 5% formic acid: methanol (90:10, v/v) solution showed values for concentration that were in an appropriate range to permit the construction

Table 4. Maximum absorption (λ), specific solution used and molar absorptivity of each pomegranate anthocyanin.

Anthocyanins	λ (nm)	Specific solution	Molar Absorptivity
Delphinidin-3,5-diglucoside	520	1% HCl in MeOH	40368*
Cyanidin-3,5-diglucoside	508.5	1% HCl in MeOH	35000**
Delphinidin-3-glucoside	543	1% HCl in MeOH	29000**
Pelargonidin-3,5-Diglucoside diglucoside	510	1% HCl in MeOH	32360**
Cyanidin-3-glucoside	530	1% HCl in MeOH	34300**
Pelargonidin-3-glucoside	496	1% HCl in H ₂ O	31620**

*Molar absorptivity calculated by spectrophotometry using a commercial analytical standard of the anthocyanin, purchased from Chromadex (USA). **Source: Giusti *et al.* (1999).




Figure 1. Chromatogram of the pomegranate juice anthocyanins (peak 1: delphinidin-3,5-diglucoside; peak 2: cyanidin-3,5-diglucoside; peak 3: delphinidin-3-glucoside; peak 4: pelargonidin-3,5-diglucoside; peak 5: cyanidin-3-glucoside; peak 6: pelargonidin-3-glucoside).

Analytical standards for the analysis of pomegranate anthocyanins by HPLC

SANTIAGO, M. C. P. A. et al.




Figure 2. Chromatogram and UV/Vis spectra at 520nm of the isolated anthocyanins: a) delphinidin-3,5-diglucoside; b) cyanidin-3,5-diglucoside; c) delphinidin-3-glucoside; d) pelargonidin-3,5-diglucoside; e) cyanidin-3-glucoside; f) pelargonidin-3-glucoside.

of highly linear calibration curves (Table 5), allowing for the quantification of the matrix evaluated.

All the six anthocyanins isolated could be used as analytical standards, since besides the high grade of purity, the identity of each was confirmed by a reliable technique, such as high resolution mass spectrometry (Table 6).

As described by Müller et al. (2012) the use of single isolated anthocyanin standards better reflects the absolute content of these compounds in a solution than the use of just one anthocyanin to calculate all their concentrations. The same authors found higher anthocyanin contents in blueberries (*Vaccinium corymbosum* L.) and bilberries (*Vaccinium myrtillus* L.), using single isolated standards

Analytical standards for the analysis of pomegranate anthocyanins by HPLC

SANTIAGO, M. C. P. A. *et al.*

Table 5. Data for the external standard calibration curves.

Anthocyanin	Calibration curve data		Concentration range ($\mu\text{g mL}^{-1}$)
	Equation	(R^2)	
Delphinidin-3,5-diglucoside	$Y=9.89 \cdot 10^6 X + 3.45 \cdot 10^4$	0.999	3.20-102.34
Cyanidin-3,5-diglucoside	$Y=8.23 \cdot 10^6 X + 3 \cdot 10^5$	0.989	25.75-103.00
Delphinidin-3-glucoside	$Y=1.09 \cdot 10^7 X + 3.35 \cdot 10^4$	0.995	3.25-13.00
Pelargonidin-3,5-diglucoside	$Y=4.59 \cdot 10^7 X + 4.61 \cdot 10^3$	0.992	0.27-3.58
Cyanidin-3-glucoside	$Y=9.55 \cdot 10^6 X + 2.65 \cdot 10^4$	0.998	1.21-21.21
Pelargonidin-3-glucoside	$Y=4.59 \cdot 10^7 X + 4.61 \cdot 10^3$	0.998	0.27-2.16

Table 6. Identification of the anthocyanins isolated by high resolution mass spectrometry.

Peak	t_R (min)	$[M]^+$ (m/z)	MS/MS (m/z)	Anthocyanin
1	1.8	627.1781	465.1274/303.0672	Delphinidin-3,5-diglucoside
2	2.6	611.1812	449.1287/287.0653	Cyanidin-3,5-diglucoside
3	2.9	465.1115	303.0608	Delphinidin-3-glucoside
4	3.6	595.2229	433.1647/271.1086	Pelargonidin-3,5-diglucoside
5	4.0	449.1287	287.0716	Cyanidin-3-glucoside
6	5.4	433.1424	271.0739	Pelargonidin-3-glucoside

Table 7. Anthocyanin concentrations in raw pomegranate juice and microcapsules obtained by spray drying.

Anthocyanin	Concentration ($\text{mg } 100\text{g}^{-1}$)*	
	Raw juice	Microcapsules
Delphinidin-3,5-diglucoside	8.32 ± 0.45	27.00 ± 0.74
Cyanidin-3,5-diglucoside	10.78 ± 0.40	57.42 ± 1.17
Delphinidin-3-glucoside	0.53 ± 0.01	3.23 ± 0.03
Pelargonidin-3,5-diglucoside	0.49 ± 0.02	1.93 ± 0.18
Cyanidin-3-glucoside	5.50 ± 0.18	20.43 ± 0.56
Pelargonidin-3-glucoside	0.46 ± 0.01	1.49 ± 0.12

*Values expressed as means of two determinations \pm standard deviation.

in the quantification step than by using cyanidin-3-glucoside equivalents.

The proposed method is an important tool for the quality control of pomegranate drinks and other pomegranate products. The possibility of this evaluation is important not only for consumers but also for industry, since it can be used to understand and optimize some of the operational parameters that could influence the stability of the bioactive compounds, in this case, more specifically, of the anthocyanins.

Using the isolated anthocyanins and the calibration curves prepared, it was possible to evaluate the contents of these compounds in different pomegranate products such as juice and the microcapsules obtained by spray drying, as shown in Table 7.

4 Conclusion

The proposed procedure showed that it was possible to obtain analytical anthocyanin standards with a high grade of purity from natural sources, and proved

to be a feasible alternative for laboratories to produce standards according to their research requirements.

Furthermore, these standards could be used in the analysis of other matrices where the same anthocyanins are present.

Acknowledgments

To CNPQ and FAPERJ for their financial support.

References

ALIGHOURCHI, H.; BARZEGAR, M. Some physicochemical characteristics and degradation kinetic of anthocyanin of reconstituted pomegranate juice during storage. **Journal of Food Engineering**, Oxford, v. 90, n. 2, p. 179-185, 2009. <http://dx.doi.org/10.1016/j.jfoodeng.2008.06.019>

BOROCHOV-NEORI, H.; JUDEINSTEIN, S.; TRIPLER, E.; HARARI, M.; GREENBERG, A.; SHOMER, I.; HOLLAND, D. Seasonal and cultivar variations in antioxidant and sensory quality of pomegranate (*Punica granatum* L.) fruit. **Journal**

Analytical standards for the analysis of pomegranate anthocyanins by HPLCSANTIAGO, M. C. P. A. *et al.*

of **Food Composition and Analysis**, San Diego, v. 22, n. 3, p. 189-195, 2009. <http://dx.doi.org/10.1016/j.jfca.2008.10.011>

BRITO, E. S.; ARAUJO, M. C. P.; ALVES, R. E.; CARKEET, C.; CLEVIDENCE, B. A.; NOVOTY, J. A. Anthocyanins present in selected tropical fruits: acerola, jambolão, jussara and guajiru. **Journal of Agriculture and Food Chemistry**, Washington, v. 55, n. 23, p. 9389-9394, 2007. PMID:17929888. <http://dx.doi.org/10.1021/jf0715020>

GIL, M.; TOMAS-BARBERAN, F. A.; HESS-PIERCE, B.; HOLCROFT, D. M.; KADER, A. A. Antioxidant activity of pomegranate juice and its relationship with phenolic composition and processing. **Journal of Agricultural and Food Chemistry**, Washington, v. 48, n. 10, p. 4581-4589, 2000. <http://dx.doi.org/10.1021/jf000404a>

GIUSTI, M. M.; RODRÍGUEZ-SAONA, L. E.; WROLSTAD, R. E. Molar absorptivity and color characteristics of acylated and non-acylated pelargonidin-based anthocyanins. **Journal of Agricultural and Food Chemistry**, Washington, v. 47, n. 11, p. 4631-4637, 1999. <http://dx.doi.org/10.1021/jf981271k>

GOUVÊA, A. C. M. S.; SANTIAGO, M. C. P. A.; PACHECO, S.; GODOY, R. L. O.; CABRAL, L. M. C. Anthocyanins standards (cyanidin-3-O-glucoside and cyanidin-3-O-rutenosideo) isolation from freeze-dried açai (*Euterpe oleraceae* Mart.) by HPLC. **Ciência e Tecnologia de Alimentos**, Campinas, v. 32, n. 1, p. 43-46, 2012. <http://dx.doi.org/10.1590/S0101-20612012005000001>

KIMURA, M.; RODRIGUEZ-AMAYA, D. B. A Scheme for obtaining standards and hplc quantification of leafy vegetable carotenoids. **Food Chemistry**, New York, v. 78, n. 3, p. 389-398, 2002. [http://dx.doi.org/10.1016/S0308-8146\(02\)00203-0](http://dx.doi.org/10.1016/S0308-8146(02)00203-0)

MÜLLER, D.; SCHANTZ, M.; RICHLING, E. High performance liquid chromatography analysis of anthocyanins in bilberries (*Vaccinium myrtillus* L.), blueberries (*Vaccinium corymbosum* L.),

and corresponding juices. **Journal of Food Science**, Chicago, v. 77, n. 4, p. 340-345, 2012. PMID:22394068. <http://dx.doi.org/10.1111/j.1750-3841.2011.02605.x>

ÖZGEN, M.; SERÇE, S.; KAYA, C. Phytochemical and antioxidant properties of anthocyanin-rich morus nigra and morus rubra fruits. **Scientia Horticulturae**, Kent, v. 119, n. 3, p. 275-279, 2009. <http://dx.doi.org/10.1016/j.scienta.2008.08.007>

PATRAS, A.; BRUNTON, N. P.; O'DONNELL, C.; TIWARI, B. K. Effect of thermal processing on anthocyanin stability in foods: mechanisms and kinetics of degradation. **Trends in Food Science and Technology**, Cambridge, v. 21, n. 1, p. 3-11, 2010. <http://dx.doi.org/10.1016/j.tifs.2009.07.004>

POMAR, F.; NOVO, M.; MASA, A. Varietal differences among the anthocyanin profile of 50 red table grape cultivars studied by high performance liquid chromatography. **Journal of Chromatography A**, Amsterdam, v. 1094, n. 1-2, p.34-41, 2005.

QU, W.; BREKSA III, A. P.; PAN, Z.; MA, H. Quantitative determination of major polyphenol constituents in pomegranate products. **Food Chemistry**, New York, v. 132, n. 3, p. 1585-1591, 2011. <http://dx.doi.org/10.1016/j.foodchem.2011.11.106>

SUMNER, M. D.; ELLIOTT-ELLER, M.; WEIDNER, G.; DAUBENMIER, J. J.; CHEW, M. H.; MARLIN, R. Effects of pomegranate juice consumption on myocardial perfusion in patients with coronary heart disease. **Journal of Cardiology**, Japan, v. 96, n. 6, p. 810-814, 2005. PMID:16169367. <http://dx.doi.org/10.1016/j.amjcard.2005.05.026>

ZHANG, Y.; WANG, D.; LEE, R. P.; HENNING, S. M.; HEBER, D. California hass avocado: profiling of carotenoids, tocopherol, fatty acid, and fat content during maturation and from different growing areas. **Journal of Agricultural and Food Chemistry**, Washington, v. 57, n. 21, p. 7395-7400, 2009. PMID:20349921. <http://dx.doi.org/10.1021/jf9010017>