

RESEARCH ARTICLE

Open Access

Regulatory *nodD1* and *nodD2* genes of *Rhizobium tropici* strain CIAT 899 and their roles in the early stages of molecular signaling and host-legume nodulation

Pablo del Cerro^{1†}, Amanda Alves Paiva Rolla-Santos^{2†}, Douglas Fabiano Gomes², Bettina Berquó Marks², Francisco Pérez-Montaño¹, Miguel Ángel Rodríguez-Carvajal³, André Shigueyoshi Nakatani², Antonio Gil-Serrano³, Manuel Megías¹, Francisco Javier Ollero¹ and Mariangela Hungria^{2*}

Abstract

Background: Nodulation and symbiotic nitrogen fixation are mediated by several genes, both of the host legume and of the bacterium. The rhizobial regulatory *nodD* gene plays a critical role, orchestrating the transcription of the other nodulation genes. *Rhizobium tropici* strain CIAT 899 is an effective symbiont of several legumes—with an emphasis on common bean (*Phaseolus vulgaris*)—and is unusual in carrying multiple copies of *nodD*, the roles of which remain to be elucidated.

Results: Phenotypes, Nod factors and gene expression of *nodD1* and *nodD2* mutants of CIAT 899 were compared with those of the wild type strain, both in the presence and in the absence of the *nod*-gene-inducing molecules apigenin and salt (NaCl). Differences between the wild type and mutants were observed in swimming motility and IAA (indole acetic acid) synthesis. In the presence of both apigenin and salt, large numbers of Nod factors were detected in CIAT 899, with fewer detected in the mutants. *nodC* expression was lower in both mutants; differences in *nodD1* and *nodD2* expression were observed between the wild type and the mutants, with variation according to the inducing molecule, and with a major role of apigenin with *nodD1* and of salt with *nodD2*. In the *nodD1* mutant, nodulation was markedly reduced in common bean and abolished in leucaena (*Leucaena leucocephala*) and siratro (*Macroptilium atropurpureum*), whereas a mutation in *nodD2* reduced nodulation in common bean, but not in the other two legumes.

Conclusion: Our proposed model considers that full nodulation of common bean by *R. tropici* requires both *nodD1* and *nodD2*, whereas, in other legume species that might represent the original host, *nodD1* plays the major role. In general, *nodD2* is an activator of *nod*-gene transcription, but, in specific conditions, it can slightly repress *nodD1*. *nodD1* and *nodD2* play other roles beyond nodulation, such as swimming motility and IAA synthesis.

Keywords: *nodD* gene, Nod factors, Nodulation, Symbiosis, Nitrogen fixation, *Rhizobium tropici*

Background

Bacteria commonly known as rhizobia are capable of establishing symbioses with several leguminous species, forming specific structures, the root nodules, where the process of biological fixation of atmospheric nitrogen takes place, bringing important contributions to agriculture and

to the environment [1-3]. Legume nodulation requires a cascade of molecular signals exchanged between the host plant and the rhizobium. This molecular dialogue begins with the exudation of flavonoids from the legume, which are recognized by the bacterium. When induced by these plant molecules, rhizobia synthesize lipochitooligosaccharides (LCOs), also known as Nod factors, responsible for launching the nodulation process [3-8]. It is noteworthy that an increasing number of reports show that Nod factors may play roles beyond the nodulation process, including

* Correspondence: mariangela.hungria@embrapa.br

†Equal contributors

²Embrapa Soja, C.P. 231, 86001-970 Londrina, Paraná, Brazil

Full list of author information is available at the end of the article

stimulation of photosynthesis, improvements in plant growth and grain yield and changes in immune responses in both legumes and non-legumes [9-12].

We consider *nodD* as the most intriguing regulatory nodulation gene; it belongs to the LysR family of transcription regulators, and it is constitutively expressed and responsible for the transcription of other nodulation genes in the presence of suitable plant inducers, usually flavonoids, thus initiating the nodulation process [8,13,14]. Furthermore, many other symbiosis-related phenotypes, such as polysaccharide production, phytohormone synthesis, motility, quorum-sensing and the activation of the type-III secretion system are directly or indirectly regulated by means of inducing flavonoids via NodD in rhizobia [15-20]. Studies of genomes of rhizobia indicate that, depending on the rhizobial species, there are one to five copies of *nodD*. In the species that possess only one copy of this gene, such as *Rhizobium leguminosarum* bv. trifolii, a mutation usually results in loss of nodulation, whereas, in the presence of multiple copies, as in *Sinorhizobium* (= *Ensifer*) *meliloti*, *Rhizobium leguminosarum* bv. phaseoli and *Bradyrhizobium japonicum*, an intricate interaction between the *nodD* genes seems to occur and the nodulation is not completely suppressed [21-23].

Rhizobium tropici strain CIAT 899 is an effective microsymbiont of common bean (*Phaseolus vulgaris* L.) in the tropical acid soils of South America. Notable properties of this strain are its high tolerance of environmental stresses and its broad legume host-range [24-26]. Another intriguing feature of CIAT 899 is its capacity for producing a large variety of Nod factors [27,28]. Interestingly, this bacterium is able to produce these key symbiotic molecules under abiotic stresses, such as acid and saline conditions, in the absence of plant-molecular signals [28-30]. In a pioneering study, five distinct *nodD*-hybridizing DNA regions were identified in CIAT 899 [31], later confirmed as five *nodD* genes in the sequenced genome of the strain [32]. The *nodD1* gene preceding the *nodABC* operon seems to play the major role in nodulation [31], but a more precise study of the regulatory functions of *nodD1* in *R. tropici* is lacking. In addition, the role of *nodD2*, present in some rhizobial species, is unclear. It has been reported to be a repressor of the *nodABC* operon, leading to a negative effect on Nod-factor production in *Sinorhizobium* (= *Ensifer*) *fredii* strain NGR 234 [33]. A suppressive role has also been observed in *B. japonicum* [23] and a negative regulation by NodD2 products was reported in *Bradyrhizobium* (*Arachis*) [34].

Here we report a study in which phenotypes, Nod factors and gene expression of mutants of nodulation genes *nodD1* and *nodD2* of strain CIAT 899 were compared with those of the wild type strain, to help to elucidate the roles of these regulatory genes.

Results and discussion

Phenotypic characterization of wild type and mutant strains

As defined in the genome of *R. tropici* strain CIAT 899 [32], *nodD1* precedes the *nodABC* operon, while *nodD2*, corresponding to *nodD5* described by van Rhijn et al. [31], is adjacent to the *nodA2* and *hsnT* genes (Additional file 1: Figure S1). *R. tropici* CIAT 899 *nodD1* mutant was obtained in a previous work by insertion of a Km^R cassette into a unique *XhoI* restriction site located on the gene [30]. As described in the Material and Methods section, the *nodD2* mutant was obtained after deletion of a 0.6 kb *PstI* fragment of the gene and the insertion in its place of the Ω interposon (Additional file 1: Figure S1).

Growth rate was not affected by mutation in *nodD1* or *nodD2* genes of *R. tropici* CIAT 899 (data not shown). However, it is known that some bacterial properties may be regulated via NodD proteins, such as EPS (exopolysaccharide) production, LPS (lipopolysaccharide) profiles, swimming and swarming motilities, biofilm formation and IAA (indole acetic acid) synthesis, among others (e.g., [15-20]). We evaluated some these properties in the wild type and mutant strains in the presence or absence of two *nod*-gene inducing molecules, apigenin (3.7 μ M) and salt (NaCl 300 mM). Results showed statistical differences only in swarming motility (Figure 1) and in the production of IAA (Figure 2).

Swarming motility is caused by rotation of single or multiple flagellae along wetted surfaces [35] and, in certain rhizobia, is altered in the presence of legume root exudates that are rich in flavonoids [36]. Our experiments showed that, under control conditions, both *nodD* mutants showed more swarming motility than did the wild type strain. Interestingly, these differences were even stronger when the medium was augmented with apigenin for the *nodD1* mutant and with NaCl for the *nodD2* mutant (Figure 1). Therefore, the results suggest a constitutive suppression of swarming by NodD1 and NodD2 proteins.

IAA is an essential plant hormone that promotes growth, including lateral-root proliferation. Previous work has demonstrated that synthesis of this molecule is regulated by NodD1 and NodD2 in *S. fredii* strain NGR234 [16]. In CIAT 899, our experiments showed an increase in the production of IAA in the presence of apigenin and NaCl (Figure 2), suggesting that both inducing molecules promote the synthesis of this phytohormone. This finding is supported by the presence of a *nod*-box upstream of the IAA operon in the genome of CIAT 899 [32]. In addition, NodD1 seems to be the main regulator in the presence of apigenin, since, in this mutant, the production of IAA was significantly lower than in the presence of NaCl. The production of IAA in the *nodD2* mutant was strongly reduced when the medium was supplemented with NaCl (Figure 2),




Figure 1 Swarming motility phenotype of the *R. tropici* CIAT 899 wild type and *nodD1* and *nodD2* mutants. Quantified swarm ring diameters of wild type strain (continuous line), the *nodD1* mutant (striped line) and the *nodD2* mutant (dotted line). Values are the averages of three swarm plates per strain. *nodD1* and *nodD2* mutant parameters were individually compared with the parental strain CIAT 899 parameters by using the Mann–Whitney non-parametric test. Values tagged by * are significantly different at the level $\alpha = 5\%$. Swarming motility in: **A.** TY medium, **B.** TY medium supplemented with 3.7 μM of apigenin, and **C.** TY medium supplemented with 300 mM of NaCl.

suggesting that this regulator may be mainly implied in the activation of the IAA operon in the presence of salt. Altogether, the results indicate a predominant role of NodD1 in activation of the IAA gene by apigenin and a predominant role of NodD2 when the inducing molecule is NaCl.

The nodulation phenotype in common bean was first evaluated in pouches bags, where it was possible to observe that a mutation in *nodD1* caused a significant decrease in nodule number of common bean; to a lesser extent, a decrease was also observed with a mutation in *nodD2* (Additional file 2: Figure S2). In both leucaena [*Leucaena leucocephala* (Lam.) de Wit] and siratro [*Macroptilium atropurpureum* (DC.) Urb.], no nodules

were observed when plants were inoculated with the *nodD1* mutant, but apparently no differences were observed when plants of both species were inoculated with the *nodD2* mutant in comparison with plants inoculated with the wild type strain (data not shown).

Nodulation of the type and mutants was confirmed by growing plants in larger pots, in Leonard jars containing sterile substrate. In common bean, a mutation in *nodD1* did not suppress nodulation, but caused a reduction of 82% in nodule number (Table 1). The absence of nodulation in both leucaena and siratro when inoculated with the *nodD1* mutant was confirmed. *Vis-à-vis* the *nodD2* mutant, nodulation of common bean was reduced by 55%, and no statistical differences in relation to the wild




Figure 2 Indole-3-acetic acid (IAA) relative production by *R. tropici* CIAT 899 wild type, and by the *nodD1* and *nodD2* mutants. Bacteria were grown in TY medium containing tryptophan in absence and presence of apigenin (3.7 μ M) or NaCl (300 mM). Supernatants were taken 96 h after the addition of flavonoid or salt. IAA production was calculated relative to the production without inducing molecules of the wild type strain by using the Mann–Whitney non-parametrical test. The asterisks indicate a significant different at the level $\alpha = 5\%$. Black bars: CIAT 899. Light gray bars: *nodD1* mutant. Dark gray bars: *nodD2* mutant.

type strain were observed in the nodulation of either leucaena or siratro. However, shoot dry weight of the leucaena plants inoculated with the *nodD2* mutant was lower than with the wild type ($\alpha = 10\%$) (Table 1).

In rhizobial species with more than one copy of the regulatory *nodD* gene, *nodD1* preceding the *nodABC* operon has been recognized as the main gene regulating nodulation e.g. [23,33,34,37–39]. However, reports show that the role of each *nodD* copy, their responses to flavonoids, and the nodulation phenotypes vary on a case-by-case basis with the rhizobium strain and the host-plant species/cultivar. An intricate pattern of responses in nodulation leads to the assumption that *S. meliloti* utilizes the three copies of *nodD* to optimize the interaction with each of its legume hosts [37,40]. A mutation in *nodD1* of *S. meliloti* delays but does not eliminate nodulation of both alfalfa (*Medicago sativa*) and sweet clover (*Melilotus alba*), and only a triple mutation of *nodD1-nodD2-nodD3* results in absence of nodules [37]. Contrarily, in the promiscuous strain *S. fredii* NGR 234,

capable of nodulating more than 110 plant species, a mutation in *nodD1* abolishes nodulation in several temperate and tropical species [41]. In addition, in *B. japonicum nodD1* is sufficient for nodulation of the putative main host plant, soybean (*Glycine max*), but the additional genes *nodVW* are required for the nodulation of mung bean (*Vigna radiata*), cowpea (*Vigna unguiculata*) and siratro [39,42]. In our study, a mutation in *nodD1* decreased, but did not suppress, nodulation of common bean; however, *nodD1* proved to be essential for the nodulation both of leucaena and of siratro (Table 1).

Still considering nodulation phenotype, in *S. meliloti* the *nodD2* gene did not have any apparent effect on nodulation of either alfalfa or sweet clover [37]. Similarly, no detectable effects were observed by inoculating siratro and cowpea with a *nodD2* mutant of *Bradyrhizobium* (*Arachis*) sp. strain NC92 [34]. Contrarily, in our study, a significant decrease in nodulation of common bean was detected with the mutation in *nodD2*, but no effects were observed in leucaena and siratro (Table 1).

Table 1 Nodule number ($n^{\circ} \text{plant}^{-1}$) and shoot dry weight (g plant^{-1}) of common bean, leucaena and siratro inoculated with *R. tropici* strain CIAT 899 and *nodD* derivatives

Strains	<i>P. vulgaris</i> ^a		<i>L. leucocephala</i> ^a		<i>M. atropurpureum</i> ^a	
	Nodule number	Shoot dry weight	Nodule number	Shoot dry weight	Nodule number	Shoot dry weight
<i>R. tropici</i> CIAT899	213 \pm 52	1.82 \pm 0.64	13 \pm 4	0.41 \pm 0.03	34 \pm 8	0.05 \pm 0.01
<i>nodD1</i> mutant	38 \pm 11*	1.42 \pm 0.35	0 \pm 0*	0.09 \pm 0.01*	0 \pm 0*	0.05 \pm 0
<i>nodD2</i> mutant	95 \pm 38*	1.03 \pm 0.27	10 \pm 3	0.36 \pm 0.04**	24 \pm 8	0.05 \pm 0
none	0 \pm 0*	0.80 \pm 0.25*	0 \pm 0*	0.09 \pm 0.01*	0 \pm 0*	0.05 \pm 0

^aData represent means \pm SD (standard deviation) of six jars, each with two plants. *nodD1* and *nodD2* mutant parameters were individually compared with the parental strain CIAT 899 parameters by using the Mann–Whitney non-parametric test. Values tagged by *and **are significantly different at the level $\alpha = 5$ and 10%, respectively.

Plants evaluated after 25 (common bean) or 42 days (leucaena and siratro) of growth under controlled conditions.

Nod factor patterns

Rhizobium tropici strain CIAT 899 is known as an interesting strain in relation to its production of a large variety of Nod factors, not only when induced by flavonoids [27,28], but also under high-salinity conditions in the absence of flavonoids [28-30].

A list of all Nod factors detected in the wild type strain in comparison to the *nodD1* and *nodD2* mutants is shown in Tables 2, 3 and 4. Unexpectedly, Nod factors were found in the B⁻ medium [43], even in the absence of inducer molecules. In this condition, around ten Nod factors were synthesized, with no significant differences among wild type CIAT 899, *nodD1* and *nodD2* mutants (Table 2). When induced by 3.7 μM apigenin, the synthesis of a variety of Nod factors was confirmed in all strains, such that numerically, 29 Nod factors were detected in the wild type CIAT 899 and 25 in the *nodD2* mutant; a slight reduction was observed with the *nodD1* mutant, but, even then, 20 Nod factors were observed (Table 3). This number is higher than in other wild type rhizobial species, e.g. the four Nod factors identified in *B. japonicum* strain USDA 138 [44]. Up to 36 Nod factors were found in CIAT 899 under saline conditions (Table 4), and in the *nodD1* and *nodD2* mutants the numbers were lower, of 20 and 18 Nod factors, respectively. These results indicate that NaCl has a stronger *nod*-induction capacity than apigenin does, and that it is affected by *nodD2* but not *nodD1*. However, one might also consider that it deserves further studies to investigate the possibility that Nod factors are more stable in a 300 mM NaCl supplemented medium.

The production of a large number of Nod factors in all conditions tested might be related to broad host promiscuity and abiotic-stress tolerance of *R. tropici* [24-26]. The promiscuous *S. fredii* strain NGR 234 also produces a larger number of Nod factors (≥18) [41], and the composition of Nod factors produced by this strain varies with the activity of host-specific nodulation genes [45]. Furthermore, one interesting feature observed in our study was that the Nod factors with structure III (C_{18:1}, NMe), IV (C_{18:1}, NMe), IV (C_{18:0}, NMe), V (C_{18:0}, NMe) and V (C_{16:0}, NMe, S) were present in the wild type and in the *nodD2* mutant, but not in the *nodD1* mutant; therefore, this structure might be implicated in host-specific nodulation, and could explain why the mutant in the *nodD1* gene is unable to induce nodules on leucaena or siratro. It is also worth mentioning that Folch-Mallol *et al.* [46] described that in CIAT 899 the sulfation of the LCOs, mediated by the *nodHPQ* genes are important for nodulation efficiency on *L. leucocephala*. A mutant in the *nodH* gene induced about half of nodules than those induced by the wild type strain [46]. Interestingly, one of the five LCOs not synthesized by the CIAT 899 *nodD1* mutant is sulphated [V (C_{16:0}, NMe, S)] (Tables 3 and 4) and may be important for nodulation on leucaena. However, because the *nodD1* mutant is unable to nodulate leucaena, other LCOs not secreted by this mutant must be important to explain its symbiotic phenotype.

In *R. tropici*, the amount and diversity of Nod factors produced are directly influenced by the conditions of bacterial growth. Our results are consistent with the

Table 2 Nod factor structure biosynthesized in control condition (B⁻ medium) by wild type CIAT 899 and the *nodD1* and *nodD2* mutants

[M + H] ⁺ (m/z)	B _i ions	Structure ^a	CIAT899 ^b	<i>nodD1</i> ^b	<i>nodD2</i> ^b
850	426, 629, 832	III (C _{18:1})	+	+	-
1027	400, 603, 806	IV (C _{16:0})	+	+	+
1041	414, 617, 820	IV (C _{16:0} , NMe)	-	-	+
1053	426, 629, 832	IV (C _{18:1})	+	+	+
1055	428, 631, 834	IV (C _{18:0})	+	-	-
1067	440, 643, 846	IV (C _{18:1} , NMe)	+	+	+
1216	386, 589, 792, 995	V (C _{14:0} , NMe)	+	-	+
1230	400, 603, 806, 1009	V (C _{16:0})	+	+	+
1244	414, 617, 820, 1023	V (C _{16:0} , NMe)	+	+	+
1256	426, 629, 832, 1035	V (C _{18:1})	+	+	+
1270	440, 643, 846, 1049	V (C _{18:1} , NMe)	+	+	+
1350	440, 643, 846, 1049, [M-80] ^{+c} = 1270	V (C _{18:1} , NMe, S)	+	+	+
1352	442, 645, 848, 1051, [M-80] ^{+c} = 1272	V(C _{18:0} , NMe, S)	-	+	-
1378	468, 671, 874, 1077, [M-80] ^{+c} = 1298	V (C _{20:1} , NMe, S)	-	+	-

^aNF structures are represented following the convention (Spaink, 1992) [43] that indicates the number of GlcNAc residues in the backbone (Roman numeral), the length and degree of unsaturation of the fatty acyl chain, and the other substituents, which are listed in the order in which they appear, moving clockwise from the fatty acid. NMe, N-methyl group at glucosamine non reducing residue; S, sulfate group at reducing glucosamine residue.

^bSymbol: + = detected; - = non detected.

^cThese ions arise by loss of a neutral with mass 80 Da, corresponding to the loss of SO₃.

Table 3 Nod factor structure biosynthesized in the presence of apigenin (3.7 μM) by wild type CIAT899 and the *nodD1* and *nodD2* mutants

[M + H] ⁺ (m/z)	B _i ions	Structure ^a	CIAT899 ^b	<i>nodD1</i> ^b	<i>nodD2</i> ^b
810	386, 589	III (C _{14:0} , NMe)	-	-	+
824	400, 603	III (C _{16:0})	-	+	+
838	414, 617	III (C _{16:0} , NMe)	+	+	+
850	426, 629	III (C _{18:1})	+	+	+
852	428, 631	III (C _{18:0})	+	-	-
864	440, 643	III (C _{18:1} , NMe)	+	-	+
999	372, 575, 778	IV (C _{14:0})	+	+	-
1011	384, 597, 790	IV (C _{14:1} , NMe)	+	-	-
1013	386, 589, 792	IV (C _{14:0} , NMe)	+	+	+
1025	398, 601, 804	IV (C _{16:1})	+	+	+
1027	400, 603, 806	IV (C _{16:0})	+	+	+
1039	412, 615, 818	IV (C _{16:1} , NMe)	+	+	+
1041	414, 617, 820	IV (C _{16:0} , NMe)	+	+	+
1053	426, 629, 832	IV (C _{18:1})	+	+	+
1055	428, 631, 834	IV (C _{18:0})	+	+	+
1067	440, 643, 846	IV (C _{18:1} , NMe)	+	-	+
1069	442, 645, 848	IV (C _{18:0} , NMe)	+	-	+
1081	454, 657, 860	IV (C _{20:1})	+	-	-
1147	440, 643, 846	IV (C _{18:1} , NMe, S)	-	-	+
1202	372, 575, 778, 981	V (C _{14:0})	+	+	-
1214	426, 629, 790, 832, 993 ^d	V (C _{18:1} , dNAC)	+	-	-
1216	386, 589, 792, 995	V (C _{14:0} , NMe)	+	+	+
1228	440, 643, 846, 1007 ^e	V (C _{18:1} , NMe, dNAC)	+	-	-
1230	400, 603, 806, 1009	V (C _{16:0})	+	+	-
1231	440, 643, 846, 1049	IV Hex-ol (C _{18:1} , NMe)	-	-	+
1242	412, 615, 818, 1021	V (C _{16:1} , NMe)	+	+	+
1244	414, 617, 820, 1023	V (C _{16:0} , NMe)	+	+	+
1256	426, 629, 832, 1035	V (C _{18:1})	+	+	+
1270	440, 643, 846, 1049	V (C _{18:1} , NMe)	+	+	+
1272	442, 645, 848, 1051	V (C _{18:0} , NMe)	+	-	+
1284	454, 657, 860, 1063	V (C _{20:1})	-	-	+
1324	414, 617, 820, 1023	V (C _{16:0} , NMe, S)	+	-	+
1336	426, 629, 832, 1035	V (C _{18:1} , S)	+	+	-
1350	440, 643, 846, 1049, [M-80] ^{+c} = 1270	V (C _{18:1} , NMe, S)	+	+	+

^aNF structures are represented following the convention (Spaink, 1992) [43] that indicates the number of GlcNAc residues in the backbone (Roman numeral), the length and degree of unsaturation of the fatty acyl chain, and the other substituents, which are listed in the order in which they appear, moving clockwise from the fatty acid. Hex-ol, hexitol (reduced terminal hexose); NMe, *N*-methyl group at glucosamine non reducing residue; S, sulfate group at reducing glucosamine residue.

^bSymbol: + = detected; - = non detected.

^cThese ions arise by loss of a neutral with mass 80 Da, corresponding to the loss of SO₃.

^dMixture of two Nod Factors, deacetylated at glucosamine residues numbers 2 and 3, respectively.

^eNod Factor deacetylated at glucosamine residue number 2.

report that CIAT 899 produces of a high number of Nod factors in the presence of *nod*-gene-inducing molecules [27-30], which provides a better understanding of the

control of Nod-factor biosynthesis, and which, in *R. tropici*, does not follow the classical pathway mediated by flavonoids.

Table 4 Nod Factor structure biosynthesized in the presence of 300 mM NaCl by wild type CIAT899 and the *nodD1* and *nodD2* mutants

[M + H] ⁺ (m/z)	B _i ions	Structure ^a	CIAT 899 ^b	<i>nodD1</i> ^b	<i>nodD2</i> ^b
824	400, 603	III (C _{16:0})	+	+	+
838	414, 617	III (C _{16:0} , NMe)	+	+	+
850	426, 629	III (C _{18:1})	+	+	+
864	440, 643	III (C _{18:1} , NMe)	+	-	+
999	372, 575, 778	IV (C _{14:0})	+	+	-
1013	386, 589, 792	IV (C _{14:0} , NMe)	+	+	-
1025	398, 601, 804	IV (C _{16:1})	+	+	+
1027	400, 603, 806	IV (C _{16:0})	+	+	+
1041	414, 617, 820	IV (C _{16:0} , NMe)	+	+	+
1053	426, 629, 832	IV (C _{18:1})	+	+	+
1055	428, 631, 834	IV (C _{18:0})	+	-	-
1067	440, 643, 846	IV (C _{18:1} , NMe)	+	+	+
1069	442, 645, 848	IV (C _{18:0} , NMe)	+	-	-
1147	440, 643, 846	IV (C _{18:1} , NMe, S)	+	-	-
1149	442, 645, 848	IV (C _{18:0} , NMe, S)	+	-	-
1202	372, 575, 778, 981	V (C _{14:0})	-	+	-
1203	414, 617, 820, 1023	IV Hex (C _{16:0} , NMe)	+	-	-
1205	414, 617, 820, 1023	IV Hex-ol (C _{16:0} , NMe)	+	-	-
1215	426, 629, 832, 1035	IV Hex (C _{18:1})	+	+	-
1216	386, 589, 792, 995	V (C _{14:0} , NMe)	+	+	+
1229	440, 643, 846, 1049	IV Hex (C _{18:1} , NMe)	+	-	-
1230	400, 603, 806, 1009	V (C _{16:0})	+	+	+
1231	440, 643, 846, 1049	IV Hex-ol (C _{18:1} , NMe)	+	-	-
1233	442, 645, 848, 1051	IV Hex-ol (C _{18:0} , NMe)	+	-	-
1242	412, 615, 818, 1021	V (C _{16:1} , NMe)	+	+	+
1244	414, 617, 820, 1023	V (C _{16:0} , NMe)	+	+	+
1256	426, 629, 832, 1035	V (C _{18:1})	+	+	+
1258	428, 631, 834, 1037	V (C _{18:0})	+	-	-
1270	440, 643, 846, 1049	V (C _{18:1} , NMe)	+	+	+
1272	442, 645, 848, 1051	V (C _{18:0} , NMe)	+	-	-
1298	468, 671, 874, 1077	V (C _{20:1} , NMe)	+	-	-
1324	414, 617, 820, 1023	V (C _{16:0} , NMe, S)	+	-	+
1336	426, 629, 832, 1035	V (C _{18:1} , S)	+	+	+
1350	440, 643, 846, 1049, [M-80] ^{+c} = 1270	V (C _{18:1} , NMe, S)	+	+	+
1352	442, 645, 848, 1051	V (C _{18:0} , NMe, S)	+	-	-
1378	468, 671, 874, 1077	V (C _{20:1} , NMe, S)	+	-	-
1380	470, 673, 876, 1079	V (C _{20:0} , NMe, S)	+	-	-

^aNF structures are represented following the convention (Spaink, 1992) [43] that indicates the number of GlcNAc residues in the backbone (Roman numeral), the length and degree of unsaturation of the fatty acyl chain, and the other substituents, which are listed in the order in which they appear, moving clockwise from the fatty acid. Hex, hexose; Hex-ol, hexitol (reduced terminal hexose); NMe, *N*-methyl group at glucosamine non reducing residue; S, sulfate group at reducing glucosamine residue.

^bSymbol: + = detected; - = non detected.

^cThese ions arise by loss of a neutral with mass 80 Da, corresponding to the loss of SO₃.

Gene expression

In various strains of rhizobia, the *nodD1* gene is the chief regulator of Nod-factor biosynthesis and symbiotic phenotype e.g. [34,37,38,47]. Contrarily, *nodD2* has been proposed as a repressor of *nod*-gene expression [33,39,48,49], affecting the bacterial Nod-factor profile. We performed gene expression studies with the wild type and *nodD1* and *nodD2* mutants, to improve our understanding of the roles of these two genes (Figure 3).

We evaluated the relative expression of the *nodC* gene (Figure 3A), which controls the elongation of the oligosaccharide chain of Nod factors and is transcribed with the activation of *nod* genes. The relative expression of *nodC* was lower for both mutants in comparison to the wild type strain, both in the apigenin and in the salt treatments (Figure 3A).


Significant expressions of the *nodD1* gene was observed in the WT strain both with salt and apigenin, while for the *nodD2* mutant it was statistically significant only when induced with apigenin, and higher than in the WT (Figure 3B).

In relation to the expression of *nodD2*, CIAT 899 WT strain significantly expressed the gene both with salt and apigenin. Contrarily, no statistically significant expression was observed for the *nodD1* mutant in none of the conditions evaluated (Figure 3C). However, we must consider that the expression levels of *nodD2* were all very low, and numerically even higher for the *nodD1* mutant in the presence of apigenin, therefore no strong conclusion can be taken from this assay at this moment (Figure 3C).

All together, these results indicate that the *nodD1* is a positive regulator gene, while *nodD2* may positively or negatively regulate the expression of the *nodD1* gene. Supporting these results is evidence of the involvement of *nodD2* in the regulation of the expression of *nodD1* by binding to *nod* box-like sequences located upstream of its coding region [33,49].

Proposal of a regulatory model for *nodD1* and *nodD2* genes of *R. tropici*

A graphic summary of the main features of the wild type CIAT 899 and *nodD1* and *nodD2* mutants is shown in Additional file 3: Figure S3. In our study a major role of *nodD1* in *R. tropici* CIAT 899 was confirmed. In the presence of the *nod*-gene-inducer apigenin, *nodD1* greatly increased the expression of *nodC* (42-fold), decreasing to 15-fold when the gene was mutated. Similar responses, but lower in magnitude, were observed under saline conditions (Figure 3A). Although confirming a major role of *nodD1*, the results also indicate that other *nodD* genes are involved in the activation of *nodC*, in the presence both of flavonoids and of salts.



Still in relation to *nodD1*, a mutation abolished nodulation in leucaena and siratro, but not in common bean. As *nodD1* gene is the chief regulator of Nod-factor biosynthesis and thus nodulation of the host plant e.g. [5,8,43,44], our results suggest that common bean might not be the main host of *R. tropici*, although it has been largely isolated from this host legume in acid soils of South America [24,26,50-52]. Indeed, doubts about common bean as the main host of *R. tropici* have been raised, giving support to the hypothesis that the species

might be an original symbiont of another indigenous legume, further “adapting” to common bean [50]. *R. tropici* has been isolated from common bean and other indigenous legumes in Europe, Africa, Australia, and North America [50], and results of some studies suggest the following as original host candidates for *R. tropici*: *Gliricidia* spp., from which the strain has been isolated in Mexico [53] and Brazil [54]; *Acaciella angustissima* in Mexico [55]; and *Mimosa* spp. in Brazil [54].

Understanding the relation between Nod factors and host specificity has been a goal of several studies, but without full success. In our research, we found that Nod factors of the following structures, III (C_{18:1}, NMe), IV (C_{18:1}, NMe), IV (C_{18:0}, NMe), V (C_{18:0}, NMe) and V (C_{16:0}, NMe, S) (Tables 3 and 4) might be related—to a greater or lesser extent—to the nodulation of the original host plant, as they are absent in the *nodD1* mutant. We have also confirmed the great variety of Nod factors produced by *R. tropici*, as reported before [27–30], even in the absence of *nod*-gene inducers [29,30] (Additional file 3: Figure S3). We propose a new, constitutive mechanism of Nod-factor synthesis that is highly enhanced when environmental conditions are stressful, such as strongly acidic pH or salinity. Some transcriptional regulators may be activated in these conditions and they could be responsible for the regulation of *nod*-gene expression via *nodD* regulators.

In various rhizobial strains, *nodD2* has been described as a repressor of the expression of *nod* genes e.g. [23,33,39,49]. For example, in *B. japonicum*, induction of *nodC* by flavonoids is virtually suppressed by elevated levels of NodD2 [23], and in *S. fredii* extra plasmid copies of *nodD2* reduced the level of *nodD1* transcripts to below the limits of detection [49]. However, there is still no evidence that the suppression by NodD2 is mediated by *nodD1*. In our study, we found that *nodD2* activated *nodC* at similar levels as those observed for *nodD1* with both apigenin and salt. However, a slight repression of *nodD1* by *nodD2* was observed in the presence of apigenin (Figure 3B). Accordingly, we hypothesize that *nodD2* is usually an activator of *nod*-gene transcription, although, in the presence of some flavonoids it may slightly repress *nodD1*. Nevertheless, if this repression is biologically significant, it remains to be determined, as no differences in nodulation were observed for leucaena or siratro in the absence of *nodD2*, whereas nodulation was decreased in common bean.

Our model contends that full nodulation of common bean by *R. tropici* requires both *nodD1* and *nodD2*, while, in other plant species that might represent the original host, *nodD1* plays the major role. *nodD2* is not a strong repressor as described in other rhizobial species, and, in general, plays a role as an activator of *nod*-gene transcription, but, in specific conditions, it may slightly

repress *nodD1*. The *nodD* regulation in *R. tropici* CIAT 899 resembles the pattern observed in *S. meliloti*—need for three copies of *nodD* to optimize the interaction with each of its legume hosts [40]. The biological significance of producing an abundance of Nod factors is not completely understood yet, but we hypothesize that represents an evolutionary strategy to avoid abiotic stresses by nodulating a range of legume species. Reports show that *nod* genes may also control other functions that contribute to nodulation, as described for *nodD2* in the exopolysaccharide synthesis of *S. fredii* [38], and chaperones and other genes by *nodD1* in *S. meliloti* [56], *inter alia*. Our results demonstrate extra roles for *nodD1* and *nodD2* of *R. tropici* in swarming motility and IAA synthesis.

Conclusions

Our model proposes that full nodulation of common bean by *R. tropici* requires both *nodD1* and *nodD2*, while in other plant species that might represent the original host *nodD1* plays the major role. Contrarily to other rhizobial species, *nodD2* of *R. tropici* is usually not a strong repressor of *nod*-gene transcription. *R. tropici* synthesizes a variety of Nod factors that might be related to the ability of nodulating a variety of legume species, representing an evolutionary strategy of the symbiosis under abiotic stressful conditions. *nodD1* and *nodD2* of *R. tropici* also play roles in swarming motility and IAA synthesis.

Methods

Bacterial strains, plasmids, media, and growth conditions

Rhizobium tropici CIAT 899 and derivative strains were grown at 28°C on tryptone yeast (TY) medium [57], B⁻ minimal medium [43] or yeast-extract mannitol (YM) medium [58], supplemented when necessary with apigenin to a final concentration of 3.7 μM or with NaCl at 300 mM. *Escherichia coli* strains were cultured on Luria-Bertani (LB) medium [59] at 37°C. When required, the media were supplemented with the appropriate antibiotics as described by Lamrabet *et al.* [60]. *R. tropici* RSP82 [30] was used as a *nodD1* mutant derivative of *R. tropici* CIAT899 (Km^R 30 μg mL⁻¹).

To obtain the *nodD2* mutant, primer pairs *nodD2*-F (5′ – GTA GGC CAT AAT GTC CAG A) and *nodD2*-R (5′ – GCG GCT TTA TAC TCA CCA) were used for amplifying the *nodD2* gene. The 1450-bp PCR product was cloned into pGEM⁺-T Easy (Promega) (Amp^R 100 μg mL⁻¹). The PCR-amplified *nodD2* fragment was then excised from the plasmid obtained with the endonuclease *EcoRI* and cloned into the vector pK18mob [61], which is suicide in rhizobia, confers resistance to kanamycin (km^R 30 μg mL⁻¹) and was previously digested also with *EcoRI*. This new plasmid was digested with the

enzyme *Pst*I, which cuts the *nodD2* gene in two sites, releasing a fragment of approximately 600 pb. The rest of the plasmid was treated with the Klenow enzyme to convert the cohesive ends to blunt ends. This treated plasmid was ligated with a 2-Kb DNA fragment containing the Ω interposon [carrying the spectinomycin resistance gene (spc^R 100 $\mu\text{g mL}^{-1}$)], which was obtained from a previous digestion of the pHP45 Ω plasmid [62] with the *Sma*I enzyme (blunt end). The resulting plasmid was transformed into the *E. coli* strain DH5 α . Plasmids were transferred from *E. coli* to *Rhizobium* strains by conjugation, as described by Simon [63], using plasmid pRK2013 [64] as helper. The plasmid generated was used for the homogenization of the mutated version of the *nodD2* gene in *R. tropici* CIAT 899 by using the methodology previously described [65]. The homogenization was confirmed by DNA-DNA hybridization. For this purpose, DNA was blotted to Hybond-N nylon membranes (Amersham, UK), and the DigDNA method of Roche (Switzerland) was employed according to the manufacturer's instructions. Additional file 1: Figure S1 displays the type of mutation realized to obtain the *nodD2* mutant.

The parental and mutant strains are deposited in the culture collection of the Department of Biology of the Universidad de Sevilla and at the Diazotrophic and Plant Growth Promoting Bacteria Culture Collection of Embrapa Soja (WFCC Collection # 1213, WDCC Collection # 1054).

Identification of Nod factors

Purification and LC-MS/MS analyses of Nod factors produced by *R. tropici* CIAT 899 and derivative strains grown in B⁻ minimal medium [43] (supplemented when required with NaCl 300 mM or apigenin 3.7 μM) were performed as described previously [30].

RNA isolation, cDNA synthesis and quantitative RT-PCR

Wild type CIAT 899 and *nodD1* and *nodD2* mutants were pre-cultured in 10-mL aliquots of TY medium at 100 rpm and 28°C in the dark. After 48 h, the three strains pre-inoculated were transferred to new media and subjected to the following conditions: control (without induction), 300 mM NaCl and apigenin 3.7 μM . These new cultures were performed in triplicate under the same conditions as for the pre-cultures, 100 rpm and 28°C in the dark, except that were grown into the exponential phase (O.D. at 600 nm of 0.5 to 0.6).

Total RNA was extracted using Trizol[®] reagent (Life Technologies) as previously described [66]. The total concentrations were estimated in a NanoDrop ND 1000 spectrophotometer (NanoDrop-Technologies, Inc., City etc. here and elsewhere) and the integrity was assessed by gel electrophoresis. Extracted RNA samples were submitted to DNaseI treatment (Invitrogen/Life Technologies, Grand Island, NY, USA) and the first stand of

cDNA was synthesized using SuperscriptIII[™] reverse transcriptase (Invitrogen[™]), according to manufacturer's protocol.

Primers for the RT-qPCR targets, genes *nodD1*, *nodD2* and *nodC*, were designed using Primer3Plus (<http://www.bioinformatics.nl/cgi-bin/primer3plus/primer3plus.cgi/>), to obtain amplicons of 50–150 bp. With the same software, a primer to 16S rRNA was obtained and applied to normalize the relative expression of the targets. To avoid unspecific alignments, the primer sequences were searched against the *R. tropici* CIAT 899 genome (<http://www.ncbi.nlm.nih.gov/nucleotide/440224888?report=genbank>). The primer sequences and sizes of the amplified fragments are available in Additional file 4: Table S1.

RT-qPCR reactions were performed in a 7500 RT-qPCR Thermocycler (Applied Biosystems/Life Technologies). The reactions were performed in triplicate for each of the three biological replicates. The Platinum[®] SYBR Green[®] Master Mix kit (Applied Biosystems) was used according to the manufacturer's instructions. Cycling conditions were as follows: 50°C for 2 min, 95°C for 10 min, 45 cycles at 95°C for 2 min, 60°C for 30 s and 72°C for 30 s, in 45 cycles. Rest2009 software package [67] was used to evaluate the data by providing a robust statistical analysis ($p < 0.05$). The normalization of cycle threshold (Ct) of RT-qPCR amplifications was performed based on the selected endogenous gene (16S rRNA).

Studies of external exopolysaccharides

The anthrone-H₂SO₄ method, which measures the total reducing sugar content in a given sample [68] was used to determine the total carbohydrate amounts of exopolysaccharide (EPS) contained in supernatants from bacterial cultures. For this purpose, *R. tropici* CIAT 899 and derivatives were grown in 5 mL of TY liquid medium on an orbital shaker (180 rpm) for 96 h at 28°C. When required, the media were supplemented with NaCl (300 mM) or apigenin (3.7 μM). Samples of 1 mL were centrifuged to remove cells. Cell-free culture supernatants were assayed for EPS content via H₂SO₄ hydrolysis in the presence of the colorimetric indicator anthrone. Every experiment was performed three times with three replicates each time. Lipopolysaccharide (LPS) extraction, separation on SDS-PAGE, and silver staining were performed as previously described using the same bacteria, medium and conditions [17].

Motility assays

Swimming and swarming phenotypes were tested on TY medium [57] (supplemented when necessary with NaCl 300 mM or apigenin 3.7 μM) agar plates containing 0.28% or 0.4%, respectively, of Bacto Agar. The strains to be assayed (wild type and mutants) were

grown in 5 mL of TY medium on an orbital shaker (180 rpm) for 96 h at 28°C. Aliquots (2 µL) of culture suspensions were drop-inoculated (swarming assay) or sink-inoculated (swimming assay) onto plates and air-dried in a laminar-flow cabinet. The inoculated plates were wrapped with parafilm and incubated for the required time at 28°C in an upright position. Every experiment was performed three times with three replicates each time.

Biofilm formation assay

The biofilm formation assay on polystyrene surfaces was performed using the method described by O'Toole and Kolter [69] with modifications [20]. CIAT 899 and mutant strains were grown on TY medium [57] (supplemented with NaCl 300 mM or apigenin 3.7 µM when required) for 7 days with gentle rocking at 28°C. Every experiment was performed three times with eight replicates each time.

Quantification of indole acetic acid (IAA) production

Quantification of an IAA-like compound from *R. tropici* strain cultures was carried out by using Salkowski colorimetric assays [70], as described previously by Fierro-Coronado *et al.* [71]. To measure IAA production, 5 mL of TY medium with tryptophan (0.4 g L⁻¹) (supplemented when required with NaCl 300 mM or apigenin 3.7 µM) were inoculated and incubated during 96 h at 28°C on an orbital shaker (180 rpm) with *R. tropici* strains. Of these cultures, samples of 1 mL were centrifuged to remove cells. Cell-free culture supernatants were assayed for IAA production. Every experiment was performed three times with eight replicates each time.

Nodulation assays

nodD1 and *nodD2* mutants were grown in YM medium until a concentration of 10⁹ cells mL⁻¹ was achieved, to be used as inoculum. Surface-sterilized seeds [58] were used for the assays with common bean (*Phaseolus vulgaris* L.), leucaena [*Leucaena leucocephala* (Lam.) de Wit] and siratro [*Macroptilium atropurpureum* (DC) Urb.]. Pre-germinated seeds (about 2 days after germination) were placed in sterilized pouches or Leonard jars containing N-free nutrient solution [58], with 1 mL of inoculum of each strain added and verified for nodulation capacity after 25 (common bean) or 42 days (leucaena and siratro) with a 16-h 25°C/18°C photoperiod and about 70% relative humidity. Shoots were dried at 65°C until constant weight was achieved, and then weighed. Experiments were performed three times.

Additional files

Additional file 1: Figure S1. Information about the *nodD1* and *nodD2* genes of *R. tropici* used in our study. A. Gene neighborhood of *nodD1* and *nodD2* genes in the genome of *R. tropici* strain CIAT 899. B. Location of primers (dark arrows) used to perform RT-qPCR experiments. C. Schematic representation of the *nodD2* mutation.

Additional file 2: Figure S2. Nodulation phenotype in common bean (*Phaseolus vulgaris*) inoculated with CIAT 899 and derivative *nodD* strains assayed in pouch bags. Experiment performed under controlled conditions of growth chamber and plants harvested at 25 days after inoculation. A. wild type strain. B. *nodD1* mutant. C. *nodD2* mutant. D. Uninoculated.

Additional file 3: Figure S3. Main properties observed in wild type (WT), *nodD1* and *nodD2* mutants of *R. tropici* strain CIAT 899.

Additional file 4: Table S1. Sequences of the primers used in the RT-qPCR and sizes of the PCR products obtained.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

Conceived and designed the experiments: MM, FJO, MH. Performed the experiments: PC, APR-S, DFG, BBM, FP-M, MAR-C, ASN, AG-S, FJO. Analyzed the data: all authors. Contributed reagents/materials/analysis tools: MM, FJO, MH. Wrote the paper: PC, APR-S, DFG, MM, FJO, MH. All authors read and approved the final manuscript.

Acknowledgements

The study was partially supported by AGL2012-38831, Embrapa, CNPq (National Council for Scientific and Technological Development), Project Science without Borders (400205/2012-5). Authors acknowledge Dr. Jesiane S.S. Batista for helping in obtaining the *nodD2* mutant and Allan R.J. Eaglesham for suggestions on the manuscript and English review. A.A.P.R.-Santos acknowledges a posdoc fellowship from project Science without Borders, D.F.Gomes a PhD fellowship from Project Repensa and A.S.Nakatani a posdoc fellowship from Fundação Araucária. M.H. is also a research fellow from CNPq. We thank the Centro de Investigación, Tecnología e Innovación (CITIUS) of the University of Seville for MS facilities. Approved for publication by the Editorial Board of Embrapa Soja as manuscript 315/2014.

Author details

¹Departamento de Microbiología, Facultad de Biología, Universidad de Sevilla, Avda. Reina Mercedes, 6 Apdo Postal 41012 Sevilla, Spain. ²Embrapa Soja, C.P. 231, 86001-970 Londrina, Paraná, Brazil. ³Departamento de Química Orgánica, Facultad de Química, Universidad de Sevilla, Apdo Postal 553, 41071 Sevilla, Spain.

Received: 29 December 2014 Accepted: 9 March 2015

Published online: 28 March 2015

References

1. Peoples MB, Brockwell J, Herridge DF, Rochester IJ, Alves BJR, Urquiaga S, et al. The contributions of nitrogen-fixing crop legumes to the productivity of agricultural systems. *Symbiosis*. 2009;48:1–17.
2. Ormeño-Orrillo E, Hungria M, Martínez-Romero E. Dinitrogen-fixing prokaryotes. In: *The prokaryotes - prokaryotic physiology and biochemistry*. Berlin-Heidelberg: Springer-Verlag; 2013. p. 427–51.
3. Van Hameren B, Hayashi S, Gresshoff PM, Ferguson BJ. Advances in the identification of novel factors required in soybean nodulation, a process critical to sustainable agriculture and food security. *J Plant Biol Soil Health*. 2013;1:6.
4. Denarié J, Debbelle F, Promé JC. *Rhizobium* lipo-chitinoligosaccharide nodulation factors: signaling molecules mediating recognition and morphogenesis. *Ann Rev Biochem*. 1996;65:503–35.
5. Schlaman HRM, Phillips DA, Kondorosi E. Genetic organization and transcriptional regulation of rhizobial nodulation genes. In: Spaink HP, Kondorosi A, Hooykaas PJJ, editors. *The Rhizobiaceae: Molecular Biology of*

- Model Plant-Associated Bacteria. Dordrecht: Kluwer Academic Publishers; 1998. p. 361–86.
6. Geurts R, Bisseling T. *Rhizobium* Nod factor perception and signalling. *Plant Cell*. 2002;14:S239–49.
 7. Brennic A, Winans SC. Detection of and response to signals involved in host-microbe interactions by plant-associated bacteria. *Microbiol Mol Biol Rev*. 2005;69:155–94.
 8. Oldroyd GE. Speak, friend, and enter: signaling systems that promote beneficial symbiotic associations in plants. *Nat Rev Microbiol*. 2013;11:252–63.
 9. Khan W, Prithiviraj B, Smith DL. Nod factor [Nod Bj V (C18:1, MeFuc)] and lumichrome enhance photosynthesis and growth of corn and soybean. *J Plant Physiol*. 2008;163:1342–51.
 10. Liang Y, Cao Y, Tanaka K, Thibivilliers S, Wan J, Choi J, et al. Nonlegumes respond to rhizobial Nod factors by suppressing the innate immune response. *Science*. 2013;341:1384–7.
 11. Marks BB, Nogueira MA, Hungria M, Megias M. Biotechnological potential of rhizobial metabolites to enhance the performance of Bradyrhizobium spp and Azospirillum brasilense inoculants with soybean and maize. *AMB Express*. 2013;3:21.
 12. Smith S, Habib A, Kang Y, Leggett M, Diaz-Zorita M. LCO applications provide improved responses with legumes and non-legumes. In: De Bruijn F, editor. *Biological Nitrogen Fixation*. New Jersey: John Wiley & Sons, Inc; 2015. p. 1073–81.
 13. Kondoros E, Gyuris J, Schmidt J, John M, Duda E, Hoffmann B, et al. Positive and negative control of *nod* gene expression in *Rhizobium meliloti* is required for optimal nodulation. *Embo J*. 1989;8:1331–40.
 14. Spaink HP. Root nodulation and infection factors produced by rhizobial bacteria. *Annu Rev Microbiol*. 2000;54:257–88.
 15. Krause A, Doerfel A, Gottfert M. Mutational and transcriptional analysis of the type III secretion system of *Bradyrhizobium japonicum*. *Mol Plant Microbe Interact*. 2002;15:1228–35.
 16. Theunis M, Kobayashi H, Broughton WJ, Prinsen E. Flavonoids, NodD1, NodD2, and Nod-Box NB15 modulate expression of the *y4wEFG* locus that is required for indole-3-acetic acid synthesis in *Rhizobium* sp. strain NGR234. *Mol Plant Microbe Interact*. 2004;17:1153–61.
 17. Vinardell JM, Lopez-Baena FJ, Hidalgo A, Ollero FJ, Bellogin R, Espuny MR, et al. The effect of FITA mutations on the symbiotic properties of *Sinorhizobium fredii* varies in a chromosomal-background-dependent manner. *Arch Microbiol*. 2004;181:144–54.
 18. López-Baena FJ, Vinardell JM, Pérez-Montaño F, Crespo-Rivas JC, Bellogin RA, Espuny Mdel R, et al. Regulation and symbiotic significance of nodulation outer proteins secretion in *Sinorhizobium fredii* HH103. *Microbiology*. 2008;154:1825–36.
 19. Pérez-Montaño F, Guasch-Vidal B, González-Barroso S, López-Baena FJ, Cubo T, Ollero FJ, et al. Nodulation-gene-inducing flavonoids increase overall production of autoinducers and expression of N-acyl homoserine lactone synthesis genes in rhizobia. *Res Microbiol*. 2011;162:715–23.
 20. Pérez-Montaño F, Jiménez-Guerrero I, Del Cerro P, Baena-Ropero I, López-Baena FJ, Ollero FJ, et al. The symbiotic biofilm of *Sinorhizobium fredii* SMH12, necessary for successful colonization and symbiosis of *Glycine max* cv Osumi, is regulated by quorum sensing systems and inducing flavonoids via NodD1. *PLoS One*. 2014;9:e105901.
 21. Hungria M, Johnston AWB, Phillips DA. Effects of flavonoids released naturally from bean (*Phaseolus vulgaris*) on *nodD*-regulated gene transcription in *Rhizobium leguminosarum* bv. *phaseoli*. *Mol Plant Microbe Interact*. 1992;5:199–203.
 22. Broughton WJ, Jabbouri S, Perret X. Keys to symbiotic harmony. *J Bacteriol*. 2000;182:5641–52.
 23. Garcia MLJ, Dunlap J, Loh J, Stacey G. Phenotypic characterization and regulation of the *nolA* gene of *Bradyrhizobium japonicum*. *Mol Plant Microbe Interact*. 1996;9:625–35.
 24. Martínez-Romero E, Segovia L, Mercante FM, Franco AA, Graham P, Pardo MA. *Rhizobium tropici*, a novel species nodulating *Phaseolus vulgaris* L. beans and *Leucaena* sp. trees. *Int J Syst Bacteriol*. 1991;41:417–26.
 25. Hernández-Lucas I, Segovia L, Martínez-Romero E, Pueppke SG. Phylogenetic relationships and host range of *Rhizobium* spp. that nodulates *Phaseolus vulgaris* L. *Appl Environ Microbiol*. 1995;61:2775–9.
 26. Hungria M, Andrade DS, Chueire LMO, Probanza A, Guitierrez-Manero FJ, Megias M. Isolation and characterization of new efficient and competitive bean (*Phaseolus vulgaris* L.) rhizobia from Brazil. *Soil Biol Biochem*. 2000;21:1515–28.
 27. Poupot R, Martínez-Romero E, Promé JC. Nodulation factors from *Rhizobium tropici* are sulfated or nonsulfated chitopentasaccharides containing an N-methyl-N-acylglucosaminyl terminus. *Biochemistry*. 1993;32:10430–5.
 28. Morón B, Soria-Díaz ME, Ault J, Verroios G, Noreen S, Rodríguez-Navarro DN, et al. Low pH changes the profile of nodulation factors produced by *Rhizobium tropici* CIAT899. *Chem Biol*. 2005;12:1029–40.
 29. Estevez J, Soria-Díaz ME, De Cordoba FF, Moron B, Manyani H, Gil A, et al. Different and new Nod factors produced by *Rhizobium tropici* CIAT899 following Na⁺ stress. *FEMS Microbiol Lett*. 2009;293:220–31.
 30. Guasch-Vidal B, Estévez J, Dardanelli MS, Soria-Díaz ME, De Córdoba FF, Balog CI, et al. High NaCl concentrations induce the *nod* genes of *Rhizobium tropici* CIAT899 in the absence of flavonoid inducers. *Mol Plant Microbe Interact*. 2013;26:451–60.
 31. van Rhijn PJS, Feys B, Verreth C, Vanderleyden J. Multiple copies of *nodD* in *Rhizobium tropici* CIAT899 and BR816. *J Bacteriol*. 1993;175:438–47.
 32. Ormeño-Orrillo E, Menna P, Gonzaga LA, Ollero FJ, Nicolas MF, Rodrigues EP, et al. Genomic basis of broad host range and environmental adaptability of *Rhizobium tropici* CIAT 899 and *Rhizobium* sp PRF 81 which are used in inoculants for common bean (*Phaseolus vulgaris* L.). *BMC Genomics*. 2012;13:735.
 33. Fellay R, Hanin M, Montorzi G, Frey J, Freiberg C, Golinowski W, et al. *nodD2* of *Rhizobium* sp. NGR234 is involved in the repression of the *nodABC* operon. *Mol Microbiol*. 1998;27:1039–50.
 34. Gillette WK, Elkan GH. *Bradyrhizobium (Arachis)* sp. strain NC92 contains two *nodD* genes involved in the repression of *nodA* and a *nolA* gene required for the efficient nodulation of host plants. *J Bacteriol*. 1996;178:2757–66.
 35. Berg HC. The rotary motor of bacterial flagella. *Annu Rev Biochem*. 2003;72:19–54.
 36. Tambalo DD, Vanderlinde EM, Robinson S, Halmillawewa A, Hynes MF, Yost CK. Legume seed exudates and *Physcomitrella patens* extracts influence swarming behavior in *Rhizobium leguminosarum*. *Can J Microbiol*. 2014;60:15–24.
 37. Honma MA, Ausubel FM. *Rhizobium meliloti* has three functional copies of the *nodD* symbiotic regulatory gene. *Proc Natl Acad Sci U S A*. 1987;84:8558–62.
 38. Appelbaum ER, Thompson DV, Ilder K, Chartrain N. *Rhizobium japonicum* USDA 191 has two *nodD* genes that differ in primary structure and function. *J Bacteriol*. 1998;170:12–20.
 39. Loh J, Stacey G. Nodulation gene regulation in *Bradyrhizobium japonicum*: a unique integration of global regulatory circuit. *Appl Environ Microbiol*. 2003;69:10–7.
 40. Honma MA, Asomaning M, Ausubel FM. *Rhizobium meliloti nodD* genes mediate host-specific activation of *nodABC*. *J Bacteriol*. 1990;172:901–11.
 41. Relić B, Fellay R, Lewin A, Perret K, Price NPJ, Rochepeau P, et al. *nod* genes and Nod Factors of *Rhizobium* species NGR 234. In: Palacios R, Mora J, Newton WE, editors. *New Horizons in Nitrogen Fixation*. Dordrecht: Kluwer Academic Publishers; 1993. p. 183–9.
 42. Göttfert M, Groß P, Hennecke H. Proposed regulatory pathway encoded by the *nodV* and *nodW* genes, determinants of host specificity in *Bradyrhizobium japonicum*. *Proc Natl Acad Sci*. 1990;87:2680–4.
 43. Spaink HP. Rhizobial lipo-oligosaccharides: answers and questions. *Plant Mol Biol*. 1992;20:977–86.
 44. Muñoz N, Soria-Díaz ME, Manyani H, Sánchez-Matamoros RC, Serrano AG, Megias M, et al. Structure and biological activities of lipochitooligosaccharide nodulation signals produced by *Bradyrhizobium japonicum* USDA 138 under saline and osmotic stress. *Biol Fertil Soils*. 2014;50:207–15.
 45. Jabbouri S, Relić B, Hanin M, Kamalaprifa P, Burger U, Promé D, et al. *nolO* and *nolE* (HsIII) of *Rhizobium* sp. NGR234 are involved in 3-O-carbamoylation and 2-O-methylation of Nod factors. *J Bio Chem*. 1998;273:12047–55.
 46. Folch-Mallol JL, Marroqui S, Sousa C, Manyani H, Lopez-Lara IM, van der Drift KM, et al. Characterization of *Rhizobium tropici* CIAT899 nodulation factors: the role of *nodH* and *nodPQ* genes in their sulfation. *Mol Plant Microbe Interact*. 1996;9(3):151–63.
 47. Vargas C, Martínez LJ, Megias M, Quinto C. Identification and cloning of nodulation genes and host specificity determinants of the broad host-range *Rhizobium leguminosarum* biovar *phaseoli* strain CIAT899. *Mol Microbiol*. 1990;4:1899–910.
 48. Machado D, Krishnan HB. *nodD* alleles of *Sinorhizobium fredii* USDA 191 differentially influence soybean nodulation, *nodC* expression, and production of exopolysaccharides. *Curr Microbiol*. 2003;47:134–7.
 49. Machado D, Pueppke SG, Vinardell JM, Ruiz-Sainz JE, Krishnan HB. Expression of *nodD1* and *nodD2* in *Sinorhizobium fredii*, a nitrogen-fixing symbiont of soybean and other legumes. *Mol Plant Microbe Interact*. 1998;11:375–82.

50. Gomes DF, Ormeno-Orrillo E, Hungria M. Biodiversity, symbiotic efficiency and genomics of *Rhizobium tropici* and related species. In: De Bruijn F, editor. *Biological Nitrogen Fixation*. New Jersey, Hoboken: Wiley-Blackwell; 2015. p. 747–56.
51. Grange L, Hungria M, Graham PH, Martínez-Romero E. New insights into the origins and evolution of rhizobia that nodulate common bean (*Phaseolus vulgaris*) in Brazil. *Soil Biol Biochem*. 2007;39(4):867–76.
52. Pinto FGS, Hungria M, Mercante FM. Polyphasic characterization of Brazilian *Rhizobium tropici* strains effective in fixing N₂ with common bean (*Phaseolus vulgaris* L.). *Soil Biol Biochem*. 2007;39(8):1851–64.
53. Acosta-Durán C, Martínez-Romero E. Diversity of rhizobia from nodules of the leguminous tree *Gliricidia sepium*, a natural host of *Rhizobium tropici*. *Arch Microbiol*. 2002;178:161–4.
54. Menna P, Hungria M, Barcellos FG, Bangel EV, Hess PN, Martínez-Romero E. Molecular phylogeny based on the 16S rRNA gene of elite rhizobial strains used in Brazilian commercial inoculants. *Syst Appl Microbiol*. 2006;29:315–32.
55. Rincón-Rosales R, Lloret L, Ponce E, Martínez-Romero E. Rhizobia with different symbiotic efficiencies nodulate *Acaciella angustissima* in Mexico, including *Sinorhizobium chiapanecum* sp. nov. which has common symbiotic genes with *Sinorhizobium mexicanum*. *FEMS Microbiol Ecol*. 2009;67:103–17.
56. Capela D, Carrere S, Batut J. Transcriptome-based identification of the *Sinorhizobium meliloti* NodD1 regulon. *Appl Environ Microbiol*. 2005;71:4910–3.
57. Beringer JE. R factor transfer in *Rhizobium leguminosarum*. *J Gen Microbiol*. 1974;84:188–98.
58. Vincent JM. The modified Fahraeus slide technique. In: Vincent JM, editor. *A manual for the practical study of root nodule bacteria*. Oxford, UK: Blackwell Scientific Publications; 1970.
59. Sambrook J, Fritsch EF, Maniatis T. *Molecular Cloning: A Laboratory Manual*. 2nd ed. Cold Spring Harbor, NY, USA: Cold Spring Harbor Laboratory Press; 1989.
60. Lamrabet Y, Bellogín RA, Cubo T, Espuny MR, Gil-Serrano A, Krishnan HB, et al. Mutation in GDP-fucose synthesis genes of *Sinorhizobium fredii* alters Nod factors and significantly decreases competitiveness to nodulate soybeans. *Mol Plant-Microbe Interact*. 1999;12:207–17.
61. Schäfer A, Tauch A, Jäger W, Kalinowski J, Thierbach G, Pühler A. Small mobilizable multi-purpose cloning vectors derived from the *Escherichia coli* plasmids pK18 and pK19: selection of defined deletions in the chromosome of *Corynebacterium glutamicum*. *Gene*. 1994;145:69–73.62.
62. Prentki P, Krisch HM. *In vitro* insertional mutagenesis with a selectable DNA fragment. *Gene*. 1984;29:303–13.
63. Simon R. High frequency mobilization of gram-negative bacterial replicons by the in vivo constructed Tn5-Mob transposon. *Mol Gen Genet*. 1984;196:413–20.
64. Figurski DH, Helinski DR. Replication of an origin-containing derivative of plasmid RK2 dependent on a plasmid function provided in trans. *Proc Natl Acad Sci U S A*. 1979;76:1648–52.
65. López-Baena FJ, Monreal JA, Pérez-Montaño F, Guasch-Vidal B, Bellogín RA, Vinardell JM, et al. The absence of Nops secretion in *Sinorhizobium fredii* HH103 increases GmPR1 expression in Williams soybean. *Mol Plant Microbe Interact*. 2009;22:1445–54.
66. Gomes DF, Batista JSS, Rolla AAP, Silva LPS, Bloch C, Galli-Terasawa LV, et al. Proteomic analysis of free-living *Bradyrhizobium diazoefficiens*: highlighting potential determinants of a successful symbiosis. *BMC Genomics*. 2014;15:643. doi: 10.1186/1471-2164-15-643.
67. Pfaffl MW, Horgan GW, Dempfle L. Relative expression software tool (REST) for group-wise comparison and statistical analysis of relative expression results in real-time PCR. *Nucleic Acids Res*. 2002;30:e36.
68. Tomlinson AD, Ramey-Hartung B, Day TV, Merritt PM, Fuqua C. *Agrobacterium tumefaciens* ExoR represses succinoglycan biosynthesis and is required for biofilm formation and motility. *Microbiology*. 2010;156:2670–81.
69. O'Toole GA, Kolter R. Initiation of biofilm formation in *Pseudomonas fluorescens* WCS365 proceeds via multiple, convergent signalling pathways: a genetic analysis. *Mol Microbiol*. 1998;28:449–61.
70. Glickmann E, Dessaux Y. A critical examination of the specificity of the Salkowski reagent for indolic compounds produced by phytopathogenic bacteria. *Appl Environ Microbiol*. 1995;61:793–6.
71. Fierro-Coronado RA, Quiroz-Figueroa FR, García-Pérez LM, Ramírez-Chávez E, Molina-Torres J, Maldonado-Mendoza IE. IAA-producing rhizobacteria from chickpea (*Cicer arietinum* L.) induce changes in root architecture and increase root biomass. *Can J Microbiol*. 2014;60:639–48.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

