

Sunflower and peanut emergency : initial development under sugarcane mulch

Nilza Patrícia Ramos¹, Maria do Carmo de Salvo Soares Novo²; Maria Regina Gonçalves Ungaro²; Antônio Augusto Lago²

¹Embrapa Meio Ambiente, Rodovia SP 340 - Km 127,5 Jaguariúna-SP, C.P. 69, CEP: 13820-000, Brasil, www.cnpma.embrapa.br, e-mail - npramos@cnpma.embrapa.br. ²Instituto Agronômico de Campinas, Avenida Theodureto de Almeida Camargo, nº1500, Campinas-SP, C.P. 28, CEP. 13001-970, Brasil, www.iac.sp.gov.br

ABSTRACT

The research aimed to evaluate the effect of residual sugarcane mulch on sunflower and peanut plant emergence and initial development. Vases of 4.0 L were disposed in a randomized experimental blocks design, with four replications, in a factorial arrangement of five mulch amounts and three cultivars of each crop. The mulch treatments consisted of four increasing amounts (5, 10, 15 e 20 t ha⁻¹) and a tester with no mulch. The sunflower cultivars were the varieties IAC-iarama and Catissol and the hybrid H-358; the peanut cultivars were IAC-Caiapó, Runner 886 and Tatu. The speed emergency index and final percentage, the plant height and shoot dry mass were evaluated. The presence of different levels of sugarcane mulch negatively influences the emergency and initial plant development mainly in peanut but also in sunflower. The negative effects are stronger especially for the runner type of peanut cultivars, while cultivar Tatu was less influenced by the mulch thickness.

Key-words: *Arachis hypogaea* L. - *Helianthus annuus* L – mulch - oilcrops - seedling development.

INTRODUCTION

The sugarcane crop expansion is a reality in Brazil due mainly by the bio-energy or renewable energy concern. This expansion is now been associated with the biodiesel agriculture chain, favored by the possibility of sugarcane rotation with oil crops. In Sao Paulo state, sugarcane rotation with peanut crop is a reality (Borsari Filho, 2006), but there is also a great potential for sunflower crop. There are increased interests in the oil crops that can be used for biodiesel production (Brasil, 2006). Thus, together with the sunflower rotation benefits on sugarcane of about 50% in the sugar yield (Ambrosano *et al.*, 2005), the energetic benefits of fuel association are expected, due to the use of the biodiesel produced by the sugarcane chain in the agriculture and transportation vehicles from the sugar mills and farmers.

With the implementation of laws that prohibited the sugarcane burning, the harvest leaves a great amount of mulch on the soil surface. The presence of mulch on the soil surface can modify physical soil characteristics like water content and thermic extent (Vasconcelos, 2002), which contribute for the soil conservation but, on the other side, can cause problems to crop management (Furlani Neto *et al.*, 1997).

The mulch layer in sugarcane areas can be as high as 10cm thickness on the soil surface, which correspond to 20 t ha⁻¹ of residues; these form a physical barrier that reduces the light incidence and modify the local climatic conditions (Velini & Negrisoni, 2000). Those alterations are able to affect the emergency and plant development due to the influence on dormancy and seed emergency processes (Trezza & Vidal, 2004). With the increased use of oilcrops in the sugarcane rotation, the study of the influence of mulch on different crops is a necessity. In this sense, the aim of the present research was to evaluate the influence of sugarcane residual mulch on the emergency and initial development of different sunflower and peanut cultivars.

MATERIAL AND METHODS

A greenhouse experiment was done at Ecophysiology and Biophysics Center of Instituto Agronômico (IAC), Campinas, SP, Brazil. Plastic vases of 4.0 L capacity, filled with 2.7 L of an argilous sieved soil, with the following chemical composition: pH (CaCl₂) = 5.2, organic matter = 25 g dm⁻³, P (resin) = 1 mg dm⁻³, K = 0.9, Ca = 23, Mg = 6, H+Al = 28, SB=29.9, CTC=57.7, expresses in mmol_c dm⁻³ and V = 52% were used. The soil was amended according to Van Raij *et al.* (1997).

Before sowing, seeds of sunflower and peanut were physiologically characterized by determining the germination, emergence (Brasil, 1992), and speed emergence index according to Maguire (1962).

The treatments were disposed as a factorial scheme (5 x 3), in a randomized block design, with four replications, combining five amounts of sugarcane mulch (0, 5, 10, 15 e 20 t ha⁻¹) and three cultivars of each crop, in separate. For sunflower were used the open pollinated cultivars IAC-iarama and Catissol and the hybrid Helio-358, while for peanut were used the runner type varieties IAC-Caiapó and Runner 88, and the erect type Tatu.

In each vase were sowed ten seeds previously treated with Thiram 0,2%, at 3cm deep, followed by the addition of the mulch of sugarcane cultivar SP 803280, which were cut into small pieces before the distribution on soil surface. The layer thicknesses on the vases were 4, 6, 9, and 10cm, which correspond to 5, 10, 15, and 20 t ha⁻¹, respectively.

The final plant emergence (EM) was evaluated 15 days after sowing (Brasil, 1992). For the speed emergence index (SEI) the number of normal seedling was daily counted until constant number, according to Maguire (1962). The initial plant development was evaluated 30 days after sowing, harvesting the plants, measuring the plant height (HP) and, after drying the aerial part in oven at 65°C through constant mass, the shoot dry mass (SDM) was obtained.

The data were analyzed using the variance analysis with F test. The data in percentage were transformed to arco seno $\sqrt{x/100}$ before the statistical analysis but the original means were described on the tables and graphics. The Duncan test was used for the cultivars comparison. For the mulch analysis a regression analysis were utilized.

RESULTS AND DISCUSSION

The initial characterization of physiological potential for sunflower and peanut cultivars (Table 1) shows that all cultivars fitted the commercialization patterns. The three sunflower cultivars presented the same physiological level while the peanut cultivar IAC-Caiapó showed slightly lower level. Those results indicated the adequate physiological quality of all cultivars to be studied.

Table 1. Characterization of physiological potential for the seeds of sunflower and peanut cultivars in relation to initial germination level (G), final emergence (EM), and speed emergence index (SEI).

	G (%)	EM (%)	SEI
Sunflower			
IAC-iarama	99a ¹	100a	1,66a
Catissol	95a	100a	1,74a
Helio 358	98a	100a	1,51a
Peanut			
IAC-Caiapó	88b	86a	0,95a
Runner 886	92a	83a	0,87a
Tatu	97a	84a	1,04a

¹Means followed by the same letter in column, for each specie, did not differ from Duncan 's test at 5% probability

For sunflower there were significant interactions between mulch amount and cultivars only for shoot dry mass (SDM), while for peanut the interactions were significant only for seedling emergence (EM). There was almost no variation for the shoot dry mass of all sunflower cultivars with the increasing amount of mulch on soil surface (Table 2). The higher data presented by Helio-358 should be associated to the genetics vigour of hybrid.

Table 2. Means of sunflower shoot dry mass affected by mulch (M) and cultivars (C). Campinas-SP, Brazil.

Cultivars	Sugarcane mulch on the soil surface (t ha ⁻¹)					Adjustment equation and coefficient of determination (%)	
	0	5	10	15	20		
Sunflower – Shoot dry mass – SDM (g)							
IAC-iarama	1.6b ¹	1.5b	1.6b	1.4b	1.4b	Y = 1.59 - 0,01x	r ² =69
Catissol	1.5b	1.4b	1.4b	1.4b	1.7b	Y = 1.51 - 0,03x + 0.002x ²	r ² =86
Helio-358	2.0a	2.1a	1.9a	2.0a	1.9a	not significant	
M x C	0.05 ^{*2}						

¹Means followed by the same letter in column, for each specie, did not differ from Duncan 's test at 5% probability

The seedling emergence of all peanut cultivars was negatively affected by the mulch presence (Table 3); the cultivar Tatu was the least affected by the mulch on the soil surface which indicates that this cultivar should be the most indicated for the plough out sugarcane areas.

Table 3. Means of peanut seedling emergence affected by mulch (M) and cultivars (C). Campinas-SP, Brazil.

Cultivars	Sugarcane mulch on the soil surface (t ha ⁻¹)					Adjustment equation and coefficient of determination (%)	
	0	5	10	15	20		
Peanut -	Seedling emergence - EM (%)						
IAC-Caiapó	64.6a ¹	37.8b	23.2a	8.4b	2.6b	Y = 51.04 - 2.20x	r ² =99
Runner 886	60.8a	27.0b	16.5a	7.2b	7.3b	Y = 50.40 - 3.91x	r ² =99
Tatu	60.5a	57.1a	25.9a	48.4a	33.9a	Y = 51.22 - 0.97x	r ² =68
M x C	442.11**						

¹Means followed by the same letter in column, for each specie, did not differ from Duncan's test at 5% probability

The isolated effect of mulch amount and cultivars interfered with the evaluated parameters both for sunflower and peanut. The sunflower seedling emergence percentage (EM) and the speed emergence index (SEI) were directly affected by the mulch increasing on the soil surface (Figure 1). The greater seedling emergence reduction occurred with the introduction of 5 t ha⁻¹ (4 cm) of mulch on soil surface in relation to the tester with no mulch at all; between 15 t ha⁻¹ and 20 t ha⁻¹ the values did not change. The SEI followed the same pattern of reduction presented by seedling emergence (Figure 1B). According to Teasdale (1996), the mulch deposition on soil surface can cause chemical, physical and biological alterations in the environment and, depending on the plant specie, it can affect the seedling emergence and plant development. Mulch deposition is responsible for delay in the soil heat absorption, which interferes in the thermic difference between day and night, leading to a delay in the speed emergence index, in same cases, like what happened in the present research.




Fig. 1. Sunflower percentage of emergence (A), speed emergence index (B) and plant height (C) affected by the presence of sugarcane mulch on the soil. Also, sunflower plant height affected by cultivars (D). Campinas-SP, Brazil.

Differently from other parameters, the plant height of sunflower (Figure 1C) showed a small increase with the mulch thickness increasing. This was an expected behavior since with the increasing of mulch thickness the seedling elongates into sunlight direction (Carvalho & Nakagawa, 2000). In field conditions, plant shading leads to less biomass accumulation and the plants become more sensible to lodging (Correia & Durigan, 2004). There were also differences between cultivars (Figure 1D) with Helio-358 presenting the highest values for plant height in comparison to IAC-iarama and Catissol, which

did not differ between them. The better performance of Helio-358 could be related to its genetic vigour. Both seedling emergency and SEI did not vary between cultivars.

In relation to peanut crop (Figure 2), there was a significant negative effect of the sugarcane mulch on SEI (Figure 3A), plant height (Figure 2C), and dry shoot mass (Figure 2E). SEI was negatively influenced by the thickness of the mulch layer; with 10cm thick the SEI was 78% lower. In peanut the mulch negative effect was much more pronounced than in sunflower probably due to the higher temperature necessary for seedling emergence because this is tropical specie which center of origin is Brazil. Also the DSM was reduced with the mulch layer increasing up to 10t ha⁻¹; being constant after this thickness. Like sunflower, the peanut plant height was positively influenced by the mulch layer increasing, with the elongation of the plants, already described by Carvalho & Nakagawa (2000) for seedlings under light deficits.




Fig. 2. Means of peanut speed emergence index (A and B), plant height (C and D) and shoot dry mass (E and F) affected respectively by sugarcane mulch (first letter) and cultivars (second letter). Campinas-SP, Brazil.

The isolated effect of cultivars also is presented in Figure 2. The cv. Tatu presented the best performance in comparison to IAC-caiapó and Runner-886 for SEI (Figure 2B), HP (Figure 2D), and DSM (Figure 2F). This superiority could be related to its growing habit classified as erect, while 'IAC-Caiapó' and 'Runner 886' are classified as runner type. The erect plants have a tendency to grow up faster than the runner type. In the present research, cultivars with different growing habits were evaluated in order to verify if the erect cultivars would present better initial performance in comparison to the runner type which have a tendency to be more productive than the erect cultivar. In the past, the erect type was the most cultivated in Brazil and maybe will become an option for the plough out sugarcane areas with the mechanical harvest obligation.

The research showed that both emergency and initial sunflower development was less negatively influenced by the presence of sugarcane mulch than peanuts. So, in the first approach, sunflower seems to

have better conditions to give a good performance in areas with high levels of sugarcane mulch. Otherwise, field evaluations would be interesting to be done in the future.

CONCLUSION

For the greenhouse conditions is possible to conclude that the presence of different levels of sugarcane mulch on the soil surface can negatively influences both sunflower and peanut emergence and initial plant development. The negative effects are stronger for peanut cultivars, especially for the runner type; the cultivar Tatu was less influenced by the mulch thickness.

REFERENCES

- Ambrosano, E.J.; Guirado, N.; Ungaro, M.R.G.; Cantarella, H.; Mendes, P.C.D.; Rossi, F.; Ambrosano, G.M.B.; Sakai, R.H.; Schammass, E.A. 2005. Vantagens da utilização da rotação com girassol e outras leguminosas em áreas de reforma de canavial em Piracicaba, São Paulo. p.92-94. IN: Documentos Embrapa Soja n.261, 2005, Londrina.
- Borsari Filho, S. 2006. Potencial da cultura do amendoim como fonte de matéria-prima para o Programa Nacional de Produção e Uso de Biodiesel. . p.43-55. IN: Câmara, G.M.S.; Heiffig, L.S. Agronegócios de plantas oleaginosas – matérias-primas para biodiesel. FEALQ - Piracicaba.
- Brasil, 2006. Ministério da Agricultura, Pecuária e Abastecimento. Secretaria de Produção de Agroenergia Plano Nacional de Agroenergia 2011- 2006. 2 ed.rev. Brasília, DF: Embrapa Informação Tecnológica.
- Brasil, 1992. Ministério da Agricultura e Reforma Agrária. Regras para análise de sementes. Brasília: Agiplan, 365p.
- Carvalho, N.M.; Nakagawa, J. 2000. Sementes: ciência, tecnologia e produção. Jaboticabal: FUNEP. 650p.
- Correia, N.M.; Durigan, J.C. 2004 Emergência de plantas daninhas em solo coberto com palha de cana-de-açúcar. Planta Daninha, Viçosa, v.22, p.11-17.
- Furlani Neto, V.L.; Ripoli, T.C.; Vila Nova, N.A. 1997. Biomassa de cana-de-açúcar: energia contida no palhicho remanescente de colheita mecânica. Stab – Açúcar, Álcool e Subprodutos, v.15, p.24-27.
- Maguire, J.D. 1962. Speed of germination aid in selection and evaluation for seedling and vigor. Crop Science, Madison, v.2, n.2, p.176-177.
- Teasdale, J.R. 1996. Contribution of cover crops to weed management in sustainable agricultural systems. Journal of Production Agriculture, v.9, p.475-479.
- Trezzi, M.M.; Vidal, R.A. 2004. Potencial de utilização de cobertura vegetal de sorgo e milho na supressão de plantas daninhas em condição de campo: II – Efeitos da cobertura morta. Planta Daninha, v.22, p. 1-10.
- Van Raij, B.; Cantarella, H.; Quaggio, J.A.; Furlani, A.M.C. 1997. Recomendações de adubação e calagem para o estado de São Paulo, 2ª ed. Campinas, Instituto Agrônomo ; Fundação IAC. 285p.
- Vasconcelos, A.C.M. 2002. Desenvolvimento do sistema radicular da parte aérea de socas de cana-de-açúcar sob dois sistemas de colheita: crua mecanizada e queimada manual. 2002. 140p. Tese (Doutorado) - Universidade Estadual Paulista, Jaboticabal.
- Velini, E.D.; Negrisoni, E. 2000. Controle de plantas daninhas em cana-crua. p.148-164. In: Congresso Brasileiro da Ciência das Plantas Daninhas, 22., 2000, Foz Do Iguaçu.