

<b>Title</b>	<b>PGC-Enriched miRNAs Control Germ Cell Development</b>
<b>Author(s)</b>	<b>Bhin, J; Jeong, HS; Kim, JS; Shin, JO; Hong, KS; Jung, HS; Kim, C; Hwang, D; Kim, KS</b>
<b>Citation</b>	<b>Molecules and Cells, 2015, v. 38, p. 895-903</b>
<b>Issued Date</b>	<b>2015</b>
<b>URL</b>	<b><a href="http://hdl.handle.net/10722/224948">http://hdl.handle.net/10722/224948</a></b>
<b>Rights</b>	<b>This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License.</b>

# PGC-Enriched miRNAs Control Germ Cell Development

Jinhyuk Bhin<sup>1,5</sup>, Hoe-Su Jeong<sup>2,5</sup>, Jong Soo Kim<sup>2</sup>, Jeong Oh Shin<sup>3</sup>, Ki Sung Hong<sup>2</sup>, Han-Sung Jung<sup>3</sup>, Changhoon Kim<sup>2</sup>, Daehee Hwang<sup>1,4,\*</sup>, and Kye-Seong Kim<sup>2,\*</sup>

**Non-coding microRNAs (miRNAs) regulate the translation of target messenger RNAs (mRNAs) involved in the growth and development of a variety of cells, including primordial germ cells (PGCs) which play an essential role in germ cell development. However, the target mRNAs and the regulatory networks influenced by miRNAs in PGCs remain unclear. Here, we demonstrate a novel miRNAs control PGC development through targeting mRNAs involved in various cellular pathways. We reveal the PGC-enriched expression patterns of nine miRNAs, including miR-10b, -18a, -93, -106b, -126-3p, -127, -181a, -181b, and -301, using miRNA expression analysis along with mRNA microarray analysis in PGCs, embryonic gonads, and postnatal testes. These miRNAs are highly expressed in PGCs, as demonstrated by Northern blotting, miRNA in situ hybridization assay, and miRNA qPCR analysis. This integrative study utilizing mRNA microarray analysis and miRNA target prediction demonstrates the regulatory networks through which these miRNAs regulate their potential target genes during PGC development. The elucidated networks of miRNAs disclose a coordinated molecular mechanism by which these miRNAs regulate distinct cellular pathways in PGCs that determine germ cell development.**

## INTRODUCTION

Primordial germ cells (PGCs) are the embryonic precursors of

the gametes of the adult. During embryonic development, PGCs differentiate to germ cells, which develop oocytes in the female or spermatogonia in the male, and ultimately produce eggs or sperm, respectively. In mice, PGCs first emerge inside the extra-embryonic mesoderm at the posterior end of the primitive streak as a cluster of cells (Kucia et al., 2006) and then migrate toward the presumptive genital ridge, entering the genital ridge to colonize developing gonads. PGCs in gonads become distinguishable morphologically at 11.5 days post-coitum (dpc) and initiate differentiation toward the spermatogenic or oogenic morphogenesis at 12.5 dpc (McLaren, 1992; 2000). They are regarded as the primordial pluripotent cells because of their unique ability to preserve the developmental totipotency (Abe, 2007; De Felici et al., 2009; McLaren, 2003). Embryonic germ cells derived from PGCs and embryonic stem cells (ESCs) from the inner cell mass (ICM) have many similarities, but PGCs and ICM exhibit distinct self-renewal capability and lineage-specific characteristics (Pal et al., 2009; Watson and Tam, 2001).

MicroRNAs (miRNAs) are ~22 nucleotide, short, non-coding RNAs that regulate gene expression by inhibiting the transcription or translation of their target mRNAs (Bartel, 2004). Most miRNAs in animals function through the inhibition of effective mRNA translation of target genes through imperfect base pairing with the 3'-untranslated region (3' UTR) of target mRNAs. Thereby, the silencing mechanism of miRNAs is mediated by the inhibitory action of the RNA-induced silencing complex that both represses the translation and facilitates the degradation of targeted mRNAs (Filipowicz et al., 2008; Jaskiewicz and Filipowicz, 2008).

MiRNAs regulate growth and development in a variety of the cells (Kane et al., 2014). Also, a number of studies show that miRNAs regulate germ cell development (Banisch et al., 2012). During mouse germ cell development, the miR-290-295 and miR-17-92 clusters are known as significantly abundant miRNAs in PGCs. PGC-specific deficiency of Dicer, which plays a key role in microRNA biogenesis, results in poor PGC proliferation (Hayashi et al., 2008). MiR-290-295 cluster deficiency in mice results in embryonic lethality and germ cell defects with impaired PGC migration and premature ovarian failure (Medeiros et al., 2011). MiR-29b promotes female PGC development by targeting de novo DNA methyltransferase Dnmt3a and Dnmt3b (Takada et al., 2009). In chicken, miR-363 is involved in gonadal development (Huang et al., 2010). Moreover, miR-181a inhibits PGC differentiation by down-regulating

<sup>1</sup>Department of Chemical Engineering, Pohang University of Science and Technology, Pohang 790-784, Korea, <sup>2</sup>Graduate School of Biomedical Science and Engineering, Department of Biomedical Science, Hanyang University, Seoul 133-791, Korea, <sup>3</sup>Division in Anatomy and Developmental Biology, Department of Oral Biology, Oral Science Research Center, BK21 PLUS Project, Yonsei University, College of Dentistry, Seoul 120-752, Korea, <sup>4</sup>Department of New Biology and Center for Plant Aging Research, Institute for Basic Science, Daegu Gyeongbuk Institute of Science & Technology, Daegu 711-873, Korea, <sup>5</sup>These authors contributed equally to this work.

\*Correspondence: ks66kim@hanyang.ac.kr (KKS); dhhwang@dgist.ac.kr (DH)

Received 26 May, 2015; revised 25 June, 2015; accepted 8 July, 2015; published online 7 October, 2015

**Keywords:** bioinformatic analysis, *in situ* hybridization, male primordial germ cells, microarray, miRNAs

HOXA1 and PGC meiosis by targeting NR6A1 (Lee et al., 2011). Although such studies demonstrate that miRNAs play an essential role in the regulation of gene expression during PGC development, the function of miRNAs and the regulatory network of their target mRNAs remain unclear.

The aim of the present study is to identify novel PGC-enriched miRNAs that play an important role in PGC development and to reveal their regulatory networks. We performed a miRNA expression profiling analysis using a miRNA microarray and found nine novel miRNAs highly expressed in PGCs at 12.5 dpc compared to gonocytes (GCs) at 15.5 dpc, spermatogonial stem cells (SSCs) at 5 days post-partum (dpp), testes at four and eight weeks as well as ESCs. To find the potential target genes whose expression may be in inverse proportion to PGC-enriched miRNA expression, we analyzed the messenger RNA expression profiling in PGCs at 12.5 dpc, SSCs at 5 dpp, and testes at 12 dpp using an mRNA expression array. Therefore, we disclosed the potential target genes by PGC-enriched miRNAs and the regulatory networks of which PGC-enriched miRNAs regulate their target genes during PGC development. Collectively, PGC-enriched miRNAs determine germ cell development through regulating a variety of germ cell pathways, including cell growth, metabolic process, signaling, morphogenesis, and transcriptional regulation.

## MATERIALS AND METHODS

### Sample preparation

Fetal gonads were obtained from pregnant female ICR mice at 12.5 and 15.5 dpc and testes were collected from male ICR mice at 5 dpp, 12 dpp, four weeks, and eight weeks (Daehan Biolink Co. Ltd., Korea). Mouse R1 ES cell line (ATCC® SCRC-1036™) was cultured with a medium supplemented with 15% fetal bovine serum, LIF, pyruvate, glutamine,  $\beta$ -mercaptoethanol, 100 U/ml penicillin, and 100  $\mu$ g/ml streptomycin (Gibco BRL) in gelatin-coated plates and passaged with trypsin. The cells were cultured at 37°C in a humidified chamber containing 5% CO<sub>2</sub>. All animal experiments were carried out in accordance with the National Institutes of Health guide for the care and use of laboratory animals (NIH Publications No. 8023, revised 1978). These animal studies were approved by the Institutional Animal Care and Use Committee of Hanyang University College of Medicine, Korea.

### miRNA microarray analysis

Total RNAs were extracted from the cultured cells using Trizol reagent (Invitrogen, USA). For preparation of cellular miRNAs, small-sized RNAs containing miRNAs were isolated from total RNA using the RNeasy MinElute Cleanup kit (Qiagen, USA) according to the manufacturer's protocol. The isolated small-sized RNAs (~1  $\mu$ g) were subjected to direct labeling with a fluorescent dye using the Platinum Bright 647 Infrared nucleic acid labeling kit (KREATECH, Netherland) according to the manufacturer's instruction. After labeling, labeled RNAs were purified from free fluorescent substrates using KREA pure columns (KREATECH, Netherland) according to the manufacturer's instruction, and used in hybridization. A hybridization assay was carried out with Genopal®-MICM miRNA arrays (Mitsubishi Rayon, Japan) overnight, where 180-oligonucleotide DNA probes are installed for detection of mouse miRNAs in 150  $\mu$ l of hybridization buffer containing 2x SSC, 0.2% SDS, and ~1  $\mu$ g of heat-denatured labeled RNAs at 50°C. After hybridization, the miRNA chips were washed twice in 2x SSC containing 0.2% SDS at 50°C for 20 min followed by washing in 2x SSC at

50°C for 10 min. Hybridization signals were examined and analyzed using a DNA chip image analyzer according to the manufacturer's instruction (Mitsubishi Rayon, Japan).

### mRNA microarray analysis

Total RNAs were amplified and purified using the Ambion Illumina RNA amplification kit (Ambion, USA) to yield biotinylated cRNA according to the manufacturer's instruction. Briefly, 550 ng of total RNAs was reverse-transcribed using an oligo (dT) primer and second-strand cRNA were synthesized and labeled with biotin-dNTP. After purification, cRNA were quantified using the ND-1000 Spectrophotometer (NanoDrop, USA). The 750 ng of labeled cRNA samples was hybridized to the Illumina MouseRef-8-v2-BeadChip overnight at 58°C according to the manufacturer's instruction (Illumina, Inc., USA). The array was scanned using the BeadStation 500 System (Illumina, USA) to acquire the signal of the probes. The data were normalized as log<sub>2</sub> intensity values for each dataset using a quantile normalization method (Bolstad et al., 2003).

### Reverse transcription-polymerase chain reaction (RT-PCR)

One  $\mu$ g of DNase I-treatment total RNAs was reverse-transcribed using Superscript II reverse transcriptase with random hexamer (Invitrogen) according to the manufacturer's instruction. The standard PCR condition was as follows: 4 min at 94°C, followed by 25-35 cycles of 30 s at 94°C, 30 s at 55°C, and 30 s at 72°C. The primer sequences used in this study are given in Supplementary Table S1.

### miRNA quantitative real-time PCR assay

TaqMan miRNA assay was performed as previously described (Chen et al., 2005). Reverse transcription (RT) reactions were run in a GeneAmp PCR 9,700 Thermocycler (Applied Biosystems, USA). RT reaction without templates or primer was used as the negative controls. Gene expression levels were quantified using the ABI 7300 RT-PCR System (Applied Biosystems). A comparative real-time PCR including no template control was performed in triplicate using each specific primer set for miR-93, miR-106b, miR-18a, miR-181a, miR-199a, miR-10b, miR-126, miR-181b, miR-301, miR-127, miR-210, miR-376a, and snoRNA234 as the loading control. The reaction was performed at 95°C for 5 min, followed by 40 cycles of 95°C for 15 s, and 60°C for 1 min. All reagents and protocols were from Applied Biosystems (USA). The expression level of each miRNA relative to snoRNA234 was determined using the 2<sup>- $\Delta\Delta$ Ct</sup> method (Livak and Schmittgen, 2001).

### Northern blot analysis

The total RNAs were isolated from the cultured cells, including R1 ESCs, Sertoli cells, and MEF cells, and ICR mouse tissue (male gonads at 12.5 dpc and 15.5 dpc, testes at 5 dpp, four weeks, and eight weeks) using a Trizol reagent (Invitrogen, USA). Northern blotting was performed as previously described (Lagos-Quintana et al., 2001). The total RNA were separated in a 12.5% polyacrylamide gel containing 7 M urea, transferred to a Hybond-N+ membrane (GE Healthcare Life Sciences), and fixed by an ultraviolet cross-linker. Probes were end-labeled with [ $\gamma$ -32P] ATP (GE Healthcare Life Sciences) using T4 polynucleotide kinase (Takara, Japan). Prehybridization, hybridization, and wash were carried out at 42°C using ExpressHyb Hybridization Solution (Clontech, USA) according to the manufacturer's instruction. Blots were exposed on BioMax MS film (Kodak) at -80°C for 48 h. The 18S and 28S ribosomal RNA bands were stained with ethidium bromide and used as a loading

control. The sequences of probes were as follows: mmu-miR-93; 5'-caaagtgctgttcgagcaggt ag-3', mmu-miR-106a; 5'-caa agtgctaacagtgagcaggt ag-3', mmu-miR-181b; 5'-aacatt cat tgctgtcgg tggt-3', mmu-miR-18; 5'-taaggtgcatctagtgca gat ag-3'.

### miRNA *in situ* hybridization assay

For miRNA *in situ* hybridization, PGCs at 12.5 dpc and testes at 5 and 12 dpp were fixed with 4% paraformaldehyde (PFA) in 0.01 MPBS (pH 7.4) overnight at 4°C. After embedding in paraffin, the tissue was sectioned in 7 mm thicknesses. The sections were then baked at 65°C, de-waxed in xylene, rehydrated through graded washes of ethanol in water, and post-fixed in 4% PFA. The sections were pre-warmed and hybridized with the 5'-digoxigenin labeled locked nucleic acid (LNA) microRNA probes in a humid chamber containing 50% formamide in 2x SSC overnight at 20-25°C (Kloosterman et al., 2006). After washing with 2x SSC, the sections, an anti-DIG-antibody in TBS containing 10% FBS was applied and incubated at 4°C overnight. After washing with TBS, the sections were incubated in NTMT, and color development was enhanced by NBT and BCIP. The probes of miR-106b, -93, -18a, and -181a, and scramble were obtained from Exiqon.

### Bioinformatics

Identification of potential target genes by miRNAs: Potential targets by PGC-enriched miRNAs were generated through three miRNAs-target prediction programs, including TargetScan (Lewis et al., 2005), mirDB (Wang, 2008), and mirBase (Griffiths-Jones et al., 2006), in regard to the genes of each mRNA expression profiles. Mouse transcripts predicted to be miRNA targets were retrieved from the Sanger microRNA database.

Identification of differentially expressed genes (DEGs); We first defined the expressed genes across ICR mouse tissues at 12.5 dpc, 5 dpp, and 13.5 dpp using a Gaussian mixture modeling method as previously described (Lee et al., 2010). Among potential candidates, DEGs were identified by following three comparisons: 1) 12.5 dpc versus 5 dpp, 2) 12.5 dpc versus 12 dpp, 3) 5 dpp versus 12 dpp using an integrative statistical method previously reported (Chae et al., 2013). This method calculated adjusted P-values (P) by combining P-values from two-sample *t*-test and log<sub>2</sub>-median ratio test using empirical distributions of T-values and log<sub>2</sub>-median ratios by random permutations between samples. For each comparison, the DEGs were defined as the genes with P < 0.05 and log<sub>2</sub>-fold-changes > 0.58. DEGs were grouped into the eighteen clusters based on their differential expression patterns over time (Supplementary Fig. S1).

Gene ontology analysis: Gene ontology enrichment analysis was performed using a DAVID software (Huang da et al., 2009). Gene ontology Biological Processes (GOBPs) with P-value < 0.1 (a default cut-off) computed from DAVID were selected as the ones enriched by the genes in each group.

### Statistical analysis

Unless stated otherwise, each experiment was carried out using at least three replicates. Data were described as mean ± SEM. Statistically significant differences between groups were evaluated by one-way analysis of variance (ANOVA) using a log-linear model in the Statistical Analysis System (SAS, USA). Values of P < 0.05 were considered statistically significant.

## RESULTS

### PGC-enriched miRNAs during germ cell development


To examine the novel miRNAs predominantly expressed in PGCs during germ cell development, we performed miRNA expression profiling arrays in PGCs at 12.5 dpc, gonads at 15.5 dpc, testes at four and eight weeks, ESCs, Sertoli cells, and MEF cells using the Genopal®-MICM miRNA arrays as described in "Materials and Methods". Among the 180 miRNAs, we selected 12 miRNAs that were expressed specifically in PGCs with criteria of a significant (P < 0.05) increase (> 1.5-fold) in PGCs compared to controls. These 12 miRNAs, including miR-10b, -18a, -93, -106b, -126-3p, -127, -181a, -181b, -301, -199a, -210, and -376a, were expressed highly in PGCs at 12.5 dpc and gonads at 15.5 dpc compared to other controls (Fig. 1A). The expression levels of these miRNAs decreased with mature germ cell development (Fig. 1A). Of the 12 miRNAs, some miRNAs, such as miR-301, -106b, -93, and -18a, were expressed highly in PGCs, whereas miR-199a, -127, -376a, and -210 were expressed more in gonads at 15.5 dpc. Therefore, 12 miRNAs meet the criteria as potential regulatory microRNA candidates that are highly and specifically expressed in PGCs.

The accuracy of the miRNA microarray analysis is affected by the short length of mature miRNA sequences leading to an increase in the false discovery rate. Therefore, we conducted miRNA qPCR analysis to validate the PGC-enriched miRNA expression (Fig. 1B). Based on the restriction criteria of miRNA expression in PGCs, we isolated nine PGC-enriched miRNAs including miR-10b, -18a, -93, -106b, -126-3p, -127, -181a, -181b, and -301, during PGC development. miR-199a, -210, and -376a were also expressed in PGCs (Supplementary Fig. S2), but the expression level of these miRNAs was smaller compared to that of the selected nine miRNAs (data not shown). Since the PGC-enriched expression of miR-199a, -210, and -376a fell below selection criteria, these miRNAs were excluded from PGC-enriched miRNAs.

A Northern blot analysis confirmed again the PGC-enriched miRNA expression of miR-18a, -93, -106b, and -181a (Fig. 1C). To narrow our conclusion, we conducted a miRNA *in situ* hybridization assay to examine the PGC-enriched miRNA expression of miR-18a, -93, -106b, and -181a. These miRNAs were expressed precisely in PGCs at 12.5 dpc (Fig. 1D), but not in other tissues and other germ stages (Supplementary Fig. S3). Together, our data indicate that nine novel miRNAs are expressed in a PGC-dependent fashion, suggestive of their roles in regulating PGC development.

### Target genes by PGC-enriched miRNAs

Our results indicate that PGC-enriched miRNAs may regulate PGC development. To further establish this, we performed an integrative analysis using the comparison of the mRNA expression profiling between PGCs at 12.5 dpc, SSCs at 5 dpp, and testes at 12 dpp. We found a total of 4,171 DEGs from three comparisons as follows: 1) 12.5 dpc versus 5 dpp (2,313 DEGs), 2) 12.5 dpc versus 12 dpp (2,177 DEGs), and 3) 5 dpp versus 12 dpp (2,226 DEGs). These DEGs were assembled into the eighteen expression patterns (C1-C18) (Supplementary Fig. S1A and Supplementary Table S2). Among them, the gene expression patterns in C1 (29 genes) and C2 (343 genes) were inversely correlated with the PGC-enriched miRNA expression (Fig. 2A and Supplementary Fig. S1B). It seems that some genes in C1 and C2 may be the potential targets by PGC-enriched miRNAs. To test the notion, we checked the potential targets of our nine miRNAs using three microRNA target prediction programs; TargetScan, mirDB, and mirBase (Griffiths-Jones et al., 2006; Lewis et al., 2005; Wang, 2008). According to our analysis, a total of 6,511




**Fig. 1.** The expression levels of miRNAs during male germ cell development. (A) Expression profiling of 12 PGC-enriched miRNAs identified by miRNA microarray in developing male germ cells. For each miRNA, the mean-centered  $\log_2$ -expression profile (mean = 0) is shown. A hierarchical clustering was done to visualize the similarity in expression patterns of the miRNAs. The colors represent increased (red) and decreased (green) expression relative to the mean expression level. (B) PGC-enriched expression of miR-10b, -18a, -93, -106b, -126-3p, -127, -181a, -181b, and -301 (red) among 12 miRNAs identified from a miRNA microarray analysis were confirmed by miRNA qPCR. (C) Northern blot analysis demonstrates that miR-18a, -93, -106b, and -181a were expressed in PGCs during male germ cell development. In the bottom, Ethidium bromide staining of 28S, 18S, and 5S rRNAs is shown as a loading control. Northern blot assay was performed as described in Materials and Methods. (D) miRNA in situ hybridization assay demonstrated the PGC-enriched expression patterns of miRNAs in PGCs at 12.5 dpc. Arrow, a seminiferous cord; black arrowhead, primordial germ cells; white arrowhead, Sertoli cells.

genes was targeted by miR-10b, -18a, -93, -106b, -127, -181a, -181b, and -301. The 127 genes could be the potential targets because of affiliating to C1 and C2 (Fig. 2B and Supplementary Table S3). As expected, the expression pattern of 127 genes exhibited a reverse relationship with that of PGC-enriched miRNAs (Fig. 2C). Interestingly, miR-106b and -93 are reported to be transcribed together as a common polycistronic transcript and to be a member of the same miRNA cluster, miR-17-92 cluster (Olive et al., 2010). Thus, miR-106b and -93 have highly similar mature sequences and may share target genes. In line with that possibility, miR-106b and -93 shared 39 common targets among a total of 49 genes (Fig. 2D and Supplementary Table S3). Additionally, miR-181a and -181b are clustered on chromosome 1 and co-transcribed. MiR-181a and -181b had 28 common targets among a total of 36 potential genes (Fig. 2D and Supplementary Table S3).

#### PGC-enriched miRNAs regulate cellular processes

PGCs arrived at the genital ridges and become incorporated into sex cords at 11.5 dpc. They continued to proliferate until 13.5-14.5 dpc and then entered a quiescent period. After birth, GCs resumed mitosis and initiated their differentiation into spermatogonia. During this process, GCs undergo a variety of possible morphological and metabolic changes including apoptosis (Fujimoto et al., 2000; Honke et al., 2002; Zhang et al., 2005). To explore the physiological effects on GC development by stage specific expression of nine candidate miRNAs, we examined the cellular processes represented by 127 target genes based on the enrichment analysis of GOBPs using DAVID software (Huang da et al., 2009). The GOBPs of the 127 target genes indicated their involvement in germ cell development mechanisms, such as cell replication, morphogenesis, apoptosis, regulatory intracellular signaling cascades, and phospholipid metabolic processes (Fig. 3A and Supplementary



**Fig. 2.** Identification of target genes by PGC-enriched miRNAs and their biological processes. (A) The expression of target genes categorized in C1 and C2 showed a reverse correlation with nine PGC-enriched miRNAs during germ cell development. The auto-scaled log<sub>2</sub>-expression profiles of each target gene (mean = 0 and standard deviation = 1) are shown (gray line). Median expression values in C1 and C2 for each sample are expressed as blue and red lines. (B) The strategy for identifying target genes by nine PGC-enriched miRNAs. 6,511 targets were generated from the three databases using the sequence pairing algorithms. Among them, 127 were selected as target genes having a reverse expression (C1 and C2) with miRNA expression. (C) A heat map showed expression patterns of the 127 target genes. For each gene, the mean-centered log<sub>2</sub>-expression profile (mean = 0) is shown. A hierarchical clustering was done to visualize the similarity in expression patterns of the genes. The colors represent increased (red) and decreased (green) expression relative to the mean expression level. (D) miR-106b and -93 shared target genes due to possession of similar seed sequence. Similarly, miR-181a and -181b shared target genes.

Table S4). Our results suggest that the miRNA-specific targets could be down-regulated after birth to prevent precocious germ cell development. Similarly, the induction of target genes coinciding with regulation of cell cycle and apoptosis after birth suggests support for the resumption of mitosis and the restriction of germ cell loss necessary for proper germ cell development as previously noted (Kondoh et al., 1991).


#### Regulatory networks affected by PGC-enriched miRNAs

MiRNAs are known to regulate germ cell development (Banisch et al., 2012). To understand the regulatory roles of miRNAs in PGC development, we demonstrated a network of which nine miRNAs appeared to regulate PGC development through changing target gene expression. Of the 127 target genes as


members of C1 and C2, 92 genes were clustered to the known functions based on their GOBP annotations according to DAVID software (Huang da et al., 2009). Thus, we created a network model containing the 12 major modules grouped by the 92 nodes (Fig. 3B), including 1) apoptosis, 2) cell cycle and proliferation, 3) differentiation and morphogenesis, 4) structure organization, 5) intracellular signaling cascades, 6) transcription regulation, 7) transport and localization, 8) cell adhesion and migration, 9) cellular metabolic process, 10) RNA processing, 11) proteolysis, and 12) immune and defense response.

To explore the information-theoretic models for complex networks, we assembled the relative contribution by each PGC-enriched miRNAs in response to the 12 cellular processes (Fig. 4A). We calculated the score as a relative contribution of target


**A**



**B**



**Fig. 3.** A network model delineates biological processes regulated by PGC-enriched miRNAs. (A) The enrichment analysis results of GOBPs by total genes in C1 and C2 (red bar, total genes) and 127 target genes (blue bar). The red line denotes the cutoff value ( $P$ -value = 0.1) of the analysis. (B) A network model shows the relationships between nine PGC-enriched miRNAs and their target genes. Colors and shapes of the nodes represent PGC-enriched miRNAs (green diamonds), target genes in C1 (purple circles) and C2 (red circles). Edge colors were used to distinguish the interactions of individual miRNAs or miRNA families. The nodes were grouped according to their gene ontologies (colored rectangles). Nodes containing the same gene ontologies belong to same group as described in “Materials and Methods”. The genes with multiple gene ontologies were comprised of two or more functional groups. The background colors represent gene ontology groups (ellipsoidal for lower-level gene ontologies and rectangle for upper-level gene ontologies).



**Fig. 4.** A heat map represents the relationship between each PGC-enriched miRNA and the twelve biological processes. (A) A heat map shows the contribution score defined by the number of target genes regulated by each miRNA within a gene ontology group divided by the total number of the target genes of the miRNA within the group. (B) RT-PCR analysis confirms expression of the representative target genes during male germ cell development.

genes by individual miRNAs. Notably, this analysis revealed that the miR-17-92 cluster, including miR-93 and -106b, plays a crucial role in the regulation of the 12 cellular processes during germ cell development. Especially, the contribution score implied that miR-93 and -106b have an anti-apoptotic role by targeting apoptotic Cidea (Figs. 3A and 4B, and Supplementary Table S3). Our prediction is consistent with the previous conclusions that miR-106b acts as a hub molecule in an apoptotic miRNA-gene network (Liu et al., 2012) and the suppression of miR-106b-93-25 cluster expression is essential for ER stress-mediated apoptosis (Gupta et al., 2012). To corroborate the above suggested correlations in light of a network model, the effect on expression levels of target genes by individual miRNAs was examined by RT-PCR (Fig. 4B). As target gene expression was indeed decreased with germ cell development, it is suggestive that these targets might be regulated by our miRNAs.

Moreover, the contribution score implicated that miR-106b and -93 have a role in specific metabolic processes, especially phospholipid metabolism, through targeting Abhd5 and Gab1 (Fig. 4B). Moreover, miR-18a, another member of miR-17 family, regulated a broad spectrum of processes for germ cell development, such as differentiation/morphogenesis, RNA processing, transcription regulation, and cell cycle, by targeting Rhox8, Klf9, Camta2, Mbnl1, Celf2, Lmo4, Foxq1, Rassf2, and Ccng2 (Fig. 4B). Our suggestion is in agree-

ment with the previous observation that miR-18a regulates male germ cell maturation (Bjork et al., 2010). Furthermore, miR-181a and -181b negatively regulated the expression of targets, including Mbnl1 and Prkra, that modulate differentiation and development based on the contribution score (Fig. 4B). Therefore, miR-10b, -18a, -181b, and -301 along with miR-181a, which is expressed highly in chicken PGCs, might cooperatively regulate germ cell development, repress PGC differentiation, and keep PGCs from entering meiosis. Finally, the contribution analysis showed that other PGC-enriched miRNAs, such as miR-10b, -126-3p, -127, and -301, may regulate diverse cellular processes during germ cell development. Based on the contribution score, 1) miR-126-3p could regulate cell adhesion/migration, transport/localization, and transcription regulation, 2) miR-10b may regulate apoptosis, RNA processing, transport/localization, and immune/defense response, 3) miR-127 might regulate differentiation/morphogenesis, and 4) miR-301 could regulate cell cycle/proliferation, structure organization, signaling cascade, differentiation/morphogenesis, and adhesion/migration. Taken together, our results demonstrate that nine PGC-enriched miRNAs play a key role in the regulation of a variety of cellular processes related to PGC development. Furthermore, our network model could extend the current understanding of the regulatory roles of miRNAs during germ cell development.

## DISCUSSION

MiRNAs play an essential role in regulation of specification, migration, and differentiation in the gametes (Dyce et al., 2010; Jessberger, 2008; Tang, 2010). Though the Surani group reported that miRNAs, such as miR-290-295 and miR-17-92 clusters, are expressed highly in PGCs along with germ cell development (Hayashiet al., 2008), a systematic analysis of PGC-enriched miRNAs has not been explored. In this study, we find that nine novel miRNAs, including miR-10b, -18a, -93, -106b, -126-3p, -127, -181a, -181b, and -301, are expressed selectively in PGCs according to our comparative study profiling miRNA expression of PGCs at 12.5 dpc, GCs at 15.5 dpc, SSCs at 5 dpp, and testes at four and eight weeks. To understand the functional roles of these miRNAs, we investigated the potential target genes by an integrative analysis with the sequence pairing information and anti-correlated expression profile data between miRNAs and target mRNAs. Using the anti-relationship between miRNAs and their target genes, we are able to present a network model delineating how these nine miRNAs regulate the PGC developmental processes, including differentiation and morphogenesis, cell proliferation and apoptosis, cell adhesion and migration, cellular signaling cascades, and other cellular metabolic processes.

A variety of miRNA target prediction algorithms are based on the sequence pairing between miRNAs and target transcripts (Griffiths-Jones et al., 2006; Lewis et al., 2005). Recently, the miRNA integrative analysis using *in silico* predictions and the data of microRNA and mRNA expression profiling shows an inverse expression correlation between miRNA and target mRNA. This method provides more precise target prediction compared to the conventional sequence pairing algorithms (Huang et al., 2007; Joung et al., 2007; Peng et al., 2009; Tran et al., 2008; Wang, 2008). Using a newly developed integrative analysis, we propose a network model to understand the regulatory roles of nine PGC-enriched miRNAs.

Of the nine miRNAs, miR-18a, -93, and -106b belonging to the miR-17-92 cluster were previously reported to be highly expressed in mouse PGCs (Hayashi et al., 2008). In addition, miR-181a was known to be highly expressed in chicken PGCs and to inhibit the somatic differentiation of PGCs by silencing homeobox A1 (Lee et al., 2011). On the other hand, five miRNAs, such as miR-10b, -126-3p, -127, -181b, and -301, have not been previously reported in relation to PGC development. miR-18a, one of the nine PGC-enriched miRNAs, may mediate male germ cell maturation by inhibiting a transcription factor HSF2 that influences a wide range of developmental process (Bjorket al., 2010). Although we do not find that HSF2 belongs to our DEGs in PGCs development, our network model suggests that some target genes, such as *Camta2*, *Mbnl1*, *Rhox8*, and *Foxq1*, targeted by miR-18a may contribute to PGC development. Two clusters of the miR-17 family, miR-17-92 and miR-106b-25 clusters, might regulate germ cell differentiation by decreasing *Bim*, *Kit*, *Socs3*, and *Stat3* (Tong et al., 2012). Moreover, *Gab1*, *Casp8*, *Foxq1*, *Lmo4*, *Pkd1*, and *Mef2c* by miR-106b and/or -93 possibly will control PGC development. *Vcam-1*, which is targeted by the miR-126-3p (Harris et al., 2008) and *Pkd1*, which is targeted by the miR-127 along with miR-93 and -106b, mediate cell adhesion/migration and differentiation/morphology, suggesting that miR-126-3p and -127 may be involved with cell adhesion/migration and differentiation/morphology in PGCs. Additionally, the miR-181 family including miR-181a and -181b expresses in PGCs of the chicken (Lee et al., 2011), as well as in human (Sirotkin et al., 2009) and

mouse (Saunders et al., 2010). Sirtuin-1 is a target of miR-181a (Zhou et al., 2012). Sirtuin-1 null mice showed to be sterile in males and sirtuin-1 also participated with growth hormone/insulin-like growth axis (Xu et al., 2010). Collectively, the miR-181 family including miR-181a and -181b play a key role in reproductive development.

During germ cell development, massive changes in gene control and expression, such as alternative splicing and mitotic-to-meiotic transition also occur (Schmid et al., 2013). In support of these, the miR-181a/b and miR-10b modulate mechanisms of RNA processing, such as spliceosome assembly and mRNA splicing site selection in PGCs (Fig. 4A and Supplementary Tables S3). Moreover, *Tbc1d9b*, which is targeted by miR-301, and *Arhgef3* and *Arhgef4*, whose expression is decreased by the miR-17 family, mediate small GTPase-mediated signaling pathways, suggesting miR-17 family and miR-301 involvement in Ras signaling in PGCs.

In summary, our study provides a demonstration and comprehensive bioinformatics analysis of nine novel miRNAs that are likely candidates to be important in PGC miRNA-mediated regulatory networks involved in development of PGCs, such as those regulating essential mechanisms of differentiation and morphogenesis, cell proliferation and apoptosis, cell adhesion and migration, and cellular signaling cascades, as well as other cellular metabolic processes.

*Note: Supplementary information is available on the Molecules and Cells website ([www.molcells.org](http://www.molcells.org)).*

## ACKNOWLEDGMENTS

We are grateful to Professor Eric B. Richardson, M.D., Hanyang University, for giving the constructive comments and editing our manuscript. This work was supported by grant from the Medical Research Center (2008-0062190) and by Institute of Basic Science (IBS-R013-G1-2015) funded by the National Research Foundation of Korea (NRF) of the Ministry of Science, ICT & Future Planning, Republic of Korea.

## REFERENCES

- Abe, K. (2007). Developmental program for pluripotential cells and primordial germ cells in mice. *Tanpakushitsu Kakusan Koso* 52, 2046-2053.
- Banisch, T.U., Goudarzi, M., and Raz, E. (2012). Small RNAs in germ cell development. *Curr. Topics Dev. Biol.* 99, 79-113.
- Bartel, D.P. (2004). MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell* 116, 281-297.
- Bjork, J.K., Sandqvist, A., Elsing, A.N., Kotaja, N., and Sistonen, L. (2010). miR-18, a member of Oncomir-1, targets heat shock transcription factor 2 in spermatogenesis. *Development* 137, 3177-3184.
- Bolstad, B.M., Irizarry, R.A., Astrand, M., and Speed, T.P. (2003). A comparison of normalization methods for high density oligonucleotide array data based on variance and bias. *Bioinformatics* 19, 185-193.
- Chae, S., Ahn, B.Y., Byun, K., Cho, Y.M., Yu, M.H., Lee, B., Hwang, D., and Park, K.S. (2013). A systems approach for decoding mitochondrial retrograde signaling pathways. *Sci. Signal.* 6, rs4.
- Chen, C., Ridzon, D.A., Broomer, A.J., Zhou, Z., Lee, D.H., Nguyen, J.T., Barbisin, M., Xu, N.L., Mahuvakar, V.R., Andersen, M.R., et al. (2005). Real-time quantification of microRNAs by stem-loop RT-PCR. *Nucleic Acids Res.* 33, e179.
- De Felici, M., Farini, D., and Dolci, S. (2009). In or out stemness: comparing growth factor signalling in mouse embryonic stem cells and primordial germ cells. *Curr. Stem Cell Res. Ther.* 4, 87-97.
- Dyce, P.W., Toms, D., and Li, J. (2010). Stem cells and germ cells: microRNA and gene expression signatures. *Histol. Histopathol.* 25, 505-513.

- Filipowicz, W., Bhattacharyya, S.N., and Sonenberg, N. (2008). Mechanisms of post-transcriptional regulation by microRNAs: are the answers in sight? *Nat. Rev. Genet.* 9, 102-114.
- Fujimoto, H., Tadano-Aritomi, K., Tokumasu, A., Ito, K., Hikita, T., Suzuki, K., and Ishizuka, I. (2000). Requirement of seminolipid in spermatogenesis revealed by UDP-galactose: Ceramide galactosyltransferase-deficient mice. *J. Biol. Chem.* 275, 22623-22626.
- Griffiths-Jones, S., Grocock, R.J., van Dongen, S., Bateman, A. and Enright, A.J. (2006). miRBase: microRNA sequences, targets and gene nomenclature. *Nucleic Acids Res.* 34, D140-144.
- Gupta, S., Read, D.E., Deepti, A., Cawley, K., Gupta, A., Oommen, D., Verfaillie, T., Matus, S., Smith, M.A., Mott, J.L., et al. (2012). Perk-dependent repression of miR-106b-25 cluster is required for ER stress-induced apoptosis. *Cell Death Dis.* 3, e333.
- Harris, T.A., Yamakuchi, M., Ferlito, M., Mendell, J.T., and Lowenstein, C.J. (2008). MicroRNA-126 regulates endothelial expression of vascular cell adhesion molecule 1. *Proc. Natl. Acad. Sci. USA* 105, 1516-1521.
- Hayashi, K., Chuva de Sousa Lopes, S.M., Kaneda, M., Tang, F., Hajkova, P., Lao, K., O'Carroll, D., Das, P.P., Tarakhovskiy, A., Miska, E.A., et al. (2008). MicroRNA biogenesis is required for mouse primordial germ cell development and spermatogenesis. *PLoS One* 3, e1738.
- Honke, K., Hirahara, Y., Dupree, J., Suzuki, K., Popko, B., Fukushima, K., Fukushima, J., Nagasawa, T., Yoshida, N., Wada, Y., et al. (2002). Paranal junction formation and spermatogenesis require sulfoglycolipids. *Proc. Natl. Acad. Sci. USA* 99, 4227-4232.
- Huang, J.C., Babak, T., Corson, T.W., Chua, G., Khan, S., Gallie, B.L., Hughes, T.R., Blencowe, B.J., Frey, B.J., and Morris, Q.D. (2007). Using expression profiling data to identify human microRNA targets. *Nat. Methods* 4, 1045-1049.
- Huang, P., Gong, Y., Peng, X., Li, S., Yang, Y., and Feng, Y. (2010). Cloning, identification, and expression analysis at the stage of gonadal sex differentiation of chicken miR-363 and 363\*. *Acta Biochim. Biophys. Sin (Shanghai)* 42, 522-529.
- Huang da, W., Sherman, B.T., and Lempicki, R.A. (2009). Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat. Protocols* 4, 44-57.
- Jaskiewicz, L., and Filipowicz, W. (2008). Role of Dicer in posttranscriptional RNA silencing. *Curr. Top Microbiol. Immunol.* 320, 77-97.
- Jessberger, R. (2008). New insights into germ cell tumor formation. *Horm. Metab. Res.* 40, 342-346.
- Joung, J.G., Hwang, K.B., Nam, J.W., Kim, S.J., and Zhang, B.T. (2007). Discovery of microRNA-mRNA modules via population-based probabilistic learning. *Bioinformatics* 23, 1141-1147.
- Kane, N.M., Thrasher, A.J., Angelini, G.D., and Emanuelli, C. (2014). Concise review: MicroRNAs as modulators of stem cells and angiogenesis. *Stem Cells* 32, 1059-1066.
- Kloosterman, W.P., Wienholds, E., de Bruijn, E., Kauppinen, S., and Plasterk, R.H. (2006). *In situ* detection of miRNAs in animal embryos using LNA-modified oligonucleotide probes. *Nat. Methods* 3, 27-29.
- Kondoh, G., Murata, Y., Aozasa, K., Yutsudo, M., and Hakura, A. (1991). Very high incidence of germ cell tumorigenesis (seminomagenesis) in human papillomavirus type 16 transgenic mice. *J. Virol.* 65, 3335-3339.
- Kucia, M., Machalinski, B., and Ratajczak, M.Z. (2006). The developmental deposition of epiblast/germ cell-line derived cells in various organs as a hypothetical explanation of stem cell plasticity? *Acta Neurobiol. Exp. (Wars)* 66, 331-341.
- Lagos-Quintana, M., Rauhur, R., Lendeckel, W., and Tuschl, T. (2001). Identification of novel genes coding for small expressed RNAs. *Science* 294, 853-858.
- Lee, H.J., Suk, J.E., Patrick, C., Bae, E.J., Cho, J.H., Rho, S., Hwang, D., Maslah, E., and Lee, S.J. (2010). Direct transfer of alpha-synuclein from neuron to astroglia causes inflammatory responses in synucleinopathies. *J. Biol. Chem.* 285, 9262-9272.
- Lee, S.I., Lee, B.R., Hwang, Y.S., Lee, H.C., Rengaraj, D., Song, G., Park, T.S., and Han, J.Y. (2011). MicroRNA-mediated posttranscriptional regulation is required for maintaining undifferentiated properties of blastoderm and primordial germ cells in chickens. *Proc. Natl. Acad. Sci. USA* 108, 10426-10431.
- Lewis, B.P., Burge, C.B., and Bartel, D.P. (2005). Conserved seed pairing, often flanked by adenosines, indicates that thousands of human genes are microRNA targets. *Cell* 120, 15-20.
- Liu, Z., Yang, D., Xie, P., Ren, G., Sun, G., Zeng, X., and Sun, X. (2012). MiR-106b and MiR-15b modulate apoptosis and angiogenesis in myocardial infarction. *Cell. Physiol. Biochem.* 29, 851-862.
- Livak, K.J., and Schmittgen, T.D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods* 25, 402-408.
- McLaren, A. (1992). Development of primordial germ cells in the mouse. *Andrologia* 24, 243-247.
- McLaren, A. (2000). Germ and somatic cell lineages in the developing gonad. *Mol. Cell. Endocrinol.* 163, 3-9.
- McLaren, A. (2003). Primordial germ cells in the mouse. *Dev. Biol.* 262, 1-15.
- Medeiros, L.A., Dennis, L.M., Gill, M.E., Houbaviy, H., Markoulaki, S., Fu, D., White, A.C., Kirak, O., Sharp, P.A., Page, D.C., et al. (2011). Mir-290-295 deficiency in mice results in partially penetrant embryonic lethality and germ cell defects. *Proc. Natl. Acad. Sci. USA* 108, 14163-14168.
- Olive, V., Jiang, I., and He, L. (2010). mir-17-92, a cluster of miRNAs in the midst of the cancer network. *Int. J. Biochem. Cell Biol.* 42, 1348-1354.
- Pal, R., Totey, S., Mamidi, M.K., and Bhat, V.S. (2009). Propensity of human embryonic stem cell lines during early stage of lineage specification controls their terminal differentiation into mature cell types. *Exp. Biol. Med.* 234, 1230-1243.
- Peng, X., Li, Y., Walters, K.A., Rosenzweig, E.R., Lederer, S.L., Aicher, L.D., Proll, S., and Katze, M.G. (2009). Computational identification of hepatitis C virus associated microRNA-mRNA regulatory modules in human livers. *BMC Genomics* 10, 373.
- Saunders, L.R., Sharma, A.D., Tawney, J., Nakagawa, M., Okita, K., Yamanaka, S., Willenbring, H., and Verdin, E. (2010). miRNAs regulate SIRT1 expression during mouse embryonic stem cell differentiation and in adult mouse tissues. *Aging* 2, 415-431.
- Schmid, R., Greltscheid, S.N., Ehrmann, I., Dalglish, C., Danilenko, M., Paronetto, M.P., Pedrotti, S., Greltscheid, D., Dixon, R.J., Sette, C., et al. (2013). The splicing landscape is globally reprogrammed during male meiosis. *Nucleic Acids Res.* 41, 10170-10184.
- Sirotkin, A.V., Ovcharenko, D., Grossmann, R., Laukova, M., and Mlyncek, M. (2009). Identification of microRNAs controlling human ovarian cell steroidogenesis via a genome-scale screen. *J. Cell. Physiol.* 219, 415-420.
- Takada, S., Berezikov, E., Choi, Y.L., Yamashita, Y., and Mano, H. (2009). Potential role of miR-29b in modulation of Dnmt3a and Dnmt3b expression in primordial germ cells of female mouse embryos. *RNA* 15, 1507-1514.
- Tang, F. (2010). Small RNAs in mammalian germline: tiny for immortal. *Differentiation* 79, 141-146.
- Tong, M.H., Mitchell, D.A., McGowan, S.D., Evanoff, R., and Griswold, M.D. (2012). Two miRNA clusters, Mir-17-92 (Mirc1) and Mir-106b-25 (Mirc3), are involved in the regulation of spermatogonial differentiation in mice. *Biol. Reprod.* 86, 72.
- Tran, D.H., Satou, K., and Ho, T.B. (2008). Finding microRNA regulatory modules in human genome using rule induction. *BMC Bioinform.* 9 Suppl 12, S5.
- Wang, X. (2008). miRDB: a microRNA target prediction and functional annotation database with a wiki interface. *RNA* 14, 1012-1017.
- Watson, C.M., and Tam, P.P. (2001). Cell lineage determination in the mouse. *Cell Struct Funct.* 26, 123-129.
- Xu, F., Gao, Z., Zhang, J., Rivera, C.A., Yin, J., Weng, J., and Ye, J. (2010). Lack of SIRT1 (Mammalian Sirtuin 1) activity leads to liver steatosis in the SIRT1<sup>+/+</sup> mice: a role of lipid mobilization and inflammation. *Endocrinology* 151, 2504-2514.
- Zhang, Y., Hayashi, Y., Cheng, X., Watanabe, T., Wang, X., Taniguchi, N., and Honke, K. (2005). Testis-specific sulfoglycolipid, seminolipid, is essential for germ cell function in spermatogenesis. *Glycobiology* 15, 649-654.
- Zhou, B., Li, C., Qi, W., Zhang, Y., Zhang, F., Wu, J.X., Hu, Y.N., Wu, D.M., Liu, Y., Yan, T.T., et al. (2012). Downregulation of miR-181a upregulates sirtuin-1 (SIRT1) and improves hepatic insulin sensitivity. *Diabetologia* 55, 2032-2043.