



Title	The influence of pre-existing health conditions on short-term mortality risks of temperature: Evidence from a prospective Chinese elderly cohort in Hong Kong
Author(s)	Sun, S; Tian, L; Qiu, H; Chan, KP; Tsang, H; Tang, HKR; Lee, RSY; Thach, TQ; Wong, CM
Citation	Environmental Research, 2016, v. 148, p. 7-14
Issued Date	2016
URL	http://hdl.handle.net/10722/224929
Rights	This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License.

1 **The influence of pre-existing health conditions on short-term mortality risks of**
2
3 **temperature:**
4
5
6 **evidence from a prospective Chinese elderly cohort in Hong Kong**
7
8
9

10
11 Shengzhi Sun ^a, Linwei Tian ^a, Hong Qiu ^a, King-Pan Chan ^a, Hilda Tsang ^a, Robert Tang ^a,
12
13 Ruby Siu-yin Lee ^b, Thuan-Quoc Thach ^a, Chit-Ming Wong ^{a,*}
14
15
16

17
18
19
20 **Authors' affiliations:**
21

22 ^a School of Public Health, The University of Hong Kong, Hong Kong SAR, China
23
24

25 ^b Elderly Health Service, Department of Health, Hong Kong SAR, China
26
27

28
29
30
31 *** Corresponding author:**
32

33 Chit-Ming Wong, PhD
34

35 School of Public Health
36

37 The University of Hong Kong
38

39 5th Floor, Faculty of Medicine Building
40

41 21 Sassoon Road, Hong Kong SAR, China
42

43 Phone: (852) 2831 5071
44

45 Fax: (852) 2855 9528
46

47 E-mail: hrmrwcm@hku.hk
48
49
50
51
52
53
54
55
56
57
58
59
60
61

1
2
3 **Abstract (word: 242)**
4
5

6 **Background:**
7

8 Both cold and hot temperatures are associated with adverse health outcomes. Less is known
9 about the role of pre-existing medical conditions to confer individual's susceptibility to
10 temperature extremes.
11
12
13
14
15

16 **Methods:**
17

18 We studied 66,820 subjects aged ≥ 65 who were enrolled and interviewed in all the 18
19 Elderly Health Centers of Department of Health, Hong Kong from 1998 to 2001, and
20 followed up for 10-13 years. The distributed lag nonlinear model (DLNM) combined with a
21 nested case-control study design was applied to estimate the nonlinear and delayed effects of
22 cold or hot temperature on all natural mortality among subjects with different pre-existing
23 diseases.
24
25
26
27
28
29
30
31
32
33
34
35

36 **Results:**
37

38 The relative risk of all natural mortality associated with a decrease of temperature from 25th
39 percentile (19.5 °C) to 1st percentile (11.3 °C) over 0-21 lag days for participants who reported
40 to have an active disease at the baseline was 2.21 (95% confidence interval (CI): 1.19, 4.10)
41 for diabetes mellitus (DM), 1.59 (1.12, 2.26) for circulatory system diseases (CSD), and 1.23
42 (0.53, 2.84) for chronic obstructive pulmonary disease (COPD), whereas 1.04 (0.59, 1.85) for
43 non-disease group (NDG). Compared with NDG, elders with COPD had excess risk of
44 mortality associated with thermal stress attributable to hot temperature, while elders with DM
45 and CSD were vulnerable to both hot and cold temperatures.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 **Conclusions:**

2
3 Elders with pre-existing health conditions were more vulnerable to excess mortality risk to
4
5
6 hot and/or cold temperature. Preventative measures should target on elders with chronic
7
8
9 health problems.
10

11
12
13
14 **Keywords:** Distributed lag nonlinear model; Elders; Medical condition; Mortality;
15
16
17 Temperature; Nested case-control study
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 **List of abbreviations and their full forms**

2

3

4

5 **Abbreviations** **Full form**

6

7 DLNM The distributed lag nonlinear model

8

9

10 CI Confidence interval

11

12 COPD Chronic obstructive pulmonary disease

13

14

15 BMI Body mass index

16

17 ICD-9 Ninth revision of the international classification of diseases

18

19

20 PM_{2.5} Particulate matter with aerodynamic diameter less than or equal to 2.5 μm

21

22 TPU Tertiary planning unit

23

24

25 AIC Akaike information criterion

26

27 DM Diabetes mellitus

28

29

30 CSD Circulatory system diseases

31

32 NDG Non-disease group

33

34

35 SD Standard deviation

36

37 RR Relative risk

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

1 **List of Tables and Figures**

2
3
4 **Table 1.** Baseline characteristics of the participants, according to pre-existing health
5 conditions in the prospective Chinese elderly cohort in Hong Kong (n=61,441).
6

7
8 **Table 2.** All natural mortality after 10-13 years of following up at the end of study in 2011 by
9 pre-existing health conditions.
10

11
12 **Table 3.** The cumulative relative risks of cold and hot effects on all natural mortality in elders
13 with pre-existing diseases along lag days.
14

15
16 **Fig. 1.** The estimated relative risk of temperature centered at median (24.7 °C) on all natural
17 mortality over 21 days for subjects with different pre-existing diseases. NDG: non-disease
18 group; DM: diabetes mellitus; CSD: circulatory system diseases; COPD: chronic obstructive
19 pulmonary disease.
20
21

22
23 **Fig. 2.** The estimated relative risk of cold temperature effects [a decrease from 25th percentile
24 of temperature (19.5 °C) to 10th percentile of temperature (16.2 °C)] and hot effects [an
25 increase from 75th percentile of temperature (27.8 °C) to 90th percentile of temperature (29.4
26 °C)] on all natural mortality over 21 days of lag. NDG: non-disease group; DM: diabetes
27 mellitus; CSD: circulatory system diseases; COPD: chronic obstructive pulmonary disease.
28 The solid lines are mean relative risks, and grey regions are 95% CIs.
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 **1. Introduction**

2
3 Climate change is likely to bring extreme temperature events more frequently with increased
4
5
6 intensity and widespread distribution. It was observed that global warming not only increased
7
8
9 the frequency of heat waves, but also led to increase in cold spells with a steeper rate than in
10
11
12 hot spells from 1981 to 2010 worldwide (Song et al., 2013). Temperature extremes had short-
13
14
15 term adverse impacts on health in that heat waves were associated with increased mortality,
16
17
18 hospital admissions, and emergency room visits (Astrom et al., 2011; Khalaj et al., 2010;
19
20
21 Knowlton et al., 2009; Williams et al., 2012). However, only a few studies have assessed the
22
23
24 effects of cold temperature (Medina-Ramon and Schwartz, 2007; Medina-Ramon et al., 2006;
25
26
27 Ou et al., 2013; Rocklov et al., 2014).

28
29
30
31 Extreme temperature events affect vulnerable populations most, amplifying social disparities
32
33
34 in health, both within and between countries (Meehl and Tebaldi, 2004). Understanding
35
36
37 susceptibility to extreme temperature is of importance in environmental public health
38
39
40 researches (Anderson and Bell, 2009; Gasparrini, 2014). However, there were few studies for
41
42
43 effects of thermal stress on susceptible subgroups (Astrom et al., 2015; Bateson and Schwartz,
44
45
46 2004; Ma et al., 2015), fewer were studies for effects of cold temperature and in persons with
47
48
49 medical conditions. This may be due to: a lack of personal information to define susceptible
50
51
52 groups in conventional studies with ecological designs (e.g. time-series, case-crossover, or
53
54
55 case-only approaches); a lack of suitable statistical methods to assess effect modifiers of
56
57
58 time-varying exposure in studies with cohort design; also the difficulties encountered in
59
60
61 analyzing nonlinear and delayed effects for the relationship between temperature and

1 mortality.

2
3
4
5
6 Among studies to explore whether medical conditions may confer susceptibility to
7
8
9 temperature, most of them did not directly assess effects in persons with pre-existing health
10
11 problems but instead indirectly through assessing mortality outcomes for specific causes
12
13 (Atsumi et al., 2013) . This approach is not appropriate, since the underlying cause of death
14
15 may not necessarily provide the information of existing medical conditions before death. For
16
17 example, people died from cardiovascular diseases may have diabetes mellitus and persons
18
19 with diabetes mellitus not be reflected in the underlying cause. Thus vulnerability of persons
20
21 with specific health problems cannot be assessed.
22
23
24
25
26
27
28
29
30

31 In the study, we aimed to assess the nonlinear and delayed mortality effects of cold or hot
32
33 temperature among elders in a prospective Chinese elderly cohort in Hong Kong, and to
34
35 comparatively assess effect modification of pre-existing diseases.
36
37
38
39
40
41

42 **2. Methods**

43 *2.1. Study subjects*

44
45 We included a total of 66,820 adults aged 65 years or older (≥ 65 years), about 9 % of people
46
47 aged ≥ 65 years in Hong Kong, who enrolled at one of the eighteen Elderly Health Centres of
48
49 the Department of Health from 1998 to 2001 and followed up until 31st of December 2011.
50
51
52

53
54 Body mass index (BMI) was collected by physical examinations, and socio-economic
55
56 conditions, lifestyle characteristics (smoking, physical activity, education), and morbidity
57
58
59
60
61

1 status were collected by face-to-face interview during enrolment and follow-up visits by
2
3 registered nurses (Schooling et al., 2014; Wong et al., 2015). Ethics approval was obtained
4
5
6 from the Ethics Committee of the Faculty of Medicine, The University of Hong Kong and of
7
8
9 the Department of Health of Hong Kong.

10
11
12
13
14 Self-reported active chronic disease was collected through face-to face interview, which
15
16
17 included hypertension, heart diseases, stroke, diabetes mellitus, and COPD and/or asthma.

18
19
20 Self-reports of chronic diseases were confirmed and supplemented by clinical diagnoses
21
22
23 based on history (Schooling et al., 2006). All natural mortality was coded as 1-799 according
24
25
26 to the International Classification of Diseases, Ninth revision (ICD-9) before 2001 or A00-
27
28
29 R99 based on ICD-10 from 2001 by the Department of Health. Subjects who died within the
30
31 first year after recruitment were excluded from our analysis.

32 33 34 35 36 *2.2. Air pollutants and meteorological data*

37
38
39 We obtained air pollutants data of ten general monitoring stations from Environmental
40
41
42 Protection Department of Hong Kong from 1999 to 2011. We took daily 24-hour average
43
44
45 concentration of particulate matter with aerodynamic diameter less than or equal to 2.5 μm
46
47
48 ($\text{PM}_{2.5}$) and nitrogen dioxide (NO_2), and daily 8-hour (10:00-18:00 hours) average
49
50
51 concentration of ozone (O_3). Data were regarded as missing if numbers of hourly
52
53
54 concentration for one particular day were less than 75 % (18 hours for $\text{PM}_{2.5}$ and NO_2 , and 6
55
56
57 hours for O_3). Daily concentrations of air pollutants were evaluated by averaging the daily
58
59
60 concentrations across all valid monitoring stations.

1
2
3 Daily mean air temperature and relative humidity in percent were measured by digital
4
5
6 thermometers with platinum resistance sensors at the Hong Kong Observatory. The historical
7
8
9 temperature data have been used in the previous population studies of temperature effects in
10
11
12 Hong Kong (Goggins et al., 2015; Yi and Chan, 2014).
13

14 15 16 17 *2.3. Statistical analysis* 18

19
20 The nested case-control design employs a case-control approach within an established cohort
21
22 to obtain estimates from a sample of the cohort that are similar to estimates obtained from
23
24 analysis of the entire cohort (Goldstein and Langholz, 1992). This design has become popular
25
26 because it allows for computationally efficient analysis of data from a cohort with substantial
27
28 savings in cost and time; it is a useful alternative for cohort analysis with studying time-
29
30 varying exposures (Essebag et al., 2005). The nested case-control design has already been
31
32 used to studying time-varying exposure, such as PM_{2.5} (Beverland et al., 2012).
33
34
35
36
37
38
39
40
41

42 In this study, we used nested case-control design to assess short-term effects of temperature.
43
44 We constructed nested case-control data sets as follows: (1) controls selected randomly from
45
46 among the cohort subjects whose follow-up time at least as long as the case for each death in
47
48 the cohort; (2) the associated date of controls (the date when the control reached the exact
49
50 follow-up time as the case) was in the same calendar year and month as the death date of case
51
52 to control long-term trend and seasonality; (3) the date of birth of controls was within 1
53
54
55
56
57
58
59
60
61
62
63
64
65

1 outside the follow-up period of the cohort. We randomly selected 9 controls to each case from
2
3 all eligible subjects in the cohort (Beverland et al., 2012). For all participants (including
4
5 persons with and without morbidity status), about 98% of cases had the full complement of 9
6
7 controls, and for diabetes mellitus (DM), circulatory system diseases (CSD), chronic
8
9 obstructive pulmonary disease (COPD), and non-disease group (NDG), about 92%, 97%,
10
11 80%, and 93.4% respectively.
12
13
14
15
16
17
18
19

20 Temperature appears a nonlinear and delayed relationship with mortality (Guo et al., 2011). A
21
22 distributed lag nonlinear model (DLNM) has been developed based on a “cross-basis”
23
24 function, which allows simultaneously estimate the nonlinear and delayed effects of
25
26 temperature on mortality or morbidity in time-series data, and this approach was expanded to
27
28 apply to cohort study recently (Gasparrini, 2014).
29
30
31
32
33
34
35

36 We used conditional logistic regression to incorporate a DLNM in a nested case-control study
37
38 design to assess the nonlinear and delayed mortality effects of temperature among subjects in
39
40 a prospective Chinese elderly cohort, and to comparatively examine the effect modification of
41
42 morbidity status and the extent of mortality displacement related to high temperature.
43
44
45
46
47
48
49

$$50 \quad \text{logit } P = T_{t,l} + b_1 X_i + b_2 X_i(t) + \text{Strata}(\text{strata}) \quad (1)$$

51
52
53
54

55 where t is the day of death for case or an associated day of each control; $T_{t,l}$ is a matrix
56
57 obtained by applying the DLNM to temperature, and l is the number of lag days; X_i is the
58
59
60
61
62
63
64
65

1 vector of time-independent variables, including individual-level factors: gender, BMI (three
2 levels), smoking (three levels), physical exercise (four levels), education (three levels),
3 tertiary planning unit (TPU)-level factors: proportion of the population ≥ 65 years of age,
4 proportion with tertiary education, and the average monthly income in each TPU, and district
5 level factor: proportion of smokers in each district; $X_i(t)$ is the vector of time-dependent
6 variables, including dummy variable of day of the week on day t , and natural cubic B-splines
7 with 3 d.f. for relative humidity (Guo et al., 2011; Stafoggia et al., 2008). β_1 and β_2 are
8 vectors of coefficients for time-independent and time-dependent variables; *Strata* (*strata*) is a
9 categorical variable of each risk set.
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27

28 $T_{t,l}$ in model (1) was the term to estimate both the nonlinear and delayed effects of
29 temperature. In order to completely capture the overall effects of temperature, we extended
30 lags up to 21 days. We placed knots of natural cubic B-splines at equal intervals on the log
31 scale of lags. We placed knots of natural cubic B-splines at equal spaces in the range of
32 temperature, which was from 1999 to 2011. Minimum Akaike Information Criterion (AIC)
33 was used to guide selection of the d.f. (knots) for temperature and lag days (Gasparrini,
34 2014). It was found that using one knot for temperature and one knot for lag days produced
35 the best model fit for all disease categories in the present study.
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52

53 To examine the extreme cold effect, we calculated the relative risk of all natural mortality
54 associated with a decrease from 25th percentile of temperature (19.5 °C) to 1st percentile of
55 temperature (11.3 °C); and the relative risk associated with a decrease from 25th (19.5 °C) to
56
57
58
59
60
61
62
63
64
65

1 10th (16.2 °C) was used to represent the moderate cold effect (Goldberg et al., 2011; Guo et al.,
2
3 2012). Meanwhile, we calculated the relative risk of all natural mortality associated with
4
5 temperature increased from 75th (27.8 °C) to 99th (30.4 °C), and from 75th (27.8 °C) to 90th
6
7 (29.4 °C) to represent the extreme and moderate hot effect, respectively. Cumulative effects
8
9 over 0-21 days and the relative risk along the lags for both cold and hot temperatures were
10
11 examined and plotted, the mortality displacement was assessed as well.
12
13
14
15
16
17
18
19

20 The following sensitivity analyses were conducted to examine the robustness of the effect
21
22 estimates:
23

- 24
25 1. Random selection of 5, 7, and 12 controls for each case from all eligible subjects in
26
27 the cohort;
28
29
- 30
31 2. Adjustment for each of the three air pollutants PM_{2.5}, NO₂, and O₃ in the main
32
33 temperature model as a natural cubic B-splines function with 3 d.f. (Guo et al., 2011);
34
35
- 36
37 3. Time-stratified case-crossover study design incorporated with DLNM to estimate
38
39 temperature mortality effects, in that controls were selected with the same year, month,
40
41 and weekday as the case.
42
43
- 44
45 4. Increasing the number of knots for both temperature and lag days to 2;
46
47
48
49

50 Mortality risks associated with temperature between subjects with pre-existing health
51
52 conditions and NDG were compared based on calculated Z-score and corresponding one-
53
54 sided *p*-Value (Altman and Bland, 2003). Other statistical tests were all two-sided, and values
55
56 of *p*<0.05 were considered statistically significant. We used the SAS version 9.2 software
57
58
59
60
61
62
63
64
65

1 package to construct the dataset of nested case-control study. We used R software (version
2
3 3.1.2; R Development core team 2014) to fit all models, with its “dlnm” package (version
4
5 2.1.3) to perform distributed lag nonlinear models (Gasparrini et al., 2010). The SAS and R
6
7 code used are available in Supplementary Material.
8
9

10 11 12 13 14 **3. Results** 15

16 17 *3.1. Descriptive statistics* 18

19
20 A total of 66,820 subjects were recruited in the prospective Chinese elderly cohort of Hong
21
22 Kong. After excluding subjects who died within first year post recruitment and subjects with
23
24 missing data, 61,441 were left in our final analysis. Table 1 shows the baseline characteristics
25
26 of participants, stratified according to pre-existing active diseases. The characteristics of the
27
28 subjects were similar in most respects across pre-existing active disease groups. But subjects
29
30 with COPD tended to be more smokers (quitted or current), older and less females when
31
32 compared with other morbidity subgroups. Mean age at enrollment was 72 years. About two
33
34 third (66%) of subjects were female; a major part of subjects had never smoked (71.3%), and
35
36 did exercise more than six times per week (72.1%).
37
38
39
40
41
42
43
44
45
46

47 [Table 1 here]
48
49
50
51
52

53 Among 61,441 subjects at recruitment, 47.3% subjects reported to have CSD, 14.2% to have
54
55 DM, 6.2% to have COPD, and 45.4% subjects reported they did not have those three diseases
56
57 (NDG). After 10-13 years of following up till the end of the study on 31 December 2011,
58
59
60
61
62
63
64
65

1 16,497 (26.9%) subjects died among which 1,855 were subjects with COPD, 3,024 were
2
3 subjects with DM, 9,070 were with CSD, and 5,809 were NDG. The corresponding death
4
5 rates were 48.4%, 34.6%, 31.2% and 20.8, respectively (Table 2).
6
7
8
9

10
11 [Table 2 here]
12
13
14
15
16

17 *3.2. Cumulative effects of cold and hot temperatures*

18

19
20 Table 3 presents the cumulative relative risks of extreme cold effects (decreasing from 25th to
21
22 1st percentile of mean temperature), moderate cold (decreasing from 25th to 10th percentile),
23
24 moderate hot (increasing from 75th to 90th percentile), and extreme hot (increasing from 75th
25
26 to 99th percentile) on all natural mortality for subjects with different pre-existing diseases.
27
28

29
30 The cumulative effects of cold temperatures over longer lags were greater than effects over
31
32 0-3 lag days for all subjects and subgroups with DM and CSD, while the cumulative effects
33
34 of hot temperatures didn't vary much along the lags, except for subjects with DM.
35
36
37
38
39
40
41

42 Compared with the temperature at 25th percentile (19.5 °C), the relative risks of extreme cold
43
44 temperature (1st percentile of temperature, 11.3 °C) over lag 0-21 days were 2.21 (95% CI:
45
46 1.19, 4.10) for DM, 1.59 (1.12, 2.26) for CSD, 1.23 (0.53, 2.84) for COPD, and 1.49 (1.15,
47
48 1.93) for all subjects respectively, while 1.04 (0.59, 1.85) for NDG. Compared with the
49
50 temperature at 75th percentile (19.5 °C), the relative risks associated with extreme hot
51
52 temperature (99th percentile of temperature, 30.4 °C) over lag 0-3 days were 1.05 (95% CI:
53
54 0.94, 1.17) for DM, 1.06 (0.99, 1.12) for CSD, 1.14 (0.98, 1.32) for COPD, and 1.01 (0.97,
55
56
57
58
59
60
61
62
63
64
65

1 1.06) for all subjects respectively. We did not observe any significant association between
2
3 cold/hot temperatures and all natural mortality in NDG.
4
5
6
7

8 9 *3.3. Comparisons with NDG*

10
11 When compared with NDG over the same lag periods, the cumulative cold effects over 0-13
12 lag days were significantly greater for elders with DM ($p<0.05$), for elders with CSD ($p<0.1$)
13 and for all subjects ($p<0.1$), while the cumulative hot effects over 0-3 lag days appeared
14 significant greater for subjects with COPD and CSD ($p<0.05$), and subjects with DM ($p<0.1$).
15
16 Results showed that subjects with DM and CSD exhibited greater vulnerability to both cold
17 and hot weather, while subjects with COPD had excess mortality risk attributable to hot
18 temperature (Table 3).
19
20
21
22
23
24
25
26
27
28
29
30

31
32
33 [Table 3 here]
34
35
36

37 *3.4. Temperature-lag-mortality relationships*

38
39 The 3-dimensional plots show the distributed lag and non-linear relationship between all
40 natural mortality and temperature for subjects with different morbidity status at all lag days
41 (see Supplementary Material, Fig. S1). The overall temperature-mortality relationships in
42 general were U-shaped over all 21 days with increased health effects in both cold and high
43 temperatures. Overall cold temperature effects were much higher than hot effects (Fig. 1).
44
45
46
47
48
49
50
51
52
53
54

55
56 [Figure 1 here]
57
58
59
60
61
62
63
64
65

1 Generally, cold temperature effects were delayed and lasted for about 20 days, while hot
2
3 effects were immediate and lasted for less than 5 days. Cold temperature effects were delayed
4
5 for subjects with COPD and CSD, but not for subjects with DM (Fig. 2). Furthermore,
6
7 mortality displacement related to hot temperature was not found in subjects with DM.
8
9
10
11
12
13
14
15
16

17 [Figure 2 here]
18
19
20
21

22 3.5. Sensitivity analysis 23

24
25 For results of sensitivity analysis, in general, the mortality effects of temperature become
26
27 stable when more than 7 controls were selected for each case (see Supplementary Material,
28
29 Table S1 to Table S3). When increasing number of knots for both temperature and lag days to
30
31 two, or controlling for each of the three air pollutants (PM_{2.5}, NO₂, and O₃) or using time-
32
33 stratified case-crossover study design to estimate temperature mortality effects, the effect
34
35 estimates remained similar (see Supplementary Material, Table S4 to Table S6).
36
37
38
39
40
41
42
43

44 4. Discussion 45

46
47 To the best of our knowledge, this is the first study to assess the nonlinear and delayed
48
49 mortality effects of cold or hot temperature for persons with baseline existing health problem
50
51 in a cohort. The major findings of this study are: (1) persons with pre-existing disease are
52
53 more vulnerable to hot and/or cold temperature; (2) cold temperature effects are more
54
55 pronounced than hot temperature effects; and (3) cold temperature effects are delayed and
56
57
58
59
60
61
62
63
64
65

1 persist for about 20 days, while hot effects are immediate and last for shorter period.
2
3
4
5

6 *4.1. Mortality risks of temperature* 7

8
9 To date several studies have examined the temperature-mortality relationship worldwide
10
11 (Basu, 2009; Gasparrini et al., 2012; Hajat et al., 2005; Hajat et al., 2010; Nordio et al., 2015),
12
13 but few focused on persons with pre-existing active diseases were based on case-only study
14
15 design (Khalaj et al., 2010; Medina-Ramon et al., 2006; Schwartz, 2005; Zanobetti et al.,
16
17 2013). The major disadvantage of case-only study to identify susceptible subgroups is that the
18
19 excess risk of mortality associated with extreme temperatures in each subpopulation cannot
20
21 be estimated on its own, but only relative to other subpopulations. Consequently, we cannot
22
23 obtain absolute effects of extreme temperature in a susceptible subgroup.
24
25
26
27
28
29
30

31
32
33 Indeed we did not find any studies that assessed absolute mortality effects of temperature for
34
35 persons with existing diseases directly. In addition, the magnitude of temperature effects
36
37 varies greatly by climate, region, and population (Curriero et al., 2002). To avoid these
38
39 variations, we compared temperature mortality effects for all subjects (including persons with
40
41 or without morbidity) with previous studies conducted in Hong Kong. We compared with the
42
43 study of Xu et al. (2013), who used the same cohort data, and reported that hot temperature
44
45 effects were non-significant ($p < 0.05$) at lag 0-1 for all natural mortality, which were
46
47 consistent with our results. Cold temperature effects are usually delayed and lasting for a few
48
49 weeks, thus we compared our results with the study of Yi and Chan (2014), who used quasi-
50
51 Poisson model combined with a DLNM in time-series study design, and reported that cold
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 effects in terms of relative risk for decrease in temperature from 1st to 25th percentile at lag 0-
2
3 13 and lag 0-21 for subjects aged older than 75 years was 1.37 (1.26, 1.47) and 1.41 (1.35,
4
5 1.51), respectively. These results were comparable with results of our study (mean age of
6
7 subjects at baseline was 72 years, and followed up for 10 to 13 years).
8
9

10
11
12 Our results suggest that persons with DM increase susceptibility to both cold and hot
13
14 temperature. These findings agree with study conducted in two Chinese cities (Harbin and
15
16 Chongqing), which showed significant associations between both extreme hot and cold
17
18 temperatures and DM mortality (Li et al., 2014). Most studies reported that people with DM
19
20 have a high risk of death (Medina-Ramon et al., 2006; Schwartz, 2005; Zanobetti et al., 2013),
21
22 hospitalizations (Knowlton et al., 2009), and emergency department visits (Khalaj et al., 2010)
23
24 during heat wave or in high temperature, but few have indicated that persons with DM are
25
26 sensitive to cold temperature (Li et al., 2014). Mortality displacement is of public health
27
28 importance, and it has been well documented in the literature that hot effects appear more
29
30 acute and followed by harvesting (Hajat et al., 2005; Qiao et al., 2015). A novel finding of our
31
32 study is that mortality displacement related to hot temperature was not found in subjects with
33
34 DM, which suggests that persons with DM are at risk of premature death associated with
35
36 hotter temperature. Temperature not only can have an impact on very frail individual but also
37
38 on relatively healthy persons who have DM.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55

56 DM is characterized by a chronic state of low-grade inflammation, endothelial dysfunction,
57
58 and hypercoagulability (Luchsinger et al., 2001), and it has already been reported that several
59
60
61
62
63
64
65

1 factors, such as oxidative stress and protein kinase C, may contribute to macrovascular injury
2
3 from hyperglycemia. When exposed to extreme temperatures, blood vessels dilate or constrict
4
5 in order to maintain body temperature. The increased demands on the circulatory system
6
7 interact with those injured vasculature would increase risk of fatal events (Hajat et al., 2005;
8
9 Idris et al., 2006). Thus, it is biological plausible that persons with DM may increase
10
11 susceptibility to both cold and hot temperature.
12
13
14
15
16
17
18
19

20 Our results showed that subjects with CSD were at higher risk of death from hot temperature,
21
22 and also suggested higher mortality risk from cold temperature, which have been consistently
23
24 confirmed by previous studies using cause-specific mortality (Gasparri et al., 2012; Ma et
25
26 al., 2011; Ma et al., 2013; Medina-Ramon et al., 2006). In order to adapt to heat when
27
28 exposed to hot temperature, cardiac output shapely increased primarily by increased heart
29
30 rate, which put a great burden to heart (Rowell, 1974). Cold temperature can increase heart
31
32 rate, blood pressure, peripheral vasoconstriction, plasma fibrinogen concentration, blood
33
34 cholesterol level, platelet viscosity, and red cell count (Keatinge et al., 1984), which changes
35
36 may predispose the subjects to arterial thrombosis and lead to heart attack and stroke or even
37
38 death (Atsumi et al., 2013; Mercer, 2003; Nayha, 2002).
39
40
41
42
43
44
45
46
47
48
49

50 In our study, we found mortality displacements related to hot effects for subjects with CSD
51
52 and COPD, but they were not statistically significant. Previous studies have indicated that
53
54 mortality displacement related to hot stress was more strongly associated with deaths from
55
56 cardiovascular and respiratory diseases than all natural mortality (Goodman et al., 2004;
57
58
59
60
61
62
63
64
65

1 Hajat et al., 2005). Qiao et al. (2015) reported mortality displacement for respiratory deaths
2
3 was more immediate and greater than cardiovascular deaths, which were also similarly shown
4
5
6 in our findings.
7
8
9

10
11 We found persons with COPD were susceptible to increased risk of mortality when exposed
12
13 to hot temperature, but not to cold temperature. This findings are consistent with findings of
14
15 Schifano et al. (2009) and Michelozzi et al. (2009) who reported that people with COPD have
16
17 increased mortality and hospital admissions when exposed to high temperature, and
18
19 consistent with findings of Medina-Ramon et al. (2006), Ma et al. (2013), and Astrom et al.
20
21 (2015) who did not find higher mortality risks when exposed to cold temperature. But our
22
23 results are conflict with findings of Schwartz (2005) who showed that persons with COPD
24
25 had higher mortality risks attributable to extreme cold temperature than persons did not have
26
27 that condition in a case-only study. One possible explanation might be that persons with
28
29 COPD were more likely to confine to bed or stay in homes during cool seasons.
30
31
32
33
34
35
36
37
38
39
40
41

42 *4.2. The nested case-control study design and DLNM*

43

44 Many study designs have been used in assessing health effects of temperature on mortality
45
46 and morbidity, such as case-crossover (Guo et al., 2011; Medina-Ramon and Schwartz, 2007),
47
48 time-series (Anderson and Bell, 2009; Wu et al., 2013), and Bayesian hierarchical models
49
50 (Anderson et al., 2013; Bobb et al., 2014). Generally, these study designs are using
51
52 aggregated data with the limitation of not being able in identifying more potential modifiers
53
54 of exposures (e.g. temperature, air pollutants). However, cohort study can record detailed
55
56
57
58
59
60
61
62
63
64
65

1 individual information (e.g. demographic characteristics, socioeconomic position, lifestyle,
2
3 social functional status), and has a great potential to thoroughly explore modifiers of
4
5 exposure effects. Basically, nested case-control design is a type of case-control design nested
6
7 in a cohort study. This study design has been widely used (Lin et al., 2015; Olsson et al.,
8
9 2010; Sawada et al., 2010) due to its being able to control for long-term trend and seasonality
10
11 by study design through matching, which is particular useful when studying time-varying
12
13 exposure. The nested case-control design is a useful alternative to Cox regression to analyze
14
15 time-varying exposure, due to similar risk estimates with superior computational efficiency
16
17 (Essebag et al., 2005).
18
19
20
21
22
23
24
25
26
27

28 One of the main advantages of DLNM is that it allows the model to contain detailed
29
30 representation of exposure-response relationship. This in turn provides an estimate of the
31
32 overall effect in the presence of delayed contributions or harvesting with constraint on lag
33
34 days to resolve the high correlation between exposures in adjacent days and the resulting
35
36 collinearity in the model (Gasparrini et al., 2010; Zanobetti et al., 2002). Therefore, applying
37
38 DLNM in a nested case-control is applicable to estimate health effects of time-varying
39
40 exposure as well as mortality displacement in a cohort.
41
42
43
44
45
46
47
48
49

50 This is the first attempt to assess mortality effects of temperature after incorporating DLNM
51
52 to a nested case-control study design. This method has potential applications to identify
53
54 modifiers of time-varying exposure (e.g. temperature, air pollutants) on morbidity and
55
56 mortality. For instance, our approach can be applied to explore whether people living in high-
57
58
59
60
61
62
63
64
65

1 polluted areas has a higher risk to thermal stress than people living in low-polluted areas.
2
3
4
5

6 Our study has some limitations. First, medical conditions were self-reported, thus it is subject
7
8
9 to misclassification. Second, our cohort subjects being older than 65 years at recruitment may
10
11
12 have multiple health problems, which may not reflect the specific modifying effect of certain
13
14
15 existing disease. However, subjects with DM may probably comorbid with CSD or
16
17
18 respiratory diseases, the higher cold risk in this group confirms that persons with diabetes
19
20
21 mellitus may be susceptible to excess mortality risk associated with temperature stress. Last
22
23
24 but not least, the subjects were volunteers; however, in Hong Kong, the enrolment to the
25
26
27 Elderly Health Service is free and open to all residents who are 65 years or older. Thus, the
28
29
30 sample would be representative of the elderly population in Hong Kong (Schooling et al.,
31
32
33 2006). Even if the enrolment is subject to volunteer bias, that bias is towards those who are
34
35
36 health conscious, our results tend to be conservative.
37
38

39 **5. Conclusions**

40
41
42 DLNM can be applied in a nested case-control study design in a cohort to estimate nonlinear
43
44
45 and delayed effects of time-varying exposure (e.g. temperature). Mortality effects of cold
46
47
48 temperature may be more pronounced than hot temperature. Elders with pre-existing health
49
50
51 conditions are more vulnerable to excess mortality risk at short-term exposure to hot and/or
52
53
54 cold temperature. Our study provides evidence to enhance protection of the vulnerable ones.
55
56
57
58
59
60
61
62
63
64
65

1 **Conflicts of interest**

2
3 The authors declare no conflicts of interest.
4
5
6
7

8
9 **Acknowledgements:**

10
11 We thank the Hong Kong Government departments including Department of Health (Elderly
12 Health Service) for the cohort data and mortality data and Environmental Protection
13
14 Department for air pollution data. This project was supported by the earmarked grant GRF
15
16
17
18
19
20 780512 of the Hong Kong Research Grants Council.
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

References

- 1
2
3 Altman, D.G., Bland, J.M., 2003. Interaction revisited: the difference between two estimates. *B.M.J.* 326, 219.
- 4 Anderson, B.G., Bell, M.L., 2009. Weather-related mortality: how heat, cold, and heat waves affect mortality in
5 the United States. *Epidemiology* 20, 205-213.
- 6
7 Anderson, G.B., Dominici, F., Wang, Y., McCormack, M.C., Bell, M.L., Peng, R.D., 2013. Heat-related
8 emergency hospitalizations for respiratory diseases in the Medicare population. *Am. J. Respir. Crit. Care Med.*
9 187, 1098-1103.
- 10
11 Astrom, D.O., Forsberg, B., Rocklov, J., 2011. Heat wave impact on morbidity and mortality in the elderly
12 population: a review of recent studies. *Maturitas* 69, 99-105.
- 13
14 Astrom, D.O., Schifano, P., Asta, F., Lallo, A., Michelozzi, P., Rocklov, J., Forsberg, B., 2015. The effect of heat
15 waves on mortality in susceptible groups: a cohort study of a mediterranean and a northern European City.
16 *Environ. Health* 14, 30.
- 17
18 Atsumi, A., Ueda, K., Irie, F., Sairenchi, T., Iimura, K., Watanabe, H., Iso, H., Ota, H., Aonuma, K., 2013.
19 Relationship between cold temperature and cardiovascular mortality, with assessment of effect modification by
20 individual characteristics. *Circ. J.* 77, 1854-1861.
- 21
22 Basu, R., 2009. High ambient temperature and mortality: a review of epidemiologic studies from 2001 to 2008.
23 *Environ. Health* 8.
- 24
25 Bateson, T.F., Schwartz, J., 2004. Who is sensitive to the effects of particulate air pollution on mortality?
26 *Epidemiology* 15, 143-149.
- 27
28 Beverland, I.J., Cohen, G.R., Heal, M.R., Carder, M., Yap, C., Robertson, C., Hart, C.L., Agius, R.M., 2012. A
29 comparison of short-term and long-term air pollution exposure associations with mortality in two cohorts in
30 Scotland. *Environ. Health Perspect.* 120, 1280-1285.
- 31
32 Bobb, J.F., Peng, R.D., Bell, M.L., Dominici, F., 2014. Heat-related mortality and adaptation to heat in the
33 United States. *Environ. Health Perspect.* 122, 811-816.
- 34
35 Curriero, F.C., Heiner, K.S., Samet, J.M., 2002. Temperature and mortality in 11 cities of the Eastern United
36 States. *Am. J. Epidemiol.* 155, 80-87.
- 37
38 Essebag, V., Platt, R.W., Abrahamowicz, M., Pilote, L., 2005. Comparison of nested case-control and survival
39 analysis methodologies for analysis of time-dependent exposure. *BMC Med. Res. Methodol.* 5, 5.
- 40
41 Gasparrini, A., 2014. Modeling exposure-lag-response associations with distributed lag non-linear models. *Stat.*
42 *Med.* 33, 881-899.
- 43
44 Gasparrini, A., Armstrong, B., Kenward, M.G., 2010. Distributed lag non-linear models. *Stat. Med.* 29, 2224-
2234.
- 45
46 Gasparrini, A., Armstrong, B., Kovats, S., Wilkinson, P., 2012. The effect of high temperatures on cause-specific
47 mortality in England and Wales. *BMC Med. Res. Methodol.* 69, 56-61.
- 48
49 Goggins, W.B., Yang, C.Y., Hokama, T., Law, L.S.K., Y., C.E.Y., 2015. Using annual data to estimate the public
50 health impact of extreme temperature. *Am. J. Epidemiol.* 182, 80-87.
- 51
52 Goldberg, M.S., Gasparrini, A., Armstrong, B., Valois, M.F., 2011. The short-term influence of temperature on
53 daily mortality in the temperate climate of Montreal, Canada. *Environ. Res.* 111, 853-860.
- 54
55 Goldstein, L., Langholz, B., 1992. Asymptotic theory for nested case-control sampling in the Cox regression
56 model. *Ann. Stat.* 20, 1903-1928.
- 57
58 Goodman, P.G., Dockery, D.W., Clancy, L., 2004. Cause-specific mortality and the extended effects of
59 particulate pollution and temperature exposure. *Environ. Health Perspect.* 112, 179-185.
- 60
61 Guo, Y., Barnett, A.G., Pan, X., Yu, W., Tong, S., 2011. The impact of temperature on mortality in Tianjin, China:

a case-crossover design with a distributed lag nonlinear model. *Environ. Health Perspect.* 119, 1719-1725.

1 Guo, Y., Punnasiri, K., Tong, S., 2012. Effects of temperature on mortality in Chiang Mai city, Thailand: a time
2 series study. *Environ. Health* 11, 36.
3

4 Hajat, S., Armstrong, B.G., Gouveia, N., Wilkinson, P., 2005. Mortality displacement of heat-related deaths.
5 *Epidemiology* 16, 613-620.
6

7 Hajat, S., O'Connor, M., Kosatsky, T., 2010. Health effects of hot weather: from awareness of risk factors to
8 effective health protection. *Lancet* 375, 856-863.
9

10 Idris, I., Thomson, G.A., Sharma, J.C., 2006. Diabetes mellitus and stroke. *Int. J. Clin. Pract.* 60, 48-56.
11

12 Keatinge, W.R., Coleshaw, S.R.K., Cotter, F., Mattock, M., Murphy, M., Chelliah, R., 1984. Increases in platelet
13 and red counts, blood viscosity, and arterial pressure during mild surface cooling: factors in mortality from
14 coronary and cerebral thrombosis in winter. *Br. Med. J. (Clin. Res. Ed.)* 289, 1405-1408.

15 Khalaj, B., Lloyd, G., Sheppard, V., Dear, K., 2010. The health impacts of heat waves in five regions of New
16 South Wales, Australia: a case-only analysis. *Int. Arch. Occup. Environ. Health* 83, 833-842.
17

18 Knowlton, K., Rotkin-Ellman, M., King, G., Margolis, H.G., Smith, D., Solomon, G., Trent, R., English, P.,
19 2009. The 2006 California heat wave: impacts on hospitalizations and emergency department visits. *Environ.*
20 *Health Perspect.* 117, 61-67.
21

22 Li, Y., Lan, L., Wang, Y., Yang, C., Tang, W., Cui, G., Luo, S., Cheng, Y., Liu, Y., Liu, J., Jin, Y., 2014.
23 Extremely cold and hot temperatures increase the risk of diabetes mortality in metropolitan areas of two Chinese
24 cities. *Environ. Res.* 134, 91-97.
25

26 Lin, H.J., Chen, S.T., Wu, H.Y., Hsu, H.C., Chen, M.F., Lee, Y.T., Wu, K.Y., Chien, K.L., 2015. Urinary
27 biomarkers of oxidative and nitrosative stress and the risk for incident stroke: a nested case-control study from a
28 community-based cohort. *Int. J. Cardiol.* 183, 214-220.
29

30 Luchsinger, J.A., Tang, M.X., Stern, Y., Shea, S., Mayeux, R., 2001. Diabetes mellitus and risk of Alzheimer's
31 disease and dementia with stroke in a multiethnic cohort. *Am. J. Epidemiol.* 154, 635-641.
32

33 Ma, W., Xu, X., Peng, L., Kan, H., 2011. Impact of extreme temperature on hospital admission in Shanghai,
34 China. *Sci. Total Environ.* 409, 3634-3637.
35

36 Ma, W., Yang, C., Chu, C., Li, T., Tan, J., Kan, H., 2013. The impact of the 2008 cold spell on mortality in
37 Shanghai, China. *Int. J. Biometeorol.* 57, 179-184.
38

39 Ma, W., Zeng, W., Zhou, M., Wang, L., Rutherford, S., Lin, H., Liu, T., Zhang, Y., Xiao, J., Zhang, Y., Wang, X.,
40 Gu, X., Chu, C., 2015. The short-term effect of heat waves on mortality and its modifiers in China: an analysis
41 from 66 communities. *Environ. Int.* 75, 103-109.
42

43 Medina-Ramon, M., Schwartz, J., 2007. Temperature, temperature extremes, and mortality: a study of
44 acclimatisation and effect modification in 50 US cities. *Occup. Environ. Med.* 64, 827-833.
45

46 Medina-Ramon, M., Zanobetti, A., Cavanagh, D.P., Schwartz, J., 2006. Extreme temperatures and mortality:
47 assessing effect modification by personal characteristics and specific cause of death in a multi-city case-only
48 analysis. *Environ. Health Perspect.* 114, 1331-1336.
49

50 Meehl, G.A., Tebaldi, C., 2004. More intense, more frequent, and longer lasting heat waves in the 21st century.
51 *Science* 305, 994-997.
52

53 Mercer, J., 2003. Cold—an underrated risk factor for health. *Environ. Res.* 92, 8-13.
54

55 Michelozzi, P., Accetta, G., De Sario, M., D'Ippoliti, D., Marino, C., Baccini, M., Biggeri, A., Anderson, H.R.,
56 Katsouyanni, K., Ballester, F., Bisanti, L., Cadum, E., Forsberg, B., Forastiere, F., Goodman, P.G., Hojs, A.,
57 Kirchmayer, U., Medina, S., Paldy, A., Schindler, C., Sunyer, J., Perucci, C.A., Group, P.C., 2009. High
58 temperature and hospitalizations for cardiovascular and respiratory causes in 12 European cities. *Am J Respir*
59 *Crit Care Med* 179, 383-389.
60
61
62
63
64
65

1 Nayha, S., 2002. Cold and the risk of cardiovascular diseases: a review. *Int. J. Circumpolar Health* 61, 373-380.

2 Nordio, F., Zanobetti, A., Colicino, E., Kloog, I., Schwartz, J., 2015. Changing patterns of the temperature-

3 mortality association by time and location in the US, and implications for climate change. *Environ. Int.* 81, 80-

4 86.

5 Olsson, A., Kromhout, H., Agostini, M., Hansen, J., Lassen, C.F., Johansen, C., Kjaerheim, K., Langard, S.,

6 Stucker, I., Ahrens, W., Behrens, T., Lindbohm, M.L., Heikkila, P., Heederik, D., Portengen, L., Shaham, J.,

7 Ferro, G., de Vocht, F., Burstyn, I., Boffetta, P., 2010. A case-control study of lung cancer nested in a cohort of

8 European asphalt workers. *Environ. Health Perspect.* 118, 1418-1424.

9

10

11 Ou, C.Q., Song, Y.F., Yang, J., Chau, P.Y., Yang, L., Chen, P.Y., Wong, C.M., 2013. Excess winter mortality and

12 cold temperatures in a subtropical city, Guangzhou, China. *PLoS One* 8, e77150.

13

14 Qiao, Z., Guo, Y., Yu, W., Tong, S., 2015. Assessment of Short- and Long-Term Mortality Displacement in Heat-

15 Related Deaths in Brisbane, Australia, 1996-2004. *Environ. Health Perspect.*

16

17 Rocklov, J., Forsberg, B., Ebi, K., Bellander, T., 2014. Susceptibility to mortality related to temperature and heat

18 and cold wave duration in the population of Stockholm County, Sweden. *Glob. Health Action* 7, 22737.

19

20 Rowell, L.B., 1974. Human cardiovascular adjustments to exercise and thermal stress. *Physiol. Rev.* 54, 75-142.

21

22 Sawada, N., Iwasaki, M., Inoue, M., Itoh, H., Sasazuki, S., Yamaji, T., Shimazu, T., Tsugane, S., Japan Public

23 Health Center Based Prospective Study, G., 2010. Plasma organochlorines and subsequent risk of prostate

24 cancer in Japanese men: a nested case-control study. *Environ. Health Perspect.* 118, 659-665.

25

26 Schifano, P., Cappai, G., De Sario, M., Michelozzi, P., Marino, C., Bargagli, A.M., Perucci, C.A., 2009.

27 Susceptibility to heat wave-related mortality: a follow-up study of a cohort of elderly in Rome. *Environ Health* 8,

28 50.

29

30 Schooling, C., Chan, W., Leung, S., Lam, T., Lee, S., Shen, C., Leung, J., Leung, G., 2014. Cohort Profile: Hong

31 Kong Department of Health Elderly Health Service Cohort. *Int. J. Epidemiol.*

32

33 Schooling, C.M., Lam, T.H., Li, Z.B., Ho, S.Y., M., C.W., Ho, K.S., THAM, M.K., Cowling, B.J., Leung, G.M.,

34 2006. Obesity, physical activity, and mortality in a prospective Chinese elderly cohort. *Arch Intern Med* 166,

35 1498-1504.

36

37 Schwartz, J., 2005. Who is sensitive to extremes of temperature? *Epidemiology* 16, 67-72.

38

39 Song, X., Zhang, Z., Chen, Y., Wang, P., Xiang, M., Shi, P., Tao, F., 2013. Spatiotemporal changes of global

40 extreme temperature events (ETEs) since 1981 and the meteorological causes. *Nat. Hazards* 70, 975-994.

41

42 Stafoggia, M., Schwartz, J., Forastiere, F., Perucci, C.A., 2008. Does temperature modify the association

43 between air pollution and mortality? A multicity case-crossover analysis in Italy. *Am. J. Epidemiol.* 167, 1476-

44 1485.

45

46 Williams, S., Nitschke, M., Weinstein, P., Pisaniello, D.L., Parton, K.A., Bi, P., 2012. The impact of summer

47 temperatures and heatwaves on mortality and morbidity in Perth, Australia 1994-2008. *Environ. Int.* 40, 33-38.

48

49 Wong, C.M., Lai, H.K., Tsang, H., Thach, T.Q., Thomas, G.N., Lam, K.B., Chan, K.P., Yang, L., Lau, A.K.,

50 Ayres, J.G., Lee, S.Y., Chan, W.M., Hedley, A.J., Lam, T.H., 2015. Satellite-Based Estimates of Long-Term

51 Exposure to Fine Particles and Association with Mortality in Elderly Hong Kong Residents. *Environ. Health*

52 *Perspect.*

53

54 Wu, W., Xiao, Y., Li, G., Zeng, W., Lin, H., Rutherford, S., Xu, Y., Luo, Y., Xu, X., Chu, C., Ma, W., 2013.

55 Temperature-mortality relationship in four subtropical Chinese cities: a time-series study using a distributed lag

56 non-linear model. *Sci. Total Environ.* 449, 355-362.

57

58 Xu, W., Thach, T.Q., Chau, Y.K., Lai, H.K., Lam, T.H., Chan, W.M., Lee, R.S., Hedley, A.J., Wong, C.M., 2013.

59 Thermal stress associated mortality risk and effect modification by sex and obesity in an elderly cohort of

60 Chinese in Hong Kong. *Environ. Pollut.* 178, 288-293.

61

62

63

64

65

1 Yi, W., Chan, A.P., 2014. Effects of temperature on mortality in Hong Kong: a time series analysis. *Int. J.*
2 *Biometeorol.*

3 Zanobetti, A., O'Neill, M.S., Gronlund, C.J., Schwartz, J.D., 2013. Susceptibility to mortality in weather
4 extremes: effect modification by personal and small-area characteristics. *Epidemiology* 24, 809-819.

5 Zanobetti, A., Schwartz, J., Samoli, E., Gryparis, A., Touloumi, G., Atkinson, R., Tertre, A.L., 2002. The
6 temporal pattern of mortality responses to air pollution: a multicity assessment of mortality displacement.
7 *Epidemiology* 13, 87-93.
8
9

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 1. Baseline characteristics of the participants, according to pre-existing health conditions in the prospective Chinese elderly cohort in Hong Kong (n=61,441).

Characteristic	Pre-existing diseases				
	NDG	DM	CSD	COPD	All subjects
Individual-level					
Age in year (mean ± SD)	71.5 ± 5.5	71.9 ± 5.2	72.4 ± 5.6	73.3 ± 5.7	72.0 ± 5.6
Sex (%)					
Male	33.2	33.0	33.3	49.3	33.9
Female	66.8	67.0	66.7	50.7	66.1
BMI (%)					
1 st (<21.6)	50.6	52.7	51.2	45.3	50.8
2 nd - 3 rd (21.6-26.3)	28.5	15.2	16.5	31.2	22.8
4 th (>26.3)	20.9	32.0	32.3	23.5	26.3
Smoking (%)					
Never	71.7	72.9	72.9	43.7	71.3
Quitted	17.2	19.5	19.8	37.9	19.2
Current	11.1	7.5	7.2	18.3	9.5
Exercise (%)					
Never	16.5	13.3	14.0	16.7	15.2
<4	7.3	6.8	6.7	6.8	7.0
4-6	5.7	5.7	5.6	5.8	5.7
>6	70.4	74.2	73.7	70.7	72.1
Education (%)					
Post-secondary	3.8	3.6	3.7	3.1	3.7
Secondary	13.3	13.6	13.7	11.7	13.5
Primary	37.0	35.8	36.5	39.5	36.9
Below primary	45.8	47.0	46.1	45.7	46.0
TPU-level					
Age ≥ 65 (%)	12.1	12.2	12.2	12.1	12.1
Tertiary education (%)	12.9	12.9	13.1	12.2	13.0
Income ≥ US\$1,923/m (%)	59.5	59.5	59.6	58.6	59.5
District-level					
Smoking rate (%)	11.5	11.6	11.6	11.6	11.6

Abbreviations: BMI, body mass index; TPU, tertiary planning unit; NDG: non-disease group; DM: diabetes mellitus; CSD, circulatory system diseases; COPD, chronic obstructive pulmonary disease; SD, standard deviation.

Table 2. All natural mortality after 10-13 years of following up at the end of study in 2011 by pre-existing health conditions.

Morbidity status	No. of subjects	No. of death	%
NDG	27890	5809	20.8
DM	8752	3024	34.6
CSD	29081	9070	31.2
Hypertension	25154	7712	30.7
Heart	8164	2899	35.5
Stroke	1745	764	43.8
COPD	3829	1855	48.4
All subjects	61441	16497	26.9

Abbreviations: NDG: non-disease group; DM: diabetes mellitus; CSD, circulatory system diseases; COPD, chronic obstructive pulmonary disease.

Table 3. The cumulative relative risks of cold and hot effects on all natural mortality in elders with pre-existing diseases along lag days.

Pre-existing disease	Lag (day)	Extreme cold ^a	Moderate cold ^b	Moderate hot ^c	Extreme hot ^d
NDG	0-1	0.98 (0.84, 1.13)	1.00 (0.95, 1.05)	0.98 (0.95, 1.02)	0.97 (0.91, 1.04)
	0-3	0.96 (0.76, 1.22)	1.00 (0.92, 1.08)	0.97 (0.91, 1.03)	0.95 (0.86, 1.06)
	0-13	0.97 (0.64, 1.49)	1.01 (0.88, 1.15)	0.96 (0.87, 1.05)	0.93 (0.79, 1.09)
	0-21	1.04 (0.59, 1.85)	1.02 (0.86, 1.22)	0.97 (0.86, 1.10)	0.96 (0.78, 1.17)
DM	0-1	1.11 (0.95, 1.29)	1.03 (0.98, 1.08)	1.02 (0.98, 1.06) [#]	1.03 (0.96, 1.11)
	0-3	1.22 (0.95, 1.56) [#]	1.06 (0.98, 1.14)	1.03 (0.96, 1.10)	1.05 (0.94, 1.17) [#]
	0-13	1.77 (1.12, 2.80)[*]	1.20 (1.04, 1.38)[*]	1.03 (0.93, 1.14)	1.06 (0.89, 1.25)
	0-21	2.21 (1.19, 4.10)[*]	1.25 (1.03, 1.51)[#]	1.12 (0.99, 1.27) [#]	1.22 (0.98, 1.50) [#]
CSD	0-1	1.05 (0.96, 1.14)	1.01 (0.98, 1.04)	1.02 (1.00, 1.05) [*]	1.04 (1.00, 1.08) [*]
	0-3	1.09 (0.95, 1.26)	1.02 (0.97, 1.07)	1.03 (0.99, 1.07) [#]	1.06 (0.99, 1.12) [*]
	0-13	1.34 (1.04, 1.74)[#]	1.11 (1.02, 1.20)[*]	0.99 (0.93, 1.05)	0.99 (0.89, 1.09)
	0-21	1.59 (1.12, 2.26)[*]	1.16 (1.04, 1.29)[*]	1.02 (0.95, 1.10)	1.03 (0.91, 1.17)
COPD	0-1	1.08 (0.88, 1.31)	1.01 (0.94, 1.07)	1.05 (1.00, 1.11) [*]	1.09 (0.99, 1.20) [*]
	0-3	1.15 (0.84, 1.58)	1.02 (0.92, 1.13)	1.08 (0.99, 1.18) [*]	1.14 (0.98, 1.32) [*]
	0-13	1.37 (0.76, 2.48)	1.10 (0.91, 1.32)	1.03 (0.89, 1.18)	1.05 (0.82, 1.33)
	0-21	1.23 (0.53, 2.84)	1.04 (0.81, 1.35)	1.07 (0.90, 1.28)	1.12 (0.83, 1.52)
All subjects	0-1	1.02 (0.96, 1.09)	1.00 (0.98, 1.03)	1.01 (0.99, 1.02) [#]	1.01 (0.98, 1.04)
	0-3	1.05 (0.95, 1.17)	1.01 (0.98, 1.05)	1.01 (0.98, 1.03)	1.01 (0.97, 1.06)
	0-13	1.30 (1.07, 1.57)[#]	1.10 (1.03, 1.17)[*]	0.98 (0.94, 1.02)	0.97 (0.90, 1.04)
	0-21	1.49 (1.15, 1.93)[*]	1.14 (1.06, 1.24)[*]	1.00 (0.95, 1.06)	1.00 (0.92, 1.10)

Abbreviations: NDG: non-disease group; DM: diabetes mellitus; CSD, circulatory system diseases; COPD, chronic obstructive pulmonary disease.

^a A decrease from 25th percentile of temperature (19.5 °C) to 1st percentile of temperature (11.3 °C);

^b A decrease from 25th percentile of temperature (19.5 °C) to 10th percentile of temperature (16.2 °C);

^c An increase from 75th percentile of temperature (27.8 °C) to 90th percentile of temperature (29.4 °C);

^d An increase from 75th percentile of temperature (27.8 °C) to 99th percentile of temperature (30.4 °C).

^{*}: Compared with NDG in the same lag period, *p*-Value < 0.05;

[#]: Compared with NDG in the same lag period, 0.05 ≤ *p*-Value < 0.1.

Figure legends

Fig. 1. The estimated relative risk of temperature centered at median (24.7 °C) on all natural mortality over 21 days for subjects with different pre-existing diseases. NDG: non-disease group; DM: diabetes mellitus; CSD: circulatory system diseases; COPD: chronic obstructive pulmonary disease.

Fig. 2. The estimated relative risk of cold temperature effects [a decrease from 25th percentile of temperature (19.5 °C) to 10th percentile of temperature (16.2 °C)] and hot effects [an increase from 75th percentile of temperature (27.8 °C) to 90th percentile of temperature (29.4 °C)] on all natural mortality over 21 days of lag. NDG: non-disease group; DM: diabetes mellitus; CSD: circulatory system diseases; COPD: chronic obstructive pulmonary disease. The solid lines are mean relative risks, and grey regions are 95% CIs.

Figure1




Figure 2

