



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

In vitro administration of sodium arsenite in mouse prepubertal testis induces germ cell loss and apoptosis

Citation for published version:

Anwar, N, Qureshi, IZ, Spears, N & Lopes, F 2020, 'In vitro administration of sodium arsenite in mouse prepubertal testis induces germ cell loss and apoptosis', *Toxicology in Vitro*.
<https://doi.org/10.1016/j.tiv.2020.104924>

Digital Object Identifier (DOI):

[10.1016/j.tiv.2020.104924](https://doi.org/10.1016/j.tiv.2020.104924)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Toxicology in Vitro

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



1 ***In vitro* administration of sodium arsenite in mouse prepubertal testis induces germ cell**
2 **loss and apoptosis**

3

4 **Naureen Anwar^a, Irfan Zia Qureshi^a, Norah Spears^b, Federica Lopes^{b,*}**

5

6 ^a Quaid-i-Azam University, Department of Animal Sciences, Laboratory of Animal and
7 Human Physiology, PO Box 45320, Islamabad, Pakistan.

8 ^b University of Edinburgh, School of Biomedical Sciences, Edinburgh, United Kingdom.

9 *Corresponding author: Centre for Reproductive Health, Queen's Medical Research Institute,
10 University of Edinburgh, 47 Little France, Edinburgh EH16 4TJ, Scotland, UK.

11 E-mail: Federica.Lopes@ed.ac.uk ; Tel: +44 (0)131 242 9126

12

13 **Abstract**

14 High levels of arsenic contamination in drinking water pose serious health risks in numerous
15 countries. The documentation reporting arsenic toxicity on reproduction and development is
16 increasing, with evidence of arsenic inducing fertility and developmental issues. Nonetheless,
17 the impact of arsenic exposure on the development of the male reproductive system is not fully
18 elucidated. In the present study, we have investigated the direct effects of arsenic on
19 prepubertal mouse testis using an *in vitro* testicular organ culture system. Culture medium was
20 supplemented with a range of concentrations of sodium arsenite, examining effects of low (0.5
21 and 1 μM) and high (10, 50, 100 μM) concentrations, in cultures of post-natal day 5 CD1 mouse
22 testis. *In vitro* exposure of low arsenic concentrations (0.5 or 1 μM) for 6 days did not cause
23 any change in the testicular morphology, germ cells density, or apoptotic marker cleaved

24 caspase 3 (CC3) expression. In contrast, exposure of prepubertal testis to high arsenic
25 concentrations (10, 50 or 100 μ M) induced drastic changes: severe destruction of testicular
26 morphology, with loss of seminiferous tubule integrity; a dose-dependent decrease in germ cell
27 density, and a hundred-fold increase in CC3 expression after 50 μ M arsenic exposure. In
28 conclusion, high arsenic treatment induced a dose-dependent induction of apoptosis and germ
29 cell loss in prepubertal mouse testis.

30

31 **Keywords:**

32 Sodium arsenite; *in vitro*; prepubertal mouse; testis; toxicity.

33

34

35 **1. Introduction**

36 Arsenic is a metalloid naturally present in food, soil, and water (Duker et al., 2005;
37 Huang et al., 2004) with a toxicity defined by its binding state; inorganic arsenic compounds
38 are more toxic than organic compounds (Maitani et al., 1987; Styblo et al., 2000). Usually,
39 natural water contains inorganic arsenic forms such as trivalent (As III) and pentavalent arsenic
40 (As V) (Feng et al., 2001). Trivalent arsenic, such as sodium arsenite, is more toxic than the
41 pentavalent form, for example sodium arsenate (Styblo et al., 2000; Vega et al., 2001). Arsenic
42 is a potent carcinogen, its chronic exposure can lead to adverse effects on skin and on the
43 cardiovascular, neurological and respiratory systems (reviewed in Abdul et al., 2015; Ratnaike,
44 2003).

45 Arsenic poisoning has become an escalating problem in many developing countries.
46 Usually the contamination of drinking waters with arsenic occurs due to natural environmental

47 sources seeping into aquifers, mining, and other industrial activities (Sulanjari et al., 2015). In
48 particular, in the past few decades its exposure through drinking water has vastly increased in
49 Bangladesh, India, China, and Cambodia (Smith, 2003; Xie, 2011). Recently, the risk of arsenic
50 exposure has also been reported to have worsened in Pakistan, exposing almost 50 million
51 people to high levels of arsenic through drinking groundwater (Podgorski et al., 2017). Overall,
52 more than 150 million people around the world are affected by drinking arsenic-contaminated
53 water (Munoz et al., 2015): its chronic exposure is therefore a major concern for public health
54 (Abhyankar et al., 2012).

55 There is a clear link between exposure to inorganic arsenic and male reproductive
56 dysfunction (Feng et al., 2001; Kim and Kim, 2015; Lima et al., 2018; Wang et al., 2006). In
57 the testis, exposure to arsenic can alter spermatogenesis, interrupt steroidogenesis, and reduce
58 gonadotrophin as well as testosterone levels (Kim and Kim, 2015). *In vivo* studies investigating
59 the reproductive toxicity of arsenic have shown effects on adult rodent reproductive system.
60 For instance, Chang et al. (2007) reported reduced testicular weight and sperm count in mice
61 after subchronic exposure to 20 or 40 mg/l sodium arsenite. Similarly, mice exposed to sodium
62 arsenite via drinking water at concentration of around 530 μ mol/l presented with a significant
63 reduction in sperm count and motility (Pant et al., 2001). A comparative study in male rats
64 exposed to arsenite or arsenate suggested a higher reproductive toxicity of the inorganic form
65 (Lima et al., 2018). A small number of *in vitro* studies have assessed effects of arsenic on
66 embryonic development and reproduction. Different concentrations of sodium arsenite exert
67 toxic effects on preimplantation development of mouse embryos, with 100 μ M arsenic
68 concentration rapidly killing two-cell embryos, while 1 μ M impairs blastocyst formation
69 (Muller et al., 1986). *In vitro* studies of arsenic on post-implantation mouse embryo
70 development confirmed that sodium arsenite possesses higher toxicity compared to sodium

71 arsenate (Chaineau et al., 1990; Tabocova et al., 1996). Moreover, Sertoli cell lines undergo
72 apoptosis after arsenic trioxide exposure (Kim et al., 2011).

73 In general, children are considered more sensitive to toxicants than adults (Scheuplein
74 et al., 2002). There is an accumulating evidence of a strong negative impact of arsenic exposure
75 in early life, includes link to increased mortality rates during both childhood (Rahman et al.,
76 2013) and also adulthood (Yuan et al., 2007). Prepubertal testicular development is a highly
77 sensitive stage and it is possible that toxicants such as arsenic may act differently on testis of
78 young compared to adult males. Presently, only a few animal studies have examined the effects
79 of arsenic on the reproductive axis of the prepubertal mammal: prepubertal rat prostate
80 development has been found to be affected (Aquino et al., 2019), but that study provides little
81 information about the testicular cell populations that could be damaged (i.e. somatic and/or
82 germ cells); in one other study, prepubertal rats exposed to sodium arsenite (0.01 or 10 mg/l)
83 for 30 days in the drinking water showed a significant disruption of normal testicular
84 morphology (da Cunha de Medeiros et al., 2019).

85 Here, we have investigated the effect of sodium arsenite exposure on immature mouse
86 testis using an *in vitro* culture system that supports the developing neonatal testis. *In vitro*
87 models have been commonly used to elucidate the metabolism of arsenic (Drobna et al., 2009),
88 and the testis culture model has already been used for other toxicological studies on the male
89 gonad (Allen et al., 2020; Lopes et al., 2016; Nakamura et al., 2019; Smart et al., 2018).

90

91 **2. Materials and Methods:**

92 2.1. Animals

93 This work was approved by the Local Ethical Review Committee of the University of
94 Edinburgh and conducted according to UK home office regulations. CD1 male mice, obtained

95 from University of Edinburgh breeding stock, were kept at 14 h:10 h light and dark
96 photoperiod, provided *ad libitum* food (expanded RM3 diet, SDS; DMB Scotland) and water.
97 All mice used in the experiments were euthanized by decapitation at post-natal day (PND) 5
98 for tissue collection or by cervical dislocation at PND 12 for age-matched control, with the day
99 of the pup delivery considered as PND 0.

100 2.2. Tissue culture

101 PND 5 testes were collected, fragmented and cultured according to Lopes et al. (2016).
102 In brief, for each culture run, testes from at least two mice were dissected, pooled together and
103 placed into dissection medium Leibovitz L-15 (Invitrogen, UK) supplemented with 3mg/ml of
104 bovine serum albumin (BSA; Sigma-Aldrich Ltd, UK) at 37°C. After removal of epididymis
105 and tunica albuginea, testes were fragmented into approximately 0.5mm³ pieces, using a scalpel
106 blade. Fragments that were homogeneous in size were randomly distributed on a 24-well
107 culture plate (Greiner Bio-one, UK), with testes obtained from two pups per litter for each
108 culture run. On the first day of culture (Day 1), each testicular fragment was placed on a floating
109 polycarbonate membrane (Whatman, Camlab Ltd, Cambridge, UK) per each well, containing
110 1 ml α -Minimal Essential Medium (MEM; Invitrogen, UK) supplemented with 10% Knockout
111 Serum Replacement (Invitrogen, UK); half of the medium was replaced with fresh medium on
112 Days 4 and 6. The culture plate was incubated at 34°C, 5% CO₂. Fragments were left untreated
113 for the first 24 h (Day 1). From Day 2 of culture, testicular fragments were exposed to different
114 concentrations of sodium arsenite prepared from a stock solution of 0.05 mol/l (0.1N)
115 Tritipur®reag. (Merck, Germany). Sodium arsenite solution was diluted in culture medium and
116 final concentrations were prepared through serial dilutions. Two sets of experiments (low and
117 high arsenic doses) were performed with cultures maintained for seven days, of which six days
118 were with arsenic exposure (Days 2-7): the *in vitro* study on mouse embryos arsenic exposure
119 was used as a reference for selection of the highest dose (Muller et al., 1986). The first

120 experiment (low arsenic doses) comprised of 0.5 and 1 μ M sodium arsenite concentrations,
121 whereas the second experiment (high arsenic doses) included concentrations of 10, 50 and 100
122 μ M arsenic. In each culture run, the arsenic effect was compared to a culture control that had
123 been maintained throughout in arsenite-free medium. Each experiment was repeated six times,
124 including six culture runs, six histological and immunohistochemistry (IHC) runs and six sets
125 of image analysis. On Day 7, tissues were fixed for 1 h in 10% neutral buffered formalin (Sigma
126 Aldrich, UK) prior to IHC, or in Bouin's fluid (Sigma Aldrich Ltd, UK) prior to histological
127 examination. Tissues were placed in 3% agar (Sigma Aldrich Ltd, UK) blocks and then
128 embedded in paraffin wax prior to sectioning.

129 2.3. Histology/ Immunohistochemistry

130 Five μ m thick serial sections were cut. A middle section from each tissue fragment was
131 dewaxed, rehydrated through a series of decreasing alcohol concentrations, stained with
132 hematoxylin and eosin (H&E), and used for morphological analysis. A serial section next to
133 the middle section used for histological examination was selected for IHC reactions. Sections
134 were dewaxed, rehydrated, and processed for antigen retrieval with citrate buffer (10mM
135 sodium citrate, pH 6, Fisher Scientific, UK). In between each step, slides were washed with
136 phosphate-buffered saline (Fisher Scientific, UK) with 0.1% Triton X (PBSTx, Sigma-Aldrich,
137 UK). Blocking used 20% normal goat serum diluted in PBSTx supplemented with 5% w/v
138 BSA for 1 h at room temperature. Primary antibodies were applied and incubated overnight in
139 a humidified environment: (i) rabbit anti-cleaved caspase 3 (CC3) antibody (9661S, 1:500
140 dilution; Cell Signaling Technology, USA) and (ii) mouse anti-DEAD-box helicase 4
141 (DDX4)/Mouse Vasa Homologue (MVH) antibody (ab27591, 1:100 dilution; Abcam, UK).
142 After washing, appropriate secondary antibodies were applied (both 1:200 dilution): (i) goat
143 anti-rabbit biotinylated (E0432, DakoCytomation, Denmark) and (ii) Alexafluor goat anti-
144 mouse IgG1 568nm (A-21124, Invitrogen, UK). Omission of primary or secondary antibodies

145 were used as technical negative controls. For CC3, the visualization reagent Streptavidin Alexa
146 Fluor 488 conjugate (S-32354, Invitrogen, UK) was subsequently applied at 1:200 dilution. All
147 sections were counterstained with 4,6-Diamidino-2-phenylindole (DAPI; Invitrogen, UK) and
148 mounted with Vectashield hard-set mounting medium (Vector Laboratories, USA).

149 2.4. Image attainment and analysis

150 H&E stained sections were photomicrographed (DMLB Leica microscope, Leica
151 Microsystems Ltd, UK), with images taken using a DFC369FX camera on a Leica DM5500B
152 microscope (Leica Microsystem Ltd, UK), using filters for DAPI, A4 (BP 360/40); for Alexa
153 Fluor 488, N3 (BP 546/12); for Alexa Fluor 568, Y5 (BP 620/60). Image J software was used
154 for image analysis, with the assessor blind as to treatment. Germ cells (MVH⁺) were analyzed
155 by manual counting and the total number given relative to the area of seminiferous tubule, in
156 order to obtain a value for germ cell density (number of cells that were MVH-positive divided
157 by seminiferous tubule area in mm²). Density was also determined for whole section area
158 (number of MVH positive cells divided by the area of section in mm²) to allow determination
159 of germ cell density even where tubule integrity had been disrupted to the extent that precluded
160 measurement of tubule area. For CC3, fluorophore area was measured as a percentage of
161 tubule, interstitium and section area (proportion of CC3-positive area in mm² relative to area
162 of seminiferous tubule, interstitium or whole section, all in mm²).

163 2.5. Statistical analysis

164 GraphPad Prism was used for statistical analyses. The statistically significance of any
165 difference between the control group and the arsenic-treated groups was determined by one-
166 way ANOVA, followed by Dunnett's *post hoc* test where the ANOVA was statistically
167 significant. For analyses including only 2 experimental groups (comparison of the control
168 group to the 10 µM sodium arsenite treatment group for: density of germ cells in the

169 seminiferous tubules, and percentage of seminiferous tubule or interstitium area positive for
170 CC3), paired T-test was performed. Data is expressed as mean \pm SEM, with $p \leq 0.05$ considered
171 statistically significant.

172


173 **3. Results**

174 3.1. Morphological analysis


175 To assess the impact of sodium arsenite on the developing testis of prepubertal (PND
176 5) mouse, testis fragments were exposed *in vitro* for six days to low or high concentrations of
177 sodium arsenite, compared with arsenite-free culture control testicular fragments in each case.
178 Morphological changes were evaluated on H&E stained sections (Fig. 1).

179 **Figure 1**


A: Uncultured control



B: Low Sodium Arsenite Doses Experiment



C: High Sodium Arsenite Doses Experiment



181

182 **Figure 1. Morphological changes of prepubertal mouse testis induced by *in vitro* exposure**
183 **to sodium arsenite.** Representative images of hematoxylin and eosin stained sections of
184 uncultured and cultured testis fragments. Pre-culture control tissue (**Ai**) obtained from PND 5
185 mouse showed germ cells only at the spermatogonial stage (**black arrow**), while testes from
186 mice age-matched to be equivalent to end of culture control testis (PND 12: **Aii**) showed germ
187 cells that have entered meiosis (spermatocytes, **white arrow**). Representative images showing
188 the effect on testis morphology of low (**B**) and high (**C**) concentrations of arsenic. Control
189 tissues in both the low dose (**Bi**) and high dose experiment (**Ci**), as well as low doses of arsenic
190 (**Bii**: 0.5 μM sodium arsenite; **Biii**: 1 μM sodium arsenite) preserved normal morphology, with
191 germ cells lining the basal membrane (**black arrow**), some cells undergoing mitosis (**black**
192 **arrow head**) and several differentiating germ cells (spermatocytes, **white arrow**)
193 accumulating in the forming lumen of well-defined seminiferous tubules. High doses of arsenic
194 caused dramatic morphological changes, starting at 10 μM sodium arsenite (**Cii**) with evidence

195 of apoptotic/necrotic cells (**white arrowhead**), and with disruption or loss of seminiferous
196 tubule integrity at 50 μM (**Ciii**) and 100 μM sodium arsenite (**Civ**). B-C: scale bars represent
197 100 μm ; A and insets: scale bars represent 50 μm .

198

199 The morphology of control testicular fragments was well preserved at the end of the six
200 day culture period for both sets of experiments: necrosis was not present in the centre of the
201 section, seminiferous tubule structure and integrity was maintained, with many germ cells lined
202 along the basement membrane of all tubules, several actively undergoing a mitotic division
203 (Fig. 1 Bi and Ci). Control tissues closely resembled the morphology of equivalent age-
204 matched PND 12 testis (Fig. 1 Aii), with mitotically active spermatogonia lining the basal
205 lamina and spermatocytes in the lumen of the seminiferous tubules, showing that the culture
206 system supports physiological testicular development, given that pre-culture control testis
207 displayed presence of germ cells only at the spermatogonial stage (Fig. 1 Ai). Exposure to low
208 sodium arsenite concentrations did not cause any morphological changes in seminiferous
209 tubules when compared to the control group: seminiferous tubules maintained their
210 physiological shape with abundant spermatogonial germ cells present in tissues treated with
211 both low arsenic doses (0.5 or 1 μM ; Fig. 1 Bii and Biii). In contrast, higher concentrations of
212 sodium arsenite severely disrupted normal testicular morphology (Fig. 1 Cii, Ciii, and Civ).
213 Apoptotic/necrotic cells were present, as evidenced by dense dark staining of the chromatin
214 and more intense pink/red staining of the cytoplasm, and seminiferous tubules were irregularly
215 defined, particularly at the 50 or 100 μM arsenic doses.


216 3.2. Germ cell density

217 Density of spermatogonial germ cells (MVH⁺) was assessed at the end of the culture
218 period for both sets of experiments, low and high arsenic doses (Figs 2 and 3 respectively).

219

220

Figure 2



221


222 **Figure 2. Effect of low sodium arsenite concentrations *in vitro* exposure on germ cell**
223 **density and apoptosis in the prepubertal mouse testis.** Photomicrographs showing
224 immunohistochemical localization of MVH (red) and CC3 (green) expression, counterstained
225 with DAPI (blue) in representative images of low dose arsenic experiment. In control tissue
226 (**Ai**), as well as in the testis fragments exposed to low doses of arsenic (**Aii**, 0.5 μ M sodium
227 arsenite; **Aiii**, 1 μ M sodium arsenite), germ (MVH⁺) cells were abundant and apoptotic (CC3⁺)

228 areas were sparse. Scale bars represent 100 μm . Graphs show germ cell density: the number of
229 MVH⁺ cells per mm^2 of seminiferous tubules (**Aiv**) and per mm^2 of section (**Av**). Density of
230 germ cells was unaffected in the low arsenic dose treated testis fragments compared to that of
231 control tissue. Data represents individual data points (N=6), mean \pm SEM.


232

233 **Figure 3**

High Sodium Arsenite Doses Experiment



MVH= germ cells CC3= apoptotic cells DAPI= cell nuclei



234

235

236 **Figure 3. Effect of high sodium arsenite concentrations *in vitro* exposure on germ cell**
 237 **density and apoptosis in the prepubertal mouse testis. Photomicrographs showing**
 238 **immunohistochemical localization of MVH (red) and CC3 (green) expression, counterstained**
 239 **with DAPI (blue) in representative images of high dose arsenic experiment. In control tissue**

240 (Ai), germ (MVH⁺) cells were abundant, and apoptotic (CC3⁺) areas were sparse. In contrast,
241 in all arsenic treated groups (Aii, 10 μM; Aiii, 50 μM; Aiv, 100 μM sodium arsenite) a lower
242 density of germ cells was present, with vast areas of CC3 expression. Graphs show germ cell
243 density: the number of MVH⁺ cells per mm² of seminiferous tubules (Av) and per mm² of
244 section (Avi). A significant reduction in germ cell density occurred in all three groups exposed
245 to high concentrations of sodium arsenite compared to control tissue. Note that it was not
246 possible to obtain data about germ cell density within seminiferous tubules in the 50 and 100
247 μM arsenic dose groups due to severe loss of tubule integrity. Data represent individual data
248 points (N=6), mean ± SEM. ** and *** indicate p < 0.01 and 0.001, respectively.

249

250 Control tissues had many MVH⁺ cells lining the seminiferous tubule basal membrane
251 (Fig. 2 Ai and 3 Ai). Similarly, germ cells were numerous in testis fragments exposed to low
252 doses of sodium arsenite (Fig. 2 Aii and Aiii). In contrast, testis exposed to high doses of arsenic
253 contained only sparsely distributed germ cells (Fig. 3 Aii, Aiii, and Aiv). Analysis confirmed
254 that, in comparison to control tissue, germ cell density per tubule area and per section area of
255 treated tissues in the low arsenic dose experiment remained unaffected (p-value 0.47 and 0.99
256 for 0.5 and 1 μM respectively; Fig. 2 Aiv and Avi); in contrast, in the high arsenic doses
257 experiment, sodium arsenite exposure caused a significant decrease in germ cell density (Fig.
258 3 Av and Avi). Compared to the control group, germ cell density per tubule area was
259 significantly reduced in 10 μM arsenic treated tissues, even though tubule integrity was still
260 maintained (p < 0.01) (Fig. 3 Av). Due to seminiferous tubule disruption in tissues exposed to
261 higher doses of sodium arsenite, it was not possible to accurately define tubule margins: as a
262 consequence, in these cases density of MVH⁺ cells was assessed only relative to testis section
263 area (Fig. 3 Avi): germ cell density per area of testis section decreased by a 2.5-fold magnitude


264 after 10 μ M arsenic treatment, while after 50 or 100 μ M exposure the decrease was 3-fold for
265 all three doses ($p < 0.001$ for all).

266 3.2. Analysis of apoptosis

267 Activation of the apoptotic marker CC3 was measured at the end of the culture period.
268 In the experiment using low arsenic doses, expression of CC3 was low overall, with arsenic
269 exposed tissues (Fig. 2 Aii and Aiii) similar to those of control tissues (Fig. 2 Ai). In contrast,
270 exposure to high doses of sodium arsenite induced increased CC3 expression (Fig. 3 Aii, Aiii,
271 and Aiv) compared to control tissue (Fig. 3 Ai).

272 Analysis of apoptosis after exposure to low arsenic doses, revealed that the treatment
273 did not affect the area of CC3 expression (Fig. 4 A): the percentage of the seminiferous tubule
274 (Fig 4 Ai), interstitium (Fig 4 Aii) and total section (Fig 4 Aiii) area expressing CC3 was similar
275 to that of control tissue ($p = 0.968, 0.88$ and 0.958 , respectively).

276



278

279

280 **Figure 4. Apoptosis of the prepubertal mouse testis was induced by *in vitro* exposure to**
 281 **high arsenic doses.** Percentage of CC3 positive area of testicular fragments was not affected
 282 by low arsenic doses exposure compared to control (**Ai**, **Aii**, **Aiii**). However, exposure to 10
 283 μM arsenic increased the CC3 positive area of seminiferous tubules (**Bi**). In the interstitium,
 284 exposure to 10 μM did not affect CC3 expression compared to control tissue (**Bii**). The
 285 percentage of CC3 expression per total section area was increased compared with control after
 286 50 μM arsenic exposure only (**Biii**). Note that it was not possible to obtain data about apoptosis
 287 within seminiferous tubules in the 50 and 100 μM arsenic dose groups due to the severe loss
 288 of tubule integrity. Data represent individual data points (N=6), mean ± SEM. * and ***
 289 indicate p < 0.05 and 0.001, respectively.

290

291 As per the analysis of germ cell density, in the high arsenic dose experiment, the loss
292 of the seminiferous tubule integrity after exposure to 50 and 100 μM sodium arsenite rendered
293 it impossible to analyze CC3 expression per unit tubule area. In the 10 μM treatment group,
294 1.5% of the seminiferous tubule area expressed CC3, significantly increased over expression
295 in the control tissue ($p < 0.05$; Fig. 4 Bi). No change was seen outside of the tubule areas, with
296 the percentage of the interstitium expressing CC3 unaffected by treatment ($p = 0.43$; Fig. 4
297 Bii). When examining the tissue as a whole, CC3 expression increased a hundred-fold in tissues
298 exposed to 50 μM sodium arsenite compared to control tissue (Fig. 4 Biii). All three of the high
299 arsenic doses exhibited increased CC3 expression compared to control tissue (3.2%, 10.5%,
300 and 2.9% of the section area CC3-positive in the 10, 50, and 100 μM treatment groups
301 respectively, compared to 0.1% of the tissue CC3-positive in the control tissue), although only
302 the 50 μM sodium arsenite group reached statistical significance ($p < 0.001$; Fig. 4 Biii).

303

304 **4. Discussion**

305 Arsenic is a well-known carcinogen present naturally in the Earth's crust. Human and
306 animal exposure to arsenic is a common consequence of consumption of contaminated food or
307 water, or arising from the use of contaminated water for crop irrigation. Arsenic is also used in
308 a variety of industrial processes, contributing to its environmental distribution as well as
309 representing a risk of occupational exposure. Both chronic and acute exposure to arsenic can
310 contribute to a spectrum of diseases in humans. Effects of arsenic on adult male reproduction
311 has been studied, with reports that arsenic inhibits steroidogenesis, causes spermatotoxicity,
312 and adversely impacts on the weight of the reproductive organs of mice *in vivo* (Pant et al.,
313 2004; Sarkar et al., 2003). Similarly, *in vivo* arsenic exposure induces biochemical and

314 morphological changes in adult rat testis (de Araujo Ramos et al., 2017). Given the marked
315 differences between the pre- and post-pubertal testis, it is important to investigate effects of
316 sodium arsenite directly on the developing male reproductive tissues; with, in contrast, little
317 information available: thirty-day administration of sodium arsenite to prepubertal rats induced
318 dysfunction of normal prostate development (Aquino et al., 2019), spermatogenesis and
319 epididymal structure (da Cunha de Medeiros et al., 2019). Here, we have used an *in vitro* culture
320 system of prepubertal mouse testis to assess the impact of arsenic on the testicular development.
321 The culture system used is capable of supporting testicular development (Sato et al., 2011) and
322 has already been used by our group (Allen et al., 2020; Lopes et al., 2016; Smart et al., 2018)
323 as well as others (Chapin et al., 2016; Nakamura et al., 2019; Park et al., 2020) for toxicological
324 studies on the prepubertal testis. In the results reported here, consistency and health of the
325 control tissue, along with the presence of active spermatogenesis within the vast majority of
326 seminiferous tubules and the absence of any necrotic central area, strengthens the suitability of
327 this culture model for the evaluation of the short-term impact of a toxicant on the immature
328 testis (Chapin et al., 2016; Nakamura et al., 2019). Use of such *in vitro* models, in addition to
329 allowing strict control over exposure conditions, also complies with regulations requiring
330 reduction of animal suffering by avoiding *in vivo* administration. This is particularly important
331 here because there is little information about the toxicological effect of arsenic upon neonatal
332 mice and because it allows examination of the effects of arsenic on the early reproductive
333 system whilst avoiding the inherent difficulties that come with administration of a toxicant to
334 early neonatal animals (Twaddle et al., 2019). Nonetheless, the absence of physiological
335 pharmacokinetics of arsenic in the body, inevitably influencing circulating and tissue arsenic
336 levels, needs to be taken into account when extrapolating data from the present work, as for
337 any *in vitro* study. It is known that some metabolic processes, such as methylation, increase
338 renal clearance, reducing arsenic toxicity, although it can also produce even more cytotoxic

339 compounds (Drobna et al., 2009; Twaddle et al., 2018a). Furthermore, a non-linear relationship
340 has been observed between dose of arsenic administered to experimental animals and its
341 metabolism and toxicity, given which it could be difficult to interpret *in vivo* experiments that
342 use doses higher than human exposure (Twaddle et al., 2018b).

343 Exposure to low sodium arsenite concentrations (0.5 or 1 μM) for six days did not
344 appear to affect mouse testis integrity, with regularly shaped seminiferous tubules lined by
345 germ cells at spermatogonial and spermatocyte phases. On the other hand, high arsenic
346 concentrations (10, 50, or 100 μM) caused significant changes in the testicular structure, with
347 a dose-response pattern: sodium arsenite at 10 μM concentration induced a significant damage
348 with a number of degenerating cells, despite leaving the seminiferous tubule morphology
349 reasonably well distinguishable, while both 50 and 100 μM sodium arsenite treatments
350 drastically compromised seminiferous tubule structure. In all three high arsenic doses, the germ
351 cell population was drastically reduced. The extensive disorganization along with loss of germ
352 cells, shape, as well as shrinkage of the seminiferous tubule observed in the present study after
353 high arsenic exposure is in line with similar observations previously reported in the adult testis
354 after arsenic exposure *in vivo* (Baltaci et al., 2016; Ferreira et al., 2012). A study on the effect
355 of arsenic on prepubertal rats showed abnormal testicular structure with impairment of Sertoli
356 cell number (da Cunha de Medeiros et al., 2019). That study revealed no dose-response pattern,
357 with the both low and high doses administered causing similar effect on the testis, in contrast
358 with results here where a difference effects are observed in low and high concentrations.
359 Nonetheless, a direct comparison between the two studies is challenging, as the *in vivo* study
360 on rats used only two arsenic doses and for an extended (30-day) period of exposure (da Cunha
361 de Medeiros et al., 2019), while our *in vitro* study exposed mouse tissues to five different
362 concentrations for six days. A dose-dependent reduction of germ cells has been already

363 reported in adult mice exposed *in vivo* to sodium arsenite both short- and long-term
364 (Sanghamitra et al., 2008; Zeng et al., 2019).

365 Results here also showed an increase in the density of apoptotic cells present within
366 seminiferous tubules after exposure to 10 μ M sodium arsenite, with an increase in apoptosis
367 also seen in the 50 μ M sodium arsenite treatment group: in the latter case, tubules were ill
368 defined, and so analysis could only examine density of apoptotic cells across the entire
369 testicular section. Paradoxically, the increased expression of CC3 after exposure to 100 μ M of
370 sodium arsenite did not reach statistical significance. One possible explanation could be that at
371 the top dose, apoptotic cells died shortly after arsenic administration and the CC3 signal had
372 disappeared relatively rapidly, long before the six days culture end point: the entire apoptotic
373 cell death process can be complete within as short a time as 2-3 hours (Elmore, 2007).
374 Apoptosis is self-destruction process which can occurs by intrinsic and/or extrinsic pathways
375 (Dua et al., 2015). Reactive oxygen species (ROS) produced by mitochondria are considered
376 as the biochemical mediators of apoptosis (reviewed in Orrenius et al., 2015), thus a high
377 enough accumulation of ROS can initiate apoptosis (reviewed in Redza-Dutordoir and Averill-
378 Bates, 2016). Arsenic is known to cause oxidative stress by increasing ROS production and
379 reducing antioxidant defense systems (Yamanaka et al., 1991). Available literature indicates
380 that *in vivo* arsenic exposure can cause an increase in testicular oxidative stress and apoptosis
381 (Das et al., 2009; Uygur et al., 2016). Here, it is possible that high arsenic exposure to the
382 prepubertal testis culture might have induced ROS generation and oxidative stress, thus
383 resulting in cleaved caspase-3-mediated apoptosis.

384 According to the World Health Organization (WHO), water is safe for drinking only
385 where there are concentrations of arsenic that are less than 10 parts per billion (ppb) (Brown
386 and Ross, 2002), but the concentrations of arsenic in the drinking water in some areas of
387 developing countries such as Pakistan, Bangladesh, Bihar and India are well above that limit

388 (Brammer and Ravenscroft, 2009; Sanjrani et al., 2017). For instance, in Bangladesh, the
389 national standard for arsenic in drinking water is set at 50 ppb, although survey results have
390 shown that 25% of wells providing domestic water exceed even that level (Ravenscroft, 2005).
391 Analysis of water from irrigation shallow tube wells in Bangladesh has revealed that most of
392 them produce water with levels above 100 ppb arsenic, in several instances even above 200
393 ppb arsenic (Islam, 2005). Top arsenic concentrations of 1891 ppb have been found in Bihar
394 and India (Ghosh, 2007), with the highest of all in Nepal, 2629 ppb arsenic (Shrestha, 2003).
395 The doses of sodium arsenite used in the present study are greater than the WHO safe limits of
396 arsenic in the drinking water, but the doses used in the low arsenic study are well within the
397 range of arsenic concentrations found in drinking water in several developing countries (Ghosh,
398 2007); our low arsenic doses of 0.5 and 1 μM correspond to 64 and 129 ppb, respectively. In
399 the present study, lower doses administered for the short period of six days did not affect
400 testicular development. These results cannot, however, exclude the possibility that chronic
401 long-term exposure to such concentrations of arsenic may be detrimental to the prepubertal
402 testis, and further longer term studies are advisable. Several studies on rodents have revealed
403 arsenic accumulation in the testis and other accessory sex organs (Dua et al., 2015; Pant et al.,
404 2001; Prathima et al., 2018). Such persistent nature after accumulation in the reproductive
405 organs can also adversely impact the developing testis in both human and animals.
406 Furthermore, it is possible that other physiological processes have been compromised outside
407 those assessed in the present study that might influence fertility in the adulthood. In the high
408 arsenic doses experiment here, the lowest concentration of 10 μM , corresponding to 1299 ppb
409 arsenic, was highly detrimental to normal testicular physiology even after six days of exposure,
410 with increased cell death and a reduction in spermatogonial germ cell density. Such dose is
411 below the top concentrations of arsenic found in irrigation shallow tube wells, wells that are
412 used as a source of drinking water for local populations in rural areas of developing countries.

413 Data here suggest a potential concerning scenario for the population living in the highly arsenic
414 contaminated rural areas. Caution needs to be exercised before extrapolating data obtained
415 from an *in vitro* mouse study such as this to the situation in human and animal populations, in
416 light of the limitations previously discussed: as such, further studies are required in order to
417 confirm the translational risk of future fertility impairment for young boys. This is likely to
418 include the need for *in vivo* studies, in order to fully evaluate the long-term impact, and the
419 effect of metabolic processes on arsenic circulating levels and toxicity. Comparative *in vivo*
420 studies would also be required to validate the *in vitro* system. Such investigation should be
421 extended to the fertility of large animals; in areas where animal husbandry is an important
422 economic factor, arsenic-induced infertility of farm animals could represent a further financial
423 burden for the unstable economy of farmers in developing countries.

424

425 In conclusion, using a short term culture system of mouse prepubertal testis, this study
426 shows that exposure to high levels of arsenic for six days caused a dose dependent reduction
427 of spermatogonial germ cells and a marked increase in testicular cell apoptosis.

428

429 **Conflict of Interest**

430 The authors declare no conflict of interest.

431

432 **Funding**

433 The present work was funded by Higher Education Commission, Pakistan.

434

435 **References**

- 436 Abdul, K.S., Jayasinghe, S.S., Chandana, E.P., Jayasumana, C., De Silva, P.M., 2015. Arsenic and human
437 health effects: A review. *Environ Toxicol Pharmacol* 40, 828-846.
- 438 Abhyankar, L.N., Jones, M.R., Guallar, E., Navas-Acien, A., 2012. Arsenic exposure and hypertension: a
439 systematic review. *Environ Health Perspect* 120, 494-500.
- 440 Allen, C.M., Lopes, F., Mitchell, R.T., Spears, N., 2020. Comparative gonadotoxicity of the
441 chemotherapy drugs cisplatin and carboplatin on prepubertal mouse gonads. *Mol Hum Reprod* 26,
442 129-140.
- 443 Aquino, A.M., Salata, G.C., Pinho, C.F., de Freitas, A., Perico, L.L., de Lion Siervo, G.E.M., Mendes, L.O.,
444 Da Cunha De Medeiros, P., Justulin, L.A., Fernandes, G.S.A., Perobelli, J.E., Scarano, W.R., 2019. Arsenic
445 exposure during prepuberty alters prostate maturation in pubescent rats. *Reprod Toxicol* 89, 136-144.
- 446 Baltaci, B.B., Uygur, R., Caglar, V., Aktas, C., Aydin, M., Ozen, O.A., 2016. Protective effects of quercetin
447 against arsenic-induced testicular damage in rats. *Andrologia* 48, 1202-1213.
- 448 Brammer, H., Ravenscroft, P., 2009. Arsenic in groundwater: a threat to sustainable agriculture in
449 South and South-east Asia. *Environ Int* 35, 647-654.
- 450 Brown, K.G., Ross, G.L., 2002. Arsenic, drinking water, and health: a position paper of the American
451 Council on Science and Health. *Regul Toxicol Pharmacol* 36, 162-174.
- 452 Chaineau, E., Binet, S., Pol, D., Chatellier, G., Meininger, V., 1990. Embryotoxic effects of sodium
453 arsenite and sodium arsenate on mouse embryos in culture. *Teratology* 41, 105-112.
- 454 Chang, S.I., Jin, B., Youn, P., Park, C., Park, J.D., Ryu, D.Y., 2007. Arsenic-induced toxicity and the
455 protective role of ascorbic acid in mouse testis. *Toxicol Appl Pharmacol* 218, 196-203.
- 456 Chapin, R.E., Winton, T., Nowland, W., Danis, N., Kumpf, S., Johnson, K., Coburn, A., Stukenborg, J.B.,
457 2016. Lost in translation: The search for an in vitro screen for spermatogenic toxicity. *Birth Defects*
458 *Res B Dev Reprod Toxicol* 107, 225-242.
- 459 da Cunha de Medeiros, P., Samelo, R.R., Silva, A.P.G., da Silva Araujo Santiago, M., Duarte, F.A., de
460 Castro, I.B., Perobelli, J.E., 2019. Prepubertal exposure to low doses of sodium arsenite impairs
461 spermatogenesis and epididymal histophysiology in rats. *Environ Toxicol* 34, 83-91.
- 462 Das, J., Ghosh, J., Manna, P., Sinha, M., Sil, P.C., 2009. Taurine protects rat testes against NaAsO(2)-
463 induced oxidative stress and apoptosis via mitochondrial dependent and independent pathways.
464 *Toxicol Lett* 187, 201-210.
- 465 de Araujo Ramos, A.T., Diamante, M.A.S., de Almeida Lamas, C., Dolder, H., de Souza Predes, F., 2017.
466 Morphological and morphometrical changes on adult Wistar rat testis caused by chronic sodium
467 arsenite exposure. *Environ Sci Pollut Res Int* 24, 27905-27912.
- 468 Drobna, Z., Styblo, M., Thomas, D.J., 2009. An Overview of Arsenic Metabolism and Toxicity. *Curr*
469 *Protoc Toxicol* 42, 4.31.31-34.31.36.
- 470 Dua, T.K., Dewanjee, S., Gangopadhyay, M., Khanra, R., Zia-Ul-Haq, M., De Feo, V., 2015. Ameliorative
471 effect of water spinach, *Ipomea aquatica* (Convolvulaceae), against experimentally induced arsenic
472 toxicity. *J Transl Med* 13, 81.
- 473 Duker, A.A., Carranza, E.J., Hale, M., 2005. Arsenic geochemistry and health. *Environ Int* 31, 631-641.
- 474 Elmore, S., 2007. Apoptosis: a review of programmed cell death. *Toxicol Pathol* 35, 495-516.
- 475 Feng, Z., Xia, Y., Tian, D., Wu, K., Schmitt, M., Kwok, R.K., Mumford, J.L., 2001. DNA damage in buccal
476 epithelial cells from individuals chronically exposed to arsenic via drinking water in Inner Mongolia,
477 China. *Anticancer Res* 21, 51-57.
- 478 Ferreira, M., Matos, R.C., Oliveira, H., Nunes, B., Pereira Mde, L., 2012. Impairment of mice
479 spermatogenesis by sodium arsenite. *Hum Exp Toxicol* 31, 290-302.
- 480 Ghosh, A.S., S.K. Bose, N. Chowdhary, S., 2007. Arsenic contaminated aquifers: a study of the Ganga
481 levee zone in Bihar, India. Symposium on Arsenic: the geography of a global problem, Royal
482 Geographical Society, London.
- 483 Huang, C., Ke, Q., Costa, M., Shi, X., 2004. Molecular mechanisms of arsenic carcinogenesis. *Mol Cell*
484 *Biochem* 255, 57-66.

485 Islam, M.R.J., M. Rahman, G.K.M.M, Miah, M.A.M. Farid, A.T.M. Panaullah, G.M. Loeppert, R.H.
486 Duxbury, J.M. Meisner, C.A. , 2005. Arsenic in paddy soils of Bangladesh: Levels, distribution and
487 contribution of irrigation and sediments. . Symposium on the Behaviour of Arsenic in Aquifers, Soils
488 and Plants: Implications for Management, Dhaka.
489 Kim, Y.J., Chung, J.Y., Lee, S.G., Kim, J.Y., Park, J.E., Kim, W.R., Joo, B.S., Han, S.H., Yoo, K.S., Yoo, Y.H.,
490 Kim, J.M., 2011. Arsenic trioxide-induced apoptosis in TM4 Sertoli cells: the potential involvement of
491 p21 expression and p53 phosphorylation. *Toxicology* 285, 142-151.
492 Kim, Y.J., Kim, J.M., 2015. Arsenic Toxicity in Male Reproduction and Development. *Dev Reprod* 19,
493 167-180.
494 Lima, G.D.A., Sertorio, M.N., Souza, A.C.F., Menezes, T.P., Mouro, V.G.S., Goncalves, N.M., Oliveira,
495 J.M., Henry, M., Machado-Neves, M., 2018. Fertility in male rats: Disentangling adverse effects of
496 arsenic compounds. *Reprod Toxicol* 78, 130-140.
497 Lopes, F., Smith, R., Nash, S., Mitchell, R.T., Spears, N., 2016. Irinotecan metabolite SN38 results in
498 germ cell loss in the testis but not in the ovary of prepubertal mice. *Mol Hum Reprod* 22, 745-755.
499 Maitani, T., Saito, N., Abe, M., Uchiyama, S., Saito, Y., 1987. Chemical form-dependent induction of
500 hepatic zinc-thionein by arsenic administration and effect of co-administered selenium in mice. *Toxicol*
501 *Lett* 39, 63-70.
502 Muller, W.U., Streffer, C., Fischer-Lahdo, C., 1986. Toxicity of sodium arsenite in mouse embryos in
503 vitro and its influence on radiation risk. *Arch Toxicol* 59, 172-175.
504 Munoz, A., Chervona, Y., Hall, M., Kluz, T., Gamble, M.V., Costa, M., 2015. Sex-specific patterns and
505 deregulation of endocrine pathways in the gene expression profiles of Bangladeshi adults exposed to
506 arsenic contaminated drinking water. *Toxicol Appl Pharmacol* 284, 330-338.
507 Nakamura, N., Sloper, D.T., Del Valle, P.L., 2019. Evaluation of an in vitro mouse testis organ culture
508 system for assessing male reproductive toxicity. *Birth Defects Res* 111, 70-77.
509 Orrenius, S., Gogvadze, V., Zhivotovsky, B., 2015. Calcium and mitochondria in the regulation of cell
510 death. *Biochem Biophys Res Commun* 460, 72-81.
511 Pant, N., Kumar, R., Murthy, R.C., Srivastava, S.P., 2001. Male reproductive effect of arsenic in mice.
512 *Biometals* 14, 113-117.
513 Pant, N., Murthy, R.C., Srivastava, S.P., 2004. Male reproductive toxicity of sodium arsenite in mice.
514 *Hum Exp Toxicol* 23, 399-403.
515 Park, H.J., Lee, W.Y., Zhang, M., Hong, K.H., Park, C., Kim, J.H., Song, H., 2020. Evaluation of Resmethrin
516 Toxicity to Neonatal Testes in Organ Culture. *Toxicol Sci* 173, 53-64.
517 Podgorski, J.E., Eqani, S., Khanam, T., Ullah, R., Shen, H., Berg, M., 2017. Extensive arsenic
518 contamination in high-pH unconfined aquifers in the Indus Valley. *Sci Adv* 3, e1700935.
519 Prathima, P., Pavani, R., Sukeerthi, S., Sainath, S.B., 2018. alpha-Lipoic acid inhibits testicular and
520 epididymal oxidative damage and improves fertility efficacy in arsenic-intoxicated rats. *J Biochem Mol*
521 *Toxicol* 32.
522 Rahman, M., Sohel, N., Yunus, M., Chowdhury, M.E., Hore, S.K., Zaman, K., Bhuiya, A., Streatfield, P.K.,
523 2013. Increased childhood mortality and arsenic in drinking water in Matlab, Bangladesh: a
524 population-based cohort study. *PLoS One* 8, e55014.
525 Ratnaik, R.N., 2003. Acute and chronic arsenic toxicity. *Postgrad Med J* 79, 391-396.
526 Ravenscroft, B.W., Ahmed K.M., Burren M. , Perrin J., 2005. Arsenic in groundwater of the Bengal
527 Basin, Bangladesh: distribution, field relations and hydrogeological setting. *Hydrogeology Journal* 13,
528 727-751.
529 Redza-Dutordoir, M., Averill-Bates, D.A., 2016. Activation of apoptosis signalling pathways by reactive
530 oxygen species. *Biochim Biophys Acta* 1863, 2977-2992.
531 Sanghamitra, S., Hazra, J., Upadhyay, S.N., Singh, R.K., Amal, R.C., 2008. Arsenic induced toxicity on
532 testicular tissue of mice. *Indian J Physiol Pharmacol* 52, 84-90.
533 Sanjrani, M., Mek, T., Sanjrani, N., Leghari, S., Moryani, H., Shabnam, A., 2017. Current situation of
534 aqueous arsenic contamination in Pakistan, focused on Sindh and Punjab Province, Pakistan: A review.
535 *J Pollut Eff Cont* 5, 2.

536 Sarkar, M., Chaudhuri, G.R., Chattopadhyay, A., Biswas, N.M., 2003. Effect of sodium arsenite on
537 spermatogenesis, plasma gonadotrophins and testosterone in rats. *Asian J Androl* 5, 27-31.

538 Sato, T., Katagiri, K., Gohbara, A., Inoue, K., Ogonuki, N., Ogura, A., Kubota, Y., Ogawa, T., 2011. In vitro
539 production of functional sperm in cultured neonatal mouse testes. *Nature* 471, 504-507.

540 Scheuplein, R., Charnley, G., Dourson, M., 2002. Differential sensitivity of children and adults to
541 chemical toxicity. I. Biological basis. *Regul Toxicol Pharmacol* 35, 429-447.

542 Shresta, B.R., Witney, J.W. and Shresta, K.B., 2003. The state of arsenic in Nepal –2003. Kathmandu:
543 National Arsenic Steering Committee. Environment and Public Health Organisation, Kathmandu, Nepal
544 126.

545 Smart, E., Lopes, F., Rice, S., Nagy, B., Anderson, R.A., Mitchell, R.T., Spears, N., 2018. Chemotherapy
546 drugs cyclophosphamide, cisplatin and doxorubicin induce germ cell loss in an in vitro model of the
547 prepubertal testis. *Sci Rep* 8, 1773.

548 Smith, J., Jankowski & Sammut, 2003. Vertical distribution of As(III) and As(V) in a coastal sandy aquifer:
549 factors controlling the concentration and speciation of arsenic in the Stuarts Point groundwater
550 system, northern New South Wales, Australia. *Applied Geochemistry* 18, 1479–1496.

551 Styblo, M., Del Razo, L.M., Vega, L., Germolec, D.R., LeCluyse, E.L., Hamilton, G.A., Reed, W., Wang, C.,
552 Cullen, W.R., Thomas, D.J., 2000. Comparative toxicity of trivalent and pentavalent inorganic and
553 methylated arsenicals in rat and human cells. *Arch Toxicol* 74, 289-299.

554 Sulanjari, D., Handryani, E., Setiyono, A., 2015. Subchronic accumulation of arsenic (Arsenic Acid) on
555 sprague dawley rats. *American-Eurasian Journal of Toxicological Sciences* 7, 88-94.

556 Tabocova, S., Hunter, E.S., 3rd, Gladen, B.C., 1996. Developmental toxicity of inorganic arsenic in
557 whole embryo: culture oxidation state, dose, time, and gestational age dependence. *Toxicol Appl*
558 *Pharmacol* 138, 298-307.

559 Twaddle, N.C., Vanlandingham, M., Beland, F.A., Doerge, D.R., 2018a. Metabolism and disposition of
560 arsenic species after repeated oral dosing with sodium arsenite in drinking water. II. Measurements
561 in pregnant and fetal CD-1 mice. *Food Chem Toxicol* 115, 178-184.

562 Twaddle, N.C., Vanlandingham, M., Beland, F.A., Fisher, J.W., Doerge, D.R., 2019. Metabolism and
563 disposition of arsenic species from oral dosing with sodium arsenite in neonatal CD-1 mice. IV.
564 Toxicokinetics following gavage administration and lactational transfer. *Food Chem Toxicol* 123, 28-
565 41.

566 Twaddle, N.C., Vanlandingham, M., Churchwell, M.I., Doerge, D.R., 2018b. Metabolism and disposition
567 of arsenic species from controlled oral dosing with sodium arsenite in adult female CD-1 mice. I. Pilot
568 study to determine dosing, analytical measurements, and sampling strategies. *Food Chem Toxicol* 111,
569 482-493.

570 Uygur, R., Aktas, C., Caglar, V., Uygur, E., Erdogan, H., Ozen, O.A., 2016. Protective effects of melatonin
571 against arsenic-induced apoptosis and oxidative stress in rat testes. *Toxicol Ind Health* 32, 848-859.

572 Vega, L., Styblo, M., Patterson, R., Cullen, W., Wang, C., Germolec, D., 2001. Differential effects of
573 trivalent and pentavalent arsenicals on cell proliferation and cytokine secretion in normal human
574 epidermal keratinocytes. *Toxicol Appl Pharmacol* 172, 225-232.

575 Wang, A., Holladay, S.D., Wolf, D.C., Ahmed, S.A., Robertson, J.L., 2006. Reproductive and
576 developmental toxicity of arsenic in rodents: a review. *Int J Toxicol* 25, 319-331.

577 Xie, X., Wang, Y., Ellis, A., Su, C., Li, J., Li, M., 2011. The sources of geogenic arsenic in aquifers at Datong
578 basin, northern China: Constraints from isotopic and geochemical data. *Journal of Geochemical*
579 *Exploration*. 110, 155-166.

580 Yamanaka, K., Hasegawa, A., Sawamura, R., Okada, S., 1991. Cellular response to oxidative damage in
581 lung induced by the administration of dimethylarsinic acid, a major metabolite of inorganic arsenics,
582 in mice. *Toxicol Appl Pharmacol* 108, 205-213.

583 Yuan, Y., Marshall, G., Ferreccio, C., Steinmaus, C., Selvin, S., Liaw, J., Bates, M.N., Smith, A.H., 2007.
584 Acute myocardial infarction mortality in comparison with lung and bladder cancer mortality in arsenic-
585 exposed region II of Chile from 1950 to 2000. *Am J Epidemiol* 166, 1381-1391.

586 Zeng, Q., Yi, H., Huang, L., An, Q., Wang, H., 2019. Long-term arsenite exposure induces testicular
587 toxicity by redox imbalance, G2/M cell arrest and apoptosis in mice. *Toxicology* 411, 122-132.
588