

Northumbria Research Link

Citation: Farsi, Dominic N., Uthumange, Dinithi, Munoz Munoz, Jose and Commane, Daniel M. (2021) The nutritional impact of replacing dietary meat with meat alternatives in the UK: a modelling analysis using nationally representative data. *British Journal of Nutrition*. ISSN 0007-1145 (In Press)

Published by: Cambridge University Press

URL: <https://doi.org/10.1017/S0007114521002750>
<<https://doi.org/10.1017/S0007114521002750>>

This version was downloaded from Northumbria Research Link:
<http://nrl.northumbria.ac.uk/id/eprint/46726/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)

The nutritional impact of replacing dietary meat with meat alternatives in the UK: a modelling analysis using nationally representative data

Dominic N Farsi¹, Dinithi Uthumange¹, Jose Munoz Munoz¹, Daniel M Commane¹

¹Applied and Health Sciences, University of Northumbria, Sutherland Building, Ellison Place, Newcastle upon Tyne, NE1 8ST, United Kingdom

Corresponding author: Dominic N Farsi, dominic.farsi@northumbria.ac.uk

Short title: Nutritional impact of meat alternatives.



This peer-reviewed article has been accepted for publication but not yet copyedited or typeset, and so may be subject to change during the production process. The article is considered published and may be cited using its DOI

10.1017/S0007114521002750

The British Journal of Nutrition is published by Cambridge University Press on behalf of The Nutrition Society

Abstract

Dietary patterns high in meat compromise both planetary and human health. Meat-alternatives may help facilitate meat reduction, however the nutritional implications of displacing meat with meat-alternatives does not appear to have been evaluated. Here, data from the 9th cycle of the National Diet and Nutrition Survey was used as the basis of models to assess the effect of meat substitution on nutritional intake. We implemented three models; model 1 progressively replaced 25%, 50%, 75%, or 100% of the current meat intake with a weighted mean of meat-alternatives available in the UK market. Model 2 compared different ingredient categories of meat-alternative; vegetable, mycoprotein, a combination of bean and pea, tofu, nut and soy. Model 3 compared fortified versus unfortified meat-alternatives. The models elicited significant shifts in nutrients. Overall, there were increases in carbohydrate, fibre, sugars and sodium, whereas reductions were found for protein, total and saturated fat, iron and B12. The greatest effects were seen for; vegetable-based (+24.63g/day carbohydrates), mycoprotein-based (-6.12g/day total fat), nut-based (-19.79g/day protein, +10.23g/day fibre; -4.80g/day saturated fat, +7.44g/day sugars), soy-based (+495.98mg/day sodium), and tofu-based (+7.63mg/day iron, -2.02µg/day B12). Our results suggest meat-alternatives can be a healthful replacement for meat if chosen correctly. Consumers should seek out meat-alternatives which are low in sodium and sugar, high in fibre, protein and with high micronutrient density, to avoid compromising nutritional intake if reducing their meat intake. Manufacturers and policy makers should consider fortification of meat-alternatives with nutrients such as iron and B12 and focus on reducing sodium and sugar content.

Keywords: Meat replacement, meat alternatives, Nutritional intake, Nutritional requirements, Public health.

Introduction

Globally, annual per capita meat consumption has increased by 20 kilograms since 1961 (1). This sector of our food system contributes substantially to greenhouse gas emissions (2), depletes natural resources (3-5), and compromises animal welfare. In the UK, meat can be a valuable source of nutrients, however, overconsumption is associated with an increased risk of cardiovascular disease, type 2 diabetes (6-8), and cancer (9-16). Reducing meat consumption may therefore promote human health, protect animal welfare and benefit the environment.

Changing dietary behaviours to elicit a transition towards alternative protein sources is challenging in part owing to strongly held taste preferences, culinary traditions, and social and cultural norms (17). Meat alternatives, resembling meat in appearance, preparation and eating experience, could therefore play an important role in enabling dietary change (18). Integration of meat alternatives into the diet is subtle and does not require an overhaul of food consumption and meal patterns (19-21). Although meat alternatives have been available for years, their popularity with consumers has increased rapidly in recent times; due in part to producers better simulating the taste, texture and functionality of traditional meat products. There has also been an increase in the direction of marketing towards meat-eating consumers, rather than just vegans and vegetarians (22).

The rapid increase in the consumption of meat alternatives has raised concerns regarding their overall healthfulness and the potential displacement of valuable nutrients. The developing meat alternative market includes an array of food products with variable nutritional quality, some of which might be considered ultra-processed (23). If these foods are to play a role in reducing meat consumption at a population level, it is essential that any population level nutritional benefits and consequences are identified early, and actions taken to mitigate undesirable effects.

Here, we model the potential nutritional implications of meat replacement by using the 9th cycle (years 2016-2017) of the nationally representative UK National Diet and Nutrition Survey rolling program (NDNS RP). We present the nutritional implications of complete and partial meat substitution, using models that either; assume replacement of meat with a range of alternative products, based on current UK purchasing data; or models that assume replacement of meat with specific meat alternative product categories. To our understanding,

this is the first study to evaluate the nutritional implications of such dietary replacement scenarios.

Methods

Study Population

Dietary intake data for the UK population was drawn from the 9th cycle of the NDNS rolling programme. The NDNS rolling programme is a Public Health England and Food Standards Agency funded survey of the food consumption, nutrient intake, and nutritional status of people aged 1.5yr and older living in private households in the UK and is representative of the UK population (24). Detailed descriptions of the NDNS RP survey design and sampling methods can be found elsewhere (25-27). The 2016-2017 survey cycle year used in the present modelling analysis included 1,253 participants aged 1.5-95yr.

Dietary Records

The NDNS RP collected habitual dietary data using four-day estimated food diaries (24-26). Data from food diaries has been aggregated for each respondent to provide publicly available daily averages of self-reported food and nutrient intake. For this analysis we excluded data sets for participants with extreme reported energy intakes (< 500kcal - > 5000kcal). Datasets from subjects who self-reported abstaining from meat, or with dietary data devoid of any meat, were also excluded. A common occurrence in the collection of habitual dietary intake data is underreporting of energy intake (28), which the NDNS RP data drawn upon here corroborates. There are different methods to approach underreporting (29); in the present analysis we adjusted nutrients to age and gender specific EARs for energy.

Sub-analyses were performed by stratifying by sex and age group, using the same stratification system as the NDNS RP (male and female; 4-10yr, 11-18yr and 19-64yr, over 65yr). Children in 1.5-3yr subgroups were not included as nutrient requirements are specific for each age in this sub-population. In addition, dietary patterns at this stage in life may not include substantial amounts of solid food such as meat. This left a total of 1,110 respondents for inclusion in the analysis.

Self-reported nutritional intake from meat

The current level of meat consumption in the UK was estimated from the NDNS dataset. Briefly, the NDNS RP reports meat intake in grams per day broken down by type (i.e. “beef, lamb, pork, offal etc”), We allocated these categories into food type groups; ‘red meat;

‘processed red meat’, ‘white meat’, ‘processed white meat’, and ‘seafood’. In this analysis seafood was included with meat as a source of animal flesh. We note that the UK dietary guidelines encourage a population level increase in fish consumption as a source of omega 3 fatty acids and other nutrients, however, most UK vegetarians describe themselves as lacto-ovo-vegetarian or vegan, and only a minority describe themselves as pescatarian.

The allocation and stratification of animal meat within the NDNS dataset is described in **Table 1**.

The total nutritional contribution of meat to the UK diet was estimated by combining the intake data with the nutritional composition data from the McCance and Widdowson composition of food tables (30). Briefly, an aggregate quantity of nutrient, per gram, for each of the meat subtypes was multiplied by the intake in grams per day, for each respondent. The totality of these nutrition scores by subtype was then used to estimate the total nutritional value of meat products to the diet for each individual respondent.

Nutritional content of meat alternatives

Products were identified through product searches on UK supermarket websites using the key words, ‘meat alternatives’, ‘meat substitutes’, ‘meat-free’, ‘plant-based’, ‘vegan’, and ‘vegetarian’, to ensure a wide capture of meat alternative products. Following data collection, the products that met these criteria were grouped into ingredient categories (**Supplementary Material, Table 1**).

The nutritional profile of meat alternatives was calculated based on product labels, information from supermarket and manufacturer websites, as well as searching nutrition databases for products. Where nutrient data was not available for a product, manufacturers were contacted to request further details. This data was used to estimate the aggregate nutritional composition per gram for each of the meat alternative ingredient categories, based on the mean nutritional composition of the individual products included in each respective category.

There was incomplete availability of nutritional data across the product categories for many of the micronutrients including iodine, zinc, and selenium. There was available nutritional data for the following nutrients: carbohydrate, protein, total fat, saturated fat, fibre, sugar, sodium, iron and B12. We deemed the data on these nutrients of good quality for inclusion in subsequent analysis

Meat Replacement Scenarios

Here, we assessed the nutritional implications of a set of scenarios for the widespread replacement of meat with meat alternatives in the UK population (**Table 2**).

Model 1: The effect of meat reduction and replacement with meat alternatives

In the first replacement model, we generated a weighted mean of the nutritional quality of meat alternatives based on the current market share for meat alternative products in the UK; vegetable 23%, mycoprotein 18%, bean 17%, soy 15%, nut 14%, and tofu 13% (31).

We then looked at the effect of progressively reducing the amount of meat in each respondent's diet by 25% (MA-25), 50% (MA-50), 75% (MA-75), and 100% (MA-100), and replacing the removed nutrients with the weighted nutritional score of the meat alternatives on a per weight basis. (**Table 2**).

Model 2: A comparison of the effects of replacing nutrients from meat across specific ingredient categories of meat alternative.

In a second model, we compared the nutritional impact of replacing meat with meat alternatives produced from a particular category of ingredient. Specifically, we replaced the total amount of meat in each individual respondent's diet with vegetable-based (Vegetable), mycoprotein-based (Mycoprotein), a combination of bean and pea-based (Legume), tofu-based (Tofu), nut-based (Nut) and soy-based (Soy) meat alternatives (**Table 2**).

Model 3: substitution with fortified versus unfortified meat alternatives

Within the food products identified, 14% were fortified with either iron, B12, or both. Taking this into account, in a third and final model, we compared the effects of replacing the total amount of meat in each person's diet with either fortified meat alternatives (Fortified) or unfortified meat alternatives (Unfortified) (**Table 2**).

We implemented each replacement strategy by removing the nutrient contribution of meat from each individual respondents' total nutritional intake, followed by replacing this with the nutritional data for the equivalent weight of meat alternatives. We adjusted the final total nutritional intake after substitution to match the energy content of each participant's diet.

Comparators and subjective model corrections

Due to the variable quality of available nutritional data, and the relevance to public health, we have elected in this analysis to focus on the macronutrients, protein, fat and carbohydrate, as well as saturated fat, total sugars and sodium due to public health guidance to limit their intake (32, 33). In addition, fibre was selected as authoritative bodies advise an intake of 30g/day for the adult population (33); which many fail to achieve (34). We hypothesised that fibre would increase in the replacement models as a result of the higher fibre content of meat alternatives compared to meat (35). We also assessed influences on iron and B12, as the intake of these nutrients may be inadequate when transitioning to a dietary pattern devoid of animal based foods (36).

We analysed the nutritional adequacy of the meat replacement models by comparing the nutrient values to the Dietary Reference Values for Food Energy and Nutrients for the UK (DRVs) (32, 33, 37). DRVs comprise a series of estimates of the amount of energy and nutrients needed by different groups of healthy people in the UK population. In accordance with the DRVs, we used the Estimated Average Requirements (EARs) as a comparator for total energy, whereas for protein, sodium, iron and B12, the Reference Nutrient Intakes (RNIs) were used. For fat and carbohydrates (including total sugars, saturated fat, and fibre) we used the UK DRVs which for total fat (35%), saturated fat (11%), and total carbohydrates (50%), are given as a percentage of daily energy intake (32, 33, 37). The current recommendations for protein is to intake 0.75g per kilogram bodyweight (33, 37), as the dataset in the current analysis contains a diverse age range which will also comprise different body types, a protein target of making up the remaining 15% of total energy intake after the contributions of fat (35%) and carbohydrates (50%) was employed instead.

Since the recommendations differ by age and sex, we calculated the average values for nutrients across the population subgroups (male and female; 4-10yr, 11-18yr, 19-64yr, over 65yr). We also incorporated a group that represented the total number of participants (both male and female, ages 4yr and above), and averaged all DRV categories to obtain a population average. This group would act as the representative of the UK population and used in the main analysis.

Statistical Analysis

Differences in nutrient intake between the meat replacement models and the current intake were compared using linear models with nutrients as the dependent variable, and the replacement models as the independent variable, using the current intake as the reference. The differences in nutrient intake between all scenarios (current and replacement models) with reference to DRVs were then compared using linear models in the same manner but including the current intake with the replacement models as the independent variable and then using DRVs used as the reference. To assess the nutrient intake between meat replacement models, an Analysis of Variance (ANOVA) was implemented, using an $\alpha = 0.05$, followed by a post hoc Tukey test to determine the significance of differences between models. All analysis was performed using R statistical software (version 4.0.0) (38).

Results

Self-reported meat intake

The average self-reported intake of both total meat and meat subgroup for the total population, as well separate population subgroups are shown in **Table 3**. For the total population, the self-reported average meat intake was 132.25g/day (95% CI, 128.02-136.48g/day). Processed meat was the largest contributor to total meat intake (35.98%), followed by white meat (27.22), red meat (19.44%), and seafood (13.36%), with processed white meat contributing the lowest (3.99%). Males aged 19-64 years consumed the greatest amount of meat (mean, 177.55g/day; 95% CI, 165.33-189.77g/day), while girls aged 4-10 years consumed the least amount of meat (mean, 98.75g/day; 95% CI, 91.28-106.23g/day).

Projected changes in nutritional intake

Overall, the implemented meat replacement scenarios elicited many differences in nutritional intake. The current nutrient intake for the total population as well as for each replacement model (MA-25, MA-50, MA-75, MA-100, Vegetable, Mycoprotein, Legume, Tofu, Nut, Soy, Fortified, Unfortified) are included in **Table 4**; while the projected differences for each replacement model in comparison to the current intake are included in **Table 5**.

Model 1: The effect of meat reduction and replacement with meat alternatives

In model 1, we explored projected changes in nutritional intake given a graded replacement of meat from replacing 25% of current meat intake, through to a 100% replacement, using weighted composite nutritional values for the meat alternatives based on consumer purchasing data. We observed a linear increase in total carbohydrate consumption with the decrease in meat, reaching an additional 19 grams per day in a 100% meat replacement scenario. This increase in projected carbohydrates was statistically significant in all scenarios except the 25% replacement model (MA-50, $P = 0.018$; MA-75, $P = <0.001$; MA-100, $P <0.001$). Importantly, the projected increased intake of carbohydrates was only associated with a small linear increase in projected sugar intake; this increase did not reach statistical significance until 100% of current meat was replaced (+04.16g/day, $P = 0.05$). In contrast, with decreasing consumption of meat and increasing consumption of meat alternatives there was a graded decrease in projected total protein intake (MA-25, -3.40g/day, $P = 0.002$; MA-50, -6.99g/day, $P = <0.001$; MA-75, -10.77g/day, $P = <0.001$; MA-100, -14.77g/day, $P <0.001$). There was also a linear decrease in the intake of fat and importantly, of saturated fat, with decreasing meat and increasing meat alternative intake. The decrease in saturated fat with complete meat substitution was equivalent to 3.96 g per day ($P <0.001$).

Total fibre intake was increased significantly under each meat replacement scenario. A 100% replacement of meat with meat alternatives was associated with an 8.15g increase in total daily fibre intake ($P <0.001$). Notably, there was also a linear increase in sodium intake, with a total replacement of meat being associated with a projected 228.03 mg increase in sodium per day ($P <0.001$). There were non-statistically significant decreases in projected intake of both iron and of vitamin B12.

Similar effects were found in sub analyses across each of the population subgroups (**Supplementary Material, Tables 3-18**).

Model 2: A comparison of the effects of replacing nutrients from meat across specific categories of alternative.

This model is based on a 100% replacement of meat with the mean nutrient values from each of the separate categories of meat alternative. Compared to the current intake, the substitution of meat, with foods from any of the meat alternative categories, led to a meaningful increase in the projected mean intake of, fibre. The greatest projected increase in total fibre was observed in models replacing meat with nut-based products, (10.23 g day⁻¹), followed closely

by substitution with vegetable (8.73 g day⁻¹) or mycoprotein based products (8.71 g day⁻¹). Substitution with tofu-based products had the least impact on projected total fibre intake (+3.14 g day⁻¹).

All product categories were associated with meaningful reductions in projected saturated fat intake. The greatest reduction in saturated fat intake is projected for substitutions with nut-based meat alternatives (-4.8 g day⁻¹, $P < 0.001$), whilst the greatest reduction in total projected fat intake was associated with substitution for mycoprotein (6.12 g day⁻¹, $P < 0.001$). Only the replacements with nut-based and tofu-based meat alternatives did not cause significant reductions in projected total fat intake.

Conversely, projected intakes of protein were lower across each meat alternative category. The greatest projected reductions in total protein intake were observed when substituting in products exclusively from the nut (-19.79 g day⁻¹), vegetable (-19.33 g day⁻¹), or legume (-15.43 g day⁻¹) categories. Tofu had the least projected impact on total protein consumption, but it was still associated with a significant reduction in daily intake (-4.87 g day⁻¹).

Replacement of meat with meat alternatives was also associated with an increase in total projected sodium consumption except for the nut-based products (-124.30mg day⁻¹). Soy based meat alternatives had the highest sodium content, and replacement with the soy product category raised projected sodium intake by almost 0.5 g day⁻¹.

Regarding micronutrients, the projected intake of vitamin B12 was shown to be statistically significantly reduced in the replacement models with tofu (-2.02ug/day; $P = 0.03$), nut (-1.88ug/day; $P = 0.04$), and soy (-1.76ug/day; $P = 0.05$), although in each case mean projected intake across the population remained in excess of the RNI.

Total iron intake was projected to increase in the substitution model with tofu (+7.63 mg day⁻¹, $P < 0.001$). It was not significantly affected by the other meat replacement scenarios.

In sub-analyses by population group (age and gender) the described influence of these substitution scenarios were replicated in each of the demographics (**Supplementary Material, Tables 3-18**).

Model 3: substitution with fortified versus unfortified meat alternatives

In this model we compared the impact of selecting iron and B12 fortified, versus unfortified, meat alternatives, assuming a 100% replacement of meat. Fortified products accounted for 14% of the meat alternative products. The projected intake of iron was significantly higher

than current intakes when choosing fortified meat alternatives (+3.12mg/day; $P < 0.001$) and would comfortably attain the UK dietary recommendations (Fortified, 16.77mg/day vs 14.8mg/day RNI for girls and women aged 11-50, 8.7mg/day RNI for remaining population). The projected intake of iron was not statistically significantly reduced from current levels of intake in the unfortified model.

The projected intake of vitamin B12 was significantly decreased compared to current intakes in the unfortified model (-1.79ug/day, $P = 0.05$), however projected mean intake did not fall below the RNI. In the fortified model projected intakes of vitamin B12 were not significantly different from current levels of intake.

Interestingly, fortified meat alternatives were associated with a much smaller projected reduction in total protein from current intakes (-3.09 g day⁻¹ versus -18.19 g day⁻¹ for the unfortified products $P = 0.005$) however, it was also associated with a greater projected increase in total sodium from current levels of intake (-482 mg day⁻¹ versus +361 mg day⁻¹ for the unfortified products $P < 0.001$).

In sub analyses by population group (age and gender) the effect of choosing fortified products described above were replicated across subgroups (**Supplementary Material, Tables 3-18**).

Discussion

We have modelled the projected nutritional impact of a shift in the culinary practices of the meat-eating UK population towards favouring meat alternative products and have identified some important nutritional benefits. Notably, our models predict significant increases in fibre intake and significant decreases in both total fat and saturated fat intake as people switch from meat to meat-alternatives. On the other hand, not all projected impacts on nutrient intake would be described as beneficial; there were notable reductions in total protein and vitamin B12, and projected increases in the intake of sodium and of sugars.

The UK dietary guidelines recommend limiting fat to no more than 35% of total energy. This target is generally met by the UK population, but further displacement of fat with fibre may have the potential to reduce total energy intake with consequences for body composition. We observed significant reductions in projected total fat intake across our model scenarios, with the exception of the nut and tofu-based alternatives. The greatest projected reduction in fat was associated with mycoprotein; if we assumed no dietary adjustment for total energy, daily replacement of meat with mycoprotein might be projected to reduce fat intake by the

equivalent of 54 kcals per day, a level of reduction more than sufficient to offset weight gain (39).

The UK guidelines set an upper limit of 11% of total energy for saturated fat due to its association with cardiovascular disease (33, 37). Actual intakes of saturated fat remain above this threshold despite long-running public health initiatives. We observed that a 100% replacement of meat with meat-alternatives would take total projected saturated fat intake very close to the 11% of food energy target. This observation was consistent across the categories of meat alternative, but with the greatest projected reductions in saturated fat being observed with the nut-based and tofu-based products categories. Nut and tofu-based meat alternatives may be of further interest as the projected displacement of saturated fat was with mono-unsaturated fatty acids more so than carbohydrate (40-42). Swapping SFA, for MUFAs and PUFAs is considered optimal for heart health (43).

The RNI for protein is greatly exceeded by most UK consumers pursuing an omnivorous, energy balanced diet. However, in our models, the projected reduction in total protein intake from a 100% replacement of meat was equivalent to almost 20 grams per day if alternative foods were drawn exclusively from the vegetable- or nut-based product categories. These losses have the potential to compromise protein status in athletes, in the very young, in the aged and in certain disease states where requirements may be above the RNI. Where protein intake is a concern, we would recommend choosing products from the higher protein, tofu-, soy- and mycoprotein-based meat-alternative categories.

Dietary patterns high in fibre are associated with improved satiety, reduced risk of obesity, improved metabolic health, and reduced risk of diverticular disease, cancer, cardiovascular disease and hypertension (44, 45); the UK dietary guidelines recommend that adults consume upwards of 30g of AOAC fibre per day (33, 37). Despite consistent public health campaigns over many years, dietary fibre consumption amongst the UK public remains significantly below this recommendation (46). Using a refined NDNS cohort, incorporating meat eaters only, we noted that current mean fibre intake is 21.86g day⁻¹. In our models, under every meat-replacement scenario, projected fibre intake was shown to increase significantly compared to current intake for the total population. Importantly, the models showing a 100% replacement of meat with meat alternatives would increase the mean projected fibre intake, at a population level, to 30 g day⁻¹ (**Table 4**).

We noted considerable differences in fibre content between, and within, our meat alternative

product categories, with the highest projected increase in fibre intake being associated with vegetable, nut, and mycoprotein based products. Not all fibre elicits the same physiological benefits (47), and therefore it is important to note that fibre composition, and functional activity, is better described for soy-based (48) and mycoprotein-based (49) meat alternatives than it is for the nut, vegetable and tofu categories of product.

Non-milk extrinsic sugar intake in the UK remains well above current guidelines which suggest limiting intake to no higher than 5% of total calories. In our meat eating NDNS cohort, sugar intake was closer to 20% of total calories. A high intake of sugars may be associated with obesity and poor dental health outcomes (50), therefore any intervention that raises sugar consumption must be viewed as problematic. We noted particularly high sugar levels in the nut-based and vegetable-based product categories and in the 100% meat displacement scenarios involving these two product categories, projected sugar intake rose to almost 22% of calories.

Due to its associations with hypertension, kidney disease and some cancers, dietary sodium reduction has been the goal of a multipronged and effective public health campaign over the last two decades (51). Intakes in the UK have decreased by clinically meaningful levels in recent years; in the meat eating NDNS cohort used in this analysis, mean sodium intake was reported as being very close to the current guidelines. Unfortunately, many of the meat alternative products were found to be high in sodium, and in the 100% meat replacement model, total projected sodium intake was increased by 0.23 grams. As with the other nutrients, there was considerable variation in the sodium content of foods from within and across alternative product categories; of note, soy-based meat alternatives were found to be particularly high in sodium, and a scenario with 100% meat displacement for soy-based products was projected to raise sodium intake by almost 0.5 grams day⁻¹. We caution that our displacement models can't fully account for salt added at the table or in the preparation of meat dishes.

Iron intake amongst girls and women in the UK is marginal with the mean intakes for both girls aged 11 to 18 years, and women aged 19 to 64 years below the RNI (56% and 76% of the RNI respectively), and further, 49% of girls aged 11 to 18 years and 25% of women aged 19 to 64 years have iron intakes below the lower RNI (52). There is also evidence of both iron-deficiency anaemia (as indicated by low haemoglobin levels) and low iron stores (plasma ferritin) in 9% of older girls, 5% of adult women and 2% of older women (52).

Replacing meat with meat alternative products did not significantly reduce projected total iron intake; in fact, replacement with either tofu, or with iron-fortified meat alternatives, was associated with a significant increase in iron intake. Therefore, if meat alternatives are integrated into the dietary habits of the UK population, fortification of these foods with iron could facilitate widespread attainment of RNI's.

In the UK population, vitamin B12 deficiency is rarely due to poor dietary intake, but it can be induced by a poorly managed vegan diet (54), and thus B12 intake should be considered in any public health approaches to reducing animal product intake. In our model scenarios we observed slight reductions in B12, approaching statistical significance with a total replacement of meat, and particularly evident when the replacement products were from the nut and tofu product categories. This reduction in projected B12 intake was largely offset in the model with fortified alternatives. In our models, the projected intakes of B12 following meat replacement remained significantly above the RNI, and therefore might be deemed of limited public health consequence. However, we caution that we did not remove milk, dairy, or eggs from our models, and we would expect further reductions in B12 under those scenarios. For individuals pursuing a vegan diet, it remains sensible to choose fortified products and to consider B12 supplementation.

To our knowledge, this is the first study to consider the projected impact of this important, relatively novel, food category at a population level in the UK. The strengths of this work are that we leverage the National Dietary Nutrition Survey dataset; this is a very well developed and informative record of food habits in the United Kingdom. We have also captured a broad picture of the nutritional quality of meat alternative products from within the UK market in 2020.

The limitations of this study are that; we are reliant upon self-reported dietary intake data from the NDNS. The 4 day food record collection tool used in the NDNS can change habitual diet, and the heavy respondent work load may precipitate socioeconomic representation bias in the cohort. We have also made subjective decisions in approaching the analysis; most notably, we chose to keep the energy content of the substitution model's constant by increasing the nutritional intake uniformly across dataset after the substitution. The energy adjustment approach is consistent with previous literature evaluating the impacts of dietary substitutions (55, 56). However, whilst we assume that individuals will maintain energy balance in a meat replacement scenario, but we can't be certain which foods they are likely to

adjust their intake of, and this could affect overall projected nutritional intake. We were further limited by the depth of nutritional data and relatively poor characterisation of many of the meat alternatives. Across the product categories, mycoprotein and soy-based products were generally well characterised, but there was a paucity of B vitamin and mineral data for many of the product categories, which has been noted previously (35). This meant that we were limited in the nutrients we could assess. Data pertaining to the nutritional profile of meat alternatives within the McCance and Widdowson dataset is scarce and there is a need for greater analytical work to be done to update and formulate databases for these foods. If meat alternatives are to be regularly consumed as part of the diet in place of meat, publication of a more comprehensive characterisation their nutritional composition would be welcome.

There are implications arising from this work for consumers, producers, and food policy makers (**Table 6**). For food consumers, it is clear there are many nutritious approaches to reducing meat intake using meat-alternatives. We note, the variable density of different nutrients across meat substitute categories as summarised in **Table 7**, none of these substitutes fully replicates the nutritional composition of meat. If choosing to use meat alternatives, we therefore recommend using a variety of products from across the meat-substitute categories. We also recommend choosing products high in protein, high in fibre, and low in saturated fat and sugar; where possible, we would advise choosing products that are good sources of iron and vitamin B12. Producers of meat alternatives could better serve consumers with stronger nutritional labelling and by focussing on sodium and sugar reduction. Producers might also consider voluntary fortification with iron and B12. Food policy makers might consider regulation around fortification, marketing and labelling to better guide consumer choice.

Conclusion.

Reducing meat consumption will be a focus of public health and ecological food policy for the foreseeable future. It is therefore anticipated that meat alternatives will play an increasingly prominent role in the UK food plate. Consumer confidence in the new and dynamic meat alternative market will be an important aspect in facilitating the level of dietary change required to protect planetary and consumer health. Going forward, periodic monitoring of the nutritional quality and healthfulness of meat alternatives will be necessary to ensure these foods compare well on a nutritional playing field, and that nutrient analytical

databases are kept up to date with this information. This will then aid better quality research and the compilation of more accurate recommendations that relate to public health policy.

Acknowledgements

We would like to thank Dr Emma Derbyshire for kindly providing a constructive proof-reading of this manuscript.

Financial Support

This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

Conflict of Interest

D.N.F is supported via a PhD studentship part funded by Marlow Foods Ltd.

D.M.C and J.M have received research funding from Marlow Foods Ltd.

This work has been conducted independently of Marlow foods and the authors have no financial or other vested interest in the outcome of the work.

Authorship

The authors' responsibilities were as follows: D.N.F was responsible for study conception, data analyses and drafting the manuscript. D.M.C aided in study conception and drafting the manuscript. D.U aided in data analyses and J. M. contributed to drafting and revising the manuscript.

References

1. Xu T. Analysis on the Meat Consumption, Dietary Transformation and Cultured Meat. *IOP Conference Series: Earth and Environmental Science*. 2020;512:012079.
2. Vermeulen SJ, Campbell BM, Ingram JSI. Climate Change and Food Systems. *Annual Review of Environment and Resources*. 2012;37(1):195-222.
3. Diaz RJ, Rosenberg R. Spreading Dead Zones and Consequences for Marine Ecosystems. *Science*. 2008;321(5891):926-9.
4. Ramankutty N, Evan AT, Monfreda C, Foley JA. Farming the planet: 1. Geographic distribution of global agricultural lands in the year 2000. *Global Biogeochemical Cycles*. 2008;22(1).
5. Shiklomanov I, Rodda J. World Water Resources at the Beginning of the Twenty-First Century. 2004;13.
6. Micha R, Michas G, Mozaffarian D. Unprocessed red and processed meats and risk of coronary artery disease and type 2 diabetes--an updated review of the evidence. *Curr Atheroscler Rep*. 2012;14(6):515-24.
7. Pan A, Sun Q, Bernstein AM, Manson JE, Willett WC, Hu FB. Changes in red meat consumption and subsequent risk of type 2 diabetes mellitus: three cohorts of US men and women. *JAMA Intern Med*. 2013;173(14):1328-35.
8. Zhong VW, Van Horn L, Greenland P, Carnethon MR, Ning H, Wilkins JT, et al. Associations of Processed Meat, Unprocessed Red Meat, Poultry, or Fish Intake With Incident Cardiovascular Disease and All-Cause Mortality. *JAMA Intern Med*. 2020.
9. Alexander DD, Mink PJ, Cushing CA, Scurman B. A review and meta-analysis of prospective studies of red and processed meat intake and prostate cancer. *Nutr J*. 2010;9:50-.
10. Bernstein AM, Song M, Zhang X, Pan A, Wang M, Fuchs CS, et al. Processed and unprocessed red meat and risk of colorectal cancer: analysis by tumor location and modification by time. *PloS one*. 2015;10(8):e0135959.
11. Bouvard V, Loomis D, Guyton KZ, Grosse Y, Ghissassi FE, Benbrahim-Tallaa L, et al. Carcinogenicity of consumption of red and processed meat. *The Lancet Oncology*. 2015;16(16):1599-600.
12. Cho E, Chen WY, Hunter DJ, Stampfer MJ, Colditz GA, Hankinson SE, et al. Red Meat Intake and Risk of Breast Cancer Among Premenopausal Women. *Archives of Internal Medicine*. 2006;166(20):2253-9.

13. Cross AJ, Leitzmann MF, Gail MH, Hollenbeck AR, Schatzkin A, Sinha R. A prospective study of red and processed meat intake in relation to cancer risk. *PLoS medicine*. 2007;4(12):e325.
14. Genkinger JM, Koushik A. Meat consumption and cancer risk. *PLoS Med*. 2007;4(12):e345-e.
15. Inoue-Choi M, Sinha R, Gierach GL, Ward MH. Red and processed meat, nitrite, and heme iron intakes and postmenopausal breast cancer risk in the NIH-AARP Diet and Health Study. *Int J Cancer*. 2016;138(7):1609-18.
16. Key TJ, Bradbury KE, Murphy N. Diet and colorectal cancer in UK Biobank: a prospective study. 2019.
17. Sanchez-Sabate R, Sabaté J. Consumer Attitudes Towards Environmental Concerns of Meat Consumption: A Systematic Review. *International Journal of Environmental Research and Public Health*. 2019;16(7):1220.
18. Kumar P, Chatli MK, Mehta N, Singh P, Malav OP, Verma AK. Meat analogues: Health promising sustainable meat substitutes. *Critical Reviews in Food Science and Nutrition*. 2017;57(5):923-32.
19. Apostolidis C, McLeay F. Should we stop meating like this? Reducing meat consumption through substitution. *Food Policy*. 2016;65:74-89.
20. Dagevos H, Voordouw J. Sustainability and meat consumption: is reduction realistic? *Sustainability: Science, Practice and Policy*. 2013;9(2):60-9.
21. Schösler H, Boer Jd, Boersema JJ. Can we cut out the meat of the dish? Constructing consumer-oriented pathways towards meat substitution. *Appetite*. 2012;58(1):39-47.
22. Hu FB, Otis BO, McCarthy G. Can Plant-Based Meat Alternatives Be Part of a Healthy and Sustainable Diet? *JAMA*. 2019;322(16):1547-8.
23. Hemler EC, Hu FB. Plant-Based Diets for Cardiovascular Disease Prevention: All Plant Foods Are Not Created Equal. *Curr Atheroscler Rep*. 2019;21(5):18.
24. NatCen Social Research; MRC Elsie Widdowson Laboratory, inventor National Diet and Nutrition Survey Years 1-9, 2008/09-2016/172019.
25. Bates B, Cox L, Nicholson S, Page P, Prentice A, Steer T, et al. National Diet and Nutrition Survey results from years 5 and 6 (combined) of the Rolling Programme (2012/2013–2013/2014). London: Public Health England. 2016.
26. Bates B, Lennox A, Prentice A, Bates CJ, Page P, Nicholson S, et al. National diet and nutrition survey: Results from years 1, 2, 3 and 4 (combined) of the rolling programme

(2008/2009-2011/2012): A survey carried out on behalf of public health England and the food standards agency: Public Health England; 2014.

27. Roberts C, Steer T, Maplethorpe N, Cox L, Meadows S, Nicholson S, et al. National Diet and nutrition survey: results from years 7 and 8 (combined) of the rolling Programme (2014/2015–2015/2016). 2018.
28. Macdiarmid J, Blundell J. Assessing dietary intake: Who, what and why of under-reporting. *Nutr Res Rev.* 1998;11(2):231-53.
29. Poslusna K, Ruprich J, de Vries JHM, Jakubikova M, van't Veer P. Misreporting of energy and micronutrient intake estimated by food records and 24 hour recalls, control and adjustment methods in practice. *British Journal of Nutrition.* 2009;101(S2):S73-S85.
30. McCance RA, Widdowson EM. McCance and Widdowson's the Composition of Foods: Royal Society of Chemistry; 2014.
31. Mintel Group Ltd. Executive Summary Meat-Free Foods. 2019.
32. Department of Health. Dietary Reference Values for Food Energy and Nutrients for the United Kingdom: Report of the Panel on Dietary Reference Values of the Committee on Medical Aspects of Food Policy: Reports of Health and Social Subjects; 1991.
33. Public Health England. Government Dietary Recommendations. In: Health Do, editor. London, Public Health England: PHE publications 2016.
34. Gibson R, Eriksen R, Chambers E, Gao H, Aresu M, Heard A, et al. Intakes and Food Sources of Dietary Fibre and Their Associations with Measures of Body Composition and Inflammation in UK Adults: Cross-Sectional Analysis of the Airwave Health Monitoring Study. *Nutrients.* 2019;11(8):1839.
35. Curtain F, Grafenauer S. Plant-Based Meat Substitutes in the Flexitarian Age: An Audit of Products on Supermarket Shelves. *Nutrients.* 2019;11(11).
36. Craig WJ. Health effects of vegan diets. *The American Journal of Clinical Nutrition.* 2009;89(5):1627S-33S.
37. Scientific Advisory Committee on Nutrition. Dietary Reference Values for Energy. In: Scientific Advisory Committee on Nutrition, editor. London 2011.
38. R Core Team. R: A language and environment for statistical computing.: R Foundation for Statistical Computing; 2020.
39. Sacks FM, Bray GA, Carey VJ, Smith SR, Ryan DH, Anton SD, et al. Comparison of weight-loss diets with different compositions of fat, protein, and carbohydrates. *N Engl J Med.* 2009;360(9):859-73.

40. Rizzo G, Baroni L. Soy, Soy Foods and Their Role in Vegetarian Diets. *Nutrients*. 2018;10(1):43.
41. Ros E. Health benefits of nut consumption. *Nutrients*. 2010;2(7):652-82.
42. Ros E, Mataix J. Fatty acid composition of nuts--implications for cardiovascular health. *Br J Nutr*. 2006;96 Suppl 2:S29-35.
43. Nettleton JA, Brouwer IA, Geleijnse JM, Hornstra G. Saturated Fat Consumption and Risk of Coronary Heart Disease and Ischemic Stroke: A Science Update. *Annals of Nutrition and Metabolism*. 2017;70(1):26-33.
44. Slavin J. Fiber and prebiotics: mechanisms and health benefits. *Nutrients*. 2013;5(4):1417-35.
45. Veronese N, Solmi M, Caruso MG, Giannelli G, Osella AR, Evangelou E, et al. Dietary fiber and health outcomes: an umbrella review of systematic reviews and meta-analyses. *Am J Clin Nutr*. 2018;107(3):436-44.
46. Stephen AM, Champ MMJ, Cloran SJ, Fleith M, van Lieshout L, Mejbourn H, et al. Dietary fibre in Europe: current state of knowledge on definitions, sources, recommendations, intakes and relationships to health. *Nutrition Research Reviews*. 2017;30(2):149-90.
47. Gibson GR, Hutkins R, Sanders ME, Prescott SL, Reimer RA, Salminen SJ, et al. Expert consensus document: The International Scientific Association for Probiotics and Prebiotics (ISAPP) consensus statement on the definition and scope of prebiotics. *Nature Reviews Gastroenterology & Hepatology*. 2017;14(8):491-502.
48. Yang L, Zhao Y, Huang J, Zhang H, Lin Q, Han L, et al. Insoluble dietary fiber from soy hulls regulates the gut microbiota in vitro and increases the abundance of bifidobacteriales and lactobacillales. *J Food Sci Technol*. 2020;57(1):152-62.
49. Harris HC, Edwards CA, Morrison DJ. Short Chain Fatty Acid Production from Mycoprotein and Mycoprotein Fibre in an In Vitro Fermentation Model. *Nutrients*. 2019;11(4).
50. Campbell R, Tasevska N, Jackson KG, Sagi-Kiss V, di Paolo N, Mindell JS, et al. Association between urinary biomarkers of total sugars intake and measures of obesity in a cross-sectional study. *PLOS ONE*. 2017;12(7):e0179508.
51. Mente A, O'Donnell M, Rangarajan S, McQueen M, Dagenais G, Wielgosz A, et al. Urinary sodium excretion, blood pressure, cardiovascular disease, and mortality: a community-level prospective epidemiological cohort study. *Lancet*. 2018;392(10146):496-506.

52. Public Health England. National Diet and Nutrition Survey (NDNS): results from years 9 to 11 (combined) – statistical summary. Published 11 December 2020.
53. Lopez A, Cacoub P, Macdougall IC, Peyrin-Biroulet L. Iron deficiency anaemia. *The Lancet*. 2016;387(10021):907-16.
54. Lederer A-K, Hannibal L, Hettich M, Behringer S, Spiekerkoetter U, Steinborn C, et al. Vitamin B12 Status Upon Short-Term Intervention with a Vegan Diet-A Randomized Controlled Trial in Healthy Participants. *Nutrients*. 2019;11(11):2815.
55. Mertens E, Biesbroek S, Dofková M, Mistura L, D'Addezio L, Turrini A, et al. Potential Impact of Meat Replacers on Nutrient Quality and Greenhouse Gas Emissions of Diets in Four European Countries. *Sustainability*. 2020;12(17):6838.
56. Springmann M, Wiebe K, Mason-D'Croz D, Sulser TB, Rayner M, Scarborough P. Health and nutritional aspects of sustainable diet strategies and their association with environmental impacts: a global modelling analysis with country-level detail. *The Lancet Planetary Health*. 2018;2(10):e451-e61.

Tables

Table 1. NDNS food groups used to calculate self-reported meat intake

Meat subtype	NDNS category*	Description of category
Red meat	"beefg"	Amount of beef consumed (g)
	"lambg"	Amount of lamb consumed (g)
	"porkg"	Amount of pork consumed (g)
	"offalg"	Amount of offal consumed (g)
	"liverdishes"	Amount of liver consumed (g)
	"otherredmeatg"	Amount of other red meat consumed (g)
Processed red meat	"burgersg"	Amount of burger consumed (g)
	"sausagesg"	Amount of sausages consumed (g)
	"baconandham"	Amount of bacon and/or ham consumed (g)
	"burgersandkebabs"	Amount of burger and/or kebab consumed (g)
Total Meat	"processedredmeatg"	Amount of other processed red meat consumed (g)
White meat	"poultryg"	Amount of poultry consumed (g)
	"gamebirdsg"	Amount of game consumed (g)
Processed white meat	"coatedchicken"	Amount of coated chicken products consumed (g)
	"processedpoultryg"	Amount of processed poultry consumed (g)
Seafood	"whitefishg"	Amount of white fish consumed (g)
	"oilyfishg"	Amount of oily fish consumed (g)
	"cannedtunag"	Amount of canned tuna consumed (g)
	"shellfishg"	Amount of shellfish consumed (g)
	"whitefishcoatedorfried"	Amount of white fish (coated or fried) consumed (g)

*As displayed in NDNS data set.

Table 2. Meat replacement models implemented in the present modelling analysis

Replacement Scenario Models	Description of model	
Progressive (Model 1)	MA-25	Partial replacement substituting 25% of the current self-reported meat intake with a composite of meat alternatives.
	MA-50	Partial replacement substituting 50% of the current self-reported meat intake with a composite of meat alternatives.
	MA-75	Partial replacement substituting 75% of the current self-reported meat intake with a composite of meat alternatives.
	MA-100	Full replacement substituting 100% of the current self-reported meat intake with a composite of meat alternatives.
Ingredient (Model 2)	Vegetable	Full replacement substituting 100% of the current self-reported meat intake with meat alternatives produced predominantly from vegetables.
	Mycoprotein	Full replacement substituting 100% of the current self-reported meat intake with meat alternatives produced predominantly from mycoprotein.
	Legume	Full replacement substituting 100% of the current self-reported meat intake with meat alternatives produced predominantly from bean and pea.
	Tofu	Full replacement substituting 100% of the current self-reported meat intake with meat alternatives produced from tofu.
	Nut	Full replacement substituting 100% of the current self-reported meat intake with meat alternatives produced predominantly from nut.
Fortification (Model 3) *	Soy	Full replacement substituting 100% of the current self-reported meat intake with meat alternatives produced predominantly from soy.
	Fortified	Full replacement substituting 100% of the current self-reported meat intake with meat alternatives that have been fortified with nutrients.
	Unfortified	Full replacement substituting 100% of the current self-reported meat intake with meat alternatives that have not been fortified with nutrients.

*Products identified as fortified if 1) Explicitly stated nutrient fortification had been applied during manufacturing process; 2) Isolated nutrients included in ingredient list. Products not meeting 1) or 2) identified as unfortified.

Table 3. Self-reported meat intake (total meat, and meat subtype) stratified by age and gender.

Subgroup	Total meat, g/day	Red meat, g/day	Contribution to total meat, %	Processed meat, g/day	Contribution to total meat, %	White meat, g/day	Contribution to total meat, %	Processed white meat, g/day	Contribution to total meat, %	Seafood, g/day	Contribution to total meat, %
Male, 4 -10yr	106.01 (96.61, 115.42)	14.76 (11.74, 17.77)	13.92 (12.16, 15.39)	47.66 (41.19, 54.13)	44.95 (42.63, 46.89)	26.27 (22.30, 30.24)	24.78 (23.08, 26.20)	6.41 (4.88, 7.95)	6.05 (5.05, 6.89)	10.92 (8.64, 13.20)	10.30 (8.95, 11.43)
Female, 4 -10yr	98.75 (91.28, 106.23)	13.72 (10.92, 16.51)	13.89 (11.97, 15.54)	40.97 (34.84, 47.11)	41.49 (38.17, 44.35)	24.47 (20.88, 28.07)	24.78 (22.87, 26.42)	6.88 (5.26, 8.49)	6.96 (5.77, 7.99)	12.71 (10.20, 15.22)	12.87 (11.18, 14.33)
Male, 11 -18yr	146.71 (134.32, 159.10)	21.4 (16.47, 26.32)	14.59 (12.26, 16.54)	60.37 (50.55, 70.18)	41.15 (37.63, 44.11)	40.79 (33.57, 48.01)	27.8 (24.99, 30.17)	9.14 (5.71, 12.57)	6.23 (4.25, 7.90)	15.01 (10.57, 19.46)	10.23 (7.87, 12.23)
Female, 11 -18yr	131.36 (117.86, 144.87)	19.87 (15.51, 24.23)	15.12 (13.16, 16.72)	50.21 (40.47, 59.94)	38.22 (34.34, 41.38)	39.70 (34.19, 45.21)	30.22 (29.01, 31.21)	9.41 (6.75, 12.07)	7.16 (5.73, 8.33)	12.18 (9.25, 15.10)	9.27 (7.85, 10.42)
Male, 19 -64yr	177.55 (165.3)	40.07 (34.21)	22.57 (20.69)	62.46 (53.53)	35.18 (32.38)	50.05 (43.67)	28.19 (26.41)	5.04 (3.34, 6.74)	2.84 (2.02)	19.92 (16.09)	11.22 (9.73)

Accepted manuscript

	3, 189.77)	45.93)	24.20)	71.40)	37.62)	, 56.44)	29.74)	3.55)	23.75)	12.51)	
Female, 19 -64yr	127.84 (119.6 3, 136.06)	26.33 (22.44, 30.21)	20.59 (18.76, 22.20)	38.94 (33.29, 44.59)	30.46 (27.83, 32.77)	39.41 (34.79 , 44.02)	30.83 (29.09, 32.36)	3.31 (2.16, 4.46)	2.59 (1.81, 3.28)	19.86 (16.76, 22.95)	15.53 (14.01, 16.87)
Male, 65yr +	151.01 (133.5 3, 168.49)	42.22 (33.65, 50.80)	27.96 (25.20, 30.15)	52.64 (40.51, 64.77)	34.86 (30.34, 38.44)	29.01 (19.20 , 38.82)	19.21 (14.38, 23.04)	1.18 (0.04, 2.41)	0.78 (0.03, 1.43)	25.95 (20.12, 31.78)	17.18 (15.07, 18.86)
Female, 65yr +	103.97 (94.29, 113.66)	26.61 (20.68, 32.54)	25.60 (21.94, 28.63)	27.29 (21.29, 33.30)	26.25 (22.58, 29.30)	21.98 (16.59 , 27.36)	21.14 (17.59, 24.08)	0.80 (0.09, 1.51)	0.77 (0.10, 1.33)	27.29 (21.66, 32.92)	26.25 (22.97, 28.96)
Total Populatio n	132.25 (128.0 2, 136.48)	25.72 (23.88, 27.55)	19.44 (18.65, 20.19)	47.58 (44.68, 50.49)	35.98 (34.90, 36.99)	36.00 (33.88 , 38.12)	27.22 (26.46, 27.93)	5.28 (4.61, 5.95)	3.99 (3.60, 4.36)	17.67 (16.31, 19.04)	13.36 (12.74, 13.95)

Data expressed as means with 95% confidence intervals in parentheses

Table 4. Mean nutrient intake of total population (n = 1110) for current and replacement models, with reference to DRVs

Nutrient	DRV*	Current	Progressive Model				Ingredient Model						Fortification Model	
			MA-25	MA-50	MA-75	MA-100	Vegetable	Mycoprotein	Legume	Tofu	Nut	Soy	Fortified	Unfortified
Energy, kcal/day	2150	-	-	-	-	-	-	-	-	-	-	-	-	-
Carbohydrate, g/day†	268.80	270.77	275.16	279.78	284.65	289.80	295.39	293.17	290.68	279.77	282.24	285.33	282.01	294.03
Protein, g/day†	80.60	86.67	83.27	79.68	75.90	71.90	67.34	77.86	71.24	81.80	66.88	79.48	83.58	68.49
Fat, g/day	83.60	81.97	81.19	80.38	79.51	78.60	77.58	75.85	78.48	79.60	83.35	77.26	77.08	77.78
Fibre, g/day	30	21.86	23.74	25.71	27.80	30.00	30.59	30.57	28.66	25.00	32.10	28.12	28.94	30.23
Sugars, g/day	90	110.52	111.48	112.49	113.56	114.68	114.94	113.30	113.93	112.66	117.96	112.71	112.20	114.63
Saturated fat, g/day	26.30	30.96	30.05	29.08	28.07	27.00	27.13	27.31	27.23	26.63	26.16	27.27	27.66	27.02
Sodium, mg/day	2400	2413.80	2466.37	2521.74	2580.14	2641.82	2720.90	2726.12	2690.65	2510.20	2289.49	2909.77	2896.42	2774.88
Iron, mg/day	8.7/14.8‡	13.65	13.59	13.52	13.45	13.38	13.17	12.68	13.26	21.28	12.74	13.61	16.77	13.10
B12, ug/day	1.5	7.47	7.07	6.65	6.21	5.74	5.72	5.98	5.77	5.45	5.59	5.71	7.14	5.68

*DRV values correspond to EAR for energy; RNI for Protein, Sodium, Iron, B12; DRV for Carbohydrate, Fat, Fibre, Sugars, Saturated fat.

† Protein target elected as 15% total energy intake and carbohydrate as 50% total energy intake in the current analysis

‡ Iron RNI for females aged 11-50

Colours: Green, meeting DRV. Red, not meeting DRV. Grey, not meeting 15% protein and/or 50% carbohydrate target elected in the current analysis. Blue, not meeting higher RNI for iron.

DRV, Dietary Reference Value; EAR, Estimated Average Requirement; RNI, Reference Nutrient Intake; MA-25, Replacement model substituting 25% of self-reported meat intake with composite of meat alternatives; MA-50, Replacement model substituting 50% of self-reported meat intake with composite of meat alternatives; MA-75, Replacement model substituting 75% of self-reported meat intake with composite of meat alternatives; MA-100, Replacement model substituting 100% of self-reported meat intake with composite of meat alternatives.

Table 5. Projected differences from current intake for meat alternatives across the total population (n = 1110)

Progressive Model													
	Current	MA-25	P	MA-50	P	MA-75	P	MA-100	P				
Energy, kcal/day	2150	-	-	-	-	-	-	-	-				
Carbohydrate, g/day	270.77	+4.39	0.25	+9.01	0.018*	+13.89	<0.001*	+19.04	<0.001*				
Protein, g/day	86.67	-3.40	0.002*	-6.99	<0.001*	-10.77	<0.001*	-14.77	<0.001*				
Fat, g/day	81.97	-0.78	0.55	-1.60	0.22	-2.46	0.06	-3.37	0.009*				
Fibre, g/day	21.86	+1.88	<0.001*	+3.85	<0.001*	+5.93	<0.001*	+8.13	<0.001*				
Sugars, g/day	110.52	+0.96	0.66	+1.97	0.36	+3.04	0.16	+4.16	0.05*				
Saturated fat, g/day	30.96	-0.91	0.09	-1.88	0.001*	-2.89	<0.001*	-3.96	<0.001*				
Sodium, mg/day	2413.80	+52.57	0.21	+107.94	0.01	+166.34	<0.001*	+228.03	<0.001*				
Iron, mg/day	13.65	-0.06	0.93	-0.13	0.85	-0.20	0.78	-0.27	0.70				
B12, ug/day	7.47	-0.40	0.66	-0.82	0.37	-1.26	0.17	-1.73	0.06				
Ingredient Model													
	Current	Vegetable	P	Mycoprotein	P	Legume	P	Tofu	P	Nut	P	Soy	P
Energy,	2150			-	-	-	-					-	-

kcal/day													
Carbohydrate, g/day	270.77	+24.63	<0.001*	+22.40	<0.001*	+19.92	<0.001*	+9.00	0.018*	+11.48	0.003*	+14.56	<0.001*
Protein, g/day	86.67	-19.33	<0.001*	-8.82	<0.001*	-15.43	<0.001*	-4.87	<0.001*	-19.79	<0.001*	-7.19	<0.001*
Fat, g/day	81.97	-4.39	0.001*	-6.12	<0.001*	-3.49	0.007*	-2.37	0.07	+1.38	0.29	-4.71	<0.001*
Fibre, g/day	21.86	+8.73	<0.001*	+8.71	<0.001*	+6.79	<0.001*	+3.14	<0.001*	+10.23	<0.001*	+6.25	<0.001*
Sugars, g/day	110.52	+4.43	0.04*	+2.78	0.20	+3.42	0.11	+2.15	0.32	+7.44	0.001*	+2.20	0.31
Saturated fat, g/day	30.96	-3.83	<0.001*	-3.65	<0.001*	-3.73	<0.001*	-4.33	<0.001*	-4.80	<0.001*	-3.69	<0.001*
Sodium, mg/day	2413.80	+307.10	<0.001*	+312.32	<0.001*	+276.85	<0.001*	+96.40	0.02*	-124.30	0.003*	+495.98	<0.001*
Iron, mg/day	13.65	-0.48	0.49	-0.97	0.16	-0.39	0.57	+7.63	<0.001*	-0.91	0.19	-0.04	0.95
B12, ug/day	7.47	-1.75	0.06	-1.49	0.10	-1.70	0.06	-2.02	0.03*	-1.88	0.04*	-1.76	0.05*
Fortification Model													
	Current	Fortified	P	Unfortified	P								
Energy, kcal/day	2150	-	-	-	-								
Carbohydrate, g/day	270.77	+11.24	0.003*	+23.26	<0.001*								
Protein, g/day	86.67	-3.09	0.005*	-18.19	<0.001								

Fat, g/day	81.97	-4.90	<0.001*	-4.19	0.001*
Fibre, g/day	21.86	+7.08	<0.001*	+8.37	<0.001*
Sugars, g/day	110.52	+1.68	0.43	+4.12	0.06
Saturated fat, g/day	30.96	-3.30	<0.001*	-3.94	<0.001*
Sodium, mg/day	2413.80	+482.63	<0.001*	+361.09	<0.001*
Iron, mg/day	13.65	+3.12	<0.001*	-0.54	0.43
B12, ug/day	7.47	-0.33	0.72	-1.79	0.05*

Blue indicates significant increase compared to Current. Gold indicates significant decrease compared to Current.

*Differences compared using regression models with significance threshold $P < 0.05$

MA-25, Replacement model substituting 25% of self-reported meat intake with composite of meat alternatives; MA-50, Replacement model substituting 50% of self-reported meat intake with composite of meat alternatives; MA-75, Replacement model substituting 75% of self-reported meat intake with composite of meat alternatives; MA-100, Replacement model substituting 100% of self-reported meat intake with composite of meat alternatives.

Table 6. Implications and recommendations

i/ <u>Food consumers</u>	Consumers choosing to reduce their meat intake should understand that meat alternative products are a culinary replacement for meat, which differ in their nutritional profile. With the exception of tofu and mycoprotein, we caution that few meat alternative products have been well evaluated for their physiological effects in the consumer. We would therefore strongly recommend selecting meat alternatives based on the level of nutritional characterisation and the availability of high-quality evidence about their healthfulness. It is also important to consider the wider balance of the diet to ensure that the intake of nutrients found in meat are not compromised when reducing intake.
ii/ <u>Producers</u>	Producers of meat alternatives should be encouraged to focus on sodium reduction, deeper nutritional characterisation, and where feasible, investigations evidencing the likely health benefits of their products. Given their market, voluntary product fortification with iron and B12 might also be considered.
iii/ <u>Policy makers</u>	Meat alternatives are intended to displace a nutritionally dense component of the diet, better labelling and nutritional characterisation would be in the interest of the food consumer. Further, in 1960 policy makers in the UK responded to the displacement of butter and animal fats from the diet with legislation instructing the mandatory fortification of margarines with vitamins A and D, there may soon be a need for a discussion about fortification, and or the regulation of nutritional quality and the healthfulness of this category of food; this discussion would be better informed by appropriate dietary intervention studies.

Table 7. Nutritional comparisons across meat substitute product category.

Ingredient Model	Carbohydrate g/day	Protein g/day	Fat g/day	Fiber g/day	Sugars g/day	Saturated fat g/day	Sodium mg/day	Iron mg/day	B12 ug/day
Vegetable	↑	↓	↓	↑	↑	↓	↑	-	-
Mycoprotein	↑	↓	↓	↑	+	↓	↑	-	-
Legume	↑	↓	↓	↑	+	↓	↑	-	-
Tofu	↑	↓	-	↑	+	↓	↑	↑	↓
Nut	↑	↓	+	↑	↑	↓	↓	-	↓
Soy	↑	↓	↓	↑	+	↓	↑	-	↓

+, non-significant increase compared to current intake; -, non-significant decrease from current intake. ↑, significant increase compared to current intake; ↓, significant decrease from current intake. Green indicates best performing model with regards to dietary recommendations for nutrient. Red indicates worst performing model.