

Combined Experimental and Theoretical Study of the Competitive Absorption of CO₂ and NO₂ by a Superbase Ionic Liquid

Adam J. Greer,* S. F. Rebecca Taylor, Helen Daly, Matthew G. Quesne, Nora H. de Leeuw, C. Richard A. Catlow, Johan Jacquemin,* and Christopher Hardacre*



Cite This: *ACS Sustainable Chem. Eng.* 2021, 9, 7578–7586



Read Online

ACCESS |



Metrics & More




Article Recommendations



Supporting Information

ABSTRACT: A superbase ionic liquid (IL), trihexyltetradecylphosphonium benzimidazolide ([P₆₆₆₁₄][Benzim]), is investigated for the capture of CO₂ in the presence of NO₂ impurities. The effect of the waste gas stream contaminant on the ability of the IL to absorb simultaneously CO₂ is demonstrated using novel measurement techniques, including a mass spectrometry breakthrough method and *in situ* infrared spectroscopy. The findings show that the presence of an industrially relevant concentration of NO₂ in a combined feed with CO₂ has the effect of reducing the capacity of the IL to absorb CO₂ efficiently by ~60% after 10 absorption–desorption cycles. This finding is supported by physical property analysis (viscosity, ¹H and ¹³C NMR, and X-ray photoelectron spectroscopy) and spectroscopic infrared characterization, in addition to density functional theory (DFT) calculations, to determine the structure of the IL–NO₂ complex. The results are presented in comparison with another flue gas component, NO, demonstrating that the absorption of NO₂ is more favorable, thereby hindering the ability of the IL to absorb CO₂. Significantly, this work aids understanding of the effects that individual components of flue gas have on CO₂ capture sorbents, through studying a contaminant that has received limited interest previously.

KEYWORDS: ionic liquids, CO₂ capture, NO₂, flue gas, competitive absorption, infrared, DFT



INTRODUCTION

Post-combustion CO₂ capture is an important requirement of many industrial processes. The high-temperature combustion of fossil fuels produces large quantities of CO₂ (10–15 vol %), as well as other impurities, such as SO₂ (0.05–0.2 vol %) and NO_x (0.15–0.25 vol %), which consists predominantly of NO and NO₂.^{1–3} The reported values are obtained before any desulfurization or denitrification technologies. In particular, NO_x is known to have a significant impact on health and the environment, causing the formation of atmospheric ozone and acid rain.⁴ It is therefore vital that NO_x emissions are regulated (40 μg·m⁻³ a year), leading to the fitting of NO_x scrubbers to power stations, comprising oxidizing and reducing agents responsible for the conversion of NO_x to N₂.⁵ Aqueous alkanolamines have been employed as CO₂ capture sorbents, but the presence of NO_x was found to result in the irreversible formation of carcinogenic nitrosamines and a decrease in CO₂ capture efficiency.^{6–8} Ionic liquids (ILs) have also been widely investigated for the capture of CO₂ as a non-volatile alternative to toxic alkanolamines. However, to date, the effect of NO₂ on the ability of an IL to capture CO₂ in a combined feed has not been investigated.


The interest in ILs stems from the ability to alter their physiochemical properties, such as thermal stability or CO₂

absorption capacity, through changing the combination of cation and anion, which enables the tuning of their properties for specific applications.^{9,10} For example, the amount of CO₂ absorbed by a particular IL has been shown to have a strong dependence on the anion, with conventional anions only physically absorbing small quantities of CO₂,^{11,12} compared with task-specific ILs that incorporate amine functionality and chemically absorb up to 1 nCO₂:nIL.^{13,14} Superbase ILs (SBILs) containing an aprotic heterocyclic anion (AHA) were developed to minimize the increase in viscosity observed in amine-functionalized ILs, and they can reversibly capture a greater than equimolar amount of CO₂.^{15–18} Extensive studies into the absorption of other acidic gases such as SO₂ and NO by SBILs have found that irreversible absorption was observed in many cases, often on multiple active sites within the IL, affecting the recyclability of the system.^{19–25}

Received: March 2, 2021

Revised: April 14, 2021

Published: May 26, 2021



The effect of impurities on the CO₂ uptake of the SBIL trihexyltetradecylphosphonium benzimidazolide, [P₆₆₆₁₄][Benzim], has previously been investigated in combined feeds. SO₂ was shown to deactivate the IL through binding to the absorption site available to CO₂, while the presence of NO exhibited little effect on the IL's capacity for the uptake of CO₂.^{26,27} For NO, the co-binding of CO₂ and NO as carbamate and NONO-ate species, respectively, was observed at different N-sites of the benzimidazolide anion. However, competition for the same binding site was reported between CO₂ and SO₂, which markedly influenced the absorption capacity and recyclability of the IL. The differing effects of SO₂ and NO on the absorption of CO₂ by [P₆₆₆₁₄][Benzim] highlight the need to assess the components of flue gas impurities, both individually and in combination with CO₂.

The capture of NO₂ by ILs has been the focus of only a few studies, where it was found that an increased uptake is observed for NO₂ compared to NO.^{28–30} To date, NO₂ has not been studied in a combined feed with CO₂, so its direct influence on the ability of a sorbent to capture CO₂ is unknown. The competitive absorption of CO₂ with industrially relevant concentrations of H₂O, SO₂, or NO, independently, has been investigated previously in [P₆₆₆₁₄][Benzim], and this IL was therefore selected for the current study to gain a comprehensive insight into more complex, multi-component feeds.^{18,26,27} The use of a recently developed analytical method utilizing mass spectrometry allows the study of this superbase IL under realistic and dry flue gas conditions, with a feed containing 14% CO₂ and 0.2% NO₂.²⁶ Further molecular-level information was provided by density functional theory (DFT) calculations and spectroscopic data (NMR, IR, and X-ray photoelectron spectroscopy).

EXPERIMENTAL SECTION

Materials. Trihexyltetradecylphosphonium chloride ([P₆₆₆₁₄]⁺Cl⁻, 97.7 wt %, CAS: 258864-54-9) was procured from IoLiTec, and benzimidazole (98 wt %, CAS: 51-17-2) was purchased from Sigma-Aldrich. [P₆₆₆₁₄][Benzim] was prepared using a two-step synthesis method reported previously.¹⁸ The halide content was determined to be <5 ppm by a silver nitrate test.³¹ The water content was measured to be <0.1 wt % using a Metrohm 787 KF Titrimo Karl Fischer machine. The structure and purity of the IL, after synthesis and post-absorption, were analyzed by ¹H NMR and ¹³C NMR with a Bruker Avance II 400 MHz Ultra shield Plus and carried out as neat ILs in the presence of a glass capillary insert containing a deuterated solvent (DMSO-*d*₆, purchased from Cambridge Isotope Laboratories Inc., CAS: 2206-27-1). Gases were obtained from BOC; argon (99.998%, CAS: 7440-37-1); carbon dioxide (99.99%, CAS: 124-38-9); nitrogen dioxide (1% in argon, CAS: 10102-44-0).

Methods. The gas absorption measurement techniques used in this work were reported in detail previously, and the same protocol was followed in this work.^{26,27} To briefly summarize this, the uptake of a single component gas feed (1% NO₂ in argon) by [P₆₆₆₁₄][Benzim] was studied gravimetrically at 22 ± 0.5 °C. A mass spectrometer-based method was utilized to study the gas phase concentrations at the outlet after the IL was exposed to a mixed gas feed of 14% CO₂ + 0.2% NO₂ in Ar. A series of cycles were performed consisting of a 2 h absorption period under feed conditions at 22 °C, followed by a 2 h desorption period under Ar at 90 °C.

Analysis. The viscosity of the IL samples was measured before and after NO₂ absorption using a TA Instruments AR2000. Elemental analysis was carried out using a Thermo Scientific Flash 2000 elemental analyzer. X-ray photoelectron spectroscopy (XPS) was performed with a Kratos AXIS Ultra DLD apparatus, with a monochromated Al K α radiation X-ray source, charge neutralizer,

and hemispherical electron energy analyzer. During data acquisition, the chamber pressure was kept below 10⁻⁹ mbar. The spectra were analyzed by CasaXPS and corrected for charging using the C 1s feature at 284.8 eV.

Attenuated total reflectance-infrared (ATR-IR) spectra were recorded in a modified *in situ* cell with a ZnSe crystal and a PIKE ATRMax II accessory housed in a Bruker Tensor II infrared spectrometer. A thin film of [P₆₆₆₁₄][Benzim] (~250 mg) coated the ZnSe crystal in the cell before the introduction of the gas feed (14% CO₂ in Ar, 0.2% NO₂ in Ar, or a mixed gas feed of 14% CO₂ with 0.2% NO₂ in Ar) with a flow rate of 15 cm³·min⁻¹ at 22 °C. Desorption was performed at 90 °C under Ar. The background for all spectra was the ZnSe crystal in the cell, and spectra were recorded with 8 scans at 4 cm⁻¹ resolution. The spectrum of the IL before introduction of the gas feed has been subtracted from all the spectra of the IL under gas absorption.

DFT Calculations. DFT calculations followed a similar protocol to previous work on this system, and while a brief overview will be included here, a more detailed description can be found in the literature.²⁷ Calculations were performed with the Gaussian09 software package³² using a combination of the hybrid functional UB3LYP and the triplet- ζ basis set 6-311+G*, as reported in previous works.^{33–35} Starting geometries for [P₃₃₃₃][Benzim] models were also informed by previous molecular dynamical studies,³⁶ with absorbates manually added using the ChemCraft software package.³⁷ Minima structures for all possible reaction mechanisms were fully optimized without constraints with transition states located by initially running geometry scans, where only the degree of freedom connecting two minima was fixed. Full transition state optimizations were subsequently performed on the highest energy structures obtained along each reaction coordinate. The verification of both minima and transition states were carried out with the aid of analytical frequencies at 1 atm and 298.15 K, whereby only positive frequencies were observed for each minima, with each transition state possessing a single imaginary frequency for the mode associated with the reaction coordinate. Corrections for long-range interactions were included with the aid of the Grimme D3 dispersion model,³⁸ while solvent effects were simulated with an implicit model of acetonitrile ($\epsilon = 35.688$) using a polarizable continuum model (PCM).

RESULTS AND DISCUSSION

The absorption of 1% NO₂ in Ar by [P₆₆₆₁₄][Benzim] was initially examined by a gravimetric technique to allow direct comparison with the literature on the uptake of individual gases. It was found that an average of 4.60 nNO₂:nIL was absorbed at saturation (Figure S1), with the higher than equimolar capacity indicating a multi-site absorption effect for NO₂, as was found for NO.²⁷ The amount of NO₂ absorbed far exceeds that found for 1% NO by the same IL (1.73 nNO:nIL), indicating a different mechanism of absorption for NO₂.²⁷ The regeneration of the IL at 90 °C under argon was studied (well below the IL's decomposition temperature of 289 °C),¹⁸ with 3.64 nNO₂:nIL remaining after 2 h. NO₂ absorption is evidently not a reversible process, with the absorbed species strongly bound to the IL, which has also been observed for caprolactam-based ILs.²⁸

The ability of NO₂ to compete with CO₂ for absorption by [P₆₆₆₁₄][Benzim] was investigated using a combined feed through a series of absorption and desorption cycles, employing a mass spectrometry-based gas absorption rig.²⁶ Realistic flue gas concentrations of 14% CO₂ and 0.2% NO₂ were selected, with the results shown in Figure 1 (and Table S1). It demonstrates that after two cycles in the gas absorption rig, the CO₂ capacity of the IL was unaffected by the presence of NO₂, with 0.78 nCO₂:nIL still absorbed. After the third cycle, a clear decrease in CO₂ capacity was observed, with 0.66 nCO₂:nIL absorbed after exposure to a calculated 0.38













Figure 1. CO₂ capacity (bars) of [P₆₆₆₁₄][Benzim], calculated from the MS (± 0.04 nCO₂:nIL) and calculated exposure to NO₂ (solid line), after multiple cycles of a 2 h absorption under a feed of 14% CO₂ and 0.2% NO₂ in argon, and a 2 h desorption at 90 °C. A dashed line depicts the 14% CO₂ only value (0.78 nCO₂:nIL).

nNO₂:nIL. The ability of [P₆₆₆₁₄][Benzim] to capture CO₂ in the presence of NO₂ continued to decrease, reaching 0.31 nCO₂:nIL after 10 absorption/desorption cycles, an ~60% reduction in capacity. It was evident that the desorption conditions (2 h under Ar at 90 °C) were unable to regenerate the original CO₂ capacity. This strong irreversible absorption of NO₂ is in contrast to the behavior observed with the NO co-feed, where 0.72 nCO₂:nIL was still absorbed after 10 cycles.²⁷

Following the 10 competitive absorption/desorption cycles with CO₂ and NO₂, the treated IL was characterized using a number of techniques. Elemental analysis showed an increase in the nitrogen content of [P₆₆₆₁₄][Benzim] from 6.10 to 7.67 wt % and an increase in the viscosity from 1087 to 1516 mPa·s (at 25 °C). These changes were assigned to the irreversible incorporation of nitrogen into the IL through the absorption of NO₂. Changes in the physical properties of the IL after exposure to NO₂ were more significant than those observed after exposure to NO (nitrogen content increased from 6.10 to 6.48% and viscosity from 1087 to 1235 mPa·s), correlating with the reduced effect of the NO impurity (0.72 nCO₂:nIL

Table 1. Reaction Landscapes Showing Intermediates (I) and Transition States (TS) for N₂O₄ and N₂O₄/CO₂ Absorption by [P₃₃₃₃][Benzim], Depicting Potential Energy Surfaces for (a) Absorption of N₂O₄ and (b) Subsequent Heterolytic Cleavage of the N–N Bond, (c) Absorption of Two Moles of N₂O₄ and (d) the Barrier to Cleavage of the N–N Bond, and (e,f) the Absorption of CO₂ by [Benzim-N₂O₄]^a

Absorbate	Intermediate/ I ₁ (kJ·mol ⁻¹)	Transition State/ TS (kJ·mol ⁻¹)	Intermediate/ I ₂ (kJ·mol ⁻¹)
N ₂ O ₄	(a) [Benzim]····N ₂ O ₄		(b) [Benzim-NO ₂] + [NO ₂] ⁻
			
	-107.82 {-85.69}	-66.23 {-65.78}	-162.29 {-137.54}
(N ₂ O ₄) ₂	(c) [Benzim]····(N ₂ O ₄) ₂		(d) [Benzim]····(N ₂ O ₄) ₂
			
	-210.53 {-174.52}	-158.64 {-154.11}	-231.11 {-215.89}
N ₂ O ₄ + CO ₂	(e) [Benzim]····N ₂ O ₄ ····CO ₂		(f) [Benzim-CO ₂] ⁻ ····N ₂ O ₄
			
	-137.39 {-102.60}	-131.28 {-99.13}	-147.32 {-131.15}

^aValues are given in kJ·mol⁻¹ with zero-point corrected gas phase and {solvent} corrected energies calculated at B3LYP/6-311+G* level of theory (pseudo bonds = physisorption, solid bonds = chemisorption).




Figure 2. ATR-IR spectra of $[P_{66614}][Benzim]$ exposed to a feed of 0.2% NO_2 in Ar from 0 to 2 min. The spectrum of the IL before introduction of NO_2 has been subtracted from all spectra recorded under the NO_2 feed. Studied at 22 °C with a flow rate of $15 \text{ cm}^3 \cdot \text{min}^{-1}$.

was still absorbed after 10 cycles under NO and CO_2).²⁷ As the irreversible absorption of NO_2 causes changes in the physical properties of the IL, NMR and XPS of the IL after exposure were performed to characterize the nature of the strongly bound NO_2 species.

The structure was initially probed by 1H NMR (Figure S2), where it was expected that changes in the spectra upon the absorption of NO_2 would be caused by a change in the environment of the protons in the heterocyclic anion, which was demonstrated by a downfield shift in the peaks at 6.34/6.91/7.36 ppm (a) to 6.57/7.10/7.65 ppm (b) after the IL was treated with 1% NO_2 for 24 h, similar to shifts observed with SO_2 and NO .^{26,27} The IL from the absorption rig after the final cycle (c) displayed a similar shift, showing that the same absorbed species was formed under different conditions. In both spectra after NO_2 exposure, new peaks were observed above 13 ppm, suggesting the formation of HNO_3 from the reaction of NO_2 with residual water in the IL. The formation of HNO_3 was further confirmed by IR spectroscopy of $[P_{66614}][Benzim]$ after 24 h exposure to 1% NO_2 (Figure S4). The ^{13}C NMR spectra (Figure S3) showed analogous changes, where small shifts were noted for the peaks attributed to the benzimidazolide anion (114–148 ppm) due to changes in aromaticity caused by NO_2 absorption.

Ex situ XPS was further used to characterize the IL post-exposure in the absorption rig. A comparison of the N 1s region showed the presence of two new N 1s photoelectron peaks at 402.0 and 406.1 eV assigned to N–O species (Figure S5). The broad peak at 402.0 eV is assigned to adsorbed NO_2 , with the peak at a higher binding energy of 406.1 eV indicating a more oxidized N species, which can be attributed to N_2O_4 absorption.^{39,40} Absorption of NO in the same IL resulted in a single photoelectron peak observed at 402.4 eV, which was attributed to N_2O_2 formation.^{27,41} These species are expected to be strongly absorbed to the IL as the high vacuum in the XPS analyzing chamber would negate the detection of any weakly absorbed gases.

ATR-IR spectra of $[P_{66614}][Benzim]$ under separate CO_2 or NO feeds have been reported previously, showing characteristic bands for the absorbed species, with additional changes in

the spectra observed due to changes in the aromaticity of the $[Benzim]^-$ anion.²⁷ For CO_2 , reversible absorption and an equimolar capacity for absorption presented a simpler system than that for multi-site absorption, which has been reported here for NO_2 and NO .²⁷ With the higher absorption capacity of $[P_{66614}][Benzim]$ for NO_2 and the formation of a strongly bound species, a combined approach by theoretical DFT calculations and FTIR was utilized to investigate the nature of the absorbed species in $[P_{66614}][Benzim]$.

DFT calculations revealed that the formation of an N_2O_4 dimer, physically adsorbed to the $[Benzim]^-$ anion, was the most thermodynamically favored absorption mode (Table 1a). The charge perturbation from the $[Benzim]^-$ anion strongly enhanced the rate of dimerization of NO_2 , pushing the equilibrium towards N_2O_4 formation by strongly absorbing the dimer, thus removing it from the gas phase. The N_2O_4 dimer was found to physisorb strongly to the $[Benzim]^-$ anion, with a zero-point corrected absorption enthalpy (E_{ZPE}) of $-107.8 \text{ kJ} \cdot \text{mol}^{-1}$ (Tables S2 and S3). Although the absorption of N_2O_4 is energetically very favorable, the absorbed geometry together with a detailed comparison of grouped Mulliken charges (Tables S4 and S5) indicates that it is a purely physical process without charge transfer from the anion, in comparison to that of chemically adsorbed CO_2 . Instead, there are extremely strong dipole interactions between the two δ^+ nitrogen centers of the adsorbate and the formal negative charge on the anion.

Furthermore, the physical adsorption of two N_2O_4 dimers at both N-sites of the $[Benzim]^-$ anion was energetically favorable, with an absorption enthalpy of $-210.5 \text{ kJ} \cdot \text{mol}^{-1}$ (Table 1c). The multi-site adsorption and greater than equimolar adsorption of NO_2 to the IL correlates with the large gravimetric uptake capacity ($4.60 \text{ nNO}_2:\text{nIL}$), and the high absorption enthalpies relate to the drop in CO_2 capacity observed. These findings are also consistent with the experimentally observed results, where only a small decrease in adsorbed NO_2 was observed after desorption, probably due to the loss of weaker, physically bound, NO_2 .




ATR-IR spectra recorded during the absorption of 0.2% NO_2 in Ar by $[P_{66614}][Benzim]$ are shown in Figure 2. It is evident that upon the introduction of NO_2 , a series of

overlapping bands in the 1300–1400 cm^{-1} region increased in intensity. Interestingly, upon initial exposure to the feed, a band at 1335 cm^{-1} was the most intense in this region, but with increasing exposure time, a band at 1315 cm^{-1} dominated the spectra. In addition, bands at 1620, 1233, and 1036 cm^{-1} increased in intensity with exposure to the NO_2 feed. These changing bands could indicate a change in the nature of the absorbed species or absorption of a NO_2 species at another N-site on the $[\text{Benzim}]^-$ anion, as was observed with NO .²⁷

Calculated vibrational spectra for one or two moles of physisorbed N_2O_4 to N-sites on the $[\text{Benzim}]^-$ anion showed indistinguishable spectral profiles, with theoretically derived bands at 1426, 1308, and 847 cm^{-1} for one mole of physically absorbed N_2O_4 (Table 2) and 1433–1427, 1313–1305, and 847 cm^{-1} for two moles of N_2O_4 . The bands that formed upon the introduction of NO_2 at 868, 1335 and 1373 cm^{-1} correspond with the calculated bands for physisorbed N_2O_4 (Table 2). Bands at 1764/68 and 1825/24 cm^{-1} were predicted for the $\nu_{\text{asym}}(\text{O}-\text{N}-\text{O})$ vibration when one/two moles of N_2O_4 were physically absorbed, but these bands were not observed experimentally, as described by the other authors.^{42–44}



While physisorbed N_2O_4 was observed initially (eq 1), the spectra showed changes with time on stream, which were not simply an increase in the intensity of the bands assigned to N_2O_4 , thus suggesting the evolution of the absorbed species

Table 2. Depicts the Experimental and [Theoretical] IR Vibrations when $[\text{P}_{66614}][\text{Benzim}]$ Is Exposed to NO_2

Species	Vibration	IR vibration (cm^{-1})
(a) $[\text{Benzim}] \dots [\text{N}_2\text{O}_4]$		
	$\nu(\text{N}-\text{N})$	868 [847]
	$\nu_s(\text{O}-\text{N}-\text{O})$	1335 [1308]
	$\nu(\text{N}_2\text{O}_4)$	1373 [1426]
	$\nu_{\text{as}}(\text{O}-\text{N}-\text{O})$	–
		[1764/1825]
(b) $[\text{Benzim}-\text{NO}_2]$		
	$\nu(\text{O}-\text{N}-\text{O})$	868 [843]
	$\nu(\text{N}-\text{N})$	1036 [1071]
	$\nu_s(\text{O}-\text{N}-\text{O})$	1308 [1312-47]
	$\nu_{\text{as}}(\text{O}-\text{N}-\text{O})$	1620 [1630-76]
(c) $[\text{P}_{\text{RRRR}}][\text{NO}_2]$		
	$\nu(\text{O}-\text{N}-\text{O})$	828 [816]
	$\nu_s(\text{O}-\text{N}-\text{O})$	1233 [1236]
	$\nu_{\text{as}}(\text{O}-\text{N}-\text{O})$	1405 [1383]

with an increasing NO_2 concentration. The N–N bond of absorbed N_2O_4 has been reported to undergo heterolytic cleavage, forming $[\text{NO}_3]^-$ and $[\text{NO}]^+$ ionic species due to perturbation by an external charged species (eq 2 and 3).^{45,46}

The energy diagrams in Table 1 show the most favored reaction pathway following the physisorption of one and two moles of N_2O_4 . In these reaction schemes, the heterolytic cleavage of the N–N bond of absorbed N_2O_4 is favorable (b), but the activation barrier for the disproportionation reaction to $\text{NO}^+/\text{NO}_3^-$ was above 100 kJ mol^{-1} and was considered too high (Figure S6). The cleavage of the N–N bond to form $[\text{NO}_2]^+$ and $[\text{NO}_2]^-$, however, can occur owing to the lower barrier (41.6 kJ mol^{-1}), forming a thermodynamically stable complex, which is the favored theoretical pathway following the initial physisorption of N_2O_4 on $[\text{P}_{66614}][\text{Benzim}]$.



The theoretical pathway involving $[\text{NO}_2]^+$ and $[\text{NO}_2]^-$ species has significant implications for the absorption capacity of the IL. The $[\text{NO}_2]^+$ ion is proposed to absorb chemically to the $[\text{Benzim}]^-$ anion, leading to a neutral $[\text{Benzim}-\text{NO}_2]$ complex (Table 2b). The neutralization of the $[\text{Benzim}]^-$ anion removes the active site for absorption, causing deactivation of the anion/IL, and provides evidence for the decreased CO_2 capacity. The $[\text{NO}_2]^-$ ion stabilizes the positively charged $[\text{P}_{3333}]^+$ cation by orienting the negatively charged oxygen atoms into a position to counter the positively charged phosphorus (Table 2c). It could be proposed that the $[\text{NO}_2]^-$ ion reacts with, or deprotonates, the acidic α -protons close to the phosphonium cation center ($\text{P}-\text{CH}_2-$), but this pathway was not observed (see Table 2c, which shows the lowest energy structure).⁴⁷ Additionally, Table 1c,d shows that when two moles of N_2O_4 are physically bound, which is more likely at increased exposure times, the heterolytic cleavage of the N–N bonds has little driving force due to similar initial and final state energies, as well as a higher barrier (51.9 kJ mol^{-1}). Thus, the formation of the neutral $[\text{Benzim}-\text{NO}_2]$ complex (Table 1d) is expected to be equilibrium-limited, leading to the gradual decrease in the CO_2 capacity of the IL.

To probe the evolution of the bands in Figure 2, the spectrum after 1 min under the NO_2 feed was subtracted from all subsequent spectra (Figure 3) to allow comparison with the theoretically determined band positions of the $[\text{Benzim}-\text{NO}_2]$ and $[\text{P}_{\text{RRRR}}][\text{NO}_2]$ complexes (Table 2). A new band was observed at 1233 cm^{-1} , attributed to the formation of a $[\text{NO}_2]^-$ anion, and related to vibrations at 1405 and 828 cm^{-1} . Evidence of a new N–N bond was observed at 1036 cm^{-1} , from the formation of the neutral $[\text{Benzim}-\text{NO}_2]$ complex, associated with bands at 1620, 1308, and 868 cm^{-1} .^{48–50} The DFT calculated species correlated well with the observed ATR spectra, and additional features were due to changes in the aromaticity of the $[\text{Benzim}]^-$ anion, as observed with $\text{CO}_2/\text{SO}_2/\text{NO}$.^{26,27} Interestingly, the changes in the anion aromaticity are observed during the physical absorption of N_2O_4 , showing that the strong initial interaction distorts the charge density of the anion.




Figure 3. ATR-IR spectra of $[P_{66614}][Benzim]$ exposed to a feed of (i) 0.2% NO_2 in Ar after (a) 1 min and (b) between 1 and 5 min with the subtraction of the spectrum recorded at 1 min; and (ii) 14% $CO_2 + 0.2\% NO_2$ in Ar for 0–2 min. Studied at 22 °C with a flow rate of $15 \text{ cm}^3 \cdot \text{min}^{-1}$. The color of the labeled bands indicates physisorbed N_2O_4 (black), changes in the aromaticity of the IL (gray), chemical absorption of CO_2 (green), $[Benzim-NO_2]$ (blue), and $[P_{66614}][NO_2]$ (red).

Combining both the DFT calculations and the FTIR spectroscopic study allowed the determination of the absorbed species and how the speciation changed with time. The DFT results indicate the thermodynamically stable intermediates, while FTIR provides evidence for the proposed intermediates and the kinetics of absorption. Together, the results indicate the strong physical absorption of N_2O_4 followed by the deactivation of the IL through the formation of $[Benzim-NO_2]/[P_{RRRR}][NO_2]$ complexes at increased exposure times, which would be expected to result in a loss of absorption sites for CO_2 and the co-absorption of CO_2 was therefore studied as well.

A co-feed of 14% CO_2 and 0.2% NO_2 in Ar was investigated, simulating conditions in the gas absorption rig (Figure 3ii), to study whether bands assigned to the neutral complex would form and influence the CO_2 absorption capacity of the IL (as observed in the absorption rig results). The chemical absorption of CO_2 results in the formation of a carbamate species, with bands at 1709 (C=O) and 1281 cm^{-1} (N-COO⁻) quickly increasing during the first minute of exposure to the feed, which is caused by the greater concentration of CO_2 in the feed.²⁶ Simultaneously, bands associated with the physical absorption of N_2O_4 grow (1373 and 1334 cm^{-1}) due to the strong absorption enthalpy of the species ($-107.8 \text{ kJ} \cdot \text{mol}^{-1}$) (Table 1a). As the exposure time increases, the band at 1709 cm^{-1} (C=O) is noted to decrease, while weak bands at 1233, 1036, 868, and 828 cm^{-1} appear, indicating the chemical absorption of NO_2^+ to the $[Benzim]^-$ anion and the formation of $[P_{66614}][NO_2]$. This correlates with the stepped reduction in CO_2 capacity observed in the gas absorption rig. Subtracted data in Figure S7 show that bands due to the heterolytic cleavage of physisorbed N_2O_4 are observed to form with and without the presence of CO_2 . The absorption of CO_2 does not hinder the absorption and subsequent heterolytic cleavage of N_2O_4 , which, however, deactivates the IL to further absorption of CO_2 .

The physical absorption of N_2O_4 at one site drastically changes the kinetics and thermodynamics of absorption at the

remaining site, where the barrier for CO_2 activation is low ($6.1 \text{ kJ} \cdot \text{mol}^{-1}$) (Table 1e,f). Calculations show that the co-absorption of N_2O_4 and CO_2 to separate N-sites is thermodynamically stable ($-147.3 \text{ kJ} \cdot \text{mol}^{-1}$), aided by the low concentration of NO_2 . After an extended period, both the deactivation of the $[Benzim]^-$ anion and the larger absorption enthalpy of the binding of two moles of N_2O_4 ($-210.5 \text{ kJ} \cdot \text{mol}^{-1}$) suggest that decreases in CO_2 capacity are observed due to the loss of absorption sites. The reaction mechanism between the IL and CO_2/NO_2 is depicted in Figure S9, showing how the binding of a second N_2O_4 dimer further deactivates $[P_{66614}][Benzim]$ for CO_2 capture, before the formation of the $[Benzim-NO_2]/[P_{RRRR}][NO_2]$ complexes occurs at increased exposure times. The irreversible nature of the bound species is further demonstrated through studying the regeneration of the IL at 90 °C (Figure S8), where it was clear that the NO_x species could not be fully desorbed from $[P_{66614}][Benzim]$.

These results differ from those when NO is in the co-feed, where the co-absorption of both CO_2 and NO was observed.²⁷ The results obtained previously with NO were rationalized by the reaction landscape depicted in Figure S10, where the reduction of CO_2 is a barrierless process, and the activation barrier to form the NONO-ate complex is relatively high, showing a preference for the absorption of CO_2 . After the formation of the carbamate, slow deactivation of the IL is observed in the presence of NO via a CO_2 bond intermediate (which enables NO to co-bind). Using *in silico* techniques, this observation was explained by the relatively high kinetic barriers for the formation of the NO_x species bound to the IL from NO, as opposed to NO_2 . Therefore, it is evident that an alternative absorption mechanism occurs, accelerating the effect that NO_2 has on the uptake of CO_2 .

A summary of the effect of the different waste gas stream impurities on $[P_{66614}][Benzim]$ studied here is shown in Figure 4. It is clear that SO_2 has the greatest effect on reducing the CO_2 uptake in ILs followed by NO_2 and finally NO, in clear accord with the calculated absorption energies of -123.9 ,²⁷




Figure 4. CO₂ uptake of [P₆₆₆₁₄][Benzim] after exposure to an increasing amount of flue gas impurity after multiple absorption/desorption cycles of a feed containing 14% CO₂ and 0.2% impurity; dashed line, 14% CO₂ only value. Green triangles, SO₂. Red solid diamonds, NO₂. Blue circles, NO. Adapted with permission from ref 26. Copyright 2018 American Chemical Society. Adapted with permission from ref 27. Copyright 2019 American Chemical Society

−107.8, and −91.1 kJ·mol^{−1}, respectively (CO₂ = −52.1 kJ·mol^{−1}). It should be noted that, whereas the formation of the NONO-ate is thermodynamically stable (from the absorption of NO), this process is kinetically slow, hindered by the high concentration of CO₂ in the gas feed.

SUMMARY AND CONCLUSIONS

The accelerated effect of a flue gas contaminant, NO₂, on the deactivation of [P₆₆₆₁₄][Benzim] was investigated and compared to NO. Industrially relevant gas concentrations were used to demonstrate how small amounts of impurities can dramatically change the capability of a sorbent to capture CO₂, and in this particular case, an ~60% decrease in CO₂ capacity was observed. This decrease was corroborated by other experimental techniques and DFT calculations, showing the importance of considering contaminants when designing ionic liquids, or new sorbents in general, for CO₂ capture. Spectroscopic results in combination with DFT calculations showed that NO₂ was predominately strongly physically adsorbed on multiple sites in the form of N₂O₄. This species was then found to undergo heterolytic cleavage resulting in the deactivation of the IL. The ability to investigate the effects of flue gas impurities, individually and competitively, on the uptake of CO₂, is an important experimental tool in the development of new sorbents and highlights the need for rigorous experimental methods, preferably in tandem with theoretical studies and the investigation of more complex, realistic multi-component feeds.

Overall, these results show the importance of investigating the effect of flue gas contaminants. Further studies into the feasibility and optimization of ILs in such a process are still required, but a consideration of impurities and sorbent recyclability is an essential factor. Significant improvements in tuning the selectivity and absorption enthalpies of ILs to reversibly capture NO_x or SO₂ at high efficiencies are challenging, and alternative methods of removing acidic impurities will still be required (gas scrubbers and traps). Additionally, tuning the basicity of the IL by selecting anions with different pK_a values presents an interesting opportunity for the pretreatment of waste gas feeds, where contaminants such as NO_x could be reversibly captured.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acssuschemeng.1c01451>.

Gravimetric results (Figure S1); absorption rig data displayed in Figure 1 (Table S1); NMR data and ¹H/¹³C NMR spectra (Figures S2 and S3, respectively); ATR-IR spectra after the absorption of 1% NO₂ (Figure S4); N 1 s XPS spectra (Figure S5); calculated absorption energies (Tables S2 and S3) and grouped Mulliken charges/spin populations for various gases to [P₃₃₃₃][Benzim] (Tables S4 and S5); potential energy landscape of oxygen transfer from bound NO₂⁺ (Figure S6); ATR-IR difference spectra of the absorption of NO₂ and NO₂ + CO₂ co-feed (Figure S7); ATR-IR spectra showing desorption of the NO₂, and NO₂ + CO₂ co-feed (Figure S8); mechanism depicting the effect of N₂O₄ adsorption on the CO₂ recyclability of [P₆₆₆₁₄][Benzim] (Figure S9); potential energy landscape of CO₂/NO absorption (Figure S10); listed Cartesian co-ordinates (PDF)

AUTHOR INFORMATION

Corresponding Authors

Adam J. Greer – School of Chemistry and Chemical Engineering, Queen's University Belfast, Belfast BT9 5AG, Northern Ireland; Department of Chemical Engineering and Analytical Science, The University of Manchester, Manchester M13 9PL, United Kingdom; orcid.org/0000-0003-1639-5433; Phone: +44 (0) 161 306 2227; Email: adam.greer@manchester.ac.uk

Johan Jacquemin – Laboratoire PCM2E, Université de Tours, 37200 Tours, France; Materials Science and Nano-Engineering, Mohammed VI Polytechnic University, Ben Guerir 43150, Morocco; orcid.org/0000-0002-4178-8629; Phone: +33 (0) 2 47 36 73 29; Email: jj@univ-tours.fr; Fax: +33 (0) 2 47 36 70 73

Christopher Hardacre – Department of Chemical Engineering and Analytical Science, The University of Manchester, Manchester M13 9PL, United Kingdom; orcid.org/0000-0001-7256-6765; Phone: +44 (0) 161 306 2672; Email: c.hardacre@manchester.ac.uk

Authors

S. F. Rebecca Taylor – Department of Chemical Engineering and Analytical Science, The University of Manchester, Manchester M13 9PL, United Kingdom; orcid.org/0000-0002-6175-2631

Helen Daly – Department of Chemical Engineering and Analytical Science, The University of Manchester, Manchester M13 9PL, United Kingdom; orcid.org/0000-0002-1019-8490

Matthew G. Quesne – School of Chemistry, Cardiff University, Cardiff CF10 3AT, United Kingdom; UK Catalysis Hub, Research Complex at Harwell, STFC Rutherford Appleton Laboratory, Didcot, Oxfordshire OX11 0FA, United Kingdom; orcid.org/0000-0001-5130-1266

Nora H. de Leeuw – School of Chemistry, Cardiff University, Cardiff CF10 3AT, United Kingdom; School of Chemistry, University of Leeds, Leeds LS2 9JT, United Kingdom

C. Richard A. Catlow – School of Chemistry, Cardiff University, Cardiff CF10 3AT, United Kingdom; UK

Catalysis Hub, Research Complex at Harwell, STFC Rutherford Appleton Laboratory, Didcot, Oxfordshire OX11 0FA, United Kingdom; Department of Chemistry, University College London, London WC1H 0AJ, United Kingdom

Complete contact information is available at: <https://pubs.acs.org/10.1021/acssuschemeng.1c01451>

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

The authors acknowledge funding from the EPSRC under grant no. EP/N009533/1, a multi-disciplinary approach to generating low carbon fuels, carried out in collaboration with the University of Manchester, Queen's University Belfast, Cardiff University, and University College London. Open access data can be found via the University of Manchester research portal. A.J.G. further acknowledges funding from the Department for Employment and Learning. Computing facilities for this work were provided by ARCCA at Cardiff University, HPC Wales, and through the membership of the UK's Materials Chemistry Consortium (MCC). Additionally, computing resources were provided by the STFC Scientific Computing Department's SCARF cluster. The MCC is funded by the EPSRC (EP/F067496). The UK Catalysis Hub is thanked for resources and support provided via the membership of the UK Catalysis Hub Consortium and funded by EPSRC (grants EP/R026815/1, EP/ K014854/1, and EP/ M013219/1).

REFERENCES

- (1) Khatri, R. A.; Chuang, S. S. C.; Soong, Y.; Gray, M. Thermal and Chemical Stability of Regenerable Solid Amine Sorbent for CO₂ Capture. *Energy Fuels* **2006**, *20*, 1514–1520.
- (2) Ertl, G.; Knoezinger, H. *Handbook of Heterogeneous Catalysis*; vol. 5. (Wiley-VCH: 1997).
- (3) Gao, J.; Wang, S.; Zhao, B.; Qi, G.; Chen, C. Pilot-scale experimental study on the CO₂ capture process with existing SO₂: Degradation, reaction rate, and mass transfer. *Energy Fuels* **2011**, *25*, 5802–5809.
- (4) Clean Air Technology Centre. *Nitrogen Oxides (NO_x), Why and How They Are Controlled*; EPA 456/F-99-006R; United States Environmental Protection Agency: Research Triangle Park, 1999.
- (5) *European Parliament Directive 2008/50/EC of 21 May 2008 on ambient air quality and cleaner air for Europe*; European Union law: [Online]. [Accessed 22 October 2018]. Available at: <https://eur-lex.europa.eu/>.
- (6) Fostås, B.; Gangstad, A.; Nenseter, B.; Pedersen, S.; Sjøvoll, M.; Sørensen, A. L. Effects of NO_x in the flue gas degradation of MEA. *Energy Procedia* **2011**, *4*, 1566–1573.
- (7) Supap, T.; Shi, H.; Idem, R.; Gelowitz, D.; Campbell, C.; Ball, M. Nitrosamine Formation Mechanism in Amine-Based CO₂ Capture: Experimental Validation. *Energy Procedia* **2017**, *114*, 952–958.
- (8) Magee, P. N.; Barnes, J. M. The production of malignant primary hepatic tumours in the rat by feeding dimethylnitrosamine. *Br. J. Cancer* **1956**, *10*, 114–122.
- (9) Rogers, R. D.; Seddon, K. R. Ionic Liquids–Solvents of the Future? *Science* **2003**, *302*, 792–793.
- (10) Greer, A. J.; Jacquemin, J.; Hardacre, C. Industrial Applications of Ionic Liquids. *Molecules* **2020**, *25*, 5207.
- (11) Blanchard, L. A.; Hancu, D.; Beckman, E. J.; Brennecke, J. F. Green processing using ionic liquids and CO₂. *Nature* **1999**, *399*, 28–29.
- (12) Muldoon, M. J.; Aki, S. N. V. K.; Anderson, J. L.; Dixon, J. K.; Brennecke, J. F. Improving carbon dioxide solubility in ionic liquids. *J. Phys. Chem. B* **2007**, *111*, 9001–9009.
- (13) Zhang, Y.; Zhang, S.; Lu, X.; Zhou, Q.; Fan, W.; Zhang, X. Dual amino-functionalised phosphonium ionic liquids for CO₂ capture. *Chemistry* **2009**, *15*, 3003–3011.
- (14) Gurkan, B. E.; de la Fuente, J. C.; Mindrup, E. M.; Ficke, L. E.; Goodrich, B. F.; Price, E. A.; Schneider, W. F.; Brennecke, J. F. Equimolar CO₂ Absorption by Anion-Functionalized Ionic Liquids. *J. Am. Chem. Soc.* **2010**, *132*, 2116–2117.
- (15) Wang, C.; Luo, X.; Luo, H.; Jiang, D. E.; Li, H.; Dai, S. Tuning the basicity of ionic liquids for equimolar CO₂ capture. *Angew. Chem., Int. Ed.* **2011**, *50*, 4918–4922.
- (16) Seo, S.; Quiroz-Guzman, M.; DeSilva, M. A.; Lee, T. B.; Huang, Y.; Goodrich, B. F.; Schneider, W. F.; Brennecke, J. F. Chemically Tunable Ionic Liquids with Aprotic Heterocyclic Anion (AHA) for CO₂ Capture. *J. Phys. Chem. B* **2014**, *118*, 5740–5751.
- (17) Gohndrone, T. R.; Bum Lee, T.; DeSilva, M. A.; Quiroz-Guzman, M.; Schneider, W. F.; Brennecke, J. F. Competing Reactions of CO₂ with Cations and Anions in Azolide Ionic Liquids. *ChemSusChem* **2014**, *7*, 1970–1975.
- (18) Taylor, S. F. R.; McCrellis, C.; McStay, C.; Jacquemin, J.; Hardacre, C.; Mercy, M.; Bell, R. G.; de Leeuw, N. H. CO₂ Capture in Wet and Dry Superbase Ionic Liquids. *J. Solution Chem.* **2015**, *44*, 511–527.
- (19) Wang, C.; Cui, G.; Luo, X.; Xu, Y.; Li, H.; Dai, S. Highly efficient and reversible SO₂ capture by tunable azole-based ionic liquids through multiple-site chemical absorption. *J. Am. Chem. Soc.* **2011**, *133*, 11916–11919.
- (20) Cui, G.; Zheng, J.; Luo, X.; Lin, W.; Ding, F.; Li, H.; Wang, C. Tuning anion-functionalized ionic liquids for improved SO₂ capture. *Angew. Chem., Int. Ed. Engl.* **2013**, *52*, 10620–10624.
- (21) Chen, K.; Lin, W.; Yu, X.; Luo, X.; Ding, F.; He, X.; Li, H.; Wang, C. Designing of anion-functionalized ionic liquids for efficient capture of SO₂ from flue gas. *AIChE J.* **2015**, *61*, 2028–2034.
- (22) Chen, K.; Shi, G.; Zhou, X.; Li, H.; Wang, C. Highly Efficient Nitric Oxide Capture by Azole-Based Ionic Liquids through Multiple-Site Absorption. *Angew. Chemie Int. Ed.* **2016**, *55*, 14364–14368.
- (23) Cao, N.; Gan, L.; Xiao, Q.; Lv, X.; Lin, W.; Li, H.; Wang, C. Highly Efficient and Reversible Nitric Oxide Capture by Functionalized Ionic Liquids through Multiple-Site Absorption. *ACS Sustainable Chem. Eng.* **2020**, *8*, 2990–2995.
- (24) Sun, Y.; Ren, S.; Hou, Y.; Zhang, K.; Zhang, Q.; Wu, W. Highly Reversible and Efficient Absorption of Low-Concentration NO by Amino-Acid-Based Ionic Liquids. *ACS Sustainable Chem. Eng.* **2020**, *8*, 3283–3290.
- (25) Zhai, R.; He, X.; Mei, K.; Chen, K.; Cao, N.; Lin, W.; Li, H.; Wang, C. Ultrahigh Nitric Oxide Capture by Tetrakis(azolyl)borate Ionic Liquid through Multiple-Sites Uniform Interaction. *ACS Sustainable Chem. Eng.* **2021**, *9*, 3357–3362.
- (26) Taylor, S. F. R.; McClung, M.; McReynolds, C.; Daly, H.; Greer, A. J.; Jacquemin, J.; Hardacre, C. Understanding the Competitive Gas Absorption of CO₂ and SO₂ in Superbase Ionic Liquids. *Ind. Eng. Chem. Res.* **2018**, *57*, 17033–17042.
- (27) Greer, A. J.; Taylor, S. F. R.; Daly, H.; Quesne, M.; Catlow, C. R. A.; Jacquemin, J.; Hardacre, C. Investigating the Effect of NO on the Capture of CO₂ Using Superbase Ionic Liquids for Flue Gas Applications. *ACS Sustainable Chem. Eng.* **2019**, *7*, 3567–3574.
- (28) Duan, E.; Guo, B.; Zhang, D.; Shi, L.; Sun, H.; Wang, Y. Absorption of NO and NO₂ in Caprolactam Tetrabutyl Ammonium Halide Ionic Liquids. *J. Air Waste Manage. Assoc.* **2011**, *61*, 1393–1397.
- (29) Li, X.; Zhang, L.; Li, L.; Hu, Y.; Liu, J.; Xu, Y.; Luo, C.; Zheng, C. NO Removal from Flue Gas Using Conventional Imidazolium-Based Ionic Liquids at High Pressures. *Energy Fuels* **2018**, *32*, 6039–6048.
- (30) Kunov-Kruse, A. J.; Thomassen, P. L.; Riisager, A.; Mossin, S.; Fehrmann, R. Absorption and Oxidation of Nitrogen Oxide in Ionic Liquids. *Chem.-A Eur. J.* **2016**, *22*, 11745–11755.

- (31) Oster, K.; Goodrich, P.; Jacquemin, J.; Hardacre, C.; Ribeiro, A. P. C.; Elsinawi, A. A new insight into pure and water-saturated quaternary phosphonium-based carboxylate ionic liquids: Density, heat capacity, ionic conductivity, thermogravimetric analysis, thermal conductivity and viscosity. *J. Chem. Thermodyn.* **2018**, *121*, 97–111.
- (32) Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Scalmani, G.; Barone, V.; Mennucci, B.; Petersson, G. A.; Nakatsuji, H.; Caricato, M.; Li, X.; Hratchian, H. P.; Izmaylov, A. F.; Bloino, J.; Zheng, G.; Sonnenberg, J. L.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Vreven, T.; Montgomery, J. A.; Peralta, J. E.; Ogliaro, F.; Bearpark, M.; Heyd, J. J.; Brothers, E.; Kudin, K. N.; Staroverov, V. N.; Kobayashi, R.; Normand, J.; Raghavachari, K.; Rendell, A.; Burant, J. C.; Iyengar, S. S.; Tomasi, J.; Cossi, M.; Rega, N.; Millam, J. M.; Klene, M.; Knox, J. E.; Cross, J. B.; Bakken, V.; Adamo, C.; Jaramillo, J.; Gomperts, R.; Stratmann, R. E.; Yazyev, O.; Austin, A. J.; Cammi, R.; Pomelli, C.; Ochterski, J. W.; Martin, R. L.; Morokuma, K.; Zakrzewski, V. G.; Voth, G. A.; Salvador, P.; Dannenberg, J. J.; Dapprich, S.; Daniels, A. D.; Farkas, Foresman, J. B.; Ortiz, J. V.; Cioslowski, J.; Fox, D. J. *Gaussian 09*; Revis. A.02, Gaussian, Inc.: Wallingford CT, 200.
- (33) Sahu, S.; Quesne, M. G.; Davies, C. G.; Dörr, M.; Ivanović-Burmazović, I.; Siegler, M. A.; Jameson, G. N. L.; De Visser, S. P.; Goldberg, D. P. Direct observation of a nonheme iron(IV)-oxo complex that mediates aromatic C-F hydroxylation. *J. Am. Chem. Soc.* **2014**, *136*, 13542–13545.
- (34) Neu, H. M.; Yang, T.; Baglia, R. A.; Yosca, T. H.; Green, M. T.; Quesne, M. G.; de Visser, S. P.; Goldberg, D. P. Oxygen-Atom Transfer Reactivity of Axially Ligated Mn(V)-Oxo Complexes: Evidence for Enhanced Electrophilic and Nucleophilic Pathways. *J. Am. Chem. Soc.* **2014**, *136*, 13845–13852.
- (35) Timmins, A.; Quesne, M. G.; Borowski, T.; de Visser, S. P. Group Transfer to an Aliphatic Bond: A Biomimetic Study Inspired by Nonheme Iron Halogenases. *ACS Catal.* **2018**, *8*, 8685–8698.
- (36) Mercy, M.; Rebecca Taylor, S. F.; Jacquemin, J.; Hardacre, C.; Bell, R. G.; De Leeuw, N. H. The addition of CO₂ to four superbases ionic liquids: a DFT study. *Phys. Chem. Chem. Phys.* **2015**, *17*, 28674–28682.
- (37) Zhurko, G.; Zhurko, D. *Chemcraft Program*. Academic version 1.5. 2004.
- (38) Grimme, S.; Antony, J.; Ehrlich, S.; Krieg, H. A consistent and accurate ab initio parametrization of density functional dispersion correction (DFT-D) for the 94 elements H-Pu. *J. Chem. Phys.* **2010**, *132*, 154104.
- (39) Polzonetti, G.; Alnot, P.; Brundle, C. R. The adsorption and reactions of NO₂ on the Ag(111) surface. I. XPS/UPS and annealing studies between 90 and 300 K. *Surf. Sci.* **1990**, *238*, 226–236.
- (40) Sysoev, V. I.; Okotrub, A. V.; Guse'nikov, A. V.; Smirnov, D. A.; Bulusheva, L. G. In situ XPS Observation of Selective NO_x Adsorption on the Oxygenated Graphene Films. *Phys. status solidi* **2018**, *255*, 17002671–17002675.
- (41) Ruiz-Soria, G.; Pérez Paz, A.; Sauer, M.; Mowbray, D. J.; Lacovig, P.; Dalmiglio, M.; Lizzit, S.; Yanagi, K.; Rubio, A.; Goldoni, A.; Ayala, P.; Pichler, T. Revealing the Adsorption Mechanisms of Nitroxides on Ultrapure, Metallicity-Sorted Carbon Nanotubes. *ACS Nano* **2014**, *8*, 1375–1383.
- (42) Koch, T. G.; Horn, A. B.; Chesters, M. A.; McCoustra, M. R. S.; Sodeau, J. R. A low-temperature reflection-absorption infrared spectroscopic study of ultrathin films of dinitrogen tetroxide and dinitrogen pentoxide on gold foil. *J. Phys. Chem.* **1995**, *99*, 8362–8367.
- (43) Wang, J.; Koel, B. E. IRAS Studies of NO₂, N₂O₃, and N₂O₄ Adsorbed on Au(111) Surfaces and Reactions with Coadsorbed H₂O. *J. Phys. Chem. A* **1998**, *102*, 8573–8579.
- (44) Cataldo, F.; Compagnini, G.; D'Urso, L.; Mita, V.; Strazzulla, G.; Ursini, O.; Angelini, G. Adsorption of dinitrogen tetroxide (N₂O₄) on multi-walled carbon nanotubes (MWCNTs). *Fullerenes Nanotubes Carbon Nanostruct.* **2008**, *16*, 154–164.
- (45) Pidko, E. A.; Mignon, P.; Geerlings, P.; Schoonheydt, R. A.; Van Santen, R. A. A periodic DFT study of N₂O₄ disproportionation on alkali-exchanged zeolites X. *J. Phys. Chem. C* **2008**, *112*, 5510–5519.
- (46) Lignell, H.; Varner, M. E.; Finlayson-Pitts, B. J.; Benny Gerber, R. Isomerization and ionization of N₂O₄ on model ice and silica surfaces. *Chem. Phys.* **2012**, *405*, 52–59.
- (47) Yeadon, D. J.; Jacquemin, J.; Plechkova, N. V.; Maréchal, M.; Seddon, K. R. Induced Protic Behaviour in Aprotic Ionic Liquids by Anion Basicity for Efficient Carbon Dioxide Capture. *ChemPhysChem* **2020**, *21*, 1369–1374.
- (48) Busca, G.; Lorenzelli, V. Infrared study of the adsorption of nitrogen dioxide, nitric oxide and nitrous oxide on hematite. *J. Catal.* **1981**, *72*, 303–313.
- (49) Weston, R. E., Jr.; Brodasky, T. F. Infrared Spectrum and Force Constants of the Nitrite Ion. *J. Chem. Phys.* **1957**, *27*, 683–689.
- (50) Gatehouse, B. M. A survey of the infrared spectra of NO₂ in metal complexes. *J. Inorg. Nucl. Chem.* **1958**, *8*, 79–86.