

## RESEARCH ARTICLE

# The time to peak blood bicarbonate ( $\text{HCO}_3^-$ ), pH, and the strong ion difference (SID) following sodium bicarbonate ( $\text{NaHCO}_3$ ) ingestion in highly trained adolescent swimmers

Josh W. Newbury<sup>1</sup>, Matthew Cole<sup>1</sup>, Adam L. Kelly<sup>1</sup>, Richard J. Chessor<sup>2</sup>, S. Andy Sparks<sup>1,3\*</sup>, Lars R. McNaughton<sup>3</sup>, Lewis A. Gough<sup>1</sup>

**1** Human Performance and Health Research Group, Centre for Life and Sport Sciences (CLaSS), Department of Sport and Exercise, Birmingham City University, Birmingham, United Kingdom, **2** Sports Science and Sports Medicine Team, British Swimming, Loughborough, Leicestershire, United Kingdom, **3** Sports Nutrition and Performance Research Group, Department of Sport and Physical Activity, Edge Hill University, Ormskirk, United Kingdom

\* [andy.sparks@edgehill.ac.uk](mailto:andy.sparks@edgehill.ac.uk)



## OPEN ACCESS

**Citation:** Newbury JW, Cole M, Kelly AL, Chessor RJ, Sparks SA, McNaughton LR, et al. (2021) The time to peak blood bicarbonate ( $\text{HCO}_3^-$ ), pH, and the strong ion difference (SID) following sodium bicarbonate ( $\text{NaHCO}_3$ ) ingestion in highly trained adolescent swimmers. *PLoS ONE* 16(7): e0248456. <https://doi.org/10.1371/journal.pone.0248456>

**Editor:** Caroline Sunderland, Nottingham Trent University, UNITED KINGDOM

**Received:** February 25, 2021

**Accepted:** June 17, 2021

**Published:** July 1, 2021

**Peer Review History:** PLOS recognizes the benefits of transparency in the peer review process; therefore, we enable the publication of all of the content of peer review and author responses alongside final, published articles. The editorial history of this article is available here: <https://doi.org/10.1371/journal.pone.0248456>

**Copyright:** © 2021 Newbury et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** We have now uploaded the data set as a [S1 Data](#).

## Abstract

The timing of sodium bicarbonate ( $\text{NaHCO}_3$ ) supplementation has been suggested to be most optimal when coincided with a personal time that bicarbonate ( $\text{HCO}_3^-$ ) or pH peaks in the blood following ingestion. However, the ergogenic mechanisms supporting this ingestion strategy are strongly contested. It is therefore plausible that  $\text{NaHCO}_3$  may be ergogenic by causing beneficial shifts in the strong ion difference (SID), though the time course of this blood acid base balance variable is yet to be investigated. Twelve highly trained, adolescent swimmers (age:  $15.9 \pm 1.0$  years, body mass:  $65.3 \pm 9.6$  kg) consumed their typical pre-competition nutrition 1–3 hours before ingesting  $0.3 \text{ g}\cdot\text{kg} \text{ BM}^{-1} \text{ NaHCO}_3$  in gelatine capsules. Capillary blood samples were then taken during seated rest on nine occasions (0, 60, 75, 90, 105, 120, 135, 150, 165 min post-ingestion) to identify the time course changes in  $\text{HCO}_3^-$ , pH, and the SID. No significant differences were found in the time to peak of each blood measure ( $\text{HCO}_3^-$ :  $130 \pm 35$  min, pH:  $120 \pm 38$  min, SID:  $98 \pm 37$  min;  $p = 0.08$ ); however, a large effect size was calculated between time to peak  $\text{HCO}_3^-$  and the SID ( $g = 0.88$ ). Considering that a difference between time to peak blood  $\text{HCO}_3^-$  and the SID was identified in adolescents, future research should compare the ergogenic effects of these two individualized  $\text{NaHCO}_3$  ingestion strategies compared to a traditional, standardized approach.

## Introduction

Sodium bicarbonate ( $\text{NaHCO}_3$ ) is recommended to athletes based upon strong evidence of a performance enhancing effect [1]. It is well acknowledged that  $\text{NaHCO}_3$  ingestion enhances buffering capacity by increasing bicarbonate ( $\text{HCO}_3^-$ ) and pH concentrations within the

**Funding:** RJC contributed to this research by providing peer review from the perspective of a world-class performance nutritionist. British Swimming as an organisation had no role within any aspect of this research project. All funding and consumables were provided by Birmingham City University, and all participants were volunteers from a local high-performance swimming team (City of Birmingham Swimming Club).

**Competing interests:** The commercial affiliation of RJC does not alter our adherence to PLOS ONE policies on sharing data and materials.

blood, although the magnitude of these increases is variable between individuals [2, 3]. Increasing blood alkalosis alters the pH gradient between the intracellular and extracellular compartments, subsequently leading to the upregulation of the lactate–hydrogen ion ( $\text{H}^+$ ) co-transporter to efflux acidic  $\text{H}^+$  from the active musculature and into circulation [4, 5]. During exercise, accelerating the removal of  $\text{H}^+$  is purported to offset fatigue since intracellular acidosis is associated with debilitating effects on the capacity to sustain muscle force production [6–10]. Specifically, cellular acidosis is suggested to inhibit muscle shortening velocity and cross-bridge cycling by reducing calcium ion ( $\text{Ca}^{2+}$ ) sensitivity and myosin ATPase activity [10, 11]. Further impairments of key glycolytic enzymes [12, 13] and the strong ion difference (SID) [7, 14, 15] are also associated with exacerbated  $\text{H}^+$  accumulation, hence limiting the available ATP substrates and action potentials necessary to maintain muscular contractions, respectively. However, each of these fatiguing mechanisms are strongly contested [16–18]. Despite this,  $\text{NaHCO}_3$  continues to provide an ergogenic benefit to exercise performance, subsequently requiring further investigation of the biochemical changes that occur following ingestion.

There is an apparent increase in the ergogenic potential of  $\text{NaHCO}_3$  when baseline blood  $\text{HCO}_3^-$  is increased by  $+5 \text{ mmol}\cdot\text{L}^{-1}$ , whereas increases above  $+6 \text{ mmol}\cdot\text{L}^{-1}$  are associated with almost certain performance enhancement [19, 20]. However, the time taken to reach these thresholds varies considerably between individuals when  $\text{NaHCO}_3$  is consumed in either capsule (range: 40–240 min [3, 21, 22]) or solution form (range: 40–125 min [2, 23–25]), therefore highlighting a potential flaw in the current dosing guidelines (i.e., 60–150 min pre-exercise for all athletes [1]). Contemporary research is addressing this issue by timing exercise to coincide with an individual peak blood  $\text{HCO}_3^-$  concentration [21, 23–25], though at present, only Boegman et al. [21] has directly compared exercise performance between  $\text{NaHCO}_3$  ingested at time to peak  $\text{HCO}_3^-$  (40–160 min pre-exercise) versus a traditional dosing protocol (60 min pre-exercise for all athletes). The individualized approach significantly improved the rowing performance (2000 m time-trial) of 18 of 23 world-class rowers (Individualized =  $367.0 \pm 10.5 \text{ s}$  vs. Traditional =  $369.0 \pm 10.3 \text{ s}$ ), however, this result occurred with only a small difference in pre-exercise  $\text{HCO}_3^-$  (Individualized =  $+6 \text{ mmol}\cdot\text{L}^{-1}$  vs. Traditional =  $+5.5 \text{ mmol}\cdot\text{L}^{-1}$ ). It is therefore plausible that another mechanism besides those associated with increased  $\text{HCO}_3^-$  is involved with the ergogenic properties of  $\text{NaHCO}_3$  supplementation.

Alternatively,  $\text{NaHCO}_3$  may mitigate fatigue by altering the intracellular and extracellular balance of strong ions such as potassium ( $\text{K}^+$ ), sodium ( $\text{Na}^+$ ), chloride ( $\text{Cl}^-$ ), and  $\text{Ca}^{2+}$  [26]. Specifically,  $\text{NaHCO}_3$  increases the influx of  $\text{K}^+$  into the muscle following ingestion and attenuates losses in intramuscular  $\text{K}^+$  during exercise, subsequently improving the capacity to sustain muscle excitability [27–30]. A concomitant increase in muscular  $\text{Cl}^-$  uptake is also observed post-ingestion [27, 29, 30], which is suspected to work synergistically with  $\text{K}^+$  to protect force and excitability when muscles begin to depolarize [31]. These effects are further strengthened by an increased plasma  $\text{Na}^+$  [2, 3, 27, 28], which together with changes in  $\text{K}^+$  and  $\text{Cl}^-$ , could indicate an upregulation of  $\text{Na}^+/\text{K}^+$ -ATPase and  $\text{Na}^+/\text{K}^+/2\text{Cl}^-$ -ATPase activity to limit depolarization and preserve excitation-contraction coupling following  $\text{NaHCO}_3$  ingestion [16, 30, 31]. Currently, only Gough et al. have reported changes in the collective SID following  $\text{NaHCO}_3$  ingestion and whole-body exercise [24, 27]. Interestingly, both studies identified an ergogenic effect of  $\text{NaHCO}_3$  (4 km cycling time-trial: 1.4% faster [24]; treadmill time-to-exhaustion: 28% further [27]) when blood  $\text{HCO}_3^-$  increases ( $+7.0 \text{ mmol}\cdot\text{L}^{-1}$  [24];  $+8.8 \text{ mmol}\cdot\text{L}^{-1}$  [27]) were identified alongside an increased SID ( $+6 \text{ mEq}\cdot\text{L}^{-1}$  [24];  $+10 \text{ mEq}\cdot\text{L}^{-1}$  [27]) prior to exercise. Given that the role of  $\text{H}^+$  accumulation is controversial within exercise fatigue [16–18], the resultant performance improvements could instead be attributed to an increased plasma SID. However, no study to date has purposely intended to identify a peak

SID measurement. If a time difference exists between the time taken to reach peak blood  $\text{HCO}_3^-$  and the SID concentrations, then it is plausible that future  $\text{NaHCO}_3$  dosing strategies based upon a time to peak SID could optimize the SID mechanisms associated with performance enhancement.

Adolescence (age: 15–19 years) is a critical time in an athlete's career, whereby competitive success is essential for continued sports participation [32]. Young athletes are therefore attracted to nutritional ergogenic aids to support their performance, though in practice, these are not always ingested in an evidence-based fashion [33]. Moreover, whether  $\text{NaHCO}_3$  supplementation is appropriate for trained adolescents is unclear, with there being little evidence to suggest that large increases in blood alkalosis occur following ingestion. For example, neither Zajac et al. ( $+3.4 \text{ mmol}\cdot\text{L}^{-1}$  [34]) or Guimarães et al. ( $+3.4 \text{ mmol}\cdot\text{L}^{-1}$  [35]) observed a blood  $\text{HCO}_3^-$  increase above the proposed  $+5 \text{ mmol}\cdot\text{L}^{-1}$  'ergogenic threshold' when measured at 60 and 90 min post-ingestion, respectively. Despite similar  $\text{HCO}_3^-$  increases, however, both of these studies differed in performance outcomes (time to complete 4 x 50 m swimming sprints: 1.3% faster [34] vs. time to complete 6 x 35 m running sprints: no change [35]). Why this discrepancy occurred remains unknown, though the authors speculated that individual factors such as genetics and gastrointestinal (GI) physiology may have had a role [35]. Indeed, both studies did not account for individual differences since standardized ingestion strategies were employed and  $\text{HCO}_3^-$  changes were reported at the group level only. In order to optimize  $\text{NaHCO}_3$  supplementation for adolescent athletes, it is therefore necessary to first understand the possible acid-base balance alterations that occur following ingestion. The purpose of this study was to explore the time course changes and peak blood concentrations of three acid-base balance variables ( $\text{HCO}_3^-$ , pH, and the SID) in highly trained adolescent swimmers following the ingestion of  $0.3 \text{ g}\cdot\text{kg BM}^{-1} \text{ NaHCO}_3$ .

## Materials and methods

### Participants

Twelve national level swimmers from a performance swim programme volunteered to participate in this study (Table 1). Three of the swimmers had recently represented Great Britain at either the European Junior Swimming Championships or the European Youth Olympic Festival in 2019, whilst seven had qualified for the senior British Swimming Championships in April 2020. At the time of the study, all swimmers were ranked within the top 25 in Great Britain in at least one event within their respected age group (mean FINA points =  $696 \pm 62$ ) and were completing a minimum of 7 x 2.5 hours pool-based and 2 x 1-hour land-based training sessions per week. The study was granted institutional ethical approval (Birmingham City

**Table 1. Characteristics of study participants ( $\pm$  SD).**

Characteristics	Males (n = 5)	Females (n = 7)	Combined (n = 12)
Age (years)	16.4 $\pm$ 1.1	15.6 $\pm$ 0.8	15.9 $\pm$ 1.0
Height (m)	1.78 $\pm$ 0.6	1.70 $\pm$ 0.4	1.73 $\pm$ 0.6
Body mass (kg)	72.4 $\pm$ 10.7	60.3 $\pm$ 4.8	65.3 $\pm$ 9.6
Sum of 8 skinfolds (mm)	55.2 $\pm$ 11.3	82.1 $\pm$ 21.5	70.9 $\pm$ 22.1
Time as a competitive junior swimmer (years)	8.2 $\pm$ 1.6	8.0 $\pm$ 0.8	8.1 $\pm$ 1.2
<b>Personal best swimming times (freestyle–long course)</b>			
50 m (sec)	26.0 $\pm$ 2.6	29.4 $\pm$ 1.5	28.0 $\pm$ 2.6
100 m (sec)	54.7 $\pm$ 3.0	61.2 $\pm$ 1.4	58.8 $\pm$ 15.6
200 m (min: sec)	1:57.7 $\pm$ 4.9	2:11.3 $\pm$ 3.6	2:05.6 $\pm$ 8.1

<https://doi.org/10.1371/journal.pone.0248456.t001>

University Ethics Committee: Newbury/3649/R(B)/2019 /Nov/HELS FAEC) and both the swimmers and their parents/guardians provided written informed consent prior to their participation in the study.

### Pre-experimental procedures

Each participant was required to attend the laboratory as per their usual training time (17:00–20:30 hrs) having eaten as they normally would prior to a competitive race (i.e., self-selected timing and meal composition). Previous studies have assessed time course changes in  $\text{HCO}_3^-$  and pH in a fasted state [2, 3, 36] or following a standardized meal [22], however, these conditions do not replicate the variance in nutritional intakes that would take place in competition. This ingestion method was selected to consider the effects of individualized nutrition on baseline acid-base balance, electrolyte status, and absorption rate that would occur in practice [37]. In addition, the ingestion of a meal has shown to reduce the occurrence and severity of potential GI side effects that are associated with  $\text{NaHCO}_3$  ingestion [38]. Each swimmer completed a 24-hour dietary recall before  $\text{NaHCO}_3$  ingestion, which was used to calculate diet composition (Table 2) using an online nutrition application (Nutritics, version 5.64, Dublin, Ireland). Similarly, water was permitted to be consumed *ad libitum* (mean intake:  $1.2 \pm 0.5$  L, range: 0.5–2.0 L), though ingestion of further nutrients were restricted once  $\text{NaHCO}_3$  had been consumed. Participants were also asked to refrain from additional exercise outside of their regular swim training programme in the 48 hours prior to participating in the study. No swimmers had ingested caffeine, although concurrent ingestion of creatine (participants 5, 8, 9, 10, and 11) and beta-alanine (participants 9 and 10) were reported. Neither of these supplements was expected to affect the blood acid-base balance alterations that would occur following  $\text{NaHCO}_3$  ingestion [39, 40]. All trials were carried out between December 2019 and February 2020 during a specific race preparation training period (weekly swimming volume =  $50.9 \pm 3.4$  km).

### Protocol and measurements

Upon arrival to the laboratory, participants engaged in five min of seated rest before a baseline capillary sample of whole blood was collected from the fingertip into a 100  $\mu\text{L}$  sodium-heparin coated glass clinitube (Radiometer Medical Ltd., Copenhagen, Denmark). Blood samples were immediately analysed using a blood gas analyser (ABL9, Radiometer Medical Ltd., Copenhagen,

**Table 2. The dietary intake of the participant cohort in the 24 hours preceding  $\text{NaHCO}_3$  ingestion.**

	24 hours pre-ingestion	1–3 hours pre-ingestion
Total energy (kcal)	2852 $\pm$ 926	655 $\pm$ 285
Relative energy (kcal·kg $\text{BM}^{-1}$ )	43.3 $\pm$ 9.8	10.0 $\pm$ 3.7
Total carbohydrate (g)	362 $\pm$ 135	75.4 $\pm$ 46.0
Relative carbohydrate (g·kg $\text{BM}^{-1}$ )	5.5 $\pm$ 1.4	1.2 $\pm$ 0.6
Total protein (g)	153 $\pm$ 37	30.9 $\pm$ 9.0
Relative protein (g·kg $\text{BM}^{-1}$ )	2.3 $\pm$ 0.4	0.5 $\pm$ 0.1
Total fat (g)	88.4 $\pm$ 31.1	25.6 $\pm$ 13.4
Relative fat (g·kg $\text{BM}^{-1}$ )	1.3 $\pm$ 0.4	0.4 $\pm$ 0.2
Total $\text{Na}^+$ (mg)	3211 $\pm$ 1110	901 $\pm$ 409
Total $\text{K}^+$ (mg)	3546 $\pm$ 1185	559 $\pm$ 560
Total $\text{Ca}^{2+}$ (mg)	1428 $\pm$ 501	276 $\pm$ 229
Total $\text{Cl}^-$ (mg)	4755 $\pm$ 1558	1374 $\pm$ 601

$\text{Na}^+$  = Dietary sodium.  $\text{K}^+$  = Dietary potassium.  $\text{Ca}^{2+}$  = Dietary calcium.  $\text{Cl}^-$  = Dietary chloride.

<https://doi.org/10.1371/journal.pone.0248456.t002>

Denmark) for measurements of  $\text{HCO}_3^-$ , pH,  $\text{K}^+$ ,  $\text{Na}^+$ ,  $\text{Ca}^{2+}$ , and  $\text{Cl}^-$ . An additional 5  $\mu\text{L}$  sample was taken for the analysis of blood lactate ( $\text{La}^-$ ) (Lactate Pro 2, Arkray, Japan) which was used in the following formula to calculate the apparent SID using a freely available spreadsheet:  $\text{K}^+ + \text{Na}^+ + \text{Ca}^{2+} - \text{Cl}^- - \text{La}^-$  [41]. This method of calculating the apparent SID has been previously utilised in similar research investigating the SID alongside  $\text{NaHCO}_3$  ingestion [24, 27]. Participants then ingested 0.3 g·kg  $\text{BM}^{-1}$   $\text{NaHCO}_3$  (Dr. Oetker, Bielefeld, Germany) contained in gelatine capsules (1 g  $\text{NaHCO}_3$  per capsule; Size 00, Bulk Powders, Colchester, UK) within a five min period prior to 165 min of quiet seated rest. Blood samples were obtained and analysed on eight more occasions (60, 75, 90, 105, 120, 135, 150, and 165 min post- $\text{NaHCO}_3$  ingestion) with further samples taken at 180 ( $n = 4$ ) and 195 min ( $n = 1$ ) to ensure a peak  $\text{HCO}_3^-$  was found. No samples were taken within the first 60 min since 0.3 g·kg $^{-1}$   $\text{BM}$   $\text{NaHCO}_3$  in capsule form was not expected to display peak  $\text{HCO}_3^-$  concentrations in this time [3, 22].

Gastrointestinal side effects were monitored using nine 200 mm visual analogue scales (VAS) for nausea, flatulence, stomach cramping, belching, stomach ache, bowel urgency, diarrhoea, vomiting and stomach bloating consistent with previous  $\text{NaHCO}_3$  research [2, 24, 25, 42]. The VAS scales were labelled “no symptom” on the left side and “severe symptom” on the right side.

## Statistical analysis

Data was normally distributed (Shapiro-Wilks) and sphericity was assumed (Mauchly) prior to statistical analysis. If sphericity was violated, Huyn-Feldt (epsilon value  $>0.75$ ) or Greenhouse-Geiser (epsilon value  $<0.75$ ) corrections were applied. A repeated measures ANOVA was conducted to establish mean differences between the time to peak of the three alkalotic variables ( $\text{HCO}_3^-$ , pH, and the SID) and group mean differences for each blood metabolite ( $\text{HCO}_3^-$ , pH, SID,  $\text{K}^+$ ,  $\text{Na}^+$ ,  $\text{Ca}^{2+}$ , and  $\text{Cl}^-$ ) at each sampling time point. Post hoc comparisons were determined by the Bonferroni correction and effect sizes are reported as partial eta squared ( $P\eta^2$ ). Additional effect sizes were calculated for each time course change using the Hedges'  $g$  bias correction, which was selected based on the small sample size ( $n < 20$ ) included in this study [43]. Effect sizes were interpreted as trivial ( $g < 0.20$ ), small ( $g = 0.20-0.49$ ), medium ( $g = 0.50-0.79$ ), and large ( $g \geq 0.80$ ) [44]. Coefficient of variation (CV) was calculated using  $\text{SD}/\text{mean} \times 100$  and described the inter-individual variance between participants for time course and absolute blood analyte changes. A Spearman's rank-order correlation ( $r_s$ ) was used to investigate associations between dietary intake (energy, macronutrients, and electrolytes consumed 24 and 3 hours before to  $\text{NaHCO}_3$  ingestion) and the time to peak, absolute change, and baseline measures of blood  $\text{HCO}_3^-$ , pH, and the SID. This non-parametric correlation was selected due to violations in the sphericity of nutritional data. All statistical tests were completed using Statistical Package for the Social Sciences (SPSS), version 25 (IBM, Chicago, USA). All data are reported as mean  $\pm$  SD with statistical significance set at  $p < 0.05$ .

## Results

### Blood analyte changes

No statistical significance was found between the time to peak concentrations of each blood acid-base variable ( $\text{HCO}_3^- = 130 \pm 34$  min, pH =  $120 \pm 38$  min, SID =  $98 \pm 37$  min;  $p = 0.077$ ,  $P\eta^2 = 0.208$ ). Despite this, a large effect size ( $g = 0.88$ ) separated the time to peak SID and  $\text{HCO}_3^-$  following a mean difference of 33 min being observed between these two measures. A moderate effect size was also observed between time to peak SID and pH (23 min,  $g = 0.58$ ), whereas the effect size was small between time to peak  $\text{HCO}_3^-$  and pH (10 min,  $g = 0.26$ ). Large inter-individual variations were present in the time to peak (CV:  $\text{HCO}_3^- = 27\%$ ;

Table 3. Individual metabolite responses to 0.3 g·kg  $\text{BM}^{-1}$  sodium bicarbonate ( $\text{NaHCO}_3$ ).

	Participant (Gender)											
	1 (M)	2 (M)	3 (F)	4 (F)	5 (M)	6 (M)	7 (F)	8 (F)	9 (F)	10 (M)	11 (F)	12 (F)
$\text{HCO}_3^-$ baseline ( $\text{mmol}\cdot\text{L}^{-1}$ )	26.5	28.0	23.3	23.9	26.0	25.6	24.8	25.9	24.7	26.4	26.4	23.6
$\text{HCO}_3^-$ peak ( $\text{mmol}\cdot\text{L}^{-1}$ )	34.3	35.3	32.3	34.5	32.8	33.9	31.4	28.3	30.5	32.7	34.8	26.4
Absolute change ( $\text{mmol}\cdot\text{L}^{-1}$ )	+7.8	+7.3	+9.0	+10.6	+6.8	+8.3	+6.6	+2.4	+5.8	+6.3	+8.4	+2.8
Time to peak (min)	90	75	120	150	150	165	75	120	150	150	135	180
pH baseline (units)	7.41	7.46	7.40	7.40	7.40	7.41	7.39	7.41	7.41	7.40	7.38	7.43
pH peak (units)	7.57	7.52	7.52	7.54	7.49	7.53	7.54	7.45	7.51	7.48	7.49	7.47
Absolute change (units)	+0.16	+0.06	+0.12	+0.14	+0.09	+0.12	+0.15	+0.04	+0.10	+0.08	+0.11	+0.04
Time to peak (min)	90	75	90	135	180	165	90	105	165	150	120	75
SID baseline ( $\text{mEq}\cdot\text{L}^{-1}$ )	33	37	33	32	33	35	33	33	31	30	34	32
SID peak ( $\text{mEq}\cdot\text{L}^{-1}$ )	36	38	38	38	37	35	38	35	35	38	38	34
Absolute change ( $\text{mEq}\cdot\text{L}^{-1}$ )	+3	+1	+5	+6	+4	0	+5	+2	+4	+8	+4	+2
Time to peak (min)	90	60	90	135	120	60	165	120	60	135	75	60

<https://doi.org/10.1371/journal.pone.0248456.t003>

pH = 32%; SID = 37%;) and absolute increase from baseline observed for each alkalotic measure ( $\text{HCO}_3^- = +6.8 \pm 2.4 \text{ mmol}\cdot\text{L}^{-1}$ , CV = 35%; pH =  $+0.10 \pm 0.04$  units, CV = 42%; SID =  $+3.7 \pm 2.2 \text{ mEq}\cdot\text{L}^{-1}$ , CV = 61%). Individual data for all blood acid-base balance variables is presented in Table 3, whereas the time course changes in  $\text{HCO}_3^-$  and the SID has been plotted in Figs 1 and 2 for four different participants to emphasise the individuality in blood responses.

Across the sampling timeframe, marked increases in acid-base variables were observed ( $\text{HCO}_3^-$ :  $p < 0.001$ ,  $P\eta^2 = 0.609$ ; pH:  $p < 0.001$ ,  $P\eta^2 = 0.529$ ; SID:  $p = 0.029$ ,  $P\eta^2 = 0.212$ ; Fig 3). Blood bicarbonate was elevated 60 min post-ingestion ( $+3.2 \pm 2.6 \text{ mmol}\cdot\text{L}^{-1}$ ,  $p = 0.046$ ) and remained  $+4$ – $6 \text{ mmol}\cdot\text{L}^{-1}$  until 165 min post-ingestion (all  $p < 0.030$ ). At 105 min mean blood  $\text{HCO}_3^-$  had increased by  $5.2 \pm 2.5 \text{ mmol}\cdot\text{L}^{-1}$  and remained elevated above the proposed





Fig 1. The time course change in blood bicarbonate ( $\text{HCO}_3^-$ ) over a 165 min period following  $\text{NaHCO}_3$  ingestion for A) participant 1, B) participant 8, C) participant 7, and D) participant 5. The dotted line represents a  $+5 \text{ mmol}\cdot\text{L}^{-1}$  increase from baseline blood  $\text{HCO}_3^-$ .

<https://doi.org/10.1371/journal.pone.0248456.g001>






**Fig 2.** The time course change in blood strong ion difference (SID) over a 165 min period following  $\text{NaHCO}_3$  ingestion for A) participant 10, B) participant 2, C) participant 12, and D) participant 3.

<https://doi.org/10.1371/journal.pone.0248456.g002>

+5  $\text{mmol}\cdot\text{L}^{-1}$  ergogenic threshold until the final 165 min time point (peak: 150 min, +5.9 ± 2.7  $\text{mmol}\cdot\text{L}^{-1}$ , +23%). An increased pH occurred at 75 min post-ingestion (+0.06 ± 0.04 units, +0.8%,  $p = 0.010$ ) and this level of increase was sustained at all remaining points in time (+0.06–0.08 units, all  $p < 0.030$ ). Peak pH occurred at 165 min post-ingestion (+0.08 ± 0.05 units, +1.1%). The SID increased by  $2.0 \pm 1.5 \text{ mEq}\cdot\text{L}^{-1}$  (+6%,  $p = 0.024$ ) at 60 min post-ingestion and peaked at 135 min post-ingestion (+2.4 ± 2.6  $\text{mEq}\cdot\text{L}^{-1}$ , +7%), however, this did not reach statistical significance.

Ingestion of  $\text{NaHCO}_3$  produced ionic shifts in the balance of strong ions ( $\text{K}^+$ :  $p = 0.005$ ,  $P\eta^2 = 0.215$ ;  $\text{Na}^+$ :  $p = 0.044$ ,  $P\eta^2 = 0.160$ ;  $\text{Ca}^{2+}$ :  $p < 0.001$ ,  $P\eta^2 = 0.562$ ;  $\text{Cl}^-$ :  $p = 0.016$ ,  $P\eta^2 = 0.306$ ). The electrolytes with the greatest absolute changes were  $\text{K}^+$  and  $\text{Cl}^-$ . Large inter-individual variations restricted the ability to detect statistical significances for all metabolites, therefore group level results are described using effect sizes (Fig 4). This analysis showed that  $\text{K}^+$  reduced gradually from baseline before reaching peak declines at 120 (−0.93 ± 1.07  $\text{mmol}\cdot\text{L}^{-1}$ , −18%,  $g = 1.26$ ) and 150 min (−1.03 ± 0.93  $\text{mmol}\cdot\text{L}^{-1}$ , −20%,  $g = 1.64$ ) post-ingestion. Large effect sizes were calculated for the reductions in  $\text{Cl}^-$  from 105 to 165 min post-ingestion (all  $g > 0.80$ ) with the lowest point also occurring at 150 min post-ingestion (−3.5 ± 1.9  $\text{mmol}\cdot\text{L}^{-1}$ , −3%,  $p = 0.002$ ). The time that each individual participant reached their lowest values of  $\text{K}^+$  (133 ± 28 min, CV = 21%),  $\text{Cl}^-$  (124 ± 27 min, CV = 21%), and  $\text{Ca}^{2+}$  (141 ± 32 min, CV = 23%) averaged within 20 min of one another, albeit with large inter-individual variances. Conversely,  $\text{Na}^+$  was largely unchanged throughout the sampling period. The absolute changes from baseline displayed high inter-individual variation for each variable ( $\text{K}^+$ : −1.28 ± 0.88  $\text{mmol}\cdot\text{L}^{-1}$ , CV = 69%;  $\text{Cl}^-$ : −4.5 ± 2.1  $\text{mmol}\cdot\text{L}^{-1}$ , CV = 47%;  $\text{Ca}^{2+}$ : −0.16 ± 0.07  $\text{mmol}\cdot\text{L}^{-1}$ , CV = 43%;  $\text{Na}^+$ : +1.6 ± 1.3  $\text{mmol}\cdot\text{L}^{-1}$ , CV = 83%).

The energy and macronutrient composition of the pre-ingestion meal had no statistically significant correlations with the increases observed in blood  $\text{HCO}_3^-$ , pH, or the SID (Table 4). However, the macronutrient consumption of the meal did affect the time course behaviour of blood  $\text{HCO}_3^-$ , with protein ( $r_s = -0.581$ ) and fat intakes ( $r_s = -0.594$ ) both having moderate, negative correlations with time to peak  $\text{HCO}_3^-$ . Conversely, protein intake had a strong,




**Fig 3.** Group mean time course changes in A) blood bicarbonate ( $\text{HCO}_3^-$ ), B) pH, and C) the strong ion difference (SID) over a 165 min period following  $\text{NaHCO}_3$  ingestion. \* = a large effect size compared to baseline ( $g > 0.80$ ). # = a large effect size compared to 60 min post-ingestion ( $g > 0.80$ ).

<https://doi.org/10.1371/journal.pone.0248456.g003>

positive correlation ( $r_s = 0.711$ ) with time to peak SID. The total energy content of the pre-ingestion also had a moderate, positive correlation with time to peak SID, although this did not reach statistical significance ( $r_s = 0.555$ ,  $p = 0.061$ ). The fat content of the pre-ingestion meal was associated with baseline blood acid-base balance, as both  $\text{HCO}_3^-$  ( $r_s = 0.609$ ) and the SID ( $r_s = 0.639$ ) displayed strong, positive correlations. The amount of  $\text{Na}^+$  consumed 1–3 hours before  $\text{NaHCO}_3$  ingestion also had a strong, positive correlation with baseline  $\text{HCO}_3^-$  ( $r_s = 0.644$ ,  $p = 0.024$ ), but not with pH ( $r_s = -0.315$ ,  $p = 0.318$ ), or the SID ( $r_s = 0.171$ ,  $p = 0.596$ ). No other correlations were found between  $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Cl}^-$ , or  $\text{Ca}^{2+}$  ingestion and time course changes in blood acid-base balance.





**Fig 4.** Group mean time course changes in A) blood sodium ( $\text{K}^+$ ), B) potassium ( $\text{Na}^+$ ), C) calcium ( $\text{Ca}^{2+}$ ), and D) chloride ( $\text{Cl}^-$ ) over a 165 min period following  $\text{NaHCO}_3$  ingestion. \* = a large effect size compared to baseline ( $g > 0.80$ ). # = a large effect size compared to 60 min post-ingestion ( $g > 0.80$ ).

<https://doi.org/10.1371/journal.pone.0248456.g004>

### Gastrointestinal discomfort

All participants reported at least one GI side effect, though the severity of these instances was rated  $\leq 5/10$  in 94% (29/31) of occurrences (Table 5). Of the reported side effects, 71% (22/31)

**Table 4.** The associations between the composition of the pre-ingestion meal with the time course changes in acid-base balance variables following  $\text{NaHCO}_3$  ingestion.

Dietary intake 1–3 hours before $\text{NaHCO}_3$ ingestion	Correlations with time to peak		
	$\text{HCO}_3^-$ (min)	pH (min)	SID (min)
Energy (kcal)	$p = 0.202, r_s = -0.397$	$p = 0.785, r_s = -0.088$	$p = 0.061, r_s = 0.555$
Carbohydrate (g)	$p = 0.506, r_s = -0.213$	$p = 0.784, r_s = -0.088$	$p = 0.194, r_s = 0.403$
Protein (g)	<b><math>p = 0.048, r_s = -0.581^*</math></b>	$p = 0.870, r_s = -0.053$	<b><math>p = 0.010, r_s = 0.711^*</math></b>
Fat (g)	<b><math>p = 0.042, r_s = -0.594^*</math></b>	$p = 0.411, r_s = -0.262$	$p = 0.539, r_s = 0.197$
Dietary intake 1–3 hours before $\text{NaHCO}_3$ ingestion	Correlations with absolute change		
	$\text{HCO}_3^-$ (mmol·L <sup>-1</sup> )	pH	SID (mEq·L <sup>-1</sup> )
Energy (kcal)	$p = 0.430, r_s = 0.252$	$p = 0.084, r_s = 0.519$	$p = 0.827, r_s = 0.071$
Carbohydrate (g)	$p = 0.372, r_s = 0.284$	$p = 0.127, r_s = 0.466$	$p = 0.814, r_s = 0.076$
Protein (g)	$p = 0.476, r_s = 0.228$	$p = 0.054, r_s = 0.569$	$p = 0.221, r_s = 0.381$
Fat (g)	$p = 0.846, r_s = 0.063$	$p = 0.316, r_s = 0.316$	$p = 0.259, r_s = -0.354$
Dietary intake 1–3 hours before $\text{NaHCO}_3$ ingestion	Correlations with baseline concentrations		
	$\text{HCO}_3^-$ (mmol·L <sup>-1</sup> )	pH	SID (mEq·L <sup>-1</sup> )
Energy (kcal)	$p = 0.285, r_s = 0.336$	$p = 0.645, r_s = -0.149$	$p = 0.569, r_s = 0.171$
Carbohydrate (g)	$p = 0.652, r_s = 0.146$	$p = 0.801, r_s = -0.082$	$p = 0.920, r_s = 0.033$
Protein (g)	$p = 0.724, r_s = 0.114$	$p = 0.114, r_s = -0.480$	$p = 0.826, r_s = 0.071$
Fat (g)	<b><math>p = 0.036, r_s = 0.609^*</math></b>	$p = 1.000, r_s = <0.001$	<b><math>p = 0.025, r_s = 0.639^*</math></b>

<https://doi.org/10.1371/journal.pone.0248456.t004>

Table 5. The three most severe symptoms of gastrointestinal discomfort experienced following the ingestion of  $0.3 \text{ g kg BM}^{-1}$  sodium bicarbonate ( $\text{NaHCO}_3$ ).

Participant (Body mass)	Caps	Symptom 1	Severity (n/10)	Peak (mins)	Symptom 2	Severity (n/10)	Peak (mins)	Symptom 3	Severity (n/10)	Peak (mins)	Aggregated
1 (90.7 kg)	28	Belching	4.9	60	Stomach bloating	4.9	60	Nausea	2.3	60	17.7
2 (69.5 kg)	21	Belching	2.7	60	Stomach bloating	1.7	135	Stomach cramp	1.1	60	6.6
3 (55.8 kg)	17	Nausea	4.1	75	Stomach bloating	1.4	60	None	0.0	N/A	5.5
4 (62.2 kg)	19	Belching	3.9	60	Stomach bloating	2.4	60	Nausea	0.7	75	6.9
5 (72.1 kg)	22	Belching	1.7	75	None	0.0	N/A	None	0.0	N/A	1.7
6 (64.6 kg)	20	Stomach ache	4.0	165	None	0.0	N/A	None	0.0	N/A	4.0
7 (60.6 kg)	19	Nausea	1.4	60	None	0.0	N/A	None	0.0	N/A	1.4
8 (54.3 kg)	17	Belching	4.9	60	Flatulence	3.9	165	Stomach ache	1.5	135	11.0
9 (57.7 kg)	18	Nausea	2.6	60	None	0.0	N/A	None	0.0	N/A	2.6
10 (65.0 kg)	20	Belching	5.0	60	None	0.0	N/A	None	0.0	N/A	5.0
11 (67.9 kg)	21	Flatulence	2.6	150	Stomach bloating	2.1	60	Bowel urgency	1.5	165	6.1
12 (63.7 kg)	20	Vomiting	10.0	105	Nausea	10.0	105	Stomach ache	1.5	90	22.3

Aggregated = Sum of the most severe incidence of all nine symptoms (n/90). Caps = Number of capsules consumed (1 g of  $\text{NaHCO}_3$  per capsule).

<https://doi.org/10.1371/journal.pone.0248456.t005>

of instances peaked in severity at 60 and 75 min post-ingestion. The most common symptom was belching (58%, 7/12), followed by stomach bloating and nausea (50%, 6/12).

## Discussion

This was the first study to compare the time to peak of three different methods of measuring peak alkalosis (i.e.,  $\text{HCO}_3^-$ , pH, and the SID) following the ingestion of  $0.3 \text{ g kg BM}^{-1}$   $\text{NaHCO}_3$  in highly trained adolescent male and female swimmers. Though no statistical significance was observed between the three measures, there was a 33 min difference between the average time that  $\text{HCO}_3^-$  and the SID peaked in the blood of trained adolescents. This study also identified large inter-individual variances in the time to peak of each alkalotic measure, whereby individual peak concentrations were identified across the sampling timeframe (between 60–180 min post-ingestion). It is possible that this difference was related to the protein content of a meal consumed 1–3 hours before  $\text{NaHCO}_3$  ingestion, with this macronutrient positively and negatively correlated with time to peak SID and  $\text{HCO}_3^-$ , respectively. Finally, this study provides justification for utilising individualised  $\text{NaHCO}_3$  dosing strategies in adolescents since some sustained  $\text{HCO}_3^-$  increases exceeding  $5 \text{ mmol}\cdot\text{L}^{-1}$  for over 60 min ( $n = 6$ ), whereas others either reached this threshold for a 30–60 min period ( $n = 4$ ), or did not reach this threshold at all ( $n = 2$ ). Based on the individuality in blood responses, future work should aim to clarify whether personalised  $\text{NaHCO}_3$  strategies (i.e., time to peak  $\text{HCO}_3^-$  and time to peak SID) have a greater ergogenic effect for adolescents than standardised dosing procedures.

The average time that  $\text{HCO}_3^-$  and pH peaked within the blood of trained adolescents occurred within 10 min of one another, supporting the relationship between these two measurements when  $\text{NaHCO}_3$  is consumed in gelatine capsules [3, 22, 38]. This discussion will therefore focus on the blood  $\text{HCO}_3^-$  time course since this measure displays less variance and

greater reliability compared to pH [2, 21]. Time to peak blood  $\text{HCO}_3^-$  varied considerably within trained adolescents (75–180 min), which is in accordance with the blood responses of recreational adults following the ingestion of  $\text{NaHCO}_3$  capsules in a fed or fasted state [3, 21, 22]. Whereas no comparisons can presently be made for the time course changes in the SID, the time to peak concentration was also highly variable between each individual (60–165 min). Interestingly, the average time that the SID peaked in trained adolescents occurred 33 min before time to peak  $\text{HCO}_3^-$ , which may have implications for  $\text{NaHCO}_3$  timing in applied practice. Why this time difference occurred is currently unknown, though meal preference in the 1–3 hours prior to  $\text{NaHCO}_3$  ingestion may be a contributing factor. Indeed, higher protein intakes were correlated ( $r_s = 0.711$ ) with a delayed time to peak SID, whereas in contrast, higher protein intakes were also correlated ( $r_s = -0.581$ ) with a faster time to peak blood  $\text{HCO}_3^-$ . Since  $\text{NaHCO}_3$  will usually be supplemented with a pre-competition meal in practice, it is plausible that these two measurements will also be separated prior to performance. However, since this research did not measure a performance outcome, future work should aim to determine whether either of these individualised approaches (i.e., time to peak SID and  $\text{HCO}_3^-$ ) has an enhanced ergogenic benefit versus a traditional, fixed ingestion method (i.e., 120 min pre-exercise).

The participants in the present study demonstrated a peak blood  $\text{HCO}_3^-$  concentration that exceeded the 5–6  $\text{mmol}\cdot\text{L}^{-1}$  threshold ( $+6.8 \pm 2.4 \text{ mmol}\cdot\text{L}^{-1}$ ), contrasting previous studies when trained adolescents ingested  $\text{NaHCO}_3$  prior to exercise [34, 35]. A possible explanation for this difference is that high-level swimmers often have an early maturation onset [45], which likely occurred in the current participants and aided their selection for the high-performance swimming programme in which they were recruited. For comparison, both the male ( $72 \pm 11 \text{ kg}$ ) and female ( $60 \pm 5 \text{ kg}$ ) swimmers from the present study carried considerably more body mass than the ‘well trained’ male swimmers ( $56 \pm 1 \text{ kg}$ ) recruited by Zajac et al. [34], despite being the same age. Consequently, the current participants may have benefitted from maturity-related differences in gut physiology (i.e., caecal pH, distal microbiota) that potentially enhanced their capacity to uptake  $\text{NaHCO}_3$  from the GI tract [46, 47]. Moreover, the advanced training age of the participants could have also enhanced muscle and blood buffering capabilities (e.g., monocarboxylate transporter density,  $\text{Na}^+/\text{HCO}_3^-$  co-transport) [48–50], which could have enabled greater blood  $\text{HCO}_3^-$  increases to occur. However, it should be noted that two participants (both aged 15 years) did not increase their blood  $\text{HCO}_3^-$  concentrations above 3  $\text{mmol}\cdot\text{L}^{-1}$ , therefore adding support to previous speculations that less biologically mature adolescents might have less capacity increase blood  $\text{HCO}_3^-$  by more than 5–6  $\text{mmol}\cdot\text{L}^{-1}$ . Though it is unclear whether this range is necessary for performance enhancement in adolescents, it is perhaps appropriate to suggest that less biologically mature individuals do not require  $\text{NaHCO}_3$  supplementation at their stage of development [51].

Highly trained adolescents also demonstrated an increased blood SID following  $\text{NaHCO}_3$  ingestion. This supports contemporary evidence by Gough et al. [24, 27] whereby an elevated SID was reported prior to exercise in trained adults. Surprisingly, both studies by Gough et al. demonstrated greater group mean increases in the SID compared to the present study (+4 vs. +6–10  $\text{mEq}\cdot\text{L}^{-1}$ ), despite not purposely seeking to identify a peak concentration. The current study also observed lower baseline (33 vs. 36–38  $\text{mEq}\cdot\text{L}^{-1}$ ) and absolute peak SID concentrations (37 vs. 42–46  $\text{mEq}\cdot\text{L}^{-1}$ ), inferring that adolescents might also be limited with their capacity for strong ion movements (e.g., reduced  $\text{Na}^+/\text{K}^+$  pump activity) [48]. Furthermore, these lower concentrations occurred despite food consumption 1–3 hours before  $\text{NaHCO}_3$  ingestion, though the energy, macronutrient, or electrolyte ( $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Cl}^-$ , and  $\text{Ca}^{2+}$ ) composition of this meal was not correlated with baseline SID or the absolute SID increases that were achieved. Since water was also permitted to be consumed *ad libitum*, it is possible that plasma

volume also increased and masked some of the ionic movements that took place [52]. For example, the absolute increases observed in plasma  $\text{Na}^+$  were mitigated (+1.6 vs. +3.0  $\text{mmol}\cdot\text{L}^{-1}$ ) in comparison to previous research that investigated  $\text{NaHCO}_3$  ingestion in a fasted state [2, 3, 27]. Nonetheless, large effect sizes were calculated for plasma reductions of  $\text{K}^+$ ,  $\text{Ca}^{2+}$ , and  $\text{Cl}^-$  between 120–150 min post-ingestion, therefore suggesting that an intramuscular uptake of these ions did occur at this time [53]. Given that this study did not consider changes in the intramuscular SID or plasma volume, however, future research is required in order to further elucidate the role of  $\text{NaHCO}_3$  on the SID during exercise performance.

Comparable to adults, each blood analyte displayed a large inter-individual variance with regards to time to peak and absolute increases that occurred from baseline concentrations [2, 3, 22]. Why this phenomenon occurs remains elusive, though it is postulated that genetic differences in the rate of gastric emptying, intestinal motility, and GI blood flow may contribute to  $\text{NaHCO}_3$  uptake characteristics [54, 55]. Despite the individual differences, the process of identifying a personal time to peak  $\text{HCO}_3^-$  measurement has recently been criticised [22]. Ingesting  $\text{NaHCO}_3$  capsules has consistently been shown to produce increases in blood  $\text{HCO}_3^-$  (>5  $\text{mmol}\cdot\text{L}^{-1}$ ) that last for over 100 min in adults [3, 22, 23], therefore producing a long lasting ‘ergogenic window’ that undermines the necessity of a single ingestion time point. However, when considering the  $\text{HCO}_3^-$  time course on an individual basis, only half ( $n = 6$ ) of the trained adolescent participants demonstrated a +5  $\text{mmol}\cdot\text{L}^{-1}$   $\text{HCO}_3^-$  increase that lasted for more than 90 min. Moreover, the time that each participant spent above the +5  $\text{mmol}\cdot\text{L}^{-1}$  threshold occurred at completely separate periods within some individuals (i.e., 60–120 min, 135–165 min post-ingestion), whereas two participants had small reductions in blood  $\text{HCO}_3^-$  (~1  $\text{mmol}\cdot\text{L}^{-1}$ ) at 60–75 min post-ingestion before concentrations increased over the next 60–75 min. Due to these erratic time course differences, it is recommended that adolescents undergo time to peak testing prior to supplementation in order to identify a personal time frame for  $\text{NaHCO}_3$  to elicit a sufficient blood  $\text{HCO}_3^-$  response. However, considering that this is an expensive procedure, adolescents should remain cautious since the effects of individualised ingestion methods on exercise performance are currently untested within this cohort.

There were minimal incidents of severe GI side effects when trained adolescents ingested  $\text{NaHCO}_3$  capsules 1–3 hours after consuming a self-selected meal. Whilst each participant reported at least one side effect, the severity of these symptoms was rated less than 5 out of 10 on 94% (29/31) of occasions. This low level of GI disturbance therefore supports the findings of Carr et al. [38], whereby the co-ingestion of  $\text{NaHCO}_3$  capsules with a meal provided the best mitigation against GI distress (vs.  $\text{NaHCO}_3$  solution/fasted state). Both capsules and meal consumption are suggested to slow the reduction of  $\text{NaHCO}_3$  to  $\text{Na}^+$  and carbonic acid ( $\text{H}_2\text{CO}_3$ ), subsequently reducing the rate of acid-base disturbances that occur in the stomach (e.g., increases in carbon dioxide and water) [56, 57]. Alternatively, the age of the participants might have contributed to the lack of GI symptoms observed in this study, therefore supporting the lack of GI disturbances in adolescents from previous  $\text{NaHCO}_3$  research [34, 35]. Whilst there is currently no clear explanation for the lack of GI disturbances in adolescents, it is plausible that a lower acidity of the GI fluids compared to adults may mediate the acid-base disturbances that occur in the stomach when an alkaline substance is ingested [58]. Another general observation is that athletes of a higher body mass (>75 kg) typically display greater GI distress than lighter individuals [42, 59]. Adolescents, being smaller in stature than their adult peers, may therefore experience less GI disturbances due to a reduced absolute  $\text{NaHCO}_3$  dosage entering the stomach. However, this study did observe vomiting in one participant, which indicates that the severity GI side effects should be assessed on an individual basis.

The study methods were applied based on the logistical (e.g., taking highly trained swimmers out of training) and ethical considerations of the participant cohort (e.g., repeated blood

analysis of young athletes). This limited the current study from including a placebo experiment or repeatability measures that could have altered the interpretation of results. Nonetheless, previous work found a placebo to cause no blood time course changes for either  $\text{HCO}_3^-$ , pH, or  $\text{Na}^+$  [2, 22], therefore it may be assumed that a placebo would not have altered baseline  $\text{HCO}_3^-$ , pH, or SID within an adolescent cohort. The repeatability of time to peak  $\text{HCO}_3^-$  is controversial, however, when  $0.3 \text{ g}\cdot\text{kg BM}^{-1}$   $\text{NaHCO}_3$  capsules are ingested in a fed state. Indeed, Boegman et al. [21] reported an excellent repeatability ( $\text{ICC} = 0.77$ ) within world-class rowers when  $\text{NaHCO}_3$  was ingested alongside a standardised snack (energy: 682 kcal; protein: 20 g; carbohydrate: 140 g). Conversely, Oliveira et al. [22] found repeatability to be poor ( $\text{ICC} = 0.34$ ) when recreational adults ingested  $\text{NaHCO}_3$  60 min following a standardised breakfast (energy: 563 kcal; protein: 9 g; carbohydrate: 90 g). Since both studies differed in meal composition, participant training status, time post-prandial, and blood analysis methods (capillary [21] vs. venous [22] samples), this leaves the repeatability of the current data unclear. However, the main purpose of this research was to identify the time course changes in three alternate blood acid-base balance variables following  $\text{NaHCO}_3$  ingestion in a highly trained adolescent cohort. Since this was the first study to suggest that a difference in time to peak  $\text{HCO}_3^-$  and SID may exist in practice, future studies should therefore aim to determine the repeatability of these two measures under different competitive scenarios (e.g., pre-competition meal, after warm-up).

## Conclusion

This study shows that highly trained adolescent swimmers have significant increases and highly individual time course changes in blood  $\text{HCO}_3^-$ , pH, and SID following the ingestion of  $\text{NaHCO}_3$  capsules. Importantly, the post-ingestion time point in which the individual time to peak occurred for  $\text{HCO}_3^-$  and the SID was separated by a large effect size. Given that the effects of acidosis on exercise performance are controversial, future studies should seek to identify  $\text{NaHCO}_3$  dosing strategies based upon a peak SID concentration could be more ergogenic than standardised ingestion strategies, or individualised approaches based upon a time to peak blood  $\text{HCO}_3^-$ . This study also suggests that highly trained adolescent athletes do not have a long lasting 'ergogenic window' following the ingestion of  $\text{NaHCO}_3$  capsules, therefore supporting the practice of identifying an individualised time point whereby a peak alkalosis occurs. It is acknowledged that a lack of control surrounding pre-ingestion food and fluids may have influenced the findings of this study, however, these conditions better replicate the actual changes that would occur when athletes are in full competition preparation.

## Supporting information

**S1 Data.**  
(XLSX)

## Acknowledgments

We gratefully acknowledge all participants who volunteered for the study and the City of Birmingham Swimming Club for facilitating the research process.

## Author Contributions

**Conceptualization:** Lars R. McNaughton.

**Data curation:** Josh W. Newbury.

**Formal analysis:** Josh W. Newbury.

**Funding acquisition:** Lewis A. Gough.

**Methodology:** Josh W. Newbury, Lewis A. Gough.

**Project administration:** Adam L. Kelly, Richard J. Chessor, Lewis A. Gough.

**Supervision:** Richard J. Chessor, S. Andy Sparks, Lars R. McNaughton, Lewis A. Gough.

**Writing – original draft:** Josh W. Newbury, Richard J. Chessor, Lewis A. Gough.

**Writing – review & editing:** Josh W. Newbury, Matthew Cole, Adam L. Kelly, Richard J. Chessor, S. Andy Sparks, Lars R. McNaughton, Lewis A. Gough.

## References

1. Maughan RJ, Burke LM, Dvorak J, Larson-Meyer DE, Peeling P, Phillips SM, et al. IOC consensus statement: dietary supplements and the high-performance athlete. *Int J Sport Nutr Exerc Metab.* 2018 Mar 1; 28(2): 104–125. <https://doi.org/10.1123/ijsnem.2018-0020> PMID: 29589768
2. Gough LA, Deb SK, Sparks AS, McNaughton LR. The reproducibility of blood acid base responses in male collegiate athletes following individualised doses of sodium bicarbonate: a randomised controlled crossover study. *Sports Med.* 2017 Oct; 47(10): 2117–2127. <https://doi.org/10.1007/s40279-017-0699-x> PMID: 28229390
3. Jones RL, Stellingwerff T, Artioli GG, Saunders B, Cooper S, Sale C. Dose-response of sodium bicarbonate ingestion highlights individuality in time course of blood analyte responses. *Int J Sport Nutr Exerc Metab.* 2016 Oct; 26(5): 445–453. <https://doi.org/10.1123/ijsnem.2015-0286> PMID: 27098290
4. Bishop D, Edge J, Davis C, Goodman C. Induced metabolic alkalosis affects muscle metabolism and repeated-sprint ability. *Med Sci Sports Exerc.* 2004 May; 36(5): 807–813. <https://doi.org/10.1249/01.mss.0000126392.20025.17> PMID: 15126714
5. Mainwood GW, Worsley-Brown P. The effects of extracellular pH and buffer concentration on the efflux of lactate from frog sartorius muscle. *J Physiol.* 1975 Aug; 250(1): 1–22. <https://doi.org/10.1113/jphysiol.1975.sp011040> PMID: 16992502
6. Fitts RH. The cross-bridge cycle and skeletal muscle fatigue. *J Appl Physiol.* 2008 Feb; 104(2): 551–558. <https://doi.org/10.1152/japplphysiol.01200.2007> PMID: 18162480
7. Fitts RH. The Role of Acidosis in Fatigue: Pro Perspective. *Med Sci Sports Exerc.* 2016 Nov; 48(11): 2335–2338. <https://doi.org/10.1249/MSS.0000000000001043> PMID: 27755382
8. Knuth ST, Dave H, Peters JR, Fitts RH. Low cell pH depresses peak power in rat skeletal muscle fibres at both 30 degrees C and 15 degrees C: implications for muscle fatigue. *J Physiol.* 2006 Sep 15; 575(Pt 3): 887–899. <https://doi.org/10.1113/jphysiol.2006.106732> PMID: 16809373
9. Sundberg CW, Fitts RH. Bioenergetic basis of skeletal muscle fatigue. *Curr Opin Physiol.* 2019 Aug; 10: 118–127. <https://doi.org/10.1016/j.cophys.2019.05.004> PMID: 31342000
10. Woodward M, Debold EP. Acidosis and phosphate directly reduce myosin's force-generating capacity through distinct molecular mechanisms. *Front Physiol.* 2018 Jul 10; 9: 862. <https://doi.org/10.3389/fphys.2018.00862> PMID: 30042692
11. Debold EP, Beck SE, Warshaw DM. Effect of low pH on single skeletal muscle myosin mechanics and kinetics. *Am J Physiol Cell Physiol.* 2008 Jul; 295(1): C173–C179. <https://doi.org/10.1152/ajpcell.00172.2008> PMID: 18480297
12. Hollidge-Horvat MG, Parolin ML, Wong D, Jones NL, Heigenhauser GJ. Effect of induced metabolic acidosis on human skeletal muscle metabolism during exercise. *Am J Physiol.* 1999 Oct; 277(4): E647–E658. <https://doi.org/10.1152/ajpendo.1999.277.4.E647> PMID: 10516124
13. Spriet LL, Lindinger MI, McKelvie RS, Heigenhauser GJ, Jones NL. Muscle glycogenolysis and H<sup>+</sup> concentration during maximal intermittent cycling. *J Appl Physiol.* 1989 Jan; 66(1): 8–13. <https://doi.org/10.1152/jappl.1989.66.1.8> PMID: 2917960
14. Street D, Nielsen JJ, Bangsbo J, Juel C. Metabolic alkalosis reduces exercise-induced acidosis and potassium accumulation in human skeletal muscle interstitium. *J Physiol.* 2005 Jul 15; 566(Pt 2): 481–489. <https://doi.org/10.1113/jphysiol.2005.086801> PMID: 15860529
15. Fitts RH. Cellular mechanisms of muscle fatigue. *Physiol Rev.* 1994 Jan; 74(1): 49–94. <https://doi.org/10.1152/physrev.1994.74.1.49> PMID: 8295935



16. Allen DG, Lamb GD, Westerblad H. Skeletal muscle fatigue: cellular mechanisms. *Physiol Rev*. 2008 Jan; 88(1): 287–332. <https://doi.org/10.1152/physrev.00015.2007> PMID: 18195089
17. Cairns SP. Lactic acid and exercise performance: culprit or friend? *Sports Med*. 2006; 36(4): 279–291. <https://doi.org/10.2165/00007256-200636040-00001> PMID: 16573355
18. Westerblad H. Acidosis is not a significant cause of skeletal muscle fatigue. *Med Sci Sports Exerc*. 2016 Nov 1; 48(11): 2339–2342. <https://doi.org/10.1249/MSS.0000000000001044> PMID: 27755383
19. Carr AJ, Hopkins WG, Gore CJ. Effects of acute alkalosis and acidosis on performance: a meta-analysis. *Sports Med*. 2011 Oct 1; 41(10): 801–814. <https://doi.org/10.2165/11591440-000000000-00000> PMID: 21923200
20. Heibel AB, Perim PHL, Oliveira LF, McNaughton LR, Saunders B. Time to optimize supplementation: modifying factors influencing the individual responses to extracellular buffering agents. *Front Nutr*. 2018 May 8; 5: 35. <https://doi.org/10.3389/fnut.2018.00035> PMID: 29868599
21. Boegman S, Stellingwerff T, Shaw G, Clarke N, Graham K, Cross R, et al. The impact of individualizing sodium bicarbonate supplementation strategies on world-class rowing performance. *Front Nutr*. 2020 Sep 9; 7: 138. <https://doi.org/10.3389/fnut.2020.00138> PMID: 33015117
22. Oliveira LF, Saunders B, Yamaguchi G, Swinton P, Artioli GG. Is individualization of sodium bicarbonate ingestion based on time to peak necessary? *Med Sci Sports Exerc*. 2020 Aug; 52(8): 1801–1808. <https://doi.org/10.1249/MSS.0000000000002313> PMID: 32102054
23. Deb SK, Gough LA, Sparks SA, McNaughton LR. Sodium bicarbonate supplementation improves severe-intensity intermittent exercise under moderate acute hypoxic conditions. *Eur J Appl Physiol*. 2018 Mar; 118(3): 607–615. <https://doi.org/10.1007/s00421-018-3801-7> PMID: 29344729
24. Gough LA, Deb SK, Brown D, Sparks SA, McNaughton LR. The effects of sodium bicarbonate ingestion on cycling performance and acid base balance recovery in acute normobaric hypoxia. *J Sports Sci*. 2019 Jul; 37(13): 1464–1471. <https://doi.org/10.1080/02640414.2019.1568173> PMID: 30668281
25. Gough LA, Deb SK, Sparks SA, McNaughton LR. Sodium bicarbonate improves 4 km time trial cycling performance when individualised to time to peak blood bicarbonate in trained male cyclists. *J Sports Sci*. 2018 Aug; 36(15): 1705–1712. <https://doi.org/10.1080/02640414.2017.1410875> PMID: 29183257
26. Siegler JC, Marshall PW, Bishop D, Shaw G, Green S. Mechanistic insights into the efficacy of sodium bicarbonate supplementation to improve athletic performance. *Sports Med Open*. 2016 Dec; 2(1): 41. <https://doi.org/10.1186/s40798-016-0065-9> PMID: 27747796
27. Gough LA, Rimmer S, Sparks SA, McNaughton LR, Higgins MF. Post-exercise supplementation of sodium bicarbonate improves acid base balance recovery and subsequent high-intensity boxing specific performance. *Front Nutr*. 2019 Oct 1; 6: 155. <https://doi.org/10.3389/fnut.2019.00155> PMID: 31632978
28. Siegler JC, Gleadow-Siddall DO. Sodium bicarbonate ingestion and repeated swim sprint performance. *J Strength Cond Res*. 2010 Nov; 24(11): 3105–3111. <https://doi.org/10.1519/JSC.0b013e3181f55eb1> PMID: 20881504
29. Siegler JC, Hirscher K. Sodium bicarbonate ingestion and boxing performance. *J Strength Cond Res*. 2010 Jan; 24(1): 103–108. <https://doi.org/10.1519/JSC.0b013e3181a392b2> PMID: 19625976
30. Sostaric SM, Skinner SL, Brown MJ, Sangkabutra T, Medved I, Medley T, et al. Alkalosis increases muscle  $\text{K}^+$  release, but lowers plasma  $[\text{K}^+]$  and delays fatigue during dynamic forearm exercise. *J Physiol*. 2006 Jan 1; 570(Pt 1): 185–205. <https://doi.org/10.1113/jphysiol.2005.094615> PMID: 16239279
31. Cairns SP, Lindinger MI. Do multiple ionic interactions contribute to skeletal muscle fatigue? *J Physiol*. 2008 Sep 1; 586(17): 4039–4054. <https://doi.org/10.1113/jphysiol.2008.155424> PMID: 18591187
32. Monteiro D, Cid L, Marinho DA, Moutão J, Vitorino A, Bento T. Determinants and reasons for dropout in swimming—systematic review. *Sports*. 2017 Jul 10; 5(3): 50. <https://doi.org/10.3390/sports5030050> PMID: 29910410
33. Jovanov P, Đorđić V, Obradović B, Barak O, Pezo L, Marić A, et al. Prevalence, knowledge and attitudes towards using sports supplements among young athletes. *J Int Soc Sports Nutr*. 2019 Jul 4; 16(1): 27. <https://doi.org/10.1186/s12970-019-0294-7> PMID: 31272457
34. Zajac A, Cholewa J, Poprzecki S, Waskiewicz Z, Langfort J. Effects of sodium bicarbonate ingestion on swim performance in youth athletes. *J Sports Sci Med*. 2009 Mar 1; 8(1): 45–50. PMID: 24150555
35. Guimarães RDS, Morais Junior AC, Schincaglia RM, Saunders B, Pimentel GD, Mota JF. Sodium bicarbonate supplementation does not improve running anaerobic sprint test performance in semi-professional adolescent soccer players. *Int J Sport Nutr Exerc Metab*. 2020 Jul 15; 30(5): 330–337. <https://doi.org/10.1123/ijnsnem.2020-0031> PMID: 32668408
36. Sparks A, Williams E, Robinson A, Miller P, Bentley DJ, Bridge C, et al. Sodium bicarbonate ingestion and individual variability in time-to-peak pH. *Res Sports Med*. 2017 Jan-Mar; 25(1): 58–66. <https://doi.org/10.1080/15438627.2016.1258645> PMID: 27934546

37. Remer T, Manz F. Potential renal acid load of foods and its influence on urine pH. *J Am Diet Assoc*. 1995 Jul; 95(7): 791–797. [https://doi.org/10.1016/S0002-8223\(95\)00219-7](https://doi.org/10.1016/S0002-8223(95)00219-7) PMID: 7797810
38. Carr AJ, Slater GJ, Gore CJ, Dawson B, Burke LM. Effect of sodium bicarbonate on  $[\text{HCO}_3^-]$ , pH, and gastrointestinal symptoms. *Int J Sport Nutr Exerc Metab*. 2011 Jun; 21(3): 189–194. <https://doi.org/10.1123/ijsnem.21.3.189> PMID: 21719899
39. Hobson RM, Harris RC, Martin D, Smith P, Macklin B, Gualano B, et al. Effect of beta-alanine, with and without sodium bicarbonate, on 2000-m rowing performance. *Int J Sport Nutr Exerc Metab*. 2013 Oct; 23(5): 480–487. <https://doi.org/10.1123/ijsnem.23.5.480> PMID: 23535873
40. Sale C, Saunders B, Hudson S, Wise JA, Harris RC, Sunderland CD. Effect of  $\beta$ -alanine plus sodium bicarbonate on high-intensity cycling capacity. *Med Sci Sports Exerc*. 2011 Oct; 43(10): 1972–1978. <https://doi.org/10.1249/MSS.0b013e3182188501> PMID: 21407127
41. Lloyd P. Strong ion calculator—a practical bedside application of modern quantitative acid-base physiology. *Crit Care Resusc*. 2004 Dec; 6(4): 285–294. PMID: 16556109
42. Cameron SL, McLay-Cooke RT, Brown RC, Gray AR, Fairbairn KA. Increased blood pH but not performance with sodium bicarbonate supplementation in elite rugby union players. *Int J Sport Nutr Exerc Metab*. 2010 Aug; 20(4): 307–321. <https://doi.org/10.1123/ijsnem.20.4.307> PMID: 20739719
43. Lakens D. Calculating and reporting effect sizes to facilitate cumulative science: a practical primer for t-tests and ANOVAs. *Front Psychol*. 2013 Nov 26; 4: 863. <https://doi.org/10.3389/fpsyg.2013.00863> PMID: 24324449
44. Cohen J. *Statistical power analysis for the behavioral sciences*. Hillsdale, NJ: L. Erlbaum Associates; 1988.
45. Baxter-Jones AD, Helms PJ. Effects of training at a young age: a review of the training of young athletes (TOYA) study. *Pediatr Exerc Sci*. 1996 Nov 1; 8(4): 310–327. <https://doi.org/10.1123/pes.8.4.310>
46. Agans R, Rigsbee L, Kenche H, Michail S, Khamis HJ, Paliy O. Distal gut microbiota of adolescent children is different from that of adults. *FEMS Microbiol Ecol*. 2011 Aug; 77(2): 404–412. <https://doi.org/10.1111/j.1574-6941.2011.01120.x> PMID: 21539582
47. Bai JPF, Burckart GJ, Mulberg AE. Literature review of gastrointestinal physiology in the elderly, in pediatric patients, and in patients with gastrointestinal diseases. *J Pharm Sci*. 2016 Feb; 105(2): 476–483. <https://doi.org/10.1002/jps.24696> PMID: 26539698
48. Juel C. Regulation of pH in human skeletal muscle: adaptations to physical activity. *Acta Physiol*. 2008 May; 193(1): 17–24. <https://doi.org/10.1111/j.1748-1716.2008.01840.x> PMID: 18267000
49. McNaughton LR, Gough L, Deb S, Bentley D, Sparks SA. Recent developments in the use of sodium bicarbonate as an ergogenic aid. *Curr Sports Med Rep*. 2016 Jul-Aug; 15(4): 233–244. <https://doi.org/10.1249/JSR.000000000000283> PMID: 27399820
50. Peart DJ, Siegler JC, Vince RV. Practical recommendations for coaches and athletes: a meta-analysis of sodium bicarbonate use for athletic performance. *J Strength Cond Res*. 2012 Jul; 26(7): 1975–1983. <https://doi.org/10.1519/JSC.0b013e3182576f3d> PMID: 22505127
51. Desbrow B, McCormack J, Burke LM, Cox GR, Fallon K, Hislop M, et al. Sports Dietitians Australia position statement: sports nutrition for the adolescent athlete. *Int J Sport Nutr Exerc Metab*. 2014 Oct; 24(5): 570–584. <https://doi.org/10.1123/ijsnem.2014-0031> PMID: 24668620
52. Brown WJ Jr, Brown FK, Krishan I. Exchangeable sodium and blood volume in normotensive and hypertensive humans on high and low sodium intake. *Circulation*. 1971 Apr; 43(4): 508–519. <https://doi.org/10.1161/01.cir.43.4.508> PMID: 4929772
53. Lühker O, Berger MM, Pohlmann A, Hotz L, Gruhlke T, Hochreiter M. Changes in acid-base and ion balance during exercise in normoxia and normobaric hypoxia. *Eur J Appl Physiol*. 2017 Nov; 117(11): 2251–2261. <https://doi.org/10.1007/s00421-017-3712-z> PMID: 28914359
54. Heading RC, Nimmo J, Prescott LF, Tothill P. The dependence of paracetamol absorption on the rate of gastric emptying. *Br J Pharmacol*. 1973 Feb; 47(2): 415–421. <https://doi.org/10.1111/j.1476-5381.1973.tb08339.x> PMID: 4722050
55. Paintaud G, Thibault P, Queneau PE, Magnette J, Berard M, Rumbach L, et al. Intraindividual variability of paracetamol absorption kinetics after a semi-solid meal in healthy volunteers. *Eur J Clin Pharmacol*. 1998 Jan; 53(5): 355–359. <https://doi.org/10.1007/s002280050393> PMID: 9516037
56. Heigenhauser GJF. Ergogenic enhancement of performance in exercise and sport: bicarbonate loading. In: Lamb DR, Williams MH, editors. *Prospective in Exercise Science and Sports Medicine*, Vol. 4. Carmel, IN: Cooper Publishing Group LLC; 1991. pp.183–212.
57. Turnberg LA, Fordtran JS, Carter NW, Rector FC. Mechanism of bicarbonate absorption and its relationship to sodium transport in the human jejunum. *J Clin Invest*. 1970 Mar 1; 49(3): 548–556. <https://doi.org/10.1172/JCI106265> PMID: 5415681

58. Merchant HA, Liu F, Orlu Gul M, Basit AW. Age-mediated changes in the gastrointestinal tract. *Int J Pharm.* 2016 Oct 30; 512(2): 382–395. <https://doi.org/10.1016/j.ijpharm.2016.04.024> PMID: 27085646
59. Kahle LE, Kelly PV, Eliot KA, Weiss EP. Acute sodium bicarbonate loading has negligible effects on resting and exercise blood pressure but causes gastrointestinal distress. *Nutr Res.* 2013 Jun; 33(6): 479–486. <https://doi.org/10.1016/j.nutres.2013.04.009> PMID: 23746564