

Article

The Relationship between Body Mass Index and Mammographic Density during a Premenopausal Weight Loss Intervention Study

Emma C. Atakpa¹, Adam R. Brentnall¹, Susan Astley^{2,3,4}, Jack Cuzick¹, D. Gareth Evans^{2,3,5,6,7} ,
Ruth M. L. Warren^{8,9}, Anthony Howell^{2,3,10} and Michelle Harvie^{2,*} 

- ¹ Centre for Cancer Prevention, Wolfson Institute of Preventive Medicine, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London EC1M 6BQ, UK; e.c.atakpa@qmul.ac.uk (E.C.A.); a.brentnall@qmul.ac.uk (A.R.B.); j.cuzick@qmul.ac.uk (J.C.)
 - ² Nightingale Breast Screening Centre & Prevent Breast Cancer Unit, Wythenshawe Hospital, Manchester University NHS Foundation Trust, Manchester M23 9LT, UK; Sue.astley@manchester.ac.uk (S.A.); Gareth.Evans@mft.nhs.uk (D.G.E.); anthony.howell@manchester.ac.uk (A.H.)
 - ³ Manchester Breast Centre, The Christie Hospital, Manchester M23 9LT, UK
 - ⁴ Division of Informatics, Imaging & Data Sciences, School of Health Sciences, Faculty of Biology, Medicine and Health, University of Manchester, Manchester M13 9PT, UK
 - ⁵ Manchester Centre for Genomic Medicine, Manchester University Hospitals NHS Foundation Trust, Manchester M23 9LT, UK
 - ⁶ Manchester Centre for Genomic Medicine, NW Genomic Laboratory Hub, Manchester University Hospitals NHS Foundation Trust, Manchester M13 9WL, UK
 - ⁷ Manchester Academic Health Science Centre, Division of Evolution and Genomic Sciences, Faculty of Biology, School of Biological Sciences, Medicine and Health, University of Manchester, Manchester M23 9LT, UK
 - ⁸ Cambridge Breast Unit, Addenbrooke's Hospital, Cambridge CB2 0QQ, UK; rmlw2@cam.ac.uk
 - ⁹ Girton College, University of Cambridge, Cambridge CB3 0JG, UK
 - ¹⁰ Manchester Academic Health Science Centre, Division of Cancer Sciences, Medicine and Health, University of Manchester, Manchester M23 9LT, UK
- * Correspondence: michelle.harvie@manchester.ac.uk



Citation: Atakpa, E.C.; Brentnall, A.R.; Astley, S.; Cuzick, J.; Evans, D.G.; Warren, R.M.L.; Howell, A.; Harvie, M. The Relationship between Body Mass Index and Mammographic Density during a Premenopausal Weight Loss Intervention Study. *Cancers* **2021**, *13*, 3245. <https://doi.org/10.3390/cancers13133245>

Academic Editors: Ranjit Manchanda and Andrea Manni

Received: 7 April 2021
Accepted: 18 June 2021
Published: 29 June 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Simple Summary: This study assessed the association between short-term weight change and mammographic density in premenopausal women losing weight through diet and exercise to reduce their risk of postmenopausal breast cancer. We aimed to understand whether a reduction in body mass index affects various components of the breast, which could indicate a potential pathway for the reduction in postmenopausal breast cancer risk seen with premenopausal weight loss. Understanding this pathway is useful for monitoring the effectiveness of prevention strategies based on lifestyle advice. We found that a short-term reduction in premenopausal body mass index through diet and exercise is associated with a reduction in breast fat, but it is unlikely to have a significant effect on the quantity of breast glandular tissue. Breast cancer risk determined by changes in breast density might not capture potential weight loss-induced breast cancer risk reduction, instead falsely ascribing an increased risk due to increased percent density.

Abstract: We evaluated the association between short-term change in body mass index (BMI) and breast density during a 1 year weight-loss intervention (Manchester, UK). We included 65 premenopausal women (35–45 years, ≥ 7 kg adult weight gain, family history of breast cancer). BMI and breast density (semi-automated area-based, automated volume-based) were measured at baseline, 1 year, and 2 years after study entry (1 year post intervention). Cross-sectional (between-women) and short-term change (within-women) associations between BMI and breast density were measured using repeated-measures correlation coefficients and multivariable linear mixed models. BMI was positively correlated with dense volume between-women ($r = 0.41$, 95%CI: 0.17, 0.61), but less so within-women ($r = 0.08$, 95%CI: -0.16 , 0.28). There was little association with dense area (between-women $r = -0.12$, 95%CI: -0.38 , 0.16; within-women $r = 0.01$, 95%CI: -0.24 , 0.25). BMI and breast fat were positively correlated (volume: between $r = 0.77$, 95%CI: 0.69, 0.84, within $r = 0.58$, 95%CI: 0.36, 0.75; area: between $r = 0.74$, 95%CI: 0.63, 0.82, within $r = 0.45$, 95%CI: 0.23, 0.63). Multivariable

models reported similar associations. Exploratory analysis suggested associations between BMI gain from 20 years and density measures (standard deviation change per +5 kg/m² BMI: dense area: +0.61 (95%CI: 0.12, 1.09); fat volume: −0.31 (95%CI: −0.62, 0.00)). Short-term BMI change is likely to be positively associated with breast fat, but we found little association with dense tissue, although power was limited by small sample size.

Keywords: mammographic density; body mass index; weight loss; breast cancer risk; breast cancer prevention; premenopausal

1. Introduction

Mammographic density (herein referred to as ‘density’) is an established risk factor for breast cancer. Women in the highest density category are at a 4- to 6-fold increased risk of breast cancer relative to those with little or no dense tissue [1]. When assessed by mammography, the breast is broadly characterised by two components: fibroglandular dense tissue and fatty non-dense tissue. Percent breast density is measured as the relative proportion of dense tissue in the breast, either in terms of area or volume depending on the measurement method. Visual assessment measures percent density with respect to the total breast area (TA) whilst automated and semi-automated methods can also measure the extent of dense and fatty tissue separately. Both absolute dense area (DA) and percentage dense area (PDA) are positively associated with risk of premenopausal (and postmenopausal) breast cancer [2–4], and absolute dense volume (DV) and percentage dense volume (PDV) have also shown positive associations [5,6]. Associations of breast fat area (FA) and volume (FV) with breast cancer risk are unclear, although there is some suggestion of an inverse relationship with premenopausal breast cancer risk [4,6].

In postmenopausal women, higher attained body mass index (BMI) is associated with a higher risk of breast cancer [7–9], with an estimated 40% increase in risk for every 10 kg/m² of BMI in never users of hormone replacement therapy [9]. This increase in risk is partly explained by increased aromatisation of androgens to oestrogen in peripheral adipose tissue, which promotes cell proliferation [10,11], carcinogenesis [10,11], and insulin resistance [12]. Whilst BMI is a widely accepted risk factor for breast cancer in postmenopausal women, there may be an inverse relationship in premenopausal women [13].

Weight gain across the premenopausal years has also been linked to an increased risk of postmenopausal breast cancer. Every 5 kg of adult weight gain is associated with an approximate 10% increase in risk amongst never or low-hormone replacement therapy users [14,15]. However, a number of studies (as summarised by Hardefeldt et al. [16]) suggest that these effects are reversible with efficient weight loss [16]. In particular, weight loss in the premenopausal years has been shown to reduce postmenopausal breast cancer risk [17,18]. Risk reductions of approximately 40% have also been seen with large weight losses as a result of bariatric surgery in populations of pre- and postmenopausal women [19].

The effects of short-term weight change on breast density are less well understood, particularly those as a result of dietary weight loss. Mammographic density is a dynamic phenotype and has the potential to respond to short-term weight changes, making density reduction a possible biomarker for reduction in risk as a result of weight loss. This study aims to explore the effect of short-term dietary weight change on density using both area-based and volumetric methods in a cohort of premenopausal women to ascertain whether the relationship between weight loss and reduced postmenopausal breast cancer risk could, in part, be mediated by reductions in mammographic tissue.

2. Materials and Methods

2.1. Study Design and Participants

The Lifestyle Study is a prospective non-randomised 1 year diet and exercise weight loss intervention study amongst 79 high-risk premenopausal women attending annual screening within the Breast Cancer Family History clinic at the Prevent Breast Cancer research unit at the Manchester University Hospital Foundation NHS Trust [20–23]. Attendees of our regional Family History Clinic, aged 35–45 years, received a mailed invitation to enter either a 12-month intensive diet and exercise weight loss programme or a usual care group receiving standard written advice only, depending on their proximity to the hospital. Eligibility required women to be premenopausal with regular menstrual cycles, non-smokers, have a self-reported adult weight gain ≥ 7 kg, and a sedentary lifestyle (<40 min moderate physical activity per week). All women had a family history of breast cancer (with lifetime risk 17–40% as assessed by the Tyrer–Cuzick model [24,25]), but were excluded if they had a known *BRCA1/2* mutation or a previous history of cancer. Women were also excluded if they were already successfully dieting or losing weight, were pregnant or planning to become pregnant over the next year, had used hormonal oral contraceptives in the last six months, or had psychiatric or physical co-morbidities that could affect their ability to take part in a diet and physical activity weight loss programme.

In the intervention group ($n = 40$), women followed a 12-month intensive supervised weight loss programme involving a 25% energy-restricted Mediterranean type diet and an individualised physical activity program (150 min moderate intensity physical activity and 40 min of resistance exercise per week). The usual care group ($n = 39$) received standard written advice about diet and physical activity but no additional support for weight loss. Women provided baseline information on alcohol intake (from a 4-day food diary) and physical activity (7-day recall from an interview questionnaire) at their baseline clinic visit. All subjects gave their informed consent for inclusion before they participated in the study. The study was conducted in accordance with the Declaration of Helsinki, and the protocol was approved by the South Manchester Ethics Committee (Reference no. 01/426).

The objective of this analysis was to assess the relationship between BMI and breast density in the entire cohort of women. All participants had changing BMI measures irrespective of the type of weight loss advice they received, hence the intervention and usual care groups were combined and treated as one cohort. Furthermore, to limit the effect of women contributing observations to an area-based measure or volumetric measure only, the cohort was restricted to those with both an area and volumetric density measurement at any one or more time points ($n = 65$, 82% of the cohort).

2.2. Mammographic Density

Mammographic films were digitised using a Kodak LS85 digitiser at a pixel size of 50 μm and with 12-bits (4096 grey levels) pixel depth. The images were then anonymised and randomised to ensure the radiologists remained unaware of the time point of each mammogram. Mammograms were analysed using three different methods: (1) a semi-automated area-based measure based on computer-assisted thresholding by a single expert user (Cumulus, Sunnybrook health sciences centre, Toronto, Canada, [26]); (2) an automated volumetric Stepwedge method developed at Manchester University [27]; and (3) a visual assessment score of percentage density read to the nearest 5% by two experienced readers and expressed as an average of the two scores to calculate PDA. Cumulus was used to calculate TA, DA, FA, and PDA, and the Manchester Stepwedge method calculated total volume (TV), DV, FV, and PDV. Density assessments were made at 3 time points: baseline, 1 year follow-up (at the end of the intervention) and 1 year after the end of the intervention. Baseline mammograms were taken at the point of entry to the study; for those women with a mammogram performed within one year of entry, their most recent mammogram within the last 12 months was used. Each woman had four mammographic views taken at each time point: Left Cranial-Caudal, Right Cranial-Caudal, Left Mediolateral-Oblique, and Right Mediolateral-Oblique, and a final mammographic score at each time point

was calculated using an average of the four views. The main analysis refers to Cumulus measured area-based density and Stepwedge measured volumetric density only to assess the effects of BMI on dense and non-dense tissue separately. Visually-assessed density had similar results to Cumulus-assessed PDA, so was included as a secondary density measure only. Results for TA and TV are also reported as secondary density measures in the Supplementary Materials.

2.3. Body Weight and Body Composition

Weight, BMI, and a variety of different measures of body composition were assessed at baseline, 1 year follow-up (at the end of the intervention), and 1 year after the end of the intervention. Weight (kg) and height (m) were determined using a calibrated beam balance and stadiometer and used to calculate BMI (kg/m^2). Other body composition assessments were also made such as waist circumference; total body fat, fat free mass and % body fat (assessed using a DXA whole body scanner (Hologic Inc., Bedford, MA, USA) and bioelectrical impedance (Tanita TBF-300A, Tanita Europe B.V., Hoogoorddreef 56E, 1101 BE Amsterdam, The Netherlands)); and intra-abdominal and abdominal subcutaneous area (assessed using a magnetic resonance imaging (MRI) scan with a single transverse scan taken at the level of the intervertebral disc between the L2 and L3 vertebrae). Weight, BMI, waist circumference, and total body fat, fat free mass, and % body fat (impedance) were recorded at all three time points. Intra-abdominal area, abdominal subcutaneous area, and total body fat, fat free mass, and % body fat (DXA) were only measured at baseline and at 1 year. Weight at age 20 years was self-reported via questionnaire, and BMI at age 20 years was calculated using weight at age 20 years and height at study entry. Long-term adult BMI gain was calculated as the difference between baseline BMI and BMI at age 20 years. We discuss BMI as the measure of body weight throughout the main analysis because BMI is a commonly used adjustment for density and it is a well-established risk factor for breast cancer. Other body composition measures gave similar correlations with density to those of BMI and were highly correlated with BMI. Therefore, other body composition measures are included as secondary analyses in the Supplementary Materials. Weight gain during the intervention was defined as $\geq +3\%$ of baseline weight, weight loss was defined as $\leq -3\%$ of baseline weight, and a weight change $> -3\%$ to $< +3\%$ of the baseline weight was defined as a stable weight [28].

2.4. Statistical Analysis

Data were visualised using custom-made ‘tadpole plots’, where each tadpole represents a woman, the head plots the woman’s BMI and density at her last time point, and the points on the tail plot her BMI and density at earlier time points. Correlation (r) between BMI and mammographic density was assessed on a cross-sectional basis (between women), and within women as their short-term BMI changed, using repeated measures methods that use all of the measurements at the same time [29,30]. Briefly, between women correlation was a weighted Pearson correlation coefficient [30], and within women correlation was based on the decomposition of sums of squares from an analysis of variance [29]. The 95% confidence intervals were estimated using an empirical bootstrap (10,000 resamples). The simultaneous association of between and within women correlations was tested using a linear mixed model adjusted for age [31] (Appendix A). To help with comparisons across different measures of breast density, the breast density values were first standardised (Appendix B). To make density measures more symmetric and approximately normally-distributed, they were transformed: a square root transformation for area measures and a cube root transformation for volumetric measures. An exploratory analysis was undertaken to assess the effect of adding BMI gain since 20 years of age to the model. An additional exploratory analysis tested whether there was an association between breast density and DXA bone density. A sensitivity analysis assessed repeated measures correlation coefficients for BMI and density stratified by intervention group.

Analysis used the statistical software R [32]. All tests were two-sided and considered significant at the 5% level.

3. Results

Baseline characteristics of the cohort are shown in Table 1. Median age was 41 years (interquartile range (IQR), 38–43 years), and the majority of women were Caucasian ($n = 60$, 92%) and parous ($n = 55$, 85%). At baseline, 27 women (42%) were classified as overweight ($\text{BMI} \geq 25 \text{ kg/m}^2$ and $<30 \text{ kg/m}^2$), 20 (31%) were obese ($\text{BMI} \geq 30 \text{ kg/m}^2$), and 18 (28%) were in the normal BMI range ($\text{BMI} \geq 18.5 \text{ kg/m}^2$ and $<25 \text{ kg/m}^2$). By the end of the 2 year study period (1 year post intervention), 22 women (34%) had lost weight, 16 (25%) had gained weight, and 26 (41%) maintained their original weight. Overall, women in the intervention group lost more weight than the usual care group (mean percentage of baseline weight at 1 year = -4.4% and 0.1% , respectively; mean percentage of baseline weight at 2 years = -2.9% and 2.0% , respectively).

Median PDA, DA, and FA of each woman's average density measure over the intervention were 37.1% (IQR, 2.5% – 71.3%), 59.9 cm^2 (IQR, 5.8 – 158.4 cm^2) and 107.3 cm^2 (IQR, 23.6 – 405.1 cm^2), respectively. For Stepwedge measures, PDV, DV, and FV were 22.7% (IQR, 6.7% – 69.4%), 191.5 cm^3 (IQR, 56.7 – 710.4 cm^3), and 573.0 cm^3 (IQR, 72.8 – 1992.1 cm^3), respectively. A flow chart detailing the availability of mammographic density measures across the intervention is shown in Figure S1 (all women had BMI available at all time-points except for one woman with missing BMI at 2 years—this data point was excluded from analyses involving BMI).

Table 2 shows the repeated measures correlations. DV was positively correlated with BMI between women ($r = 0.41$, 95%CI 0.17 to 0.61) but less so within women ($r = 0.08$, 95%CI -0.16 to 0.28). There was little association between DA and BMI (between women $r = -0.12$, 95%CI -0.38 to 0.16; within women $r = 0.01$, 95%CI -0.24 to 0.25). PDV was inversely associated with BMI between and within women (between $r = -0.48$, 95%CI -0.64 to -0.33 ; within $r = -0.36$, 95%CI -0.54 to -0.12), and PDA was inversely associated with BMI between women ($r = -0.58$, 95%CI -0.72 to -0.42), but less so within women ($r = -0.22$, 95%CI -0.44 to 0.01). FV and FA were positively correlated with BMI between and within women (volume: between $r = 0.77$, 95%CI 0.69 to 0.84, within $r = 0.58$, 95%CI 0.36 to 0.75; area: between $r = 0.74$, 95%CI 0.63 to 0.82, within $r = 0.45$, 95%CI 0.23 to 0.63). The magnitude of correlations was stronger between women than within women. These associations were also seen in Figure 1 when data were visually assessed using tadpole plots (trends in the tadpole heads represented the between women correlations and trends in the tadpole tails represented within women correlations).

Results for repeated measures correlation coefficients were similar when evaluated in a sensitivity analysis stratifying the cohort by intervention group. Within women associations for BMI and FA or FV were slightly stronger for women following the supervised weight loss programme compared with the usual care group, but there was little association (within women) for BMI and DA or DV in both intervention groups (Table S6).

Other body fat composition measures were highly correlated with BMI (Table S3), and the associations between breast density and other body fat compositions were similar to those with BMI (Tables S1 and S2). The correlations between various mammographic density measures are also reported in the Supplementary Materials (Table S4).

Table 1. Participant characteristics at study entry.

| Characteristic | All | Intervention | Usual Care |
|--|------------------|------------------|------------------|
| Total | 65 | 33 | 32 |
| Age * (years) | 41 (38–43) | 41 (39–43) | 40 (38–42) |
| Baseline BMI * (kg/m ²) | 27.1 (24.7–33.4) | 27.1 (25.1–31.9) | 27.0 (24.4–34.0) |
| Baseline BMI categories # (kg/m ²) | | | |
| Normal (≥18.5 to <25) | 18 (28%) | 7 (21%) | 11 (34%) |
| Overweight (≥25 to <30) | 27 (42%) | 16 (48%) | 11 (34%) |
| Obese (≥30) | 20 (31%) | 10 (30%) | 10 (31%) |
| BMI gain since 20 years * (kg/m ²) | 5.8 (4.7–9.4) | 6.3 (4.7–10.0) | 5.7 (4.6–8.9) |
| Height * (m) | 1.64 (1.60–1.68) | 1.63 (1.60–1.68) | 1.65 (1.59–1.68) |
| Age at menarche * (years) | 12 (12–13) | 12 (12–13) | 12 (12–13) |
| Number of live births # | | | |
| Nulliparous | 10 (15%) | 6 (18%) | 4 (13%) |
| 1–2 | 41 (63%) | 20 (61%) | 21 (66%) |
| 3–4 | 12 (18%) | 6 (18%) | 6 (19%) |
| ≥5 | 2 (3%) | 1 (3%) | 1 (3%) |
| Age first live birth * (years) | 27 (22–29) | 27 (24–31) | 26 (22–29) |
| Ethnicity # (% Caucasian) | 60 (92%) | 29 (88%) | 31 (97%) |
| Previous smoker # | | | |
| Never | 54 (83%) | 29 (88%) | 25 (78%) |
| Ever | 11 (17%) | 4 (12%) | 7 (22%) |
| Previous oral contraception use # | | | |
| Never | 5 (8%) | 3 (9%) | 2 (6%) |
| Ever | 58 (89%) | 29 (88%) | 29 (91%) |
| Unknown | 2 (3%) | 1 (3%) | 1 (3%) |
| Breastfed # | | | |
| Never | 22 (34%) | 12 (36%) | 10 (31%) |
| Ever | 41 (63%) | 21 (64%) | 20 (63%) |
| Unknown | 2 (3%) | 0 (0%) | 2 (6%) |
| 10-year Tyrer–Cuzick risk * (%) | 4 (3–5) | 5 (4–6) | 3 (3–4) |
| Alcohol intake ^{a,*} (units/week) | 11 (3–24) | 11 (3–22) | 10 (1.5–26) |
| Physical activity ^{b,*} ((kJ/kg)/week) | 974 (945–999) | 968 (941–999) | 978 (953–1007) |
| Weight change from baseline to 1 year, categories # | | | |
| Loss | 26 (40%) | 20 (61%) | 6 (19%) |
| Stable | 27 (42%) | 9 (27%) | 18 (56%) |
| Gain | 12 (18%) | 4 (12%) | 8 (25%) |
| Weight change from baseline to 1 year ** | −2.2 (5.4) | −4.4 (5.0) | 0.1 (4.8) |
| Weight change from baseline to 2 years, categories # | | | |
| Loss | 22 (34%) | 16 (48%) | 6 (19%) |
| Stable | 26 (41%) | 13 (39%) | 13 (42%) |
| Gain | 16 (25%) | 4 (12%) | 12 (39%) |
| Weight change from baseline to 2 years ** | −0.5 (7.1) | −2.9 (6.2) | 2.0 (7.1) |

BMI: Body mass index. # N (%); * Median (interquartile range); ** Mean (standard deviation) % of baseline weight (kg). ^a Alcohol from a 4-day food diary; ^b Physical activity from 7-day recall. Weight loss defined as ≤−3% of baseline weight (kg); Stable weight defined as >−3% to <+3% of baseline weight (kg); Weight gain defined as ≥+3% of baseline weight (kg).

Table 2. Repeated measures between women and within women correlations for mammographic density and body mass index.

| Field | VAS (95%CI) (sqrt%) | PDA (95%CI) (sqrt%) | PDV (95%CI) (cbirt%) | FA (95%CI) (sqrt) | FV (95%CI) (cbirt) | DA (95%CI) (sqrt) | DV (95%CI) (cbirt) |
|--------------------------------------|---------------------------|---------------------------|----------------------------|----------------------|-----------------------|-----------------------|-----------------------|
| Cross-sectional BMI (between women) | −0.62 (−0.74 to −0.47) | −0.58 (−0.72 to −0.42) | −0.48 (−0.64 to −0.33) | 0.74 (0.63 to 0.82) | 0.77 (0.69 to 0.84) | −0.12 (−0.38 to 0.16) | 0.41 (0.17 to 0.61) |
| Short-term BMI change (within women) | −0.27 (−0.48 to −0.05) | −0.22 (−0.44 to 0.01) | −0.36 (−0.54 to −0.12) | 0.45 (0.23 to 0.63) | 0.58 (0.36 to 0.75) | 0.01 (−0.24 to 0.25) | 0.08 (−0.16 to 0.28) |

VAS: Visual assessment score; PDA: percent dense area; PDV: percent dense volume; FA: fat area; FV: fat volume; DA: dense area; DV: dense volume; sqrt: square root transformed; cbirt: cube root transformed; BMI: body mass index; 95%CI: 95% confidence interval. Area-based measures from Cumulus; volumetric measures from Manchester Stepwedge. Within women correlations represent trends over the entire 2 year period.




Figure 1. Tadpole plots showing body mass index (BMI) and density measures across the 2 year follow-up. Each tadpole represents a woman: the tadpole head shows BMI and density (if density is available) at her last follow-up and the points on the tail show BMI and density (if density is available) at her earlier follow-ups. (a) Dense volume; (b) Fat volume; (c) Dense area; (d) Fat area.

The between and within women associations for density and BMI measures were similar when estimated jointly in an age-adjusted linear mixed model (Table 3). In a sensitivity analysis, the same model was fit using weight instead of BMI, but it had a worse model fit for almost all density measures (Table S5).

Table 3. Multivariable linear mixed model fit results for mammographic density on body mass index (between and within women), adjusted for age (A1).

| Density Outcome | Intercept (95%CI) | Age (95%CI) (Per 10 Years) | BMI (95%CI) [between] (Per 5 kg/m ²) | BMI (95%CI) [within] (Per 5 kg/m ²) |
|-----------------|------------------------|-------------------------------|---|--|
| VAS (sqrt%) | 3.75 (1.88 to 5.61) | −0.19 (−0.56 to 0.19) | −0.51 (−0.68 to −0.35) | −0.27 (−0.44 to −0.10) |
| PDA (sqrt%) | 2.87 (0.57 to 5.17) | −0.05 (−0.53 to 0.43) | −0.46 (−0.63 to −0.30) | −0.32 (−0.59 to −0.05) |
| PDV (cbt%) | 1.73 (−1.07 to 4.53) | 0.12 (−0.48 to 0.71) | −0.39 (−0.57 to −0.21) | −0.85 (−1.32 to −0.39) |
| FA (sqrt) | −3.63 (−5.25 to −2.02) | 0.04 (−0.28 to 0.36) | 0.60 (0.46 to 0.74) | 0.43 (0.27 to 0.58) |
| FV (cbt) | −3.46 (−5.27 to −1.64) | −0.04 (−0.42 to 0.34) | 0.63 (0.50 to 0.76) | 0.79 (0.56 to 1.03) |
| DA (sqrt) | 0.57 (−2.13 to 3.27) | −0.03 (−0.59 to 0.53) | −0.08 (−0.28 to 0.11) | 0.01 (−0.30 to 0.33) |
| DV (cbt) | −2.39 (−5.11 to 0.33) | 0.09 (−0.48 to 0.66) | 0.35 (0.16 to 0.53) | 0.16 (−0.24 to 0.55) |

VAS: Visual assessment score; PDA: percent dense area; PDV: percent dense volume; FA: fat area; FV: fat volume; DA: dense area; DV: dense volume; sqrt: square root transformed; cbrt: cube root transformed; BMI: body mass index; 95%CI: 95% confidence interval. Area-based measures from Cumulus; volumetric measures from Manchester Stepwedge. Between women BMI calculated as the mean BMI for each woman; within women BMI calculated as the difference between each woman's BMI and her mean BMI. Density measures are standardised (see Appendix B). One woman with missing BMI at age 20 years excluded. Within women effects represent trends over the entire 2 year period.

When a term for BMI gain since age 20 years was added to the linear mixed model, the model fit improved for PDA, PDV, FV, and DA (all $\Delta LR-\chi^2 p < 0.05$) (Table 4). After including BMI gain since age 20 years, between women associations for BMI became more strongly inversely associated with percent density (approximately −0.5 to −0.8), more strongly positively associated with breast fat (approximately 0.6 to 0.8), more strongly inversely associated with DA (−0.1 to −0.5), and less strongly positively associated with DV (0.4 to 0.2). Within women effects of BMI on density were almost unchanged when including BMI gain since age 20 years. BMI gain from age 20 years (adjusted for attained BMI) was positively associated with DA, PDA, and PDV (5 kg/m² increase in BMI gain since age 20 years was associated with 0.61 (95%CI 0.12 to 1.09), 0.61 (95%CI 0.21 to 1.02), and 0.47 (95%CI 0.05 to 0.88) standard deviation increase in breast density (β), respectively), and inversely associated with FV ($\beta = -0.31$, 95%CI −0.62 to 0.00), but less association was seen with DV ($\beta = 0.15$, 95%CI −0.29 to 0.59) and FA ($\beta = -0.32$, 95%CI −0.67 to 0.03).

Finally, in tests of association between breast and bone density, there was some indication of a positive between women correlation for bone density and FV ($r = 0.26$, 95%CI, 0.00 to 0.50), DV ($r = 0.33$, 95%CI, 0.09 to 0.54), and TV ($r = 0.31$, 95%CI, 0.06 to 0.54) (Table S1), but we found little correlation within women (Table S2).

Table 4. Multivariable linear mixed model fit results for mammographic density on body mass index (between and within women) and body mass index gain since 20 years of age, adjusted for age (A2).

| Density Outcome | Intercept (95%CI) | Age (95%CI) (Per 10 Years) | BMI (95%CI) (between) (Per 5kg/m ²) | BMI (95%CI) (within) (Per 5kg/m ²) | BMI Gain Since 20 Years of Age (95%CI) (Per 5kg/m ²) | Δ LR- χ^2 p-Value (A1 vs. A2) |
|-----------------|------------------------|----------------------------|---|--|--|---|
| VAS (sqrt%) | 5.47 (3.34 to 7.60) | −0.25 (−0.61 to 0.12) | −0.92 (−1.23 to −0.62) | −0.27 (−0.45 to −0.10) | 0.59 (0.20 to 0.97) | 0.0031 |
| PDA (sqrt%) | 4.90 (2.34 to 7.46) | −0.16 (−0.63 to 0.31) | −0.89 (−1.22 to −0.57) | −0.32 (−0.59 to −0.06) | 0.61 (0.21 to 1.02) | 0.0033 |
| PDV (cbt%) | 3.35 (0.30 to 6.40) | 0.01 (−0.57 to 0.60) | −0.71 (−1.05 to −0.38) | −0.85 (−1.32 to −0.39) | 0.47 (0.05 to 0.88) | 0.0267 |
| FA (sqrt) | −4.59 (−6.49 to −2.69) | 0.08 (−0.24 to 0.40) | 0.82 (0.54 to 1.10) | 0.43 (0.28 to 0.59) | −0.32 (−0.67 to 0.03) | 0.0704 |
| FV (cbt) | −4.42 (−6.44 to −2.40) | 0.01 (−0.37 to 0.38) | 0.84 (0.59 to 1.09) | 0.79 (0.56 to 1.03) | −0.31 (−0.62 to 0.00) | 0.0476 |
| DA (sqrt) | 2.58 (−0.48 to 5.64) | −0.14 (−0.70 to 0.41) | −0.51 (−0.90 to −0.12) | 0.01 (−0.31 to 0.32) | 0.61 (0.12 to 1.09) | 0.0145 |
| DV (cbt) | −1.90 (−4.96 to 1.15) | 0.06 (−0.51 to 0.64) | 0.24 (−0.12 to 0.60) | 0.16 (−0.24 to 0.55) | 0.15 (−0.29 to 0.59) | 0.4967 |

VAS: Visual assessment score; PDA: percent dense area; PDV: percent dense volume; FA: fat area; FV: fat volume; DA: dense area; DV: dense volume; sqrt: square root transformed; cbrt: cube root transformed; BMI: body mass index; 95%CI: 95% confidence interval. Area-based measures from Cumulus; volumetric measures from Manchester Stepwedge. Between women BMI calculated as the mean BMI for each woman; within women BMI calculated as the difference between each woman's BMI and her mean BMI; BMI gain from age 20 years calculated as the difference between each woman's BMI at baseline and her BMI at age 20 years. Density measures are standardised (see Appendix B). One woman with missing BMI at age 20 years excluded. Δ LR- χ^2 represents the difference in likelihood ratio for A1 versus A2. Within women effects represent trends over the entire 2 year period. All variables adjusted for each other in the multivariable model, therefore BMI gain since 20 years of age is adjusted for current BMI through the variable for between women BMI.

4. Discussion

The data in this study provide some support for the two main findings. First, it is possible that the higher a premenopausal woman's BMI, the higher her breast fat and dense tissue (in particular, dense volume), and the lower her percent density. Second, the data suggested that as a premenopausal woman loses weight, her breast fat reduces, dense tissue remains relatively unchanged, and percent dense tissue increases. Effective weight loss during premenopausal years has been associated with a reduced risk of postmenopausal breast cancer [16–18], but our study data suggest that risk reduction is unlikely to be mediated by a short-term reduction in dense breast tissue. This is likely to mean that incorporation of change in percent breast density into risk algorithms will not capture potential weight loss-induced breast cancer risk reduction and may falsely ascribe an increased risk due to increased percent density. Therefore, risk prediction models need to consider how best to incorporate changes in weight and mammographic density when predicting breast cancer risk.

The between women associations of attained premenopausal BMI and breast density observed in this study were consistent with previous studies. High BMI is associated with high dense volume [33–35], but the correlation between BMI and dense area is less strong, and often close to zero [36–39]. These differences are likely to be a result of volumetric measures representing breast tissue more accurately than area-based methods by accounting for breast thickness and overlapping tissue. Additionally, since the breast is a deposit for adipose tissue, high attained BMI is strongly associated with high levels of breast fat area [36–39] and breast fat volume [33,34], which in turn leads to an inverse association between BMI and both percent dense area [36–41] and percent dense volume [33–35,42,43].

There have been very few studies to assess the effect of dietary weight loss on breast density in premenopausal women. Boyd et al. reported reductions in total and dense area alongside modest weight change within an intervention trial of women on a 2-year low-fat, high-carbohydrate diet [44]. In particular, a 5.4% decrease in dense area was seen in premenopausal women in the low-fat diet group with a 0.1kg/m² BMI reduction ($n = 249$) compared with a 2.5% decrease in the control group with a 0.3kg/m² BMI gain ($n = 264$). These reductions may be associated with the large reductions in dietary fat (55 to 35 g/day) and saturated fat (19 to 12 g/day) rather than weight loss in this study. This was considerably higher than those advised and achieved in the current reported study (total fat reduced from 77 to 60 g/day and saturated fat reduced from 28 to 21 g/day). Other trials have also assessed the effect of lifestyle interventions for weight loss on breast density, although in postmenopausal women only. In the ALPHA trial, postmenopausal women on a 1-year aerobic exercise intervention lost on average 39 cm³ more breast fat than the controls, but there was little difference in the change in dense tissue between the two groups [45]. Furthermore, the DAMA trial reported a reduction in volumetric percent density of approximately 14% for postmenopausal women following a 2-year diet or exercise intervention when compared with the controls [46]. Large weight loss with bariatric surgery is also associated with large reductions in breast fat alongside relatively smaller reductions in dense tissue, and an increase in percent density [47–49].

As an exploratory analysis, we also found an association between increased BMI gain since 20 years of age and higher dense tissue and percent density. It is possible that this is a pathway for the increased risk of postmenopausal breast cancer seen with adult weight gain [7,15,50–52]. However, this association is likely to reflect the inverse association seen in previous studies between adolescent body adiposity and dense tissue in later life [38,40,53–55], since, given the adjustment for current BMI, women with greater gain in BMI will have had lower BMI at 20 years of age. This interesting observation requires further investigation in larger datasets of women. Additionally, exploratory analysis of bone density found little association with breast density, which is in agreement with previous studies [56].

Strengths of this study include the various measures of breast density including Cumulus and the Stepwedge method, which allowed for the assessment of dense and fatty

tissue separately as well as various measures of body weight to assess adiposity. The study also assessed breast density as an area-based measure and volumetrically; both of which have similar abilities for breast cancer risk prediction [57]. Additionally, all women were encouraged to lose weight, which produced data with large within women variation in BMI, in turn increasing the potential to see an effect of changing BMI on mammographic density. Furthermore, the Lifestyle Study provided a data source to assess premenopausal weight loss and density associations; something that is not possible in studies involving routine screening data. This also provided a greater ability to capture the effects of weight loss on density because this cohort of premenopausal women were likely to have had higher dense tissue at baseline (with greater ability to decrease) than screening populations involving postmenopausal women [58]. Finally, the use of repeated measures over a 2-year period allowed us to assess the association between BMI and breast density longitudinally, whilst making use of all available data simultaneously.

Limitations of the study include the small sample size, which limits statistical power. This is particularly relevant for volumetric measures, which had a moderate amount of missing data at the baseline. In addition, the study design was not powered for the analysis of mammographic density, which was a secondary analysis (the study was powered for salivary oestradiol). This was a relatively small study, and ideally, a larger study with sufficient power would be run to verify our evidence. Another limitation is the analysis of BMI gain since 20 years of age relies on self-reported information on weight at age 20 years. This may be less accurate than the measured values. However, validation studies show that self-reported BMI is highly correlated with independently measured BMI, and the mean difference between self-reported and measured weight is minimal [59,60]. Finally, breast thickness is likely to have changed whilst women lost weight during the intervention. Volumetric measures are influenced by breast thickness [61], hence there might have been larger variation in the serial compared with stable volumetric measurements, resulting in reduced ability to capture the within women effects of BMI on dense tissue volumetrically.

5. Conclusions

This study suggests that premenopausal weight loss reduces breast fat but that it does not reduce dense tissue. Short-term premenopausal weight loss is likely to be linked to lower postmenopausal breast cancer risk through reductions in adipose tissue, not fibroglandular tissue. This means that a potential breast cancer risk reduction as a result of weight loss might not be captured by changes in breast density, and the resulting increase in percent density may falsely ascribe an increase in risk. However, the study was limited by the small sample size, and more studies are required to provide evidence to confirm these results.

Supplementary Materials: The following are available online at <https://www.mdpi.com/article/10.3390/cancers13133245/s1>, Table S1: Complete results for repeated measures between women correlations for mammographic density and body composition measures; Table S2: Complete results for repeated measures within women correlations for mammographic density and body composition measures; Table S3: Complete results for repeated measures between women correlations for different body composition measures; Table S4: Complete results for repeated measures between women correlations for different mammographic density measures; Table S5: Multivariable linear mixed model fit results for A1 using either body mass index or weight; Table S6: Repeated measures between women and within women correlations for mammographic density and body mass index, stratified by intervention group; Figure S1: Flow chart of women included in the analysis and availability of mammographic density data.

Author Contributions: Conceptualization, A.H. and M.H.; Formal Analysis, E.C.A. and A.R.B.; Investigation, S.A., J.C., D.G.E., R.M.L.W., A.H., and M.H.; Data Curation, S.A., D.G.E., R.M.L.W., A.H., and M.H.; Writing—Original Draft Preparation, E.C.A., A.R.B., S.A., J.C., D.G.E., R.M.L.W., A.H., and M.H.; Writing—Review & Editing, E.C.A., A.R.B., S.A., J.C., D.G.E., R.M.L.W., A.H., and M.H. All authors have read and agreed to the published version of the manuscript.

Funding: E.C.A., A.R.B. and J.C. are supported by Cancer Research UK (C569/A16891 to J.C.); S.A., D.G.E., A.H., and M.H. are supported by the NIHR Manchester Biomedical Research Centre (IS-BRC-1215-20007 to D.G.E.). The funders had no role in the design of the study; in the collection, analysis and interpretation of data; in the writing of the manuscript, or in the decision to submit for publication.

Institutional Review Board Statement: The study was conducted according to the guidelines of the Declaration of Helsinki and was approved by the South Manchester Ethics Committee on 28 January 2002 (Reference no 01/426).

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: The dataset used and analysed during the current study is available from the corresponding author on reasonable request.

Acknowledgments: We thank Hilary Graff for contributions to the collection and assembly of data (University of Manchester). We are particularly grateful to the women who participated in this study and the entire medical and administrative staff who worked on the Lifestyle Study.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

Abbreviations

| | |
|-----|-------------------------|
| DA | dense area |
| PDA | percent dense area |
| FA | fat area |
| TA | total area |
| DV | dense volume |
| PDV | percent dense volume |
| FV | fat volume |
| TV | total volume |
| BMI | body mass index |
| VAS | visual assessment score |

Appendix A

Linear mixed model for mammographic density on body mass index and age. A linear mixed model was used to model density and body mass index (BMI) associations in Table 3. This model allows for repeated measures and uses all of the available data (missing pairs of density and BMI are excluded). Breast density y_{ij} for woman $i = 1, \dots, n$ at time $j = \{1, 2, 3\}$ is modelled as:

$$y_{ij} = \alpha + \beta age_{ij} + \gamma \bar{x}_i + \delta(x_{ij} - \bar{x}_i) + u_i + e_{ij}; \quad (A1)$$

where α is an overall intercept; age_{ij} is the age at baseline for woman i at time j ; β is the slope for age; \bar{x}_i is mean BMI for woman i ; γ is the between women slope; x_{ij} is the BMI of woman i at time j ; δ is the within women slope; and e_{ij} is an independent random error. Another term that allows for differences between women in their overall density level is the independent random intercept u_i for woman i . The model is completed by assuming normal distributions for u_i and e_{ij} with zero mean, unknown variances, and zero covariance. The model was fitted by maximum likelihood. To aid interpretation of the estimates across different measures of density, the density values were standardised (see Appendix B). To test $\gamma = 0$ (between women correlation) and $\delta = 0$ (within women correlation), a Wald test was applied.

The model was extended to consider BMI gain from age 20 years in Table 4:

$$y_{ij} = \alpha + \beta age_{ij} + \gamma \bar{x}_i + \delta(x_{ij} - \bar{x}_i) + u_i + \epsilon z_i + e_{ij}; \quad (A2)$$

where z_i is the BMI gain since age 20 years for woman i : calculated as the difference between baseline BMI for woman i and BMI at age 20 years for woman i , and ε is the slope for BMI gain since age 20 years. To test $\varepsilon = 0$, a Wald test was applied.

Appendix B

Standardisation of each mammographic density measure:

$$\bar{x} = \frac{\sum_{i=1}^n \bar{x}_i}{n}$$

$$\sigma = \sqrt{\frac{\sum_{i=1}^n (\bar{x}_i - \bar{x})^2}{n - 1}}$$

$$z_{ij} = \frac{d_{ij} - \bar{x}}{\sigma}$$

where \bar{x}_i is the mean density for woman $i = 1, \dots, n$; d_{ij} is the density measure for woman $i = 1, \dots, n$ at time point $j = \{1, 2, 3\}$; and z_{ij} is the standardised density measure for woman i at time point j .

References

- McCormack, V.A.; dos Santos Silva, I. Breast density and parenchymal patterns as markers of breast cancer risk: A meta-analysis. *Cancer Epidemiol. Biomark. Prev.* **2006**, *15*, 1159–1169. [[CrossRef](#)] [[PubMed](#)]
- Boyd, N.F.; Byng, J.W.; Jong, R.A.; Fishell, E.K.; Little, L.E.; Miller, A.B.; Lockwood, G.A.; Tritchler, D.L.; Yaffe, M.J. Quantitative Classification of Mammographic Densities and Breast Cancer Risk: Results from the Canadian National Breast Screening Study. *J. Natl. Cancer Inst.* **1995**, *87*, 670–675. [[CrossRef](#)]
- Byrne, C.; Schairer, C.; Wolfe, J.; Parekh, N.; Salane, M.; Brinton, L.A.; Hoover, R.; Haile, R. Mammographic Features and Breast Cancer Risk: Effects with Time, Age, and Menopause Status. *J. Natl. Cancer Inst.* **1995**, *87*, 1622–1629. [[CrossRef](#)] [[PubMed](#)]
- Pettersson, A.; Hankinson, S.E.; Willett, W.C.; Lagiou, P.; Trichopoulos, D.; Tamimi, R.M. Nondense mammographic area and risk of breast cancer. *Breast Cancer Res.* **2011**, *13*, R100. [[CrossRef](#)]
- Eng, A.; Gallant, Z.; Shepherd, J.; McCormack, V.; Li, J.; Dowsett, M.; Vinnicombe, S.; Allen, S.; Dos-Santos-Silva, I. Digital mammographic density and breast cancer risk: A case-control study of six alternative density assessment methods. *Breast Cancer Res.* **2014**, *16*, 439. [[CrossRef](#)] [[PubMed](#)]
- Brentnall, A.R.; Cohn, W.F.; Knaus, W.A.; Yaffe, M.J.; Cuzick, J.; Harvey, J.A. A Case-Control Study to Add Volumetric or Clinical Mammographic Density into the Tyrer-Cuzick Breast Cancer Risk Model. *J. Breast Imaging* **2019**, *1*, 99–106. [[CrossRef](#)]
- Kyrgiou, M.; Kalliala, I.; Markozannes, G.; Gunter, M.J.; Paraskevidis, E.; Gabra, H.; Martin-Hirsch, P.; Tsilidis, K.K. Adiposity and cancer at major anatomical sites: Umbrella review of the literature. *BMJ* **2017**, *356*, j477. [[CrossRef](#)] [[PubMed](#)]
- Renehan, A.G.; Tyson, M.; Egger, M.; Heller, R.F.; Zwahlen, M. Body-mass index and incidence of cancer: A systematic review and meta-analysis of prospective observational studies. *Lancet* **2008**, *371*, 569–578. [[CrossRef](#)]
- Reeves, G.K.; Pirie, K.; Beral, V.; Green, J.; Spencer, E.; Bull, D. Cancer incidence and mortality in relation to body mass index in the Million Women Study: Cohort study. *BMJ* **2007**, *335*, 1134. [[CrossRef](#)]
- Van Kruijsdijk, R.C.M.; Van Der Wall, E.; Visseren, F. Obesity and Cancer: The Role of Dysfunctional Adipose Tissue. *Cancer Epidemiol. Biomark. Prev.* **2009**, *18*, 2569–2578. [[CrossRef](#)]
- Travis, R.C.; Key, T.J. Oestrogen exposure and breast cancer risk. *Breast Cancer Res.* **2003**, *5*, 239–247. [[CrossRef](#)] [[PubMed](#)]
- Kim, B.-K.; Chang, Y.; Ahn, J.; Jung, H.-S.; Kim, C.-W.; Yun, K.E.; Kwon, M.-J.; Suh, B.-S.; Chung, E.C.; Shin, H.; et al. Metabolic syndrome, insulin resistance, and mammographic density in pre- and postmenopausal women. *Breast Cancer Res. Treat.* **2015**, *153*, 425–434. [[CrossRef](#)]
- Schoemaker, M.J.; Nichols, H.B.; Wright, L.B.; Brook, M.N.; Jones, M.E.; O'Brien, K.M.; Adami, H.O.; Baglietto, L.; Bernstein, L.; Bertrand, K.A.; et al. Association of Body Mass Index and Age with Subsequent Breast Cancer Risk in Premenopausal Women. *JAMA Oncol.* **2018**, *4*, e181771. [[CrossRef](#)] [[PubMed](#)]
- Amadou, A.; Ferrari, P.; Muwonge, R.; Moskal, A.; Biessy, C.; Romieu, I.; Hainaut, P. Overweight, obesity and risk of premenopausal breast cancer according to ethnicity: A systematic review and dose-response meta-analysis. *Obes. Rev.* **2013**, *14*, 665–678. [[CrossRef](#)] [[PubMed](#)]
- Keum, N.; Greenwood, D.C.; Lee, D.H.; Kim, R.; Aune, D.; Ju, W.; Hu, F.B.; Giovannucci, E.L. Adult weight gain and adiposity-related cancers: A dose-response meta-analysis of prospective observational studies. *J. Natl. Cancer Inst.* **2015**, *107*, djv088. [[CrossRef](#)]
- Hardefeldt, P.J.; Penninkilampi, R.; Edirimanne, S.; Eslick, G.D. Physical Activity and Weight Loss Reduce the Risk of Breast Cancer: A Meta-analysis of 139 Prospective and Retrospective Studies. *Clin. Breast Cancer* **2018**, *18*, e601–e612. [[CrossRef](#)] [[PubMed](#)]

17. Harvie, M.; Howell, A.; Vierkant, R.A.; Kumar, N.; Cerhan, J.R.; Kelemen, L.E.; Folsom, A.R.; Sellers, T.A. Association of Gain and Loss of Weight before and after Menopause with risk of postmenopausal breast cancer in the Iowa Women's Health Study. *Cancer Epidemiol. Biomark. Prev.* **2005**, *14*, 656–661. [[CrossRef](#)] [[PubMed](#)]
18. Trentham-Dietz, A.; Newcomb, P.A.; Egan, K.M.; Titus-Ernstoff, L.; Baron, J.A.; Storer, B.E.; Stampfer, M.; Willett, W.C. Weight change and risk of postmenopausal breast cancer (United States). *Cancer Causes Control* **2000**, *11*, 533–542. [[CrossRef](#)]
19. Winder, A.A.; Kularatna, M.; MacCormick, A.D. Does Bariatric Surgery Affect the Incidence of Breast Cancer Development? A Systematic Review. *Obes. Surg.* **2017**, *27*, 3014–3020. [[CrossRef](#)]
20. Harvie, M.; Cohen, H.; Mason, C.; Mercer, T.; Malik, R.; Adams, J.; Evans, D.G.R.; Hopwood, P.; Cuzick, J.; Howell, A. Adherence to a Diet and Exercise Weight Loss Intervention amongst Women at Increased Risk of Breast Cancer. *Open Obes. J.* **2010**, *2*, 71–80.
21. Graffy, H.; Harvie, M.; Warren, R.; Boggis, C.; Astley, S.; Evans, G.; Adams, J.; Howell, A. Abstract P4-13-09: The effect of weight change on breast adipose and dense tissue. *Poster Sess. Abstr.* **2012**, *72*, P4-13-09. [[CrossRef](#)]
22. Patel, H.G.; Astley, S.M.; Hufton, A.P.; Harvie, M.; Hagan, K.; Marchant, T.E.; Hillier, V.; Howell, A.; Warren, R.; Boggis, C.R.M. Automated breast tissue measurement of women at increased risk of breast cancer. In *Digital Mammography; IWDM, 2006; Lecture Notes in Computer, Science; Astley, S.M., Brady, M., Rose, C., Zwiggelaar, R., Eds.; Springer: Heidelberg, Germany, 2006; Volume 4046*.
23. Harvie, M.; Mercer, T.; Humphries, G.; Hopwood, P.; Adams, J.; Evans, G.; Sumner, H.; Astley, S.; Hayes, L.; Cooley, J.; et al. The effects of weight loss and exercise on biomarkers of breast cancer risk—Rationale and study design. *Recent Res. Dev. Nutr.* **2002**, *5*, 91–110.
24. Tyrer, J.; Duffy, S.W.; Cuzick, J. A breast cancer prediction model incorporating familial and personal risk factors. *Stat. Med.* **2004**, *23*, 1111–1130. [[CrossRef](#)]
25. IBIS Breast Cancer Risk Evaluation Tool. Available online: <http://www.ems-trials.org/riskevaluator/> (accessed on 22 December 2020).
26. Byng, J.W.; Boyd, N.F.; Fishell, E.; Jong, R.A.; Yaffe, M.J. The quantitative analysis of mammographic densities. *Phys. Med. Biol.* **1994**, *39*, 1629–1638. [[CrossRef](#)] [[PubMed](#)]
27. Diffey, J.; Hufton, A.; Astley, S. A new step-wedge for the volumetric measurement of mammographic density. In *Digital Mammography; IWDM, 2006; Lecture Notes in Computer, Science; Astley, S.M., Brady, M., Rose, C., Zwiggelaar, R., Eds.; Springer: Heidelberg, Germany, 2006; Volume 4046*, pp. 1–9.
28. Stevens, J.; Truesdale, K.P.; McClain, J.E.; Cai, J. The definition of weight maintenance. *Int. J. Obes.* **2005**, *30*, 391–399. [[CrossRef](#)]
29. Bland, J.M.; Altman, D.G. Statistics notes: Calculating correlation coefficients with repeated observations: Part 1—correlation within subjects. *BMJ* **1995**, *310*, 446. [[CrossRef](#)]
30. Bland, J.M.; Altman, D.G. Statistics notes: Calculating correlation coefficients with repeated observations: Part 2—correlation between subjects. *BMJ* **1995**, *310*, 633. [[CrossRef](#)]
31. Crowder, M.J.; Hand, D.J. *Analysis of Repeated Measures*, 1st ed.; Monographs on Statistics and Applied Probability; Chapman & Hall: London, UK, 1990.
32. R Core Team. *R: A Language and Environment for Statistical Computing*; R Foundation for Statistical Computing: Vienna, Austria, 2015. Available online: <https://www.r-project.org> (accessed on 28 July 2019).
33. Hart, V.; Reeves, K.W.; Sturgeon, S.R.; Reich, N.G.; Sievert, L.L.; Kerlikowske, K.; Ma, L.; Shepherd, J.; Tice, J.; Mahmoudzadeh, A.P.; et al. The Effect of Change in Body Mass Index on Volumetric Measures of Mammographic Density. *Cancer Epidemiol. Biomark. Prev.* **2015**, *24*, 1724–1730. [[CrossRef](#)]
34. Alimujiang, A.; Appleton, C.; Colditz, G.A.; Toriola, A.T. Adiposity during early adulthood, changes in adiposity during adulthood, attained adiposity, and mammographic density among premenopausal women. *Breast Cancer Res. Treat.* **2017**, *166*, 197–206. [[CrossRef](#)]
35. Pereira, A.; Garmendia, M.L.; Uauy, R.; Neira, P.; Lopez-Arana, S.; Malkov, S.; Shepherd, J. Determinants of volumetric breast density in Chilean premenopausal women. *Breast Cancer Res. Treat.* **2017**, *162*, 343–352. [[CrossRef](#)] [[PubMed](#)]
36. Sung, J.; Song, Y.-M.; Stone, J.; Lee, K.; Kim, S.-Y. Association of Body Size Measurements and Mammographic Density in Korean Women: The Healthy Twin Study. *Cancer Epidemiol. Biomark. Prev.* **2010**, *19*, 1523–1531. [[CrossRef](#)] [[PubMed](#)]
37. Boyd, N.; Lockwood, G.; Byng, J.; Little, L.; Yaffe, M.; Tritchler, D. The relationship of anthropometric measures to radiological features of the breast in premenopausal women. *Br. J. Cancer* **1998**, *78*, 1233–1238. [[CrossRef](#)]
38. Harris, H.R.; Tamimi, R.M.; Willett, W.C.; Hankinson, S.E.; Michels, K.B. Body Size Across the Life Course, Mammographic Density, and Risk of Breast Cancer. *Am. J. Epidemiol.* **2011**, *174*, 909–918. [[CrossRef](#)] [[PubMed](#)]
39. Tseng, M.; Byrne, C. Adiposity, adult weight gain and mammographic breast density in US Chinese women. *Int. J. Cancer* **2011**, *128*, 418–425. [[CrossRef](#)] [[PubMed](#)]
40. Pollán, M.; Spain, D.; Lopez-Abente, G.; Miranda-García, J.; García, M.; Casanova, F.; Sánchez-Contador, C.; Santamariña, C.; Moreo, P.; Vidal, C.; et al. Adult weight gain, fat distribution and mammographic density in Spanish pre- and post-menopausal women (DDM-Spain). *Breast Cancer Res. Treat.* **2012**, *134*, 823–838. [[CrossRef](#)]
41. Samimi, G.; Colditz, G.A.; Baer, H.J.; Tamimi, R.M. Measures of energy balance and mammographic density in the Nurses' Health Study. *Breast Cancer Res. Treat.* **2008**, *109*, 113–122. [[CrossRef](#)]

42. Dorgan, J.F.; Klifa, C.; Shepherd, J.A.; Egleston, B.L.; Kwiterovich, P.O.; Himes, J.H.; Gabriel, K.P.; Van Horn, L.; Snetselaar, L.G.; Stevens, V.J.; et al. Height, adiposity and body fat distribution and breast density in young women. *Breast Cancer Res.* **2012**, *14*, R107. [[CrossRef](#)] [[PubMed](#)]
43. Jeffreys, M.; Warren, R.; Highnam, R.; Smith, G.D. Breast cancer risk factors and a novel measure of volumetric breast density: Cross-sectional study. *Br. J. Cancer* **2007**, *98*, 210–216. [[CrossRef](#)] [[PubMed](#)]
44. Boyd, N.F.; Greenberg, C.; Lockwood, G.; Little, L.; Martin, L.; Tritchler, D.; Byng, J.; Yaffe, M. Effects at Two Years of a Low-Fat, High-Carbohydrate Diet on Radiologic Features of the Breast: Results from a Randomized Trial. *J. Natl. Cancer Inst.* **1997**, *89*, 488–496. [[CrossRef](#)]
45. Woolcott, C.G.; Courneya, K.S.; Boyd, N.F.; Yaffe, M.J.; Terry, T.; McTiernan, A.; Brant, R.; Ballard-Barbash, R.; Irwin, M.L.; Jones, C.A.; et al. Mammographic Density Change with 1 Year of Aerobic Exercise among Postmenopausal Women: A Randomized Controlled Trial. *Cancer Epidemiol. Biomark. Prev.* **2010**, *19*, 1112–1121. [[CrossRef](#)]
46. Masala, G.; Assedi, M.; Sera, F.; Ermini, I.; Occhini, D.; Castaldo, M.; Pierpaoli, E.; Caini, S.; Bendinelli, B.; Ambrogetti, D.; et al. Can Dietary and Physical Activity Modifications Reduce Breast Density in Postmenopausal Women? The DAMA Study, a Randomized Intervention Trial in Italy. *Cancer Epidemiol. Biomark. Prev.* **2019**, *28*, 41–50. [[CrossRef](#)]
47. Hassinger, T.E.; Mehaffey, J.H.; Knisely, A.T.; Contrella, B.N.; Brenin, D.R.; Schroen, A.T.; Schirmer, B.D.; Hollowell, P.T.; Harvey, J.A.; Showalter, S.L. The impact of bariatric surgery on qualitative and quantitative breast density. *Breast J.* **2019**, *25*, 1198–1205. [[CrossRef](#)]
48. Williams, A.D.; So, A.; Synnestvedt, M.; Tewksbury, C.M.; Kontos, D.; Hsiehm, M.-K.; Pantalone, L.; Conant, E.F.; Schnall, M.; Dumon, K.; et al. Mammographic breast density decreases after bariatric surgery. *Breast Cancer Res. Treat.* **2017**, *165*, 565–572. [[CrossRef](#)] [[PubMed](#)]
49. Vohra, N.A.; Kachare, S.D.; Vos, P.; Schroeder, B.F.; Schuth, O.; Suttle, D.; Fitzgerald, T.L.; Wong, J.H.; Verbanac, K.M. The Short-Term Effect of Weight Loss Surgery on Volumetric Breast Density and Fibroglandular Volume. *Obes. Surg.* **2016**, *27*, 1013–1023. [[CrossRef](#)] [[PubMed](#)]
50. Magnusson, C.; Baron, J.; Persson, I.; Wolk, A.; Bergstrom, R.; Trichopoulos, D.; Adami, H.O. Body size in different periods of life and breast cancer risk in post-menopausal women. *Int. J. Cancer* **1998**, *76*, 29–34. [[CrossRef](#)]
51. Eliassen, A.H.; Colditz, G.; Rosner, B.; Willett, W.C.; Hankinson, S.E. Adult Weight Change and Risk of Postmenopausal Breast Cancer. *JAMA* **2006**, *296*, 193–201. [[CrossRef](#)]
52. Ahn, J.; Schatzkin, A.; Lacey, J.V.; Albanes, D.; Ballard-Barbash, R.; Adams, K.F.; Kipnis, V.; Mouw, T.; Hollenbeck, A.R.; Leitzmann, M.F. Adiposity, adult weight change, and postmenopausal breast cancer risk. *Arch. Intern. Med.* **2007**, *167*, 2091–2102. [[CrossRef](#)]
53. Bertrand, K.A.; Baer, H.J.; Orav, E.J.; Klifa, C.; Shepherd, J.A.; Van Horn, L.; Snetselaar, L.; Stevens, V.J.; Hylton, N.M.; Dorgan, J.F. Body fatness during childhood and adolescence and breast density in young women: A prospective analysis. *Breast Cancer Res.* **2015**, *17*, 1–10. [[CrossRef](#)] [[PubMed](#)]
54. Hopper, J.L.; Nguyen, T.L.; Stone, J.; Aujard, K.; Matheson, M.C.; Abramson, M.J.; Burgess, J.A.; Walters, E.H.; Dite, G.S.; Bui, M.; et al. Childhood body mass index and adult mammographic density measures that predict breast cancer risk. *Breast Cancer Res. Treat.* **2016**, *156*, 163–170. [[CrossRef](#)]
55. Boyd, N.F.; Martin, L.J.; Sun, L.; Guo, H.; Chiarelli, A.; Hislop, G.; Yaffe, M.; Minkin, S. Body Size, Mammographic Density, and Breast Cancer Risk. *Cancer Epidemiol. Biomark. Prev.* **2006**, *15*, 2086–2092. [[CrossRef](#)]
56. Lee, J.M.; Holley, S.; Appleton, C.; Toriola, A.T. Is There an Association Between Bone Mineral Density and Mammographic Density? A Systematic Review. *J. Women's Health* **2017**, *26*, 389–395. [[CrossRef](#)]
57. Astley, S.M.; Harkness, E.F.; Sergeant, J.C.; Warwick, J.; Stavrinou, P.; Warren, R.; Wilson, M.; Beetles, U.; Gadde, S.; Lim, Y.; et al. A comparison of five methods of measuring mammographic density: A case-control study. *Breast Cancer Res.* **2018**, *20*, 1–13. [[CrossRef](#)]
58. Burton, A.; Maskarinec, G.; Perez-Gomez, B.; Vachon, C.; Miao, H.; Lajous, M.; López-Ridaura, R.; Rice, M.; Pereira, A.; Garmendia, M.L.; et al. Mammographic density and ageing: A collaborative pooled analysis of cross-sectional data from 22 countries worldwide. *PLoS Med.* **2017**, *14*, e1002335. [[CrossRef](#)] [[PubMed](#)]
59. Renehan, A.G.; Pegington, M.; Harvie, M.N.; Sperrin, M.; Astley, S.M.; Brentnall, A.R.; Howell, A.; Cuzick, J.; Evans, D.G. Young adulthood body mass index, adult weight gain and breast cancer risk: The PROCAS Study (United Kingdom). *Br. J. Cancer* **2020**, *122*, 1552–1561. [[CrossRef](#)] [[PubMed](#)]
60. Rimm, E.B.; Stampfer, M.J.; Colditz, G.A.; Chute, C.G.; Litin, L.B.; Willett, W.C. Validity of Self-Reported Waist and Hip Circumferences in Men and Women. *Epidemiology* **1990**, *1*, 466–473. [[CrossRef](#)] [[PubMed](#)]
61. Diffey, J.; Hufton, A.; Beeston, C.; Smith, J.; Marchant, T.; Astley, S. Quantifying Breast Thickness for Density Measurement. In *Digital Mammography*; IWDM, 2008; Lecture Notes in Computer, Science; Krupinski, E.A., Ed.; Springer: Heidelberg, Germany, 2008; Volume 5116, pp. 651–658.