

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



Frota, CC; Lima, LNC; Rocha, AD; Suffys, PN; Rolim, BN; Rodrigues, LC; Barreto, ML; Kendall, C; Kerr, LRS (2012) Mycobacterium leprae in six-banded (*Euphractus sexcinctus*) and nine-banded armadillos (*Dasypus novemcinctus*) in Northeast Brazil. *Memorias do Instituto Oswaldo Cruz*, 107. pp. 209-213. ISSN 0074-0276 DOI: <https://doi.org/10.1590/s0074-02762012000900029>

Downloaded from: <http://researchonline.lshtm.ac.uk/612245/>

DOI: [10.1590/s0074-02762012000900029](https://doi.org/10.1590/s0074-02762012000900029)

Usage Guidelines

Please refer to usage guidelines at <http://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by-nc-nd/2.5/>

***Mycobacterium leprae* in six-banded (*Euphractus sexcinctus*) and nine-banded armadillos (*Dasypus novemcinctus*) in Northeast Brazil**

Cristiane Cunha Frota^{1/+}, Luana Nepomuceno Costa Lima¹, Adalgiza da Silva Rocha², Philip Noel Suffys², Benedito Neilson Rolim³, Laura Cunha Rodrigues⁴, Maurício Lima Barreto⁵, Carl Kendall⁶, Ligia Regina Sansigolo Kerr⁷

¹Departamento de Patologia e Medicina Legal ²Departamento de Saúde Comunitária, Faculdade de Medicina, Universidade Federal do Ceará, Fortaleza, CE, Brasil ³Laboratório de Biologia Molecular Aplicada a Micobactérias, Instituto Oswaldo Cruz-Fiocruz, Rio de Janeiro, RJ, Brasil ⁴Controle de Zoonoses/Vetores, Secretaria de Saúde do Estado do Ceará, Fortaleza, CE, Brasil ⁵London School of Hygiene and Tropical Medicine, London, UK ⁶Instituto de Saúde Coletiva, Universidade Federal da Bahia, Salvador, BA, Brasil ⁷Department of Global Community Health and Behavioral Sciences, Center for Global Health Equity, Tulane University School of Public Health and Tropical Medicine, New Orleans, LA, USA

Human beings are the main reservoir of the causative agent of leprosy, Mycobacterium leprae. In the Americas, nine-banded armadillos (Dasypus novemcinctus) also act as a reservoir for the bacillus. In the state of Ceará (CE), which is located in Northeast Brazil and is an endemic area of leprosy, there are several species of armadillos, including D. novemcinctus and Euphractus sexcinctus (six-banded armadillo). Contact between humans and armadillos occur mainly through hunting, cleaning, preparing, cooking and eating. This study identified M. leprae DNA in the two main species of armadillos found in Northeast Brazil. A total of 29 wild armadillos (27 D. novemcinctus and 2 E. sexcinctus) were captured in different environments of CE countryside. Samples from the ear, nose, liver and spleen from each of these animals were tested by a nested M. leprae-specific repetitive element polymerase chain reaction assay. The samples that tested positive were confirmed by DNA sequencing. M. leprae was detected in 21% (6/29) of the animals, including five D. novemcinctus and one E. sexcinctus. This is the first Brazilian study to identify the presence of a biomarker of M. leprae in wild armadillos (D. novemcinctus and E. sexcinctus) in a leprosy hyperendemic area where there is continuous contact between humans and armadillos.

Key words: *Euphractus sexcinctus* - *Dasypus novemcinctus* - *Mycobacterium leprae* - eco-epidemiology - leprosy

Mycobacterium leprae, the causative agent of leprosy, is not cultivable in vitro. The lack of growth on standard mycobacterial isolation media differentiates this organism from other mycobacterial pathogens. Human beings are the only known reservoir of infection, except in the southern United States of America (USA), where nine-banded armadillos (*Dasypus novemcinctus*) are believed to also provide a reservoir (Truman et al. 2011).

In 1960, Shepard introduced the footpad mouse model to study experimental leprosy. In 1971, nine-banded armadillos were successfully infected with the bacillus and developed clinical symptoms and pathologies similar to the human disease (Kirchheimer et al. 1972, Kirchheimer & Sanchez 1977). Armadillos provide a much more effective animal model than the mouse footpad, producing 10⁶ acid-fast bacilli/mL (Shepard 1985). Because armadillos do not breed well in captivity, the animals to be used in the laboratory need to be captured in the wild. While searching for these animals in 1975, Walsh et al. (1977) found armadillos that were naturally infected with

M. leprae in the wild for the first time. The authors proposed that armadillos might have acquired leprosy infections from untreated human patients in the USA.

The exact mode of transmission of leprosy between humans and armadillos is not known, though cross-reactivity between IgM antibodies against phenolic glycolipid-I of humans and armadillos has been reported (Truman et al. 1991, Job et al. 1992). Infected nine-banded armadillos have been identified in the states of Texas and Louisiana and in Central and South America (Smith et al. 1983, Amezcua et al. 1984, Stallknecht et al. 1987, Zumarraga et al. 2001). Additionally, biomarkers of armadillo infection have been detected in Colombia and Brazil (Deps et al. 2007, Cardona-Castro et al. 2009). Several studies have shown an association between armadillo exposure through hunting, cleaning and eating the meat and the development of leprosy (Clark et al. 2008, Deps et al. 2008, Truman 2008). More than half of the leprosy cases that have been reported in the southeastern USA have described some direct or indirect exposure to armadillos (Bruce et al. 2000, Truman et al. 2011) and other studies have raised the hypothesis that exposure to these animals could be a significant risk factor for leprosy in Brazil (Kerr-Pontes et al. 2006, Deps et al. 2008).

Unlike Brazil, where there are other species of armadillos, only *D. novemcinctus* is found in the USA. Among the 21 species of armadillos found in Brazil, *Euphractus sexcinctus* (six-banded armadillo) is commonly found. This species is known to eat animal carcasses, justifying

Financial support: CNPq (308539/2006-0), FUNCAP (303/05), CAPES (PROCAD 0204056)

+ Corresponding author: cristianefrota71@gmail.com

Received 3 April 2012

Accepted 11 July 2012

the local name of “vulture or gravedigger” and, because of this, hunters keep the animal alive and caged for several days before eating (Dalponte & Tavares-Filho 2004, Armadillo Online! 2012). This type of “quarantine” is believed to “clean” the animal, but instead enhances its contact with humans. In the state of Ceará (CE), located in Northeast Brazil, armadillos are used as a source of meat and hunting is seen as a leisure pursuit.

Brazil is a high leprosy-burdened country, with 34,894 new cases detected in 2010 (WHO 2011). More than half (53.5%) of the cases are concentrated in areas where only 17.5% of the population live, which reflects that leprosy is a rural disease (MS/SVS 2008). A study conducted with cases reported to the Brazilian Ministry of Health between 1990–2007 shows that leprosy is geographically concentrated, and a spatial analysis shows 29 clusters of higher prevalence. In these clusters, the mean rate of detection was more than the double the rate of the rest of the country (56.2 vs. 20.6 cases per 100,000 inhabitants). CE is one of the poorest states in Brazil and more than half of the municipalities in CE (46 out of 84) reported local transmission of new cases. CE was also included in one of the 29 Brazilian leprosy clusters (Penna et al. 2009). However, in Brazil, epidemiological studies are still controversial in terms of whether armadillo meat intake and direct animal contact are associated with leprosy infection (Kerr-Pontes et al. 2006, Deps et al. 2008, Schmitt et al. 2010). Aiming to investigate the question further, we used molecular diagnostic tools to search for *M. leprae* DNA in the two main species of wild armadillos found in CE (*D. novemcinctus* and *E. sexcinctus*).

MATERIALS AND METHODS

Armadillos - A total of 29 wild armadillos from two species (27 *D. novemcinctus* and 2 *E. sexcinctus*), consisting of males and females weighing from 2.6–3.8 kg, were captured between July–August 2007. The animals were captured by local hunters under the supervision of a veterinarian in rural sites of 12 selected endemic municipalities from CE (Figure). Biopsy samples from the ear, nose, liver and spleen of each of these animals were studied.


Euthanasia - Animal captures were authorised by the Brazilian Institute of Environmental and Renewable Natural Resources. Before euthanasia, armadillos were anaesthetised with tiletamine and zolazepam (5.0 mg/kg/1.M) (Virbac, Brazil). Euthanasia was conducted in the place of capture and the animals were kept in ice until they arrived at the laboratory in Fortaleza, the capital city of CE.

Sample biopsies and DNA extraction - Stringent precautions were necessary to avoid cross-contamination. Clean protective clothing was worn and gloves were changed frequently. Before the tissue dissection, the animals were carefully rinsed with distilled water. For each desired tissue, a sterile blade was used to cut. Ear, nose, liver and spleen samples were kept frozen at -20°C in separate sterile plastic storage bags until DNA extraction was performed. The sampling extraction was conducted batch-wise, four samples at a time. The frozen sections were incubated with 50 μL of 100 mM Tris-HCl, pH 8.5, containing Tween-20 and 60 μg of proteinase K per mL

for 18 h at 60°C . The digestion was conducted at 97°C for 15 min (de Wit et al. 1991). The extract was purified once with phenol/chloroform/isoamyl alcohol (25:24:1) and then precipitated with ethanol.

Polymerase chain reaction (PCR) and nucleotide sequencing - The *M. leprae*-specific repetitive element (RLEP) PCR was amplified in a nested PCR reaction. The primers for RLEP2-1 (5'-ATATCGATGCAGGCGT-GAG-3') and RLEP2-2 (5'-GGATCATCGATGCACT-GTTC-3') amplified a 282-bp sequence of the RLEP element. The second set of inner primers, RLEP2-3 (5'-GGGTAGGGGCGTTTTAGTGT-3') and RLEP2-2, amplified a 238-bp product. A 1 μL aliquot of the isolated DNA was added to 24 μL of PCR mix, which contained 15 mM Tris-HCl (pH 8.0), 50 mM KCl, 1.5 mM MgCl_2 , 0.2 mM dNTP, 5% DMSO, 1.25 units of *Taq* DNA Polymerase and 0.2 μM of each primer. The mixture was denatured at 94°C for 4 min, followed by 35 PCR cycles (30 s at 94°C , 30 s at 59.6°C and 1 min at 72°C), with a final extension at 72°C for 10 min. Each run included negative and positive controls. For the nested PCR, 0.5 μL of product was used as the DNA template. The amplification reactions were visualised on a 1.5% agarose gel. Each PCR sample was double-blind tested by different researchers in Fortaleza and Rio de Janeiro. When the results for the same sample were different, a third PCR was performed for confirmation. Different amounts of purified DNA from *M. leprae* (kindly donated by Dr R Truman, Louisiana State University, USA) were added to all negative PCR samples to assess the presence of inhibitory substances. A standard curve was constructed by serial dilution of purified *M. leprae* DNA ranging from 10 fg–1 μg . Purified *M. leprae* DNA was also used as a positive control for the amplifications.

The *M. leprae gyrA* region was amplified using primers *gyrAF* (5'-CCCGGACCGTAGCCACGCTAAGTC-3') and *gyrAR* (5'-CATCGCTGCCGGTGGGTCATTA-3'). The thermal profile involved an initial denaturation at 94°C for 5 min and six cycles of 94°C for 45 s, 68 to 63°C



Distribution map of the 12 places where armadillos were collected in the state of Ceará, Brazil.

for 45 s and 72°C for 90 s, followed by 35 cycles of 94°C for 45 s, 62°C for 45 s and 72°C for 90 s. The final extension was for 10 min at 72°C. The 187-bp PCR products were purified using the Invitrogen ChargeSwitch PCR Clean-Up kit prior to sequencing in an Applied Biosystems DNA sequencer (Perkin-Elmer Applied Biosystems) using a BigDye Terminator Cycle Sequencing kit. The sequences were identified using SecScape software v2.7 (Applied Biosystems). A reference *gyrA* sequence (GenBank accession NC002677) was used to align the sequences.

False-positive amplifications were addressed by using individual sterile section-cutting blades and sterile glassware for each biopsy sample. The armadillo biopsies and extracted DNA samples were carefully identified and kept in separate boxes. A simple DNA extraction protocol was established to minimise the risk of contamination. The extraction method used one purification step to reduce the accumulation of impurities that would inhibit the polymerase reaction.

Ethical considerations - This project was submitted to and approved by the Ethical Committee of the Federal University of Ceará.

RESULTS

A total of 116 liver, spleen, ear and nose tissue samples from 29 armadillos were tested using the nested RLEP PCR assay. *M. leprae* was detected in six (21%) of the animals; five were from the species *D. novemcinctus* (samples 8, 21, 22, 23 and 25) and one was from the species *E. sexcinctus* (sample 9) (Table I). *M. leprae* DNA was amplified in the ear biopsy samples of all six animals, but in only five of the liver or nose biopsy samples and three of the spleen biopsy samples (Table II).

The PCR amplification of *M. leprae* with the primers RLEP2-3 and RLEP2-2 was species-specific, targeting the same region of the primers developed by Donoghue et al. (2001). It generated a single band of 238 bp, a larger amplicon compared to the Donoghue primers that amplified a 99-bp product. In addition, the sequencing of the *gyrA* region (coding sequence location from 7515 to 11451 kb) was confirmatory for the presence of *M. leprae* DNA in the armadillos analysed. The detection limit of the nested PCR for *M. leprae* was 1 pg/μL (data not shown). All negative samples were amplified after adding 1 pg of purified *M. leprae* DNA. The exception was for one sample that was only amplified after a 1000-fold dilution, therefore confirming the presence of substances that were inhibitory to the PCR. The assay was repeated for the few isolates for which there were discrepancies between the researchers.

DISCUSSION

This is the first study to identify *M. leprae* in two species of wild armadillos (*D. novemcinctus* and *E. sexcinctus*) in Brazil. The tested armadillos came from an endemic leprosy area where the prevalence rate was reported to be 2.99/100,000 inhabitants in 2008 and where there is continuous contact between humans and these animals (MS/SVS 2011). In this region, the hunting and eating of armadillos is a popular and frequent practice, a situation similar to that described in Colombia (Cardona-Castro et al. 2009). It is noteworthy that previous studies

in the Americas and in the USA (Louisiana), Mexico, Colombia and Brazil had only reported *M. leprae* infection in armadillos of the species *D. novemcinctus* (Meyers et al. 1977, Truman et al. 1990, Deps et al. 2002).

Clinically, most of the animals exhibited nodule-like lesions indicative of leprosy or other degenerative diseases, as demonstrated by a clinical exam conducted by an expert veterinarian. However, no histopathological study was conducted, as the tissue samples were not formalin-fixed for anatomopathological examination. Additionally, no blood samples were taken from the animals. Therefore, we cannot conclude that the PCR-positive armadillos had clinical leprosy.

The genome sequencing of the Tamil Nadu *M. leprae* strain and other strains was conducted with a typing system based on single-nucleotide polymorphism (SNP) differences allowing continent distribution of the leprosy bacilli, which were classified as SNP types 1 to 4 (Monot et al. 2005). In addition, a correlation was observed between a mutation in the *gyrA* gene and the SNP types, which were clustered in *gyrAC* and *gyrAT* populations; the T SNP represents SNP type 3, while the C nucleotide represents the other three types. The results from the sequencing data demonstrated that the analysed samples from the armadillos in CE belong to the *gyrAT* (SNP type 3) population, which was also identified in samples from wild armadillos in Louisiana (Monot et al. 2005) and in humans in Brazil (Fontes et al. 2009). Although this is not yet definitive evidence that armadillos act as a source of

TABLE I
Polymerase chain reaction (PCR) results for the 29 wild armadillos analyzed according to the armadillo species in the state of Ceará, Brazil

Armadillo species	Total (n)	RLEP PCR positive n (%)
<i>Dasyus novemcinctus</i>	27	5 (19)
<i>Euphractus sexcinctus</i>	2	1 (50)
Total	29	6 (21)

RLEP: *Mycobacterium leprae*-specific repetitive element.

TABLE II

Distribution of positive polymerase chain reaction (PCR) results using the *Mycobacterium leprae*-specific repetitive element repetitive sequence primer pairs in different biopsy tissues from wild armadillos in the state of Ceará, Brazil

Armadillo species	Biopsy source			
	Liver	Spleen	Ear	Nose
<i>Dasyus novemcinctus</i> (n = 5)	4	2	5	4
<i>Euphractus sexcinctus</i> (n = 1)	1	1	1	1

infection for humans, this contributes to the evidence supporting the hypothesis of zoonotic transmission, as suggested by other authors (Job et al. 1986, Walsh et al. 1988, Cardona-Castro et al. 2009, Truman & Fine 2010).

Because *M. leprae* cannot be cultivated in vitro, its detection is based on the histopathological demonstration of the bacilli in dermal nerves, mouse footpad cultivation and PCR assays of the selective amplification of *M. leprae* DNA (Truman & Fine 2010). This study used the RLEP repetitive element sequence of *M. leprae*, which is reported to be specific for *M. leprae* and is not present in other mycobacterial or bacterial species. In addition, the use of the repetitive sequence as the PCR target DNA provides the advantage of higher sensitivity over other targets because it is present at multiple sites in the genomic DNA (Donoghue et al. 2001, Truman et al. 2008). It has been suggested that many homologous sequences of the RLEP may be present in other environmental *Mycobacterium* species that have not been thoroughly investigated, which might generate false-positive results. Despite this fact, Martinez et al. (2011) found that the RLEP PCR assay can be used as a more specific and sensitive diagnostic test to detect *M. leprae* infection compared to the ones based on gene targets *Ag 85B*, *sodA* and 16S rRNA. Because we used several strategies to minimise false-positive amplifications and contamination, we are confident that these are real infections.

In conclusion, the presence of *M. leprae* DNA in wild armadillos (*D. novemcinctus* and *E. sexcinctus*) in a leprosy transmission area in Brazil provides additional evidence supporting the hypothesis that armadillos can play a role as an environmental reservoir for the bacillus in this area. Moreover, the finding supports the idea that intensive contact with these animals may increase the risk of infection in CE. The current global control strategy depends on treating all human cases, but a definitive identification of an animal reservoir, as suggested by Truman et al. (2011), could challenge this strategy and may partially explain the continuing and growing presence of leprosy in the studied area. More detailed molecular studies will be useful in monitoring and confirming the transmission of *M. leprae* between wild armadillos and humans and in guiding new strategies for prevention.

REFERENCES

- Amezcuca ME, Escobar-Gutierrez A, Storrs EE, Dhople AM, Burchfield HP 1984. Wild Mexican armadillo with leprosy-like infection. *Int J Lepr Other Mycobact Dis* 52: 254-255.
- Armadillo Online! 2012. [homepage on the Internet]. Minneapolis. Armadillos: Biology, Ecology and Images. Michigan State University Department of Zoology. [cited 2012 March 13]. Available from: msu.edu/~nixonjos/armadillo/species.html.
- Bruce S, Schroeder TL, Ellner K, Rubin H, Williams T, Wolf Jr JE 2000. Armadillo exposure and Hansen's disease: an epidemiologic survey in southern Texas. *J Am Acad Dermatol* 43: 223-228.
- Cardona-Castro N, Beltran-Alzate JC, Romero-Montoya IM, Melendez E, Torres F, Sakamuri RM, Li W, Vissa V 2009. Identification and comparison of *Mycobacterium leprae* genotypes in two geographical regions of Colombia. *Lepr Rev* 80: 316-321.
- Clark BM, Murray CK, Horvath LL, Deye GA, Rasnake MS, Longfield RN 2008. Case-control study of armadillo contact and Hansen's disease. *Am J Trop Med Hyg* 78: 962-967.
- Dalponte JC, Tavares-Filho JA 2004. Diet of the yellow armadillo, *Euphractus sexcinctus*, in south-central Brazil. *Edentata* 6: 37-41.
- de Wit MY, Faber WR, Krieg SR, Douglas JT, Lucas SB, Montreewasuwat N, Pattyn SR, Hussain R, Ponnighaus JM, Hartskeerl RA 1991. Application of a polymerase chain reaction for the detection of *Mycobacterium leprae* in skin tissues. *J Clin Microbiol* 29: 906-910.
- Deps PD, Alves BL, Gripp CG, Aragao RL, Guedes B, Filho JB, Andreatta MK, Marcari RS, Prates I, Rodrigues LC 2008. Contact with armadillos increases the risk of leprosy in Brazil: a case control study. *Indian J Dermatol Venereol Leprol* 74: 338-342.
- Deps PD, Antunes JM, Tomimori-Yamashita J 2007. Detection of *Mycobacterium leprae* infection in wild nine-banded armadillos (*Dasypus novemcinctus*) using the rapid ML Flow test. *Rev Soc Bras Med Trop* 40: 86-87.
- Deps PD, Santos AR, Yamashita-Tomimori J 2002. Detection of *Mycobacterium leprae* DNA by PCR in blood sample from nine-banded armadillo: preliminary results. *Int J Lepr Other Mycobact Dis* 70: 34-35.
- Donoghue HD, Holton J, Spigelman M 2001. PCR primers that can detect low levels of *Mycobacterium leprae* DNA. *J Med Microbiol* 50: 177-182.
- Fontes AN, Sakamuri RM, Baptista IM, Ura S, Moraes MO, Martinez AN, Sarno EN, Brennan PJ, Vissa VD, Suffys PN 2009. Genetic diversity of *Mycobacterium leprae* isolates from Brazilian leprosy patients. *Lepr Rev* 80: 302-315.
- Job CK, Drain V, Truman R, Deming AT, Sanchez RM, Hastings RC 1992. The pathogenesis of leprosy in the nine-banded armadillo and the significance of IgM antibodies to PGL-1. *Indian J Lepr* 64: 137-151.
- Job CK, Harris EB, Allen JL, Hastings RC 1986. A random survey of leprosy in wild nine-banded armadillos in Louisiana. *Int J Lepr Other Mycobact Dis* 54: 453-457.
- Kerr-Pontes LR, Barreto ML, Evangelista CM, Rodrigues LC, Heukelbach J, Feldmeier H 2006. Socioeconomic, environmental and behavioural risk factors for leprosy in Northeast Brazil: results of a case-control study. *Int J Epidemiol* 35: 994-1000.
- Kirchheimer WF, Sanchez RM 1977. Quantitative aspects of leprosy in armadillos. *Lepr India* 49: 48-53.
- Kirchheimer WF, Storrs EE, Binford CH 1972. Attempts to establish the Armadillo (*Dasypus novemcinctus* linn.) as a model for the study of leprosy. II. Histopathologic and bacteriologic post-mortem findings in lepromatoid leprosy in the Armadillo. *Int J Lepr Other Mycobact Dis* 40: 229-242.
- Martinez AN, Ribeiro-Alves M, Sarno EN, Moraes MO 2011. Evaluation of qPCR-based assays for leprosy diagnosis directly in clinical specimens. *PLoS Negl Trop Dis* 5: e1354.
- Meyers WM, Walsh GP, Brown HL, Rees RJ, Convit J 1977. Naturally acquired leprosy-like disease in the nine-banded armadillo (*Dasypus novemcinctus*): reactions in leprosy patients to lepromins prepared from naturally infected armadillos. *J Reticuloendothel Soc* 22: 369-375.
- Monot M, Honore N, Garnier T, Araoz R, Coppee JY, Lacroix C, Sow S, Spencer JS, Truman RW, Williams DL, Gelber R, Virmond M, Flageul B, Cho SN, Ji B, Paniz-Mondolfi A, Convit J, Young S, Fine PE, Rasolofy V, Brennan PJ, Cole ST 2005. On the origin of leprosy. *Science* 308: 1040-1042.
- MS/SVC - Ministério da Saúde/Secretaria de Vigilância em Saúde 2008. Situação epidemiológica da hanseníase no Brasil. Programa Nacional de Controle de Hanseníase. [cited 2012 March 13]. Available from: portal.saude.gov.br/portal/arquivos/pdf/bol-letim_novembro.pdf.

- MS/SVC - Ministério da Saúde/Secretaria de Vigilância em Saúde 2011. Taxa de prevalência de hanseníase 2011. [updated 2011 June; cited 2012 March 13]. Available from: tabnet.datasus.gov.br/cgi/idx2010/matriz.htm#morb.
- Penna ML, de Oliveira ML, Penna GO 2009. The epidemiological behaviour of leprosy in Brazil. *Lepr Rev* 80: 332-344.
- Schmitt JV, Dechandt IT, Dopke G, Ribas ML, Cerci FB, Viesi JMZ, Marchioro HZ, Zunino MMB, Miot HA 2010. pArmadillo meat intake was not associated with leprosy in a case control study, Curitiba (Brazil). *Mem Inst Oswaldo Cruz* 105: 857-862.
- Sheppard CC 1960. The experimental disease that follows the injection of human leprosy bacilli into footpads of mice. *J Expect Med* 112: 445.
- Sheppard CC 1985. Experimental leprosy. In RC Hastings (ed.), *Leprosy*, Longman Group Limited, New York, p. 269-286.
- Smith JH, Long EG, Crouse DT, Christie JD, Folse DS, Imaeda T, Barksdale L 1983. Leprosy in wild armadillos (*Dasybus novemcinctus*) of the Texas Gulf Coast. *Acta Leprol* 2: 311-318.
- Stallknecht DE, Truman RW, Hugh-Jones ME, Job CK 1987. Surveillance for naturally acquired leprosy in a nine-banded armadillo population. *J Wildl Dis* 23: 308-310.
- Truman R 2008. Armadillos as a source of infection for leprosy. *South Med J* 101: 581-582.
- Truman RW, Andrews PK, Robbins NY, Adams LB, Krahenbuhl JL, Gillis TP 2008. Enumeration of *Mycobacterium leprae* using real-time PCR. *PLoS Negl Trop Dis* 2: e328.
- Truman RW, Fine PE 2010. "Environmental" sources of *Mycobacterium leprae*: issues and evidence. *Lepr Rev* 81: 89-95.
- Truman RW, Job CK, Hastings RC 1990. Antibodies to the phenolic glycolipid-1 antigen for epidemiologic investigations of enzootic leprosy in armadillos (*Dasybus novemcinctus*). *Lepr Rev* 61: 19-24.
- Truman RW, Kumaresan JA, McDonough CM, Job CK, Hastings RC 1991. Seasonal and spatial trends in the detectability of leprosy in wild armadillos. *Epidemiol Infect* 106: 549-560.
- Truman RW, Singh P, Sharma R, Busso P, Rougemont J, Paniz-Mondolfi A, Kapopoulou A, Brisse S, Scollard DM, Gillis TP, Cole ST 2011. Probable zoonotic leprosy in the southern United States. *N Engl J Med* 364: 1626-1633.
- Walsh GP, Meyers W, Binford CH 1977. Naturally acquired leprosy-like disease in the nine-banded armadillo (*Dasybus novemcinctus*): recent epizootiologic findings. *J Reticuloendothel Soc* 22: 363-367.
- Walsh GP, Meyers WM, Binford CH, Gormus BJ, Baskin GB, Wolf RH, Gerone PJ 1988. Leprosy as a zoonosis: an update. *Acta Leprol* 6: 51-60.
- WHO - World Health Organization 2011. Weekly epidemiological record. Available from: who.int/wer/2011/wer8636.pdf.
- Zumarraga MJ, Resoagli EH, Cicuta ME, Martinez AR, Oritiz de Rott MI, de Millan SG, Caimi K, Gioffre A, Alito A, Bigi F, Cataldi AA, Romano MI 2001. PCR-restriction fragment length polymorphism analysis (PRA) of *Mycobacterium leprae* from human lepromas and from a natural case of an armadillo of Corrientes, Argentina. *Int J Lepr Other Mycobact Dis* 69: 21-25.