

Original citation:

Cheng, Wei, Rolls, Edmund T., Qiu, Jiang, Liu, Wei, Tang, Yanqing, Huang, ChuChung, Wang, XinFa, Zhang, Jie, Lin, Wei, Zheng, Lirong, Pu, JunCai, Tsai, Shih-Jen, Yang, Albert C., Lin, Ching-Po, Wang, Fei, Xie, Peng and Feng, Jianfeng. (2016) Medial reward and lateral non-reward orbitofrontal cortex circuits change in opposite directions in depression. *Brain* . doi: 10.1093/brain/aww255

Permanent WRAP URL:

<http://wrap.warwick.ac.uk/81765>

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions. Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Publisher's statement:

This is a pre-copyedited, author-produced PDF of an article accepted for publication in *Brain* following peer review. The version of record Cheng, Wei, Rolls, Edmund T., Qiu, Jiang, Liu, Wei, Tang, Yanqing, Huang, ChuChung, Wang, XinFa, Zhang, Jie, Lin, Wei, Zheng, Lirong, Pu, JunCai, Tsai, Shih-Jen, Yang, Albert C., Lin, Ching-Po, Wang, Fei, Xie, Peng and Feng, Jianfeng. (2016) Medial reward and lateral non-reward orbitofrontal cortex circuits change in opposite directions in depression. *Brain* is available online at:

<http://dx.doi.org/10.1093/brain/aww255>

A note on versions:

The version presented here may differ from the published version or, version of record, if you wish to cite this item you are advised to consult the publisher's version. Please see the 'permanent WRAP url' above for details on accessing the published version and note that access may require a subscription.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk

Medial reward and lateral non-reward orbitofrontal cortex circuits change in opposite directions in depression

Wei Cheng^{1, #}, Edmund T. Rolls^{2,3, #}, Jiang Qiu^{4,5, #}, Wei Liu^{4,5, #}, Yanqing Tang^{6, #}, Chu-Chung Huang^{7, #}, XinFa Wang^{8,9,10, #}, Jie Zhang¹, Wei Lin¹, Lirong Zheng^{14,15}, JunCai Pu^{8,9,10}, Shih-Jen Tsai¹¹, Albert C Yang¹¹, Ching-Po Lin^{7, *}, Fei Wang^{6, *}, Peng Xie^{8,9,10, *}, Jianfeng Feng^{1, 2, 12, 13, *}

1. School of Mathematical Sciences and Centre for Computational Systems Biology, Fudan University, Shanghai, 200433, PR China
2. Department of Computer Science, University of Warwick, Coventry CV4 7AL, UK
3. Oxford Centre for Computational Neuroscience, Oxford, UK
4. Key Laboratory of Cognition and Personality (SWU), Ministry of Education, Chongqing, China
5. Department of Psychology, Southwest University, Chongqing, China
6. Department of Psychiatry, The First Affiliated Hospital, China Medical University, Shenyang 110001, Liaoning, PR China
7. Institute of Neuroscience, National Yang-Ming University, Taipei, Taiwan
8. Institute of Neuroscience, Chongqing Medical University, Chongqing, China
9. Chongqing Key Laboratory of Neurobiology, Chongqing, China
10. Department of Neurology, 1st Affiliated Hospital, Chongqing Medical University, Chongqing, China
11. Department of Psychiatry, Taipei Veterans General Hospital, Taipei, Taiwan
12. School of Mathematical Sciences, School of Life Science and the Collaborative Innovation Center for Brain Science, Fudan University, Shanghai, 200433, PR China
13. School of Life Sciences, Fudan University, Shanghai, 200433, PR China
14. School of Information Science and Engineering, Fudan University, Shanghai, China
15. KTH - Royal Institute of Technology, Sweden.

These authors contributed equally to this work.

Short title: orbitofrontal cortex and depression

Keywords: depression; orbitofrontal cortex; functional connectivity; resting state functional neuroimaging; medial temporal lobe; precuneus

* Corresponding author information:

Professor Jianfeng Feng

Centre for Computational Systems Biology,

School of Mathematical Sciences, Fudan University, Shanghai 200433, China, and

Department of Computer Science, University of Warwick, Coventry CV4 7AL, UK

Tel. 86-21-65643621

Email: jianfeng64@gmail.com

Brain (2016) doi 10.1093/brain/aww255

Abstract

The first brain-wide voxel-level resting state functional-connectivity neuroimaging analysis of depression is reported, with 421 patients with major depressive disorder and 488 controls. Resting state functional connectivity between different voxels reflects correlations of activity between those voxels and is a fundamental tool in helping to understand the brain regions with altered connectivity and function in depression.

One major circuit with altered functional connectivity involved the medial orbitofrontal cortex BA 13, which is implicated in reward, and which had reduced functional connectivity in depression with memory systems in the parahippocampal gyrus and medial temporal lobe, especially involving the perirhinal cortex BA 36 and entorhinal cortex BA 28. The Hamilton Depression Rating Scale scores were correlated with weakened functional connectivity of the medial orbitofrontal cortex BA 13. Thus in depression there is decreased reward-related and memory system functional connectivity, and this is related to the depressed symptoms. The lateral orbitofrontal cortex BA 47/12, involved in non-reward and punishing events, did not have this reduced functional connectivity with memory systems.

Second, the lateral orbitofrontal cortex BA 47/12 had increased functional connectivity with the precuneus, the angular gyrus, and the temporal visual cortex BA 21. This enhanced functional connectivity of the non-reward/punishment system (BA 47/12) with the precuneus (involved in the sense of self and agency), and the angular gyrus (involved in language) is thus related to the explicit affectively negative sense of the self, and of self-esteem, in depression. A comparison of the functional connectivity in 185 depressed patients not receiving medication and 182 patients receiving medication showed that the functional connectivity of the lateral orbitofrontal cortex BA 47/12 with these three brain areas was lower in the medicated than the unmedicated patients. This is consistent with the hypothesis that the increased functional connectivity of the lateral orbitofrontal cortex BA 47/12 is related to depression.

Relating the changes in cortical connectivity to our understanding of the functions of different parts of the orbitofrontal cortex in emotion helps to provide new insight into the brain changes related to depression, which are considered in the Discussion.

Introduction

Major depressive disorder is ranked by the World Health Organization as the leading cause of years-of-life lived with disability (Drevets, 2007, Gotlib and Hammen, 2009, Hamilton *et al.*, 2013). Major depressive episodes, found in both major depressive disorder and bipolar disorder are pathological mood states characterized by persistently sad or depressed mood. Major depressive disorders are generally accompanied by: (a) altered incentive and reward processing, evidenced by amotivation, apathy, and anhedonia; (b) impaired modulation of anxiety and worry, manifested by generalized, social and panic anxiety, and oversensitivity to negative feedback; (c) inflexibility of thought and behavior in association with changing reinforcement contingencies, apparent as ruminative thoughts of self-reproach, pessimism, and guilt, and inertia toward initiating goal-directed behavior; (d) altered integration of sensory and social information, as evidenced by mood-congruent processing biases; (e) impaired attention and memory, shown as performance deficits on tests of attention set-shifting and maintenance, and autobiographical and short-term memory; and (f) visceral disturbances, including altered weight, appetite, sleep, and endocrine and autonomic function (Drevets, 2007, Gotlib and Hammen, 2009).

The ability to measure brain function using non-invasive neuroimaging techniques has greatly enhanced our understanding of this illness (Rigucci *et al.*, 2010). Patients with depression show impairments in the coordinated activity of several brain regions considered to be important for several domains of mental functioning such as emotional processing (amygdala, subgenual anterior cingulate and pallidum) (Disner *et al.*, 2011, Sheline *et al.*, 2010), self-referential processes (medial prefrontal cortex (MPFC), precuneus and posterior cingulate cortex) (Kuhn and Gallinat, 2013, Price and Drevets, 2010, Sheline *et al.*, 2010), cognitive functions such as memory (hippocampus, parahippocampal cortex) (Lorenzetti *et al.*, 2009), visual processing (fusiform gyrus, lingual gyrus and lateral temporal cortex) (Veer *et al.*, 2010), and attention processing (dorsolateral prefrontal cortex, anterior cingulate cortex, thalamus and insula) (Hamilton *et al.*, 2012).

Research into the pathophysiology of depression has included the analysis of possible differences in the functional connectivity of different brain areas to elucidate some of the brain changes that may relate to depression. Resting-state fMRI provides a task-free approach that removes some performance-related confounds, and provides a reliable measure of ‘baseline’ brain activity and connectivity (Gusnard *et al.*, 2001). The functional connectivity is measured by the correlation between the fMRI BOLD signals in different brain areas when in the resting state, that is when no task is being performed. The concept is that the correlations may reveal evidence about which brain systems may interact differently in neuropsychiatric disorders (Deco and Kringelbach, 2014). There have been a number of resting state functional connectivity studies on depression (Iwabuchi *et al.*, 2015, Wang *et al.*, 2012). Most studies do not include large numbers of participants, and therefore there are insufficient data to allow voxel-level analysis, though this can be very important in helping to reveal exactly which cortical systems may be connected differently in mental disorders (Cheng *et al.*, 2015). There is an urgent need to use methods that will allow large-scale pooling of data to increase the statistical power to obtain voxel-level analysis, as well as to reduce the impact of heterogeneity in the patient population. A meta-analysis of previous investigations of resting state functional connectivity in depression was based on seed based studies each with tens of participants, and concluded as follows (Kaiser *et al.*, 2015). “Major depressive disorder was characterized by hypoconnectivity within the frontoparietal network, a set of regions involved in cognitive control of attention and emotion regulation, and hypoconnectivity between frontoparietal systems and parietal regions of the dorsal attention network involved in attending to the

external environment. Major depressive disorder was also associated with hyperconnectivity within the default network, a network believed to support internally oriented and self-referential thought, and hyperconnectivity between frontoparietal control systems and regions of the default network. Finally, the MDD groups exhibited hypoconnectivity between neural systems involved in processing emotion or salience and midline cortical regions that may mediate top-down regulation of such functions.” For comparison, the present study included almost as many participants as this meta-analysis, was not forced because of small numbers of participants to rely on *a priori*, seed-based analyses, and was able given the voxel-based approach to focus on particular brain regions, rather than brain systems identified for example as the ‘default mode network’ or ‘fronto-parietal control systems’.

Given this background, the objective of this investigation is to perform the first investigation using a voxel-based unbiased brain-wide association study (BWAS) approach on resting-state functional magnetic resonance imaging (fMRI) data in patients with major depressive disorder. The BWAS approach is modeled along the lines of genome-wide association studies (GWAS) where large genetic data sets are pooled to identify significant genetic variations in specific disorders. An aim of this investigation is to include a large number of participants in this neuroimaging research, to enable voxel-level accuracy, and robustness of the findings (Button *et al.*, 2013). In this investigation, the voxel-level resolution of the functional connectivity enabled differences of functional connectivity to be measured in nearby but functionally different parts of the orbitofrontal, and to reveal to exactly which voxels in other brain areas there was altered functional connectivity. The voxel-level analysis enabled this advance to be made. The value of the unbiased approach was that it enabled the functional connectivity between every pair of voxels in the brain to be measured, so that the findings were not limited by prior hypotheses.

Methods

Participants

There were 421 patients with a diagnosis of major depression, and 488 controls. The patients were from Taiwan (Veteran General Hospital, Taipei), Dongei (Department of Psychiatry, First Affiliated Hospital of China Medical University and the Mental Health Center of Shenyang, China) and Xinan (First Affiliated Hospital of Chongqing Medical School in Chongqing, China). All participants were diagnosed according to the Diagnostic and Statistical Manual of Mental Disorder-IV criteria for major depressive disorder. Depression severity and symptomatology were evaluated by the Hamilton Depression Rating Scale (HAMD, 17 items) (Hamilton, 1960) and the Beck Depression Inventory (BDI) (Beck and Beamesderfer, 1974). 185 of the patients were not receiving medication at the time of the neuroimaging. Table 1 provides a summary of the demographic information and the psychiatric diagnosis (showing how they were diagnosed) of the participants. Further details are provided in the Supplementary Material.

Image Acquisition and Preprocessing

Data for resting state functional connectivity analysis were collected in 3T MRI scanners in an 8 min period in which the participants were awake in the scanner not performing a task using standard protocols described in the Supplementary Material.

Data preprocessing was performed using DPARSF (Chao-Gan and Yu-Feng, 2010) (<http://restfmri.net>) which is a toolbox based on the SPM8 software package. The first 10 EPI scans were discarded to suppress equilibration effects. The remaining scans of each subject

underwent slice timing correction by sinc interpolating volume slices, motion correction for volume to volume displacement, spatial normalization to standard Montreal Neurological Institute (MNI) space using affine transformation and nonlinear deformation with a voxel size of $3 \times 3 \times 3\text{mm}^3$, followed by spatial smoothing (8 mm Full Width Half Maximum FWHM). To remove the sources of spurious correlations present in resting-state BOLD data, all fMRI time-series underwent band-pass temporal filtering (0.01-0.1 Hz), nuisance signal removal from the ventricles, and deep white matter, and regressing out any effects of head motion using the Friston et al 24 head motion parameters procedure (Friston *et al.* , 1996). Finally, we implemented additional careful volume censoring (“scrubbing”) movement correction as reported by Power et al. (Power *et al.* , 2014) to ensure that head-motion artifacts are not driving observed effects. The mean framewise displacement (FD) was computed with FD threshold for displacement being 0.5. In addition to the frame corresponding to the displaced time point, 1 preceding and 2 succeeding time points were also deleted to reduce the ‘spill-over’ effect of head movements. Subjects with >10% displaced frames flagged were completely excluded from the analysis as it is likely that such high-level of movement would have had an influence on several volumes. Global signals were not regressed out (see Supplementary Material, where the results with global signal removal are referred to for completeness).

Any effects of gender ratio, years of education, and age between the patient and control groups were regressed out in the analysis. In fact, there were no differences in the gender ratios, though the number of years of education was lower in the patients than controls. Additional analyses showed for males vs females that the overall pattern of functional connectivity differences for patients vs controls were similar, and that the correlation of the functional connectivity changes between males and females was high (0.89, $p < 0.0001$). Further, none of the functional connectivity link differences found between patients and controls was correlated significantly (FDR $p < 0.05$) with the number of years of education. We also note that the Taiwanese sample included patients with depression in remission while under antidepressant treatment, and thus their scores on the Hamilton Depression Rating Scale (HAMD) assessment were in the low range.

Voxel-wise brain-wide association studies (vBWAS)

Step 1: analysis within each imaging centre.

In the present study, each resting-state fMRI image included 47,619 voxels, which is based on the automated anatomical labeling (AAL2) atlas (Rolls *et al.* , 2015). For each pair of voxels, the time series were extracted and their correlation was calculated for each subject followed by z-transformation. Two-tailed, two-sample t-tests were performed on the 1,133,760,771 ($47619 \times 47618 / 2$) Fisher’s z-transformed correlation coefficients to identify significantly altered functional links in patients with depression compared to controls within each imaging centre. The effect of age, gender ratios, education and head motion (mean FD) were regressed within each dataset in this step.

Step 2: combination of results from all imaging centres.

The Liptak-Stouffer z score method (Liptak, 1958) which is a well-validated method for multi-site datasets and has previously been used widely in multi-site MRI data analysis (Glahn *et al.* , 2008, Yu *et al.* , 2011) was then used to combine the results from the individual datasets. Specifically, the p-value of each functional connectivity result from two-sample t-test in step 1 was converted to its corresponding z score. This was calculated firstly as in equation: $z_i = \Phi^{-1}(1 - p_i)$, where Φ is the standard normal cumulative distribution function and i

represent the i site. Next, a combined z score for a functional connectivity was calculated using the Liptak-Stouffer formula:

$$Z = \frac{\sum_{i=1}^k w_i z_i}{\sqrt{\sum_{i=1}^k w_i^2}}$$

which follows a standard normal distribution under the null hypothesis; where $w_i = \sqrt{\text{sample size}}$ is the weight of the i dataset. Finally, The Z is transformed into its corresponding p-value and a FDR procedure was used to correct for multiple comparisons.

Step 3: calculating a measure for the association of voxels.

A measure for the association (MA) between a voxel i and the brain disorder was then defined as: $MA = N_\alpha$, where N_α is the number of links between voxel i and every other voxel in the brain that have a p-value of less than α (which in the present study with FDR correction was $p < 0.01$, corresponding to a p threshold of 2.52×10^{-7}) in t-tests. A larger value of MA implies a more significant difference in functional connectivity.

For the functional connectivity of a voxel to be treated as significantly different ($p < 0.01$) after FDR correction from another voxel, the significance level uncorrected had to be $p < 2.52 \times 10^{-7}$. The smallest p value found was approximately 10^{-13} . Clusters with less than 10 voxels were not included to reduce the possibility of false positive results and to ensure that only consistent differences in functional connectivity were considered, following earlier practice (Hart *et al.*, 2012, Konrad *et al.*, 2006, Wittmann *et al.*, 2005).

Although the voxel-level brain-wide association study (BWAS) identifies all altered voxel-wise different functional connectivities in patients with depression, it is difficult to describe and show all of these changed links. Accordingly, to facilitate the description of the voxel-wise results, we conducted post-hoc cluster-wise analyses from each cluster of voxels returned by BWAS. It should be noted that all cluster-wise analyses are based on the findings of BWAS, and that it is the BWAS statistics only on which we rely. The cluster analyses just simplify the description of the different functional connectivities in depression.

Clinical correlates

We also investigated whether the differences in functional connectivity between patients and controls were correlated with clinical variables (the Hamilton depression rating Scale (HAMD) (Hamilton, 1960), Beck Depression Inventory (BDI) (Beck and Beamesderfer, 1974), and illness duration (Bell-McGinty *et al.*, 2002, de Diego-Adelino *et al.*, 2014). We used the Liptak-Stouffer z score method (Liptak, 1958) to combine the data from the different neuroimaging sites for this analysis, for this provides a principled way to take into consideration possible differences in these measures between sites. Specifically, we calculated the partial correlation between the strength of each altered functional connection with HAMD, BDI score, and with illness duration after removing the effect of sex and age, in each individual centre, then we used the Liptak-Stouffer z score method to combine the results from the individual datasets.

Results

The fMRI resting state functional connectivity analyses were performed with 421 patients with a diagnosis of major depression, and 488 controls, and this large population was sufficient to allow voxel-level analysis with fully corrected statistics.

A Brain-Wide Association Study (BWAS) of voxels with different functional connectivity in depressed patients. As shown in Figure 1 and Table 2, there were a number of voxel clusters with different functional connectivity (FC) in patients with depression compared to controls.

A large cluster of voxels (274) was in the left medial temporal lobe (MedTL) with peak at [X Y Z -27 -9 -36]. This cluster was in the parahippocampal gyrus, extending into the fusiform gyrus and inferior temporal and temporal pole areas, and indeed these voxels were classified as in areas in the AAL2 atlas (Rolls *et al.* , 2015) that included left hippocampus, parahippocampal gyrus, fusiform gyrus, middle temporal gyrus, temporal pole (middle temporal gyrus), and inferior temporal gyrus (Table 2). The voxel with the highest measure of association (*MA*, the number of voxels with which a voxel has a functional connectivity difference significant at $p < 0.01$ (FDR correction)) was 284 (Table 2). It is clear from Fig. 1 that the majority of these voxels were in the perirhinal cortex BA 36, and the entorhinal cortex BA 28. There was a corresponding cluster on the right (with peak at [24 12 -39], Table 2).

A second large cluster of voxels was in the medial orbitofrontal cortex Brodmann Area 13 with peak at [-15 12 -15] (Table 2 and Fig. 1). In the AAL2 atlas this included left Olfactory, Rectus, OFCmed, OFCant, OFCpost, Insula, but the locations of these voxels as shown in Fig. 1 was in area 13 just extending anteriorly in area 11, according to the cytoarchitectonic designation (Öngür *et al.* , 2003) (see e.g. Fig. 4 in that paper, which shows that area 13 extends quite far posterior towards what topologically is close to the anterior ventral insula). There was a corresponding cluster on the right (with peak at [15 24 -18]).

A third cluster, with a high peak *MA* value of 136, was in lateral orbitofrontal cortex area 47/12 (Öngür *et al.* , 2003) with peak at [36 36 -12]. In Fig. 1 it can be seen in slices at Y=32, 35 and 37. It was in AAL2 region Frontal_Inf_Orb_2_R.

A fourth cluster was in the anterior cingulate cortex in the just supracallosal part with peak at [-6 36 21] (Fig. 1 and Table 2).

A fifth cluster was in the precuneus and adjoining posterior cingulate cortex with peak at [-6 -54 30] (Fig. 1 and Table 2).

A sixth cluster was in the left angular gyrus, with peaks at [-48 -60 21] and [-48 -69 42].

A seventh cluster was in the middle temporal gyrus, in a part that is BA 21 and is a high order temporal cortical visual area involved in processing objects and faces.

Other minor clusters in the thalamus and postcentral cortex are indicated in Table 2.

Analysis of the functional connectivity links that were different in patients with depression. To investigate the brain areas between which there was different functional connectivity in depression, and whether it was increased or decreased, the functional connectivity (FC) of the voxels with significant differences of FC (after FDR correction at $p < 0.05$, and within the voxel clusters shown in Table 2) were measured for each of the AAL2 regions within which the voxels were located. In this way, 23 regions of interest were identified, and the functional connectivity differences between the significantly different voxels in these regions are shown in Fig. 2 as a connectivity matrix.

First the altered functional connectivity of the medial orbitofrontal cortex, area 13, brain region is considered. The relevant voxels are in regions such as OFCmed, OFCant, OFCpost, Olfactory, Rectus, and Insula in the AAL2 atlas on the left and the right (Table 2). The voxels within this area 13 cluster have high positive correlations between them, and have generally the same pattern of altered functional connectivity in depression, so is described in the remainder of this paper as OFC13. As shown in Fig. 2, OFC13 has very significantly reduced functional connectivity with the Parahippocampal, Fusiform, Temporal-Pole and temporal_Inf

areas (and many of the voxels in these areas are in the perirhinal cortex area 36 and the entorhinal cortex area 28, as shown in Fig. 1), which again act similarly in this investigation, so are referred to as the Medial Temporal Lobe (MedTL) cluster in the remainder of this paper. OFC13 also has reduced functional connectivity between the voxels in its different AAL2 regions (Fig. 2). Some parts of OFC13, in particular OFCPost, has increased functional connectivity with the precuneus.

Second, OFC47/12 on the right (Frontal_Inf_Orb_2_R in the AAL2 atlas) has *increased* functional connectivity with the left Angular gyrus, left Precuneus, and left and right Temporal_Mid (e.g. [66 -15 -18], which is Brodmann area 21, temporal visual cortex). OFC47/12 does *not* have reduced functional connectivity with the parahippocampal areas (Fig. 2). This lateral orbitofrontal cortex network is therefore very different indeed in its change in functional connectivity in depression from the OFC13 network in the medial orbitofrontal cortex.

Third, the anterior cingulate cortex (in which the voxels are just supracallosal) has reduced functional connectivity in depressed patients with some temporal cortex areas including the fusiform gyrus, with the Angular cortex, and no difference in functional connectivity with most medial orbitofrontal areas (OFC13) (apart from a small increase with OFC_Med_R).

Other functional connectivity changes include increased thalamic connectivity with some medial orbitofrontal, and parahippocampal/temporal cortex regions, and the postcentral gyrus (Fig. 2).

Clinical symptom correlates of the altered circuits. As can be seen from Table 3, there were significant correlations ($P < 0.05$ uncorrected) between some of the region of interest-wise functional links and the symptom severity scores and illness duration. Specifically, the HAMD (Hamilton Depression Rating Scale (Hamilton, 1960)) score was correlated with weakened functional connectivity between some of the areas within the medial orbitofrontal cortex (OFC13). Analysis of the subscores of the HAMD showed that links involving OFC13 were correlated with most of the 17 subscores apart from 10 and 15. Interestingly, links involving OFC47/12 were negatively correlated with H6 Insomnia (waking up early), and with H17 Insight.

The Beck Depression Inventory score (available only for the Xinan dataset of 183 patients) was also correlated with decreased functional connectivity between several of the medial orbitofrontal cortex OFC13 subregions and several of the Parahippocampal / Temporal subregions; and also with decreased functional connectivity of the temporal areas (Temporal_Inf_L and Temporal_Pole_L), and of Temporal_Mid-R with the PostCentral gyrus (Table 3). (For the Xinan dataset, which includes both HAMD and BDI scores, the Pearson correlation coefficient between each ROI-wise Functional Connectivity and either the HAMD or the BDI reached 0.41 ($p=1.6e-11$).

The illness duration (Table 3) was negatively correlated with functional connectivity between the lateral orbitofrontal cortex OFC47/12 voxels and the left Angular gyrus and Temporal-Mid_L cluster of Table 3. The illness duration was also correlated with weaker functional connectivity between the posterior part of the medial orbitofrontal cortex OFC13 cluster (specifically the part within Insula_L in the AAL2 atlas) and some Parahippocampal/Temporal areas (Table 3). Illness duration was also correlated with weaker functional connectivity between the left Angular gyrus and the (supracallosal) Anterior Cingulate cortex. These correlations strengthen the interpretation of the changes in functional

connectivity in these regions found in patients with depression, in that these functional connectivities were related to the depression that was measured in these patients.

Comparison of functional connectivity in medicated and unmedicated patients with depression. Within the depressed group, 185 were not receiving medication, and 236 patients were receiving medication. Although it was not a primary aim of this investigation, and following a suggestion, the effects of medication were assessed by comparing the functional connectivity in 185 patients not receiving medication, and 182 patients receiving medication, from the Dongbei and Xinan datasets (see Supplementary Material for details including demographic and clinical details, and limitations). The medication consisted in most cases of selective serotonin reuptake inhibitors (SSRIs) including fluoxetine, paroxetine, sertraline, citalopram and escitalopram; or serotonin-norepinephrine reuptake inhibitors (SNRIs) such as venflaxine, or a tetracyclic antidepressant such as mirtazepine. The overall pattern of functional connectivity differences between patients and controls is similar for the unmedicated (Fig. S1A) and the medicated (Fig. S1B) subgroups of patients, providing evidence that the main differences between patients and controls shown in Figs. 1-3 were found in depressed patients whether or not they were receiving medication.

Although the overall pattern of functional connectivity is similar in the subgroup without medication (Fig. S1A) and with medication (Fig. S1B), to test for significant differences, a t-test was performed between these two functional connectivity matrices, with the results shown in Fig. S1C. As one of the main findings for differences of functional connectivity between patients and controls was increased functional connectivity of the lateral orbitofrontal cortex BA 47/12 with the pre-cuneus, angular gyrus and Mid-temporal gyrus (Figs. 2 and 3), and to relieve the burden of multiple comparisons, we tested in a pre-planned comparison whether these three functional connectivity links were weaker in medicated than in un-medicated patients. For the Frontal_Inf_Orb_2R with precuneus link there is a significantly smaller functional connectivity in the medicated than the unmedicated group ($t=2.17$, $p=0.015$, one-tailed test of the specific prediction in all 3 cases). For the Frontal_Inf_Orb_2R with angular gyrus link there is a significantly smaller functional connectivity in the medicated than the unmedicated group ($t=2.55$, $p=0.005$). For the Frontal_Inf_Orb_2R with temporal_Mid_L link there is a significantly smaller functional connectivity in the medicated than the unmedicated group ($t=1.76$, $p=0.039$). The results overall are thus consistent with the hypothesis that the increased functional connectivity of the lateral orbitofrontal cortex BA 47/12 with the precuneus, angular gyrus, and mid-temporal gyrus is related to depression, and that treatment with medication reduces the functional connectivity of these three links.

Discussion

The main findings are first that one major circuit with altered functional connectivity in depression involves the medial orbitofrontal cortex BA 13, which is implicated in reward, and which had reduced functional connectivity in depression with memory systems in the parahippocampal gyrus and medial temporal lobe, involving especially the perirhinal cortex BA 36 and entorhinal cortex BA 28. The lateral orbitofrontal cortex BA 47/12, involved in non-reward and punishing events, did not have this reduced functional connectivity with memory systems, so that there is an imbalance in depression towards decreased reward-related memory system functionality. Second, BA 47/12 had increased functional

connectivity with the precuneus, the angular gyrus, and the temporal visual cortex BA 21. This enhanced functional connectivity of the non-reward/punishment system (BA 47/12) with the precuneus (involved in the sense of self and agency), and the angular gyrus (involved in language) is thus related to the explicit affectively negative sense of the self, and of self-esteem, in depression.

In a further analysis, it was shown that the overall pattern of functional connectivity differences between patients and controls is similar for the unmedicated (Fig. S1A) and the medicated (Fig. S1B) subgroups of patients, providing evidence that the main differences between patients and controls shown in Figs. 1-3 were found in depressed patients whether or not they were receiving medication. In preplanned comparisons, it was further shown that the functional connectivities of the right lateral orbitofrontal cortex BA 47/12 (Frontal_Inf_Orb_2R) with the precuneus, angular gyrus, and mid-temporal gyrus, the links highlighted in Fig. 3B, were reduced in the medicated patients compared to the unmedicated patients (Fig. S1C, with the statistics in the Supplementary Material). The results overall are thus consistent with the hypothesis that the increased functional connectivity of the lateral orbitofrontal cortex BA 47/12 with the precuneus, angular gyrus, and mid-temporal gyrus is related to depression, and that treatment with medication reduces the functional connectivity of these three links. In addition to these preplanned tests, it is notable that the medicated patients had a lower functional connectivity between the lateral orbitofrontal cortex and the medial orbitofrontal cortex (see Supplementary Material). These parts of the orbitofrontal cortex have a reciprocal relation with respect to their activations by rewards (medially) and by non-reward or loss laterally, and the smaller functional connectivity in medicated patients than unmedicated patients may be related to a change in this reciprocal relation. It is noted that limitations of this analysis of the effects of medication are that this was not a main aim of this investigation, that this is a cross-sectional not longitudinal comparison, and that the mean illness duration was 28.5 months in the unmedicated group and 58.6 months in the medicated group.

We now place these findings on functional connectivity differences in depression in the context of the known functions of the brain regions implicated in this investigation which include emotion-related, non-reward and punishment-related regions of the orbitofrontal cortex (Rolls, 2014, Rolls, 2016b), and of previous investigations into depression. The theory that depression is associated with the maladaptive responses to non-reward and punishment and hyposensitivity to reward has been extensively investigated (Eshel and Roiser, 2010, Rolls, 2016b, Russo and Nestler, 2013, Whitton *et al.* , 2015). At the psychological level, Beck's psychological theory of depression (Beck, 1979) and Seligman's learned helplessness model (Seligman, 1972), both focused around punishment and reward, and brain areas related to punishment and reward have become primary targets in psychotherapy (Beck, 2008). At the neural level, networks related to punishment and reward have been related to depression, and in some cases to monoamines (Felger *et al.* , 2015, Harmer and Cowen, 2013, Huys *et al.* , 2015, McCabe *et al.* , 2012).

Given this background, we first consider voxels in the medial parts of the orbitofrontal cortex shown in Fig. 1 that are within the OFC13 cluster. Fig. 2 shows that in depression the voxels with altered functional connectivity in these areas have reduced functional connectivity with the Parahippocampal/Temporal lobe/Fusiform cortical left and right clusters, especially involving the perirhinal cortex BA 36 and entorhinal cortex BA 28 as shown in Fig. 1. These parahippocampal areas are involved in memory, and inter alia provide a gateway to and from

the hippocampal memory system (Kesner and Rolls, 2015). Indeed, the medial and mid orbitofrontal cortex BA13 has direct reciprocal connections with the perirhinal cortex BA 36 (Kondo *et al.* , 2005), which in turn connects via the entorhinal cortex BA 28 to the hippocampus (Kesner and Rolls, 2015). There is extensive evidence that the human medial orbitofrontal cortex areas, including OFC13, is activated by rewarding stimuli that are subjectively pleasant (including pleasant odors, pleasant touch, pleasant flavor, and monetary reward) (Grabenhorst and Rolls, 2011, O'Doherty *et al.* , 2001, Rolls, 2014). The connections between the medial and mid orbitofrontal cortex BA 13 and the perirhinal cortex BA 36 (which in turn connects to the entorhinal cortex and thus to the hippocampus) provides a route for reward / emotion-related information to reach the hippocampus to become part of an episodic memory; and during later recall for the reward / emotion-related part of an episodic memory to be recalled to the orbitofrontal cortex (Kesner and Rolls, 2015, Rolls, 2014, Rolls, 2016a). It has therefore been suggested that there are weaker functional connectivity links in depression between brain areas involved in pleasant feelings and rewards with memory systems, and that this may be part of the mechanism of depression (Rolls, 2016b). This hypothesis is strengthened by the correlation between the symptoms of depression and the weakening of links between the medial orbitofrontal cortex OFC13 system and the parahippocampal / medial temporal lobe memory system areas as shown in Table 3. Consistent with the hypothesis, the anhedonia of depression can be related to decreased effects of pleasant rewarding stimuli in the medial orbitofrontal cortex during depression, effects that can be restored by antidepressants (Ma, 2015). Consistent with the importance of the orbitofrontal cortex in depression, gray matter volume reductions are found in this area in patients with depression (Ballmaier *et al.* , 2004).

Second, the lateral orbitofrontal cortex cluster OFC47/12 is in a region that is activated by many types of non-reward and unpleasant stimuli (Grabenhorst and Rolls, 2011, Rolls, 2014), including losing money (O'Doherty *et al.* , 2001), not receiving an expected social reward (Kringelbach and Rolls, 2003), and unpleasant odors. This region has very different changed functional connectivity in depression, with *increased* functional connectivity with the precuneus, angular gyrus, and middle temporal gyrus BA 21. The precuneus is a parietal region implicated in the sense of self and agency (Cavanna and Trimble, 2006), and the left (not right) angular gyrus / middle temporal gyrus is implicated in language processing (Cabeza and Nyberg, 2000). This has led to the hypothesis that this lateral non-reward / punishment system in OFC47/12 with its increased functional connectivity with self- and language-related systems relates to some of the symptoms of depression (Rolls, 2016b). The BA 21 region is high order visual cortex corresponding to the macaque inferior temporal visual cortex where faces and objects are represented (Rolls, 2012), and the increased functional connectivity of OFC47/12 with BA 21 may result in more affectively negative processing of visual inputs (Rolls, 2016b). Indeed, this increased functional connectivity between the inferior temporal visual cortex area 21 and the lateral orbitofrontal cortex OFC47/12 non-reward / punishment system may lead to depressed patients having difficulty in categorizing happy face stimuli as happy (Harmer and Cowen, 2013).

Consistent with the hypothesis of disturbed function of the orbitofrontal cortex in depression, there is increased regional cerebral blood flow in the ventrolateral orbitofrontal cortex area 47/12 in depression (Drevets *et al.* , 1992, Drevets *et al.* , 2004, Price and Drevets, 2010). In addition, overgeneral autobiographical memory manifests in individuals with major depressive disorder (MDD) tested during depressed (dMDD) or remitted phases (rMDD), and healthy individuals at high-risk for developing MDD. During specific autobiographical

memory recall, high risk individuals have increased activity relative to rMDDs and healthy controls in the ventrolateral prefrontal cortex (VLPFC) and lateral orbitofrontal cortex (Young *et al.*, 2015). The increased functional connectivity of the lateral orbitofrontal cortex (involved in non-reward and aversive processing), the precuneus (involved in the sense of self), and the angular gyrus (involved in language) in depression is of interest, for a sign of the start of a depressive episode may be negative thoughts about the self and low self-esteem, all expressed explicitly in language (Wegener *et al.*, 2015). It is notable that OFC47/12, the non-reward punishment area, has increased functional connectivity with each of these areas, but that they do not have increased connectivity with each other. The common hub to this system is the lateral orbitofrontal cortex OFC47/12.

In comparing the medial OFC13 (reward) and lateral OFC47/12 (non-reward) systems, there is evidence that they are very different systems, for the correlation between the functional connectivities within the different AAL2 regions of OFC13 was typically high ($r=0.6-0.9$), and the functional connectivities of each of these areas with OFC47/12 were typically low ($r=0.23-0.37$) ($p<10^{-14}$). Both systems though may contribute to the lack of motivation that is frequent in depression. The medial orbitofrontal cortex / medial temporal lobe memory system reduced functional connectivity may contribute by making remembered rewards, the goals for action, less rewarding, and therefore less motivating (Rolls, 2014, Rolls, 2016b). The lateral orbitofrontal non-reward system with its increased functional connectivity may make non-reward more potent, and this facilitation would also be expected to check motivation by enhancing the inhibiting effects on behavior of non-reward and expected non-reward (Rolls, 2014, Rolls, 2016b).

The anterior cingulate cortex (in which the voxel clusters are just supracallosal) had reduced functional connectivity in depressed patients with some temporal cortex areas including the fusiform gyrus, and with the Angular cortex (Figs. 2a and 3a), and no difference in functional connectivity with most medial orbitofrontal areas (OFC13) (apart from a small increase with OFC_Med_R). This supracallosal part of the far anterior cingulate cortex is at the anterior end of a supracallosal cingulate region in which many unpleasant stimuli are represented (Grabenhorst and Rolls, 2011, Rolls, 2014), and is therefore implicated in mood (Rolls, 2014). This region is just above and behind the pregenual cingulate cortex area in which a few additional voxels with significantly different functional connectivity were found in the depressed group, and this pregenual cingulate region has representations of pleasant and rewarding stimuli (Grabenhorst and Rolls, 2011, Rolls, 2014), and is thereby also implicated in mood (Rolls, 2014). Interestingly, functional connectivity changes were not found in the subcallosal cingulate cortex including the subgenual cingulate cortex (with the area found here more ventral, in OFC13), though a few voxels with altered functional connectivity were found in the amygdala, with both these regions showing increased cerebral blood flow in depression (Drevets *et al.*, 1997, Price and Drevets, 2010).

We now consider the changes in functional connectivity in depression in other brain areas not typically associated with mood and emotion. The thalamus had increased functional connectivity with a number of cortical areas, as shown in Fig. 2, with the coordinate ([9 -27 9]) indicating that this is part of the medial pulvinar, which has temporal lobe connections including to visual temporal cortex areas (Johansen-Berg *et al.*, 2005). The medial thalamus has increased cerebral blood flow in depression (Price and Drevets, 2010).

It is interesting to relate these changes in functional connectivity to the level of activity in these different brain areas in patients with depression. Hyperactivation during affective

processing tasks has been described in the thalamus and parahippocampal gyrus (Miller *et al.* , 2015), and increased cerebral blood flow in patients with major depressive disorder has been found in the medial as well as the lateral orbitofrontal cortex (Drevets *et al.* , 1992, Price and Drevets, 2010).

Although changes have been found in some of these regions in previous studies in depression including the precuneus, angular gyrus, and hippocampal system (Sundermann *et al.* , 2014), the present study is statistically more powerful because of the large number of participants involved (421 patients with a diagnosis of major depression, and 488 controls), and therefore allows analysis at the voxel level, which as we have seen greatly facilitates the interpretation of the findings by enabling the functional connectivity to be related to the different functions of even nearby brain regions such as the medial and lateral orbitofrontal cortex.

Contributors

Jianfeng Feng, Edmund T. Rolls, Peng Xie, Fei Wang and Ching-Po Lin contributed to the design of the study. Wei Liu, Yanqing Tang, Chu-Chung Huang, XinFa Wang, JunCai Pu, Shih-Jen Tsai and Albert C Yang contributed to the collection of the data. Wei Cheng, Edmund T. Rolls, Jiang Qiu, Yanqing Tang, Wei Liu, Jie Zhang, Lirong Zheng and Wei Lin contributed to the analysis of the data and the preparation of the manuscript. Edmund T. Rolls, Wei Cheng, and Jianfeng Feng participated in writing the paper. All collaborators had an opportunity to contribute to the interpretation of the results and to the drafting of the manuscript.

Declaration of interests.

All authors declare no competing interests.

Acknowledgements

J.Feng is a Royal Society Wolfson Research Merit Award holder. J.Feng is also partially supported by the National High Technology Research and Development Program of China (No. 2015AA020507) and the key project of Shanghai Science & Technology Innovation Plan (No. 15JC1400101). The research was partially supported by the National Centre for Mathematics and Interdisciplinary Sciences (NCMIS) of the Chinese Academy of Sciences, Key Program of National Natural Science Foundation of China (No. 91230201), and the Shanghai Soft Science Research Program (No. 15692106604). Wei Cheng is partly supported by grants from the National Natural Sciences Foundation of China (No.11471081, 11101429 and 71661167002). J.Zhang is supported by National Science Foundation of China (NSFC 61104143 and 61573107), and special Funds for Major State Basic Research Projects of China (2015CB856003). CP.Lin was supported in part by funding from Ministry of Science and Technology, Taiwan (NSC100-2911-I-010-010, NSC101-2911-I-010-009, NSC100-2628-E-010-002-MY3, NSC102-2321-B-010-023, and NSC103-2911-I-010-501), National Health Research Institutes (NHRI-EX103-10310EI), Ministry of Health and Welfare of Taiwan (DOH102-TD-PB-111-NSC006), and Academia Sinica, Taipei, Taiwan. J.Qiu was supported by the National Natural Science Foundation of China (31271087; 31470981; 31571137; 31500885), National Outstanding young people plan, the Program for the Top Young Talents by Chongqing, the Fundamental Research Funds for the Central Universities (SWU1509383), Natural Science Foundation of Chongqing (cstc2015jcyjA10106), General Financial Grant from the China Postdoctoral Science Foundation (2015M572423). P.Xie is supported by

National Science Foundation of China (NSFC 31271189). W. Lin was supported by the NSFC under Grant Nos. 61273014 and 11322111.

Table 1. A summary of the demographic information and the psychiatric diagnosis in the present study.

Sites	Group	Age (years)	Sex (male/female)	Education (years)	Medication (yes / no)	HAMD	Duration of illness	Mean FD
Taiwan	Healthy	49.18±8.58	60 / 36	15.04 ± 3.83	/	/	/	0.133 ± 0.054
	Patient	52.64±14.86	33 / 21	12.66 ± 3.95	54 / 0	9.34 ± 6.99	8.63 ± 6.92	0.116 ± 0.056
	Statistic (t / p)	-1.810 / 0.072	0.028 / 0.866	3.60 / 4.3e-4		/	/	1.833 / 0.0687
Dongbei	Healthy	29.90 ± 11.89	87 / 51	14.22 ± 3.40				0.101 ± 0.040
	Patient	29.02 ± 10.49	61 / 24	11.80 ± 3.18	25 / 60	20.9 ± 8.79	1.04 ± 1.67	0.098 ± 0.039
	Statistic (t / p)	0.554 / 0.580	1.611 / 0.204	5.25 / 3.7e-7		/		0.591 / 0.555
Xinan	Healthy	39.65 ± 15.80	166 / 88	13.01 ± 3.89	/	/	/	0.133 ± 0.063
	Patient	38.74 ± 13.65	183 / 99	11.91 ± 3.58	157 / 125	20.8 ± 5.87	4.16 ± 5.51	0.125 ± 0.054
	Statistic (t / p)	0.719 / 0.472	0.013 / 0.911	3.41 / 6.9e-4	/	/	/	1.729 / 0.084

Table 2. Coordinates of the peaks of the voxel clusters with different functional connectivity in patients with depression. Brodmann areas are provided where useful.

Areas	Abbreviation	# Voxels	Peak MA value	MNI coordinates (Peak)		
Hippocampus_L, ParaHippocampal_L, Fusiform_L, Temporal_Mid_L, Temporal_Pole_Mid_L, Temporal_Inf_L	MedTL_L	274	284	-27	-9	-36
ParaHippocampal_R, Fusiform_R, Temporal_Pole_Mid_R	MedTL_R	115	141	24	12	-39
Olfactory_R, Rectus_R, OFCmed_R, OFCpost_R (BA 13)	OFC13_R	140	209	15	24	-18
Olfactory_L, Rectus_L, OFCmed_L, OFCant_L, OFCpost_L, Insula_L (BA 13)	OFC13_L	206	296	-15	12	-15
OFCmed_R (BA 13)	OFC13_R_2	12	50	12	60	-21
Temporal_Mid_R (BA 21)	MidTG21_R	14	43	66	-15	-18
Frontal_Inf_Orb_2_R (BA 47/12)	OFC47/12_R	11	136	36	36	-12
Thalamus_R	Thal_R	17	82	9	-27	9
Cingulate_Ant_L (BA 24)	ACC_L	43	86	-6	36	21
Angular_L, Temporal_Mid_L (BA 39)	Angular_L	15	18	-48	-60	21
Postcentral_L	Postcentral_L	20	21	-60	-12	18
Cingulate_Post_L, Precuneus_L	Precuneus_L	24	15	-6	-54	30
Angular_L (BA 39)	Angular_L_2	22	36	-48	-69	42

Table 3. Correlations between the functional connectivity links and the depression symptom severity scores.

Functional connectivity		Clinical variable	P value	Correlation value
Olfactory_R	OFCpost_R	BDI	0.022249	-0.14572
Olfactory_R	Temporal_Pole_Mid_R	BDI	0.041603	-0.13001
OFCmed_R	ParaHippocampal_R	BDI	0.044183	-0.12843
OFCpost_R	ParaHippocampal_R	BDI	0.044118	-0.12846
OFCpost_R	Temporal_Pole_Mid_R	BDI	0.048206	-0.12610
Fusiform_L	Temporal_Mid_R	BDI	0.030704	-0.13782
Postcentral_L	Temporal_Mid_R	BDI	0.0027053	-0.14063
Temporal_Pole_Mid_L	Temporal_Inf_L	BDI	0.027426	-0.11088
Olfactory_L	Rectus_L	HAMD	0.0094015	-0.13148
Olfactory_L	Rectus_R	HAMD	0.026386	-0.11273
Olfactory_L	OFCmed_L	HAMD	0.0046205	-0.14343
Rectus_R	OFCmed_L	HAMD	0.038747	-0.10504
Frontal_Inf_Orb_2_R	Angular_L	Illness duration	0.01825	-0.12418
Frontal_Inf_Orb_2_R	Temporal_Mid_L	Illness duration	0.00077913	-0.17688
Insula_L	ParaHippocampal_R	Illness duration	0.016651	-0.12658
Insula_L	Fusiform_L	Illness duration	0.021031	-0.12198
Insula_L	Temporal_Inf_L	Illness duration	0.046867	-0.10538
Cingulate_Ant_L	Angular_L	Illness duration	0.046772	-0.10438
Postcentral_L	Temporal_Pole_Mid_L	Illness duration	0.036223	-0.11027

BDI: Beck Depression Inventory

HAMD: Hamilton Depression Rating Scale

Figure 1. Anatomical location of consistently different functional connectivity in depression obtained from voxel-based BWAS. A) Voxels showing the largest number of whole brain voxel-level functional connectivity differences in patients with depression. Clusters of voxels containing more than 10 significant voxels are shown. The color bar represents the measure of association (*MA*) given by the number of significantly different functional connectivity links relating to each voxel. The right of the brain is on the right of each slice. Some of the different clusters are in the following range of Y values for the slices shown: MedOFC13 +37 to +8; LatOFC47/12_R +37 to +32; medial temporal lobe MTL_L +17 to -16; thalamus -25 to -31; precuneus -57 to -59; angular gyrus -57 to -59. B) Manhattan plot of voxel-based BWAS results with voxels grouped in accordance with the AAL2 atlas labels (Rolls *et al.* , 2015). The order of the bars is as shown in Table S1, and the width of each bar reflects the number of voxels in each AAL2 region.




Figure 2. The functional connectivity differences between the voxels that are significantly different in the depressed and the control group, separated by AAL2 region in which the significant voxels were located. A) Links with increased functional connectivities in patients are shown in yellow-red, and decreased connectivities are in blue. The color bar shows the $-\log_{10}$ of the p value for the difference of the functional connectivity. Entries in the matrix are provided where $p < 0.05$ (FDR correction). The matrix itself contains rows and columns for all cases in which there were 10 or more significant voxels within an AAL2 region. B) The links are shown in red if they are significantly stronger in the patient group, and in blue if they are significantly weaker in the patient group. The thickness of the lines indicates the degree of alteration of the functional connectivity. The anatomical abbreviations are for the areas in the automated anatomical atlas, with abbreviations shown in Table S1. The brain regions in the left hemisphere are in the left semicircle of the diagram.





Figure 3. The medial and lateral orbitofrontal cortex networks that show different functional connectivity in patients with depression. A decrease in functional connectivity is shown in blue, and an increase in red. MedTL –medial temporal lobe from the parahippocampal gyrus to the temporal pole; MidTG21_R – middle temporal gyrus area 21 right; OFC13 – medial orbitofrontal cortex area 13; OFC47/12_R – lateral orbitofrontal cortex area 47/12 right. The lateral orbitofrontal cortex cluster in OFC47/12 is visible on the ventral view of the brain anterior and lateral to the OFC13 clusters.



References

- Ballmaier M, Toga AW, Blanton RE, Sowell ER, Lavretsky H, Peterson J, et al. Anterior cingulate, gyrus rectus, and orbitofrontal abnormalities in elderly depressed patients: an MRI-based parcellation of the prefrontal cortex. *Am J Psychiatry*. 2004 Jan;161(1):99-108.
- Beck AT, Beamesderfer A. Assessment of depression: the depression inventory. *Mod Probl Pharmacopsychiatry*. 1974;7(0):151-69.
- Beck AT. *Cognitive Therapy of Depression*: Guilford Press; 1979.
- Beck AT. The evolution of the cognitive model of depression and its neurobiological correlates. *Am J Psychiatry*. 2008 Aug;165(8):969-77.
- Bell-McGinty S, Butters MA, Meltzer CC, Greer PJ, Reynolds CF, 3rd, Becker JT. Brain morphometric abnormalities in geriatric depression: long-term neurobiological effects of illness duration. *Am J Psychiatry*. 2002 Aug;159(8):1424-7.
- Button KS, Ioannidis JP, Mokrysz C, Nosek BA, Flint J, Robinson ES, et al. Power failure: why small sample size undermines the reliability of neuroscience. *Nat Rev Neurosci*. 2013 May;14(5):365-76.
- Cabeza R, Nyberg L. Imaging cognition II: An empirical review of 275 PET and fMRI studies. *J Cogn Neurosci*. 2000 Jan;12(1):1-47.
- Cavanna AE, Trimble MR. The precuneus: a review of its functional anatomy and behavioural correlates. *Brain*. 2006 Mar;129(Pt 3):564-83.
- Chao-Gan Y, Yu-Feng Z. DPARSF: a MATLAB toolbox for “pipeline” data analysis of resting-state fMRI. *Front Syst Neurosci*. 2010;4:13.
- Cheng W, Rolls ET, Gu H, Zhang J, Feng J. Autism: reduced functional connectivity between cortical areas involved in face expression, theory of mind, and the sense of self. *Brain*. 2015;138:1382-93.
- de Diego-Adelino J, Pires P, Gomez-Anson B, Serra-Blasco M, Vives-Gilabert Y, Puigdemont D, et al. Microstructural white-matter abnormalities associated with treatment resistance, severity and duration of illness in major depression. *Psychol Med*. 2014 Apr;44(6):1171-82.
- Deco G, Kringelbach ML. Great expectations: using whole-brain computational connectomics for understanding neuropsychiatric disorders. *Neuron*. 2014 Dec 3;84(5):892-905.
- Disner SG, Beevers CG, Haigh EA, Beck AT. Neural mechanisms of the cognitive model of depression. *Nat Rev Neurosci*. 2011 Aug;12(8):467-77.
- Drevets WC, Videen TO, Price JL, Preskorn SH, Carmichael ST, Raichle ME. A functional anatomical study of unipolar depression. *J Neurosci*. 1992 Sep;12(9):3628-41.
- Drevets WC, Price JL, Simpson JRJ, Todd RD, Reich T, Vannier M, et al. Subgenual prefrontal cortex abnormalities in mood disorders. *Nature*. 1997;386:824-7.
- Drevets WC, Gadde K, Krishnan KRR. Neuroimaging studies of mood disorder. In: Charney DS, Nestler EJ, editors. *Neurobiology of Mental Illness*. New York: Oxford

University Press; 2004. p. 461-80.

Drevets WC. Orbitofrontal cortex function and structure in depression. *Ann N Y Acad Sci.* 2007 Dec;1121:499-527.

Eshel N, Roiser JP. Reward and punishment processing in depression. *Biol Psychiatry.* 2010 Jul 15;68(2):118-24.

Felger JC, Li Z, Haroon E, Woolwine BJ, Jung MY, Hu X, et al. Inflammation is associated with decreased functional connectivity within corticostriatal reward circuitry in depression. *Mol Psychiatry.* 2015 Nov 10.

Friston KJ, Williams S, Howard R, Frackowiak RS, Turner R. Movement-related effects in fMRI time-series. *Magn Reson Med.* 1996 Mar;35(3):346-55.

Glahn DC, Laird AR, Ellison-Wright I, Thelen SM, Robinson JL, Lancaster JL, et al. Meta-analysis of gray matter anomalies in schizophrenia: application of anatomic likelihood estimation and network analysis. *Biol Psychiatry.* 2008 Nov 1;64(9):774-81.

Gotlib IH, Hammen CL, editors. *Handbook of Depression.* New York: Guilford Press; 2009.

Grabenhorst F, Rolls ET. Value, pleasure, and choice in the ventral prefrontal cortex. *Trends Cogn Sci.* 2011;15:56-67.

Gusnard DA, Raichle ME, Raichle ME. Searching for a baseline: functional imaging and the resting human brain. *Nat Rev Neurosci.* 2001 Oct;2(10):685-94.

Hamilton JP, Etkin A, Furman DJ, Lemus MG, Johnson RF, Gotlib IH. Functional neuroimaging of major depressive disorder: a meta-analysis and new integration of base line activation and neural response data. *Am J Psychiatry.* 2012 Jul;169(7):693-703.

Hamilton JP, Chen MC, Gotlib IH. Neural systems approaches to understanding major depressive disorder: an intrinsic functional organization perspective. *Neurobiol Dis.* 2013 Apr;52:4-11.

Hamilton M. A rating scale for depression. *J Neurol Neurosurg Psychiatry.* 1960 Feb;23:56-62.

Harmer CJ, Cowen PJ. 'It's the way that you look at it'--a cognitive neuropsychological account of SSRI action in depression. *Philos Trans R Soc Lond B Biol Sci.* 2013;368(1615):20120407.

Hart H, Radua J, Mataix-Cols D, Rubia K. Meta-analysis of fMRI studies of timing in attention-deficit hyperactivity disorder (ADHD). *Neurosci Biobehav Rev.* 2012 Nov;36(10):2248-56.

Huys QJ, Daw ND, Dayan P. Depression: a decision-theoretic analysis. *Annu Rev Neurosci.* 2015 Jul 8;38:1-23.

Iwabuchi SJ, Krishnadas R, Li C, Auer DP, Radua J, Palaniyappan L. Localized connectivity in depression: a meta-analysis of resting state functional imaging studies. *Neurosci Biobehav Rev.* 2015 Apr;51:77-86.

Johansen-Berg H, Behrens TE, Sillery E, Ciccarelli O, Thompson AJ, Smith SM, et al. Functional-anatomical validation and individual variation of diffusion tractography-based segmentation of the human thalamus. *Cereb Cortex.* 2005 Jan;15(1):31-9.

Kaiser RH, Andrews-Hanna JR, Wager TD, Pizzagalli DA. Large-Scale Network Dysfunction in Major Depressive Disorder: A Meta-analysis of Resting-State

Functional Connectivity. *JAMA Psychiatry*. 2015 Jun;72(6):603-11.

Kesner RP, Rolls ET. A computational theory of hippocampal function, and tests of the theory: new developments. *Neurosci Biobehav Rev*. 2015 Jan;48:92-147.

Kondo H, Saleem KS, Price JL. Differential connections of the perirhinal and parahippocampal cortex with the orbital and medial prefrontal networks in macaque monkeys. *J Comp Neurol*. 2005 Dec 26;493(4):479-509.

Konrad K, Neufang S, Hanisch C, Fink GR, Herpertz-Dahlmann B. Dysfunctional attentional networks in children with attention deficit/hyperactivity disorder: evidence from an event-related functional magnetic resonance imaging study. *Biol Psychiatry*. 2006 Apr 1;59(7):643-51.

Kringelbach ML, Rolls ET. Neural correlates of rapid reversal learning in a simple model of human social interaction. *Neuroimage*. 2003;20:1371-83.

Kuhn S, Gallinat J. Resting-state brain activity in schizophrenia and major depression: a quantitative meta-analysis. *Schizophr Bull*. 2013 Mar;39(2):358-65.

Liptak T. On the combination of independent tests. *Magyar Tud Akad Mat Kutato Int Kozl*. 1958;3:171-97.

Lorenzetti V, Allen NB, Fornito A, Yucel M. Structural brain abnormalities in major depressive disorder: a selective review of recent MRI studies. *J Affect Disord*. 2009 Sep;117(1-2):1-17.

Ma Y. Neuropsychological mechanism underlying antidepressant effect: a systematic meta-analysis. *Mol Psychiatry*. 2015 Mar;20(3):311-9.

McCabe C, Woffindale C, Harmer CJ, Cowen PJ. Neural processing of reward and punishment in young people at increased familial risk of depression. *Biol Psychiatry*. 2012 Oct 1;72(7):588-94.

Miller CH, Hamilton JP, Sacchet MD, Gotlib IH. Meta-analysis of Functional Neuroimaging of Major Depressive Disorder in Youth. *JAMA Psychiatry*. 2015 Oct 1;72(10):1045-53.

O'Doherty J, Kringelbach ML, Rolls ET, Hornak J, Andrews C. Abstract reward and punishment representations in the human orbitofrontal cortex. *Nat Neurosci*. 2001;4:95-102.

Öngür D, Ferry AT, Price JL. Architectonic division of the human orbital and medial prefrontal cortex. *J Comp Neurol*. 2003;460:425-49.

Power JD, Mitra A, Laumann TO, Snyder AZ, Schlaggar BL, Petersen SE. Methods to detect, characterize, and remove motion artifact in resting state fMRI. *Neuroimage*. 2014 Jan 1;84:320-41.

Price JL, Drevets WC. Neurocircuitry of mood disorders. *Neuropsychopharmacology*. 2010 Jan;35(1):192-216.

Rigucci S, Serafini G, Pompili M, Kotzalidis GD, Tatarelli R. Anatomical and functional correlates in major depressive disorder: the contribution of neuroimaging studies. *World J Biol Psychiatry*. 2010 Mar;11(2 Pt 2):165-80.

Rolls ET. Invariant visual object and face recognition: neural and computational bases, and a model, VisNet. *Front Comput Neurosci*. 2012;6, 35:1-70.

Rolls ET. *Emotion and Decision-Making Explained*. Oxford: Oxford University Press; 2014.

Rolls ET, Joliot M, Tzourio-Mazoyer N. Implementation of a new parcellation of the orbitofrontal cortex in the automated anatomical labeling atlas. *Neuroimage*. 2015 Aug 1;122:1-5.

Rolls ET. *Cerebral Cortex: Principles of Operation*. Oxford: Oxford University Press; 2016a.

Rolls ET. A non-reward attractor theory of depression. *Neurosci Biobehav Rev*. 2016b;68:47-58.

Russo SJ, Nestler EJ. The brain reward circuitry in mood disorders. *Nat Rev Neurosci*. 2013 Sep;14(9):609-25.

Seligman ME. Learned helplessness. *Annu Rev Med*. 1972;23:407-12.

Sheline YI, Price JL, Yan Z, Mintun MA. Resting-state functional MRI in depression unmasks increased connectivity between networks via the dorsal nexus. *Proc Natl Acad Sci U S A*. 2010 Jun 15;107(24):11020-5.

Sundermann B, Olde Lutke Beverborg M, Pfliegerer B. Toward literature-based feature selection for diagnostic classification: a meta-analysis of resting-state fMRI in depression. *Front Hum Neurosci*. 2014;8:692.

Veer IM, Beckmann CF, van Tol MJ, Ferrarini L, Milles J, Veltman DJ, et al. Whole brain resting-state analysis reveals decreased functional connectivity in major depression. *Front Syst Neurosci*. 2010;4.

Wang L, Hermens DF, Hickie IB, Lagopoulos J. A systematic review of resting-state functional-MRI studies in major depression. *J Affect Disord*. 2012 Dec 15;142(1-3):6-12.

Wegener I, Geiser F, Alfter S, Mierke J, Imbierowicz K, Kleiman A, et al. Changes of explicitly and implicitly measured self-esteem in the treatment of major depression: evidence for implicit self-esteem compensation. *Compr Psychiatry*. 2015 Apr;58:57-67.

Whitton AE, Treadway MT, Pizzagalli DA. Reward processing dysfunction in major depression, bipolar disorder and schizophrenia. *Curr Opin Psychiatry*. 2015 Jan;28(1):7-12.

Wittmann BC, Schott BH, Guderian S, Frey JU, Heinze HJ, Duzel E. Reward-related FMRI activation of dopaminergic midbrain is associated with enhanced hippocampus-dependent long-term memory formation. *Neuron*. 2005 Feb 3;45(3):459-67.

Young KD, Bellgowan PS, Bodurka J, Drevets WC. Functional neuroimaging correlates of autobiographical memory deficits in subjects at risk for depression. *Brain Sci*. 2015;5(2):144-64.

Yu KK, Cheung C, Chua SE, McAlonan GM. Can Asperger syndrome be distinguished from autism? An anatomic likelihood meta-analysis of MRI studies. *J Psychiatry Neurosci*. 2011 Nov;36(6):412-21.