

<http://mc.manuscriptcentral.com/jph>

**Associations between neighbourhood environmental factors
and the uptake and effectiveness of a brief intervention to
increase physical activity**

Journal:	<i>Journal of Public Health</i>
Manuscript ID	JPH-15-0357.R2
Manuscript Type:	Original Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Goyder, Elizabeth; The University of Sheffield, Public Health, SCHARR; Maheswaran, Ravi; University of Sheffield, SCHARR Read, Simon; University of Sheffield, SCHARR
Keywords:	Public health, Health promotion, Physical activity

SCHOLARONE™
Manuscripts

1
2
3 **Associations between neighbourhood environmental factors and the uptake and**
4 **effectiveness of a brief intervention to increase physical activity: findings from**
5 **deprived urban communities in an English city.**
6
7

8
9
10
11 Prof Goyder, E.C., Professor of Public Health, School of Health and Related
12 Research, University of Sheffield¹
13

14
15
16 Prof Maheswaran, R., Professor of Epidemiology & Public Health, Public Health GIS
17 Unit, School of Health and Related Research, University of Sheffield¹
18
19

20
21 Dr Read, S., Research Associate, Public Health GIS Unit, School of Health and
22 Related Research, University of Sheffield¹
23
24

25
26 ¹School of Health and Related Research, University of Sheffield, Regent Court, 30
27 Regent Street, Sheffield, S1 4DA, UK
28
29

30
31 On behalf of the BOOSTER Study team
32
33

34
35
36
37 *Corresponding author. E-mail: e.goyder@sheffield.ac.uk
38
39

40
41
42 Running head: Uptake and effectiveness of a brief intervention
43
44

45
46
47 Keywords: physical activity; brief intervention; inequalities; deprivation
48
49

50
51 Word count (excluding references): 2872
52
53

54
55
56 Date of submission: 28 November 2015
57
58
59
60

Abstract

Background: Evidence suggests behavioural interventions may exacerbate health inequalities, potentially due to differences in uptake or effectiveness. We used a physical activity intervention targeting deprived communities to identify neighbourhood-level factors that might explain differences in programme impact.

Methods: Individuals aged 40 to 65 were sent a postal invitation offering a brief intervention to increase physical activity. We used postcodes linkage to determine whether neighbourhood indicators of deprivation, housing, crime and proximity to green spaces and leisure facilities predicted uptake of the initial invitation or an increase in physical activity level in those receiving the brief intervention.

Results: 4134 (6.8%) individuals responded to the initial invitation and of those receiving the intervention and contactable after three months, 486 (51.6%) reported an increase in physical activity. Area deprivation scores linked to postcodes predicted intervention uptake, but not intervention effectiveness. Neighbourhood indicators did not predict either uptake or intervention effectiveness.

Conclusions: The main barrier to using brief intervention invitations to increase physical activity in deprived, middle aged populations was the low uptake of an intervention requiring significant time and motivation from participants. Once individuals have taken up the intervention offer, neighbourhood characteristics did not appear to be significant barriers to successful lifestyle change.

(200 words)

Introduction

There are clear socioeconomic gradients in physical activity levels and cardiovascular mortality, with lower physical activity levels and higher cardiovascular and all-cause mortality rates in more deprived areas.^{1,2} There is also a growing body of evidence suggesting environmental and neighbourhood characteristics may explain some of the variations in physical activity related to socioeconomic circumstances.³

Whilst there is evidence that brief interventions increase physical activity levels in the short term⁴, it is less clear to what extent these interventions might reduce or exacerbate individual or neighbourhood inequalities in physical activity levels.

Modelling suggests that this may be a crucial issue in determining cost effectiveness but, as Gulliford et al acknowledge, there is a lack of empirical data about exactly how and why specific interventions might reduce or exacerbate inequalities due to differences in participation or to differences in the efficacy of the intervention for different population groups or in different settings or neighbourhoods.⁵

We therefore used the evaluation of a brief intervention programme delivered in deprived communities in the English city of Sheffield (undertaken as part of a larger project which included a subsequent randomised trial⁶) to address this question. The intervention included a motivational DVD sent by post, information on local leisure facilities and activity programmes, such as Health Walks, and two follow up phone calls at monthly intervals, to check they had received the DVD, to answer and questions and to encourage them to use the DVD and local information.⁶

The aim was to investigate if there were neighbourhood-level factors for which local data were available and which could potentially influence uptake and effectiveness of a brief intervention to increase physical activity.^{7,8} We selected two types of factors for which there was a plausible potential causal relationship with increases in physical activity levels: firstly the proximity of relevant leisure facilities or green spaces which might facilitate active pursuits⁹, and secondly the general quality of the local environment (as reflected by crime and housing indicators) which might be a barrier to individuals being more active in their neighbourhood¹⁰.

1
2
3 The overall aim was to examine if the overall degree of deprivation or these two
4 types of more specific neighbourhood level factors (access to green spaces, gyms and
5 swimming pools or housing and crime rates) were associated with either uptake or
6 effectiveness of a brief intervention offered by postal invitation in relatively
7 socioeconomically deprived neighbourhoods.
8
9

10 11 12 13 **Methods**

14
15
16 There is a striking West-East gradient in area level deprivation across Sheffield, with
17 deprived areas situated to the east of the city.¹¹ As mailouts, described further below,
18 were sent out to postcodes in deprived neighbourhoods, we used postcodes as the
19 geographical unit of analysis in this study.
20
21
22

23
24 The design of the study is shown in the Consort diagram (Figure 1). Initially a mailout
25 was sent to all people aged 40-64 years living in deprived neighbourhoods in
26 Sheffield. More deprived neighbourhoods were selected on the basis of having an
27 above average Index of Multiple Deprivation (IMD) score, a widely used national
28 indicator of deprivation in England.¹² This mailout was done in six rounds in 2009-
29 10, including an initial pilot round targeting only one neighbourhood in order to test
30 the feasibility of the approach and estimate the likely response rate and five rounds
31 each targeting a different group of neighbourhoods. Although the total mailout was in
32 excess of 70 000, the last round was incompletely mailed out. For this analysis, we
33 therefore limited the data used to the first five mailouts, which totalled 60 429.
34 Prepaid envelopes were enclosed with the mailouts. People who sent in the replies
35 accepting the invitation to participate were then contacted and provided with the
36 initial brief intervention, as described in detail in the full report, if they were deemed
37 eligible.⁶ The main reason for ineligibility was currently being physical active (and
38 just wanting support for activities), since the intervention was targeting currently
39 sedentary individuals.
40
41
42
43
44
45
46
47
48
49
50

51
52
53 Self-reported physical activity was collected at recruitment, and then re-assessed
54 three months after sending the DVD, using the Scottish Physical Activity
55 Questionnaire (SPAQ), to see if they had increased their physical activity. We used
56 an increase from pre-intervention activity levels of at least 30 minutes a week as an
57
58
59
60

1
2
3 indicator of effective change. This was a pragmatic choice based on what was
4 considered feasible for previously sedentary participants and chosen to reflect whether
5 it was likely the intervention had led to a clinically significant change.
6
7

8
9
10 We used the IMD income domain from 2010 as the indicator of socioeconomic
11 deprivation at the small area level.¹² The Income Domain score was available at the
12 lower super-output area (LSOA) level. We assigned the LSOA score to all postcodes
13 within each LSOA. On average, there were approximately 25 residential postcodes
14 per LSOA (representing about 1500 individuals).
15
16
17

18
19
20 With regard to access to green space for physical activity, we used municipal green
21 spaces managed by Sheffield City Council. We only included green spaces that were
22 open to the public for use for physical activity. We also considered green spaces
23 surrounding Sheffield and included a park managed by Rotherham Borough Council
24 as this was accessible to residents to the east of Sheffield.
25
26
27

28
29
30 With regard to access to gyms and swimming pools, we included all such facilities
31 that were open to the public. The detailed specification of green space, gym and
32 swimming pool facilities included, and the rationale for selection of these spaces and
33 facilities, have been provided elsewhere.⁶
34
35
36

37
38 We used two indicators of housing from 2010 that were provided by Sheffield City
39 Council at the postcode level. These were the percentage of vacant domestic
40 properties in each postcode and the percentage of households that were classed as
41 being in multiple occupancy in each postcode. Both these indicators were used locally
42 as indicators of the housing environment.
43
44
45

46
47
48 We obtained geo-referenced data on crime and anti-social behaviour from the national
49 police website (<http://police.uk>). We calculated the total number of crimes reported
50 (which exclude anti-social behaviour) and the number of anti-social behaviour
51 incidents recorded within a 1km radius of each postcode centroid in 2011.
52
53
54

55
56 Access to green space, gyms and swimming pools was calculated using pedestrian
57 network information within a GIS. The shortest distance along the network from a
58
59
60

1
2
3 postcode centroid to the edge of a green space or the geo-location of a leisure facility
4 was calculated.⁶
5
6
7

8 Uptake was assessed in relation to the response to the initial postal invitation, which is
9 one relatively common way of inviting participation in public health programmes
10 targeted at specific geographical areas. Effectiveness was assessed in relation to the
11 self-reported change in physical activity, three months after the “brief intervention”
12 which is an outcome previously used to assess effectiveness of brief interventions⁴.
13
14
15
16
17

18 Statistical analysis was carried out using logistic regression, modelling the odds of a
19 positive response at each stage of the multiphase study. IMD income, distances to the
20 nearest green space, gym and swimming pool, and crime and anti-social behaviour
21 incident rates were entered as continuous variables. Because of the very small
22 percentage of postcodes which had one or more vacant properties and households in
23 multiple occupancy, these were entered as dichotomous variables (>0 vs 0). Although
24 in some phases, a positive response was not likely to be potentially associated with all
25 the predictors, for consistency we assessed associations with all predictors in all
26 phases. As response rates varied by mailout, the five mailouts were included as
27 categorical variables in the analysis to adjust for this variation. For the continuous
28 variables, the odds ratios are expressed per inter-quartile range (IQR) increase in the
29 predictor variable.
30
31
32
33
34
35
36
37
38

39 **Results**

40
41
42
43 The positive response counts at each stage of the study are given in Figure 1. Of the
44 60 429 invitations sent, only 4164 people responded to this initial invitation by mail,
45 giving a response rate of 7%. Only 54% of these individuals were contactable by
46 phone and of these only 77% were sufficiently sedentary to be considered eligible for
47 the brief intervention. Of the 941 individuals who could be contacted three months
48 after receiving the DVD, 486 (52%) had managed to increase their level of physical
49 activity.
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Table 1 shows the distribution of the postcode level predictor variables used in the
4 analysis, while Table 2 shows the adjusted odds ratios for associations between each
5 of the predictor variables and a positive response at each stage.
6
7

8
9
10 Initial response rates varied by mailout from 6% to 9%. Statistical analyses were
11 therefore adjusted for mailout round to take into account this variation.
12

13
14 There was a significant association between the level of deprivation and response to
15 the mailout. The odds ratio for a positive response for an IQR increase in deprivation
16 was 0.92 (95% CI 0.87-0.96, $p=0.0006$). There was also a significant association
17 between the level of deprivation and the odds of managing to contact those who had
18 responded to the mailout (0.85, 95% CI 0.77-0.94, $p=0.001$), indicating that both
19 using postal invitations for initial contact and telephone for subsequent follow up may
20 exacerbate inequalities in uptake of similar interventions in relatively deprived
21 communities. In contrast, there were no significant associations between deprivation
22 and the odds of response from those successfully recruited to receive the brief
23 intervention.
24
25

26
27 With regard to the other variables, there were generally no significant associations
28 apart from two relatively weak associations between being contactable three months
29 after the intervention and distance to a gym or houses in multiple occupancy (Table
30 2). These are of questionable relevance, as these variables did not predict either
31 response to the initial invitation or intervention effectiveness.
32
33

34
35 We also examined whether there were any seasonal effects on uptake or effectiveness
36 of the intervention, as this might have influenced results but no significant
37 association was identified.
38
39

40 41 42 43 44 45 46 **Discussion**

47 48 49 *Summary of main findings*

50
51
52 The response to initial postal invitation to participate in an intervention to increase
53 physical exercise sent to people aged 40-64 years living in deprived areas was very
54 low, with only 7% responding to the invitation. The finding that more than half of
55 those contacted after receiving the intervention reported an increase in physical
56
57
58
59
60

1
2
3 activity after three months, suggests that these were a highly selected group of
4 individuals already motivated to make relevant lifestyle changes.
5
6

7
8 Only area level deprivation showed a clear association with response to the initial
9 invitation, with lower response rates in more deprived areas, with this association seen
10 even within the range of deprivation observed in this analysis. The association with
11 deprivation was largely apparent only in the initial response to the postal invitation,
12 with no significant effect of deprivation on effectiveness in those receiving the
13 intervention and contactable after three months. We also found no evidence linking
14 access to green space and leisure facilities, or to neighbourhood crime rates or
15 housing indicators with uptake or effectiveness.
16
17
18
19
20
21

22 ***What is already known***

23
24
25
26 Evidence from the NHS Health Check programme suggests both that socioeconomic
27 status may influence uptake of preventive programmes and that mode of invitation is
28 a significant predictor of uptake, with postal invitation showing lower uptake than
29 telephone or verbal invitations.^{13,14}
30
31
32

33
34 *Deprivation and uptake of physical activity interventions:* Gidlow et al examined
35 uptake at different stages of the referral process for exercise referral schemes¹³ and
36 also found that uptake was lower with increasing area level deprivation at the area
37 level in the initial stages of referral. Deprivation was a predictor of uptake but not of
38 completion rates, consistent with our findings. In contrast, Harrison et al found that
39 area level deprivation did not influence attendance following referral to exercise
40 referral schemes¹⁵ and Sowden et al found that uptake of such schemes and
41 completion rates were not associated with deprivation status.¹⁶
42
43
44
45
46
47
48

49
50 *Environmental factors and physical activity:* With regard to neighbourhood level
51 environmental factors, Bauman et al found in a review that physical activity levels
52 were generally correlated with proximity to recreation facilities and with
53 environmental aesthetics, including greenness.⁸ However, they found no consistent
54 evidence of correlation in older adults. Humpel et al, in another review⁹, found that
55
56
57
58
59
60

1
2
3 high neighbourhood crime levels and fear of crime were associated with decreased
4 physical activity in some studies but not several others.
5
6

7
8 A related issue is the equity of access to leisure facilities. Hillsdon et al¹⁰ examined
9 the geographical distribution of leisure facilities, using a database of all public access
10 indoor exercise facilities in England .They found that the availability of these
11 facilities, which included gyms and swimming pools, declined with increasing levels
12 of area-level deprivation. They used the IMD score at LSOA level as their indicator of
13 socioeconomic deprivation.
14
15
16
17

18
19 Spatial autocorrelation, that is the tendency for spatial units close to each other to
20 have similar characteristics, may complicate statistical analysis. We have previously
21 carried out methodological analysis to examine this aspect using data from the initial
22 phase of the study .¹⁷ We found that the association with deprivation held after a
23 complex Bayesian spatial model was used to taken account of spatial autocorrelation.
24
25
26
27

28 29 ***What this study adds*** 30

31
32
33 Whilst there are a number of reasons why interventions to promote physical activity
34 might have lower uptake and be less effective in more deprived communities, the
35 empirical evidence is sparse and inconsistent. The large scale mailing across a large
36 number of urban neighbourhoods provided a unique opportunity to examine whether,
37 and to what extent, neighbourhood characteristics which might be barriers or
38 facilitators to physical activity, would predict uptake, and subsequently the
39 effectiveness, of a brief intervention.
40
41
42
43
44

45
46 We found that even within the deprived areas targeted in this study, there was an
47 association between deprivation and uptake. However, we found no strong evidence
48 to indicate that proximity to leisure facilities and green space or that neighbourhood
49 crime rates and housing indicators were important determinants of uptake.
50
51
52

53
54 Of practical relevance to those delivering similar programmes, are the very low
55 response rates using postal invitations with telephone follow up in relatively deprived
56
57
58
59
60

1
2
3 urban neighbourhoods. This proved to be a very inefficient way of recruiting the
4 target group of sedentary middle-aged residents.
5
6

7
8 These findings also have important implications for ensuring interventions to promote
9 physical activity do not exacerbate health inequalities. This may be a particular
10 concern when using postal invitations, since even when relatively deprived urban
11 areas are targeted for an intervention, the more deprived the neighbourhood, the lower
12 the response to a postal invitation.
13
14
15

16
17
18 The findings also suggest that the major challenge may lie in getting sufficient uptake
19 for an individual level intervention to have an impact in deprived neighbourhoods.
20 However if individuals are sufficiently motivated to take up an intervention in
21 deprived areas, access to facilities or other characteristics of the local environment do
22 not seem to have an additional impact on success, presumably because these are
23 relatively highly motivated respondents and the individual support helps to overcome
24 the environment factors that might otherwise encourage less active lifestyles.
25
26
27
28
29
30

31 *Limitations*

32
33
34

35 A number of potential limitations to our study need to be considered. We used an
36 ecological study design and this is susceptible to ecological bias that is the situation in
37 which associations at the area level may not be representative of associations which
38 exist at the individual level. However, we used the deprivation index at a small-area
39 level and used postcode level data for all the other variables. Use of very small
40 geographical units overcomes several of the limitation of traditional ecological studies
41 as explanatory factors tend to be more homogenous in small geographical areas. We
42 did not specifically address spatial autocorrelation in our analysis. However, in our
43 previous work, residual spatial autocorrelation was not a significant problem once
44 explanatory factors were taken into account.¹⁷
45
46
47
48
49
50

51
52
53 We did not take quality of green space, which may have influenced use of green
54 space, into account. The database of facilities we used may have contained some
55 inaccuracies. We only took into account physical access in terms of network walking
56 distances, and other dimensions of access (e.g. cost, access for people with
57
58
59
60

1
2
3 disabilities, opening times) could have influenced use of these facilities. Other means
4 of transport e.g. private cars and public transport could have been used to access
5 leisure facilities and green spaces. Multiple occupancy could also be due to student
6 accommodation but the areas included in our study did not have large resident student
7 populations. Whilst vacant properties were used as a proxy for run down areas, new
8 housing estates would also have been captured by this indicator. Crimes may have
9 been under-recorded and there may have been geo-location errors regarding where
10 these crimes were recorded as having occurred. Also, fear of crime rather than actual
11 crime rates may exert a stronger influence on behaviour. Give the potential limitations
12 of our analysis the results need to be interpreted with caution.
13
14
15
16
17
18
19
20

21 ***Conclusions***

22
23
24 We found that uptake of a brief intervention was significantly lower in more deprived
25 areas but there was no strong evidence to indicate that other environmental factors
26 such as access to gyms, swimming pools and green space and neighbourhood crime
27 rates and housing deprivation influenced uptake. There was also no evidence that
28 socioeconomic and neighbourhood characteristics had a significant impact on the
29 effectiveness of the intervention. These findings suggest that the main reason that
30 individual brief interventions will have a limited impact on inequalities in the domain
31 of physical activity is the lower levels of uptake, rather than specific neighbourhood
32 characteristics and further research should consider how to more effectively target and
33 recruit sedentary individuals to programmes of proven effectiveness.
34
35
36
37
38
39
40
41
42

43 **Funding**

44
45
46 This project was funded by the NIHR Health Technology Assessment programme
47 (HTA 07/25/02) and has been published in full in the Health Technology Assessment
48 journal series: www.journalslibrary.nihr.ac.uk/hta/volume-18/issue-13. The views and
49 opinions expressed therein are those of the authors and do not necessarily reflect those
50 of the Department of Health.
51
52
53
54

55 **Acknowledgements**

1
2
3 We gratefully acknowledge the following: Sarah Nickson at Activity Sheffield;
4 Jeremy Wight, Dawn Lockley, Shazia Idris, Paula Mackintosh, Fatima Musa, Stella
5 Mekonnen, Joanna Rutter, Kath Horner, Sam Davey, Sarah Hepworth, Lesley
6 Lockwood, Ellie Brown, Chetna Patel, Karen Harrison, Lyn Brandon, Tahira Faiz,
7 Elise Gilwhite, Carole Circuit, Gina Harris, Margot Jackson, Chris Nield, Elaine
8 Muscoft, Gary McCulloch, Permjeet Dhoot, Elaine Goddard;, Gareth Johnston, Anne
9 Richardson at Sheffield Primary Care Trust; Guy Weston at Southey and Owlerton
10 Area Regeneration Trust; Colin Havard and the team at Sharrow Shipshape on
11 London Rd; Miriam Densham at the Zest Centre; and, all the other people who have
12 helped us to implement this study.
13
14
15
16
17
18
19
20

21 **Booster Study team:** Elizabeth Goyder (ScHARR); Daniel Hind (CTRU);
22 Munyaradzi Dimairo (CTRU); Emma Scott (CTRU); Jeff Breckon (CSES); Robert
23 Copeland (CSES); Stephen J Walters (ScHARR); Helen Crank (CSES); Cindy
24 Cooper (CTRU); Karen Collins (Centre for Health and Social Care Research); Emma
25 Everson-Hock (ScHARR); Kim Horspool (CSES); Liam Humphreys (CSES);
26 Andrew Hutchison (CSES); Sue Kesterton (CSES); Nick Latimer (ScHARR); Jon
27 Minton (ScHARR); Simon Read (ScHARR); Peter Swaile (CSES); Rebecca Wood
28 (CSES).
29
30
31
32
33
34
35
36

37 References

- 38
39
40 1. Public Health England. Obesity Knowledge and Intelligence Observatory.
41 *Social and Economic Inequalities in Diet and Physical Activity*. London:
42 Public Health England, 2013.
43 [http://www.noo.org.uk/uploads/doc/vid_19253_Social_and_economic_inequal](http://www.noo.org.uk/uploads/doc/vid_19253_Social_and_economic_inequalities_in_diet_and_physical_activity_04.11.13.pdf)
44 [ities_in_diet_and_physical_activity_04.11.13.pdf](http://www.noo.org.uk/uploads/doc/vid_19253_Social_and_economic_inequalities_in_diet_and_physical_activity_04.11.13.pdf) (date last accessed 14 June
45 2015).
46
47
48
49
50
51 2. Scholes S, Bajekal M, Love H, Hawkins N, Raine R, O'Flaherty M, et al.
52 (2012). Persistent socioeconomic inequalities in cardiovascular risk factors in
53 England over 1994-2008: a time-trend analysis of repeated cross-sectional
54 data. *BMC Public Health* 2012;**12**:129
55
56
57
58
59
60

- 1
2
3 3. Kwarteng JL, Schulz AJ, Mentz GB, Zenk SN, Opperman AA. (2014).
4 Associations between observed neighborhood characteristics and physical
5 activity: findings from a multiethnic urban community. *J Public Health*
6 2014;**36**:358-367.
7
8
9
- 10
11 4. Orrow G, Kinmonth A-L, Sanderson S et al. Effectiveness of physical activity
12 promotion based in primary care: systematic review and meta-analysis of
13 randomised controlled trials. *BMJ* 2012;**344**:e1389.
14
15
16
17
- 18 5. Gulliford M, Charlton J, Bhattarai N, Rudisill C. Social and material
19 deprivation and the cost-effectiveness of an intervention to promote physical
20 activity: cohort study and Markov model. *J Public Health* 2014;**36**:674-683.
21
22
23
24
- 25 6. Goyder E, Hind D, Breckon J, Dimairo M, Minton J, Everson-Hock E et al. A
26 randomised controlled trial and cost-effectiveness evaluation of 'booster'
27 interventions to sustain increases in physical activity in middle-aged adults in
28 deprived urban neighbourhoods. *Health Technol Assess* 2014;**18**:1-210.
29
30
31
32
- 33 7. Gidlow C, Johnston LH, Crone D, Morris C, Smith A, Foster C, James DV.
34 Socio-demographic patterning of referral, uptake and attendance in Physical
35 Activity Referral Schemes. *J Public Health* 2007;**29**:107-13.
36
37
38
39
- 40 8. Bauman AE, Reis RS, Sallis JF, Wells JC, Loos RJ, Martin BW; Lancet
41 Physical Activity Series Working Group. Correlates of physical activity: why
42 are some people physically active and others not? *Lancet* 2012;**380**:258-71.
43
44
45
46
- 47 9. Humpel N, Owen N, Leslie E. Environmental factors associated with adults'
48 participation in physical activity: a review. *Am J Prev Med* 2002;**22**:188-99.
49
50
51
52
- 53 10. Hillsdon M, Panter J, Foster C, Jones A. Equitable access to exercise facilities.
54 *Am J Prev Med* 2007;**32**:506-8.
55
56
57
- 58 11. Maheswaran R, Pearson T, Strong M, Clifford P, Brewins L, Wight J.
59 Assessing the impact of selective migration and care homes on geographical
60

1
2
3 inequalities in health--a total population cohort study in Sheffield. *Spat*
4 *Spatiotemporal Epidemiol* 2014;**10**:85-97.
5
6
7

- 8
9 12. Department for Communities and Local Government. English indices of
10 deprivation. 2012 ([https://www.gov.uk/government/collections/english-](https://www.gov.uk/government/collections/english-indices-of-deprivation)
11 [indices-of-deprivation](https://www.gov.uk/government/collections/english-indices-of-deprivation)).
12
13
14 13. Gidlow C, Ellis N, Randall J, Cowap L, Smith G, Iqbal Z, Kumar J. (2015).
15 Method of invitation and geographical proximity as predictors of NHS Health
16 Check uptake. *J Public Health*, 2015;**37**:195-201.
17
18
19 14. Attwood S, Morton K, Sutton S. Exploring equity in uptake of the NHS
20 Health Check and a nested physical activity intervention trial. *J Public Health*
21 2015;fdv070.
22
23
24 15. Harrison RA, McNair F, Dugdill L. Access to exercise referral schemes -- a
25 population based analysis. *J Public Health* 2005;**27**:326-30.
26
27
28 16. Sowden SL, Breeze E, Barber J, Raine R. Do general practices provide
29 equitable access to physical activity interventions? *Br J Gen Pract*
30 2008;**58**:e1-8.
31
32
33 17. MacNab YC, Read S, Strong M, Pearson T, Maheswaran R, Goyder E.
34 Bayesian hierarchical modelling of noisy spatial rates on a modestly large and
35 discontinuous irregular lattice. *Stat Methods Med Res* 2014;**23**:552-71.
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1: CONSORT diagram for brief intervention uptake showing overall number of participants at each stage




Table 1. Distribution of postcode level variables in the dataset of 3662 postcodes used in this analysis of uptake and effectiveness

Variables	Distribution	
IMD 2010 Income Domain score (Percentage of population who are classified as income deprived)	Median (IQR)	0.30 (0.21 - 0.36)
Walking distance to nearest gym from postcode centroid (km)	Median (IQR)	1.09 (0.67 - 1.59)
Walking distance to nearest swimming pool from postcode centroid (km)	Median (IQR)	1.70 (1.00 - 2.36)
Walking distance to nearest municipal green space from postcode centroid (km)	Median (IQR)	0.60 (0.31 - 0.91)
Percentage of postcodes containing one or more vacant domestic properties	Percentage	39%
Percentage of postcodes containing one or more households in multiple occupation	Percentage	5%
Crime incidents recorded within a 1km radius of the postcode centroid in one year	Median (IQR)	1232 (897 – 1633)
Antisocial behaviour incidents recorded within a 1km radius of the postcode centroid in one year	Median (IQR)	1583 (1188 – 1931)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Table 2: Association of socioeconomic and environmental factors with uptake and effectiveness of the brief intervention

Variable	Odds ratio (95% CI) for an IQR increase (except where indicated otherwise)				
	Respondents (4134) vs. non-respondents (56295)	Contactable respondents (2252) vs. non-contactable (1882)	Eligible (1730) vs. non-eligible (522)	Contactable three months later: yes (941) vs. no (789)	Increased physical activity (486) vs. did not increase physical activity (455)
IMD 2010	0.92 (0.87 - 0.96) p=0.0006	0.85 (0.77 - 0.94) p=0.001	1.02 (0.87 - 1.20)	0.97 (0.83 - 1.13)	0.91 (0.74 - 1.12)
Walking distance to nearest gym	1.06 (0.99 - 1.13)	1.078 (0.94 - 1.23)	0.87 (0.70 - 1.08)	0.81 (0.66 - 1.00) p=0.04	1.02 (0.77 - 1.36)
Walking distance to nearest swimming pool	0.94 (0.88 - 1.00)	1.01 (0.89 - 1.14)	1.12 (0.91 - 1.39)	1.11 (0.92 - 1.35)	0.95 (0.73- 1.24)
Walking distance to nearest municipal green space	1.00 (0.95 - 1.05)	1.04 (0.93 - 1.16)	1.02 (0.87 - 1.19)	1.04 (0.89 - 1.22)	0.92 (0.74 - 1.13)
Vacant domestic properties (Odds ratio for >0%(1) vs 0% (0))	0.95 (0.89 - 1.01)	0.99 (0.87 - 1.13)	1.17 (0.96 - 1.43)	1.09 (0.90 - 1.32)	1.30 (1.00 - 1.70)
Domestic properties in multiple occupancy (Odds ratio for >0%(1) vs 0% (0))	1.07 (0.91 - 1.25)	0.76 (0.55 - 1.04)	0.79 (0.46 - 1.34)	0.52 (0.30 - 0.88) p=0.02	1.49 (0.66 - 3.36)

Crimes reported within a 1km radius of postcode centroid	0.97 (0.89 - 1.06)	0.88 (0.74 - 1.05)	0.89 (0.66 - 1.21)	0.80 (0.60 - 1.07)	1.04 (0.68 - 1.58)
Antisocial behaviour incidents recorded within a 1km radius of postcode centroid	1.05 (0.94 - 1.18)	1.14 (0.91 - 1.44)	1.20 (0.82 - 1.75)	1.38 (0.96 - 1.99)	0.97 (0.58 - 1.63)

Mailout round was included as a categorical variable in all analyses above.